

Contents lists available at [ScienceDirect](http://ScienceDirect)

# Biochimica et Biophysica Acta

journal homepage: [www.elsevier.com/locate/bbamcr](http://www.elsevier.com/locate/bbamcr)

## Review

# The ubiquitin proteasome system – Implications for cell cycle control and the targeted treatment of cancer<sup>☆</sup>



Florian Bassermann<sup>a,\*</sup>, Ruth Eichner<sup>a</sup>, Michele Pagano<sup>b,c,\*\*</sup>

<sup>a</sup> Department of Medicine III, Klinikum rechts der Isar, Technische Universität München, Ismaninger Strasse 22, 81675 Munich, Germany

<sup>b</sup> Department of Pathology, NYU Cancer Institute, New York University School of Medicine, 522 First Avenue, SRB 1107, New York, NY 10016, USA

<sup>c</sup> Howard Hughes Medical Institute, USA

## ARTICLE INFO

### Article history:

Received 17 December 2012

Received in revised form 7 February 2013

Accepted 22 February 2013

Available online 1 March 2013

### Keywords:

Ubiquitin proteasome system

Cell cycle

SCF complex

F-box

APC/C

Cancer

## ABSTRACT

Two families of E3 ubiquitin ligases are prominent in cell cycle regulation and mediate the timely and precise ubiquitin–proteasome-dependent degradation of key cell cycle proteins: the SCF (Skp1/Cul1/F-box protein) complex and the APC/C (anaphase promoting complex or cyclosome). While certain SCF ligases drive cell cycle progression throughout the cell cycle, APC/C (in complex with either of two substrate recruiting proteins: Cdc20 and Cdh1) orchestrates exit from mitosis (APC/C<sup>Cdc20</sup>) and establishes a stable G1 phase (APC/C<sup>Cdh1</sup>). Upon DNA damage or perturbation of the normal cell cycle, both ligases are involved in checkpoint activation. Mechanistic insight into these processes has significantly improved over the last ten years, largely due to a better understanding of APC/C and the functional characterization of multiple F-box proteins, the variable substrate recruiting components of SCF ligases. Here, we review the role of SCF- and APC/C-mediated ubiquitylation in the normal and perturbed cell cycle and discuss potential clinical implications of SCF and APC/C functions. This article is part of a Special Issue entitled: Ubiquitin–Proteasome System. Guest Editors: Thomas Sommer and Dieter H. Wolf.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

The mammalian cell cycle is a strictly regulated process controlled by the oscillating activities of cyclin-dependent kinases (CDKs), which are activated by cyclins and inhibited by CDK inhibitors (CKIs). Many cyclins and CDKs have been described, with each cyclin associating with one or more CDKs, and most CDKs interacting with one or more cyclins [1]. The prototypical cyclins driving cell cycle progression are cyclins A, B, and E, which are expressed in a cell cycle dependent manner and associate with Cdk1 or Cdk2 to mediate downstream events, such as DNA replication and mitosis. Specifically, cyclin E–Cdk2 is activated in late G<sub>1</sub> phase to promote S-phase entry and DNA replication, cyclin A–Cdk2 and cyclin A–Cdk1 are involved in S-phase progression and the G<sub>2</sub>/M transition respectively, while cyclin B–Cdk1 is activated to promote mitotic entry.

This oscillating activity of CDKs is regulated by diverse mechanisms, starting from the transcriptional and translational levels, to posttranslational modifications (phosphorylation in particular), and the periodic

degradation of cyclins and CKIs by the ubiquitin–proteasome system [2,3]. Contrary to reversible modifications, such as phosphorylation or association with CKIs, ubiquitin-mediated proteasomal degradation is an irreversible mechanism that assures the strict unidirectionality of the cell cycle, and it plays a key role in cell cycle regulation by mediating the precise spatial and temporal proteolysis of the main components of the cell cycle machinery.

Ubiquitylation, the covalent attachment of the small 76-amino-acid polypeptide ubiquitin to a target protein, occurs through a three-step enzymatic cascade. Ubiquitin is first bound and activated by the E1 ubiquitin activating enzyme in an ATP dependent manner. It is subsequently transferred to the E2 ubiquitin conjugating enzyme, before the E3 ubiquitin ligase enzyme specifically binds the substrate protein to mediate the transfer of ubiquitin to a lysine residue in the target [4]. Several rounds of ubiquitin conjugation produce long ubiquitin chains in which each ubiquitin molecule is covalently bound to a specific lysine of the previous ubiquitin moiety (polyubiquitylation). In the case of polyubiquitylation via lysine 48 or lysine 11, the polyubiquitylated substrate is committed to association with, and subsequent degradation by, the 26S proteasome. However, there are different types of ubiquitylation, which can lead to other molecular consequences, depending on the nature of the E2. Indeed, substrates can be monoubiquitylated or polyubiquitylated through five other internal lysine residues (K6, K27, K29, K33, and K63) or the N-terminus of ubiquitin [5–7]. While monoubiquitylation or K63-linked polyubiquitylation specifies non-proteolytic fates for a substrate, the implications of other chain linkages are less understood [5,7,8]. Ubiquitylation itself can be reversed by specific deubiquitinating

<sup>☆</sup> This article is part of a Special Issue entitled: Ubiquitin–Proteasome System. Guest Editors: Thomas Sommer and Dieter H. Wolf.

\* Correspondence to: F. Bassermann, Department of Medicine III, Klinikum rechts der Isar, Technische Universität München, Ismaninger Strasse 22, 81675 Munich, Germany. Tel.: +498941405038.

\*\* Correspondence to: M. Pagano, Department of Pathology, NYU Cancer Institute, New York University School of Medicine, 522 First Avenue, SRB 1107, New York, NY 10016, USA. Tel.: +12122635332.

E-mail addresses: [florian.bassermann@lrz.tum.de](mailto:florian.bassermann@lrz.tum.de) (F. Bassermann), [michele.pagano@nyumc.org](mailto:michele.pagano@nyumc.org) (M. Pagano).

enzymes (DUBs) leading to a model in which ubiquitylation is controlled by the balance between an E3 ligase and a DUB [9].

To allow for high specificity in the ubiquitylation of target proteins, more than 600 different E3 ligases exist in the human genome [10]. Depending on their homology domains, these ligases can be subdivided in two major classes, the HECT (homologous to E6-AP C-terminus) family E3 ligases and the RING (really interesting new gene) family E3 ligases [11]. The approximately 30 different HECT ligases form transient covalent linkages with ubiquitin during the ubiquitylation process, while RING ligases only mediate the transfer of ubiquitin from the E2 directly to the substrate. RING E3 ligases can be further subcategorized into those in which one protein contains both the RING and the substrate adaptor domain or multi-subunit complexes, in which these domains are part of distinct proteins within the complex [11]. One of the best described E3 families within the multi-subunit RING ligases is the cullin RING ligase (CRL) superfamily [12], which includes the CRL1 (better known as SCF, standing for Skp1–Cul1–F-box protein complex) and the APC/C (anaphase promoting complex/cyclosome, which is also referred to as a CRL-like ligase) E3 ligases.

The SCF complex consists of Cul1, the scaffold of the complex, Rbx1, the RING protein, which eventually recruits the E2, and Skp1 (S-phase kinase associated protein 1), which serves as an adaptor to bind the F-box protein, the substrate-binding subunit (Fig. 1) [12–14]. F-box proteins are named after the F-box domain, a 40-amino-acid motif initially identified in cyclin F (aka Fbxo1), which binds Skp1 [15]. They contain an additional motif, which constitutes the substrate binding domain. Depending on these domains, they are classified into three groups. F-box proteins containing WD-40 domains are called Fbxws, while those with leucine-rich repeats are called Fbxls. The remaining F-box proteins, with other motifs (e.g., tetratricopeptide repeats, kelch repeats, proline-rich motifs) are named Fbxos [13,16,17].

Sixty-nine different F-box proteins are encoded by the human genome, and individual F-box proteins recognize different substrates. Therefore, these proteins determine the broad functional bandwidth of SCF complexes. Importantly, specific substrates or biological activities have so far been assigned to only a few F-box proteins, and each of these plays a key role in essential cellular processes, such as cell cycle control, apoptosis, DNA damage responses, gene transcription, or translation [3,18].

The APC/C is a prominent E3 ubiquitin ligase involved in cell cycle regulation. It consists of thirteen different subunits and, in somatic

cells, has two co-activators: Cdc20 and Cdh1 (also known as Fzr1), which define substrate specificity and associate with the APC/C core at defined stages of the cell cycle (Fig. 1) [19]. While SCF ligases can be active throughout the cell cycle, APC/C activity is restricted to the time between metaphase and the end of the next  $G_1$ , and during this time, APC/C<sup>Cdc20</sup> initiates anaphase and mitotic exit, while APC/C<sup>Cdh1</sup> contributes to mitotic exit and establishment of a stable  $G_1$  state [3,20–22].

In the following sections, we outline the current understanding of how SCF and APC/C E3 ligases regulate cell cycle progression, their response to environmental cues in each phase of the cell cycle, and potential clinical and therapeutic implications of SCF and APC/C biology.

## 2. The $G_0$ and $G_1$ phases

### 2.1. Quiescence and $G_1$


In response to mitogen withdrawal (or after exit from mitosis), a stable  $G_0$  or  $G_1$  state has to be established. The APC/C<sup>Cdh1</sup> complex is the central player in this task, and it is activated in late mitosis to complete mitotic exit and reset the cell cycle by targeting a variety of proteins involved in DNA replication, cycle progression, and mitosis. Among the targets that promote DNA replication is Cdc6, which binds to the origin recognition complex (ORC) to form pre-replication complexes (preRCs). Cdc6 is kept at low levels during early  $G_1$  through APC/C<sup>Cdh1</sup>-mediated degradation (Fig. 2A) [23]. APC/C<sup>Cdh1</sup> also negatively regulates proliferative signal transduction by targeting Ets2 (Fig. 2A), a transcription factor involved in Ras–Raf–MAPK signaling that induces the expression of cyclin D1 [24,25]. APC/C<sup>Cdh1</sup> activity also precludes early accumulation of positive cell cycle regulators [21,26]. Specifically, Skp2 (S phase kinase-associated protein 2, aka Fbl1) is targeted for proteolysis to avert premature formation of the SCF<sup>Skp2</sup> complex, which promotes cell cycle progression by mediating the degradation of CKIs (see below and Fig. 2A) [26–28]. Importantly, the Cdc25A phosphatase, which promotes S-phase entry and mitosis by dephosphorylating and thus activating Cdk2 and Cdk1, is also kept at low levels during early  $G_1$  by APC/C<sup>Cdh1</sup> (Fig. 2A) [29]. Finally, APC/C<sup>Cdh1</sup> targets mitotic proteins for proteasomal degradation [19–21,30]. Among the most prominent targets are cyclin A and cyclin B (Fig. 2A), the main drivers of mitotic entry [31,32]. Furthermore, Aurora A, Aurora B, and polo-like kinase 1 (Plk1), important mitotic kinases involved in centrosome maturation, spindle assembly, and chromosome separation, are additional targets of APC/C<sup>Cdh1</sup> (Fig. 2A) [33–36].

Hence, by targeting protooncogenic proteins, such as Skp2 and Ets2, APC/C<sup>Cdh1</sup> exerts an anti-proliferative activity. In fact, APC/C<sup>Cdh1</sup> acts as a tumor suppressor in mice, where *Fzr1* heterozygous mice have elevated rates of spontaneous tumor formation [37].


In response to mitogen withdrawal, cells must also switch to a state of nutritional saving and restrict protein translation and cell growth to sustain survival. SCF<sup>Fbxo9</sup> contributes to this process via differential regulation of mTORC1 and mTORC2 signaling. Specifically, the Tel2 (telomere maintenance 2) and Tti1 (Tel2 interacting protein 1) proteins, integral components of both the mTORC1 and mTORC2 complexes, are phosphorylated by CKII to target them for degradation by SCF<sup>Fbxo9</sup> upon growth factor withdrawal. This pathway specifically targets Tel2 and Tti1 in the mTORC1 complex (Fig. 2B) [38]. As a consequence, mTORC1 signaling is attenuated to restrain protein translation and cell growth, but via relief of mTORC1-mediated feedback inhibition, mTORC2 signaling is sustained to promote survival [38].

### 2.2. The growth factor response

If growth factors and nutrients are available, cells respond with a number of events that lead to growth and, eventually, cell division. Certain SCF ligases are acutely involved in the orchestration of growth factor responses during  $G_1$  phase. For example, the tumor suppressor



**Fig. 1.** Structure of the SCF and the APC/C ubiquitin ligases. The SCF and APC/C E3 ligases are both members of the Cullin-RING ligase (CRL) superfamily. Cul1 and Apc2 (light grey) are the scaffold proteins of SCF and APC/C, respectively. On one end, they bind to a RING finger protein, Rbx1 or Apc11 (dark grey), which recruits the E2 ubiquitin conjugating enzyme (UBC). On their other end, they connect to the substrate specific unit via an adaptor molecule (red). In the case of SCF, the F-box proteins are the variable substrate binding components (orange), while Cdh1 and Cdc20 (in somatic cells), together with Apc10, recruit substrates to the APC/C. The scheme illustrates the relationship between APC/C and SCF components and does not represent the topology of APC/C subunits.



**Fig. 2.** SCF and APC/C mediated degradation processes in G<sub>1</sub> phase and the response to mitogen stimulation. **A.** APC/C<sup>Cdh1</sup> maintains a stable G<sub>1</sub> phase by targeting mitotic proteins like Aurora A, Aurora B, Plk1, cyclin A, and cyclin B for degradation. In addition, APC/C<sup>Cdh1</sup> prevents premature accumulation of positive cell cycle regulators, such as Ets2, Cdc6, and Cdc25A. The CDK inhibitors p21 and p27 are stabilized via APC/C<sup>Cdh1</sup>-dependent degradation of Skp2 to keep CDK activity low during G<sub>1</sub>. **B.** Upon mitogen stimulation, SCF<sup>βTrCP</sup> mediates the degradation of inhibitory or proapoptotic proteins, such as PDCD4, DEPTOR, and BimEL, promoting cell growth. Upon mitogen withdrawal, SCF<sup>Fbxo9</sup> targets Tel2 and Tti1 for degradation within mTORC1 to attenuate mTORC1 activity and sustain mTORC2 signaling. **C.** At the G<sub>1</sub>/S transition, APC/C<sup>Cdh1</sup> activity is inhibited through several mechanisms, such as inhibitory phosphorylation by CDKs and binding to Emi1. Skp2-dependent degradation of p21 and p27 leads to activation of cyclin E-Cdk2, which initiates S phase before being degraded via SCF<sup>Fbw7</sup>. See main text for more details.

PDCD4 (programmed cell death protein 4), which inhibits the translation initiation factor eIF4A, is degraded in response to mitogens in a two-step mechanism. First, PDCD4 is phosphorylated on serine 67 (and possibly adjacent serine residues) by S6K1, and it is subsequently ubiquitinated via SCF<sup>βTrCP</sup> (Fig. 2B), resulting in increased protein synthesis and cell growth [39]. In response to mitogens and survival signals, SCF<sup>βTrCP</sup> furthermore promotes survival by targeting the proapoptotic BH3-only protein BimEL for degradation in an Rsk1/2- and Erk1/2-dependent manner (Fig. 2B) [40]. Another important target of SCF<sup>βTrCP</sup> in growth factor responses is the mTOR inhibitor DEPTOR (Fig. 2B). After mitogen stimulation, DEPTOR is phosphorylated by mTORC1 or mTORC2 and CK1α, which directs ubiquitination and degradation, relieving mTOR inhibition [41,42]. Thus, in cooperation with SCF<sup>βTrCP</sup>, mTOR signaling promotes further mTOR activation via this auto-amplification loop.

### 2.3. Post-restriction point G<sub>1</sub> and the G<sub>1</sub> to S phase transition

With sufficient mitogen stimulation and cell growth, cells commit to cell division, passing a point of G<sub>1</sub> known as “restriction point”, after which cells will proceed through a round of cell division even if mitogens are removed. The transition from G<sub>1</sub> to S phase results from decreasing APC/C<sup>Cdh1</sup> activity, decreasing CKIs levels, rising cyclin expression, increasing CDK activity, phosphorylation and inactivation of Rb (retinoblastoma) protein family members, and activation of E2F transcription factor family members.

Inactivation of APC/C<sup>Cdh1</sup> during progression through the late G<sub>1</sub> phase occurs via different mechanisms. First, UbcH10, the APC/C specific E2 ubiquitin conjugating enzyme, is ubiquitinated by APC/C<sup>Cdh1</sup>

(Fig. 2C), thereby providing a negative feedback loop limiting APC/C<sup>Cdh1</sup> activity [43]. Likewise, Cdh1 initiates autoubiquitination within APC/C (Fig. 2C), thus limiting its activity [44]. Furthermore, rising levels of CDK activity lead to phosphorylation of Cdh1, disrupting its binding to APC/C (Fig. 2C) [3,45]. Later, in S-Phase, an unidentified SCF complex contributes to Cdh1 degradation (Fig. 2C), either directly or indirectly, and this process may depend on previous Cdh1-phosphorylation [46]. Finally, at the G<sub>1</sub>/S transition, the transcription factor E2F induces expression of Emi1 (an F-box protein also known as Fbxo5), which binds to APC/C<sup>Cdh1</sup> as a pseudosubstrate inhibitor, inhibiting APC/C<sup>Cdh1</sup> throughout S and G<sub>2</sub> (Fig. 2C) [47,48].

Rising CDK activity during G<sub>1</sub> phase occurs through several different mechanisms. Primarily, the expression of cyclins is induced on a transcriptional level, with cyclin E and cyclin A expression being initiated by E2F (Fig. 2C) [49,50]. However, the majority of cyclin-Cdk complexes are inhibited via association with CDK inhibitors (CKIs), such as p21 or p27. In late G<sub>1</sub>, Cdk4 and Cdk6 bound to cyclin D1 catalyze the phosphorylation (and consequent inactivation) of Rb family proteins (pRb, p107, and p130), which bind and inhibit E2F transcription factors, preventing their promotion of cell cycle progression. With decreasing APC/C<sup>Cdh1</sup> activity in late G<sub>1</sub>, Skp2 accumulates and forms the SCF<sup>Skp2</sup> complex (Fig. 2C), which targets cell cycle inhibitors for degradation, promoting cell cycle progression throughout S, G<sub>2</sub>, and M [22]. Importantly, SCF<sup>Skp2</sup> binds p27 and mediates its proteasomal degradation, liberating cyclin E-Cdk2 from p27 inhibition (Fig. 2C) [51–54]. Ubiquitination of p27 depends on phosphorylation of threonine 187 by cyclin E-Cdk2 and the co-factor Cks1 [55–59]. Other CKIs ubiquitinated by SCF<sup>Skp2</sup> include p21 and p57, whose degradation further reinforces cyclin-Cdk1/2 activation [60,61]. By analogy to p27,

phosphorylation by cyclinE-Cdk2 and binding of Cks1 are likely prerequisites for SCF<sup>Skp2</sup> dependent ubiquitylation of p21 and p27 [60,61]. These functions distinguish Skp2 as an E3 ligase with oncogenic properties, which is further supported by the frequent overexpression of Skp2 in many tumors [62].

### 3. S and G<sub>2</sub> phases of the cell cycle

#### 3.1. S phase


Cyclin E-Cdk2 and (later) cyclin A-Cdk2 are the two main CDK complexes in S phase and mediate the initiation of DNA and centrosome duplication. At the same time, cyclin A-Cdk2 further inhibits APC/C<sup>Cdh1</sup> via Cdh1 phosphorylation to ensure cyclin stability (Fig. 3A) [43]. Upon successful S-phase entry, cyclin D1 is phosphorylated by GSK3β, targeting it for degradation by an SCF ligase containing Fbxo4 and the small heat-shock protein α/B-crystallin (the latter being a substrate adaptor) [63]. As cells progress through S and approach G<sub>2</sub>, the levels of S phase cyclins decrease again. Notably, cyclin E abundance is regulated via a negative feedback loop. Specifically, as Cdk2-activity increases, Cdk2-bound cyclin E is phosphorylated on threonine 380 in a Cdk2-dependent manner, whereupon phosphorylated cyclin E is targeted for degradation by the SCF<sup>Fbw7</sup> complex (Figs. 2C and 3A) [64,65].

#### 3.2. DNA damage checkpoint


During S and G<sub>2</sub> phases, it is crucial for cells to assure accurate and complete DNA replication. Therefore, DNA damage detection mechanisms and checkpoints provide precise surveillance. Upon DNA damage, ATM activates the checkpoint kinase Chk2 and ATR activates Chk1 in a Claspin-dependent process (Fig. 4). Chk1 and Chk2 then mediate inhibitory phosphorylation of Cdc25A on serine 216 [66,67]. Cdc25A promotes cell cycle activation by virtue of its ability to remove inhibitory phosphorylations from Cdk1 and Cdk2, but Cdc25A phosphorylation by Chk1/2 results in ubiquitylation by SCF<sup>βTrCP</sup> and degradation, attenuating CDK activity (Fig. 4) [68]. Degradation of Cdc25A leads to a pause in cell cycle progression during S or G<sub>2</sub>, depending on the timing of the DNA damage.

Another target of the DNA damage checkpoint in G<sub>2</sub> phase is Plk1 [69,70]. In response to DNA damage, the phosphatase Cdc14B is released from the nucleolus to the nucleoplasm, where it dephosphorylates Cdh1 (Fig. 4). As a consequence, APC/C<sup>Cdh1</sup> becomes reactivated to target Plk1 for degradation (Fig. 4). In this way, Claspin and Wee1 are stabilized to activate the checkpoint and inhibit cell cycle progression (Fig. 4) [70]. Notably, other known targets of APC/C<sup>Cdh1</sup> are likely protected during this process by DUBs, particularly USP28 [70,71].

SCF<sup>βTrCP</sup> also plays a role during recovery from the checkpoint. Claspin, which mediates ATR-induced activation of Chk1 upon DNA damage, is phosphorylated by Plk1 during checkpoint recovery and



**Fig. 3.** SCF and APC/C mediated degradation processes in S and G<sub>2</sub> phases of the cell cycle. A. In early G<sub>2</sub>, APC/C<sup>Cdh1</sup> remains inactivated, SCF<sup>Skp2</sup> maintains cyclin A-Cdk1/2 activity via degradation of p21 and p27, SCF<sup>Fbw7</sup> promotes cyclin E degradation, and SCF<sup>NIPA</sup> prevents premature accumulation of nuclear cyclin B. Other S phase proteins (e.g., RRM2 and CP110) are also degraded via cyclin F to stop dNTP production and assure centrosomes are only replicated once per cell cycle. B. At the G<sub>2</sub>/M transition, SCF<sup>Skp2</sup> continues to target p21 and p27, while Wee1 is degraded in a Plk1- and SCF<sup>βTrCP</sup>-dependent manner to release cyclin B-Cdk1 from inhibition. See main text for more details.



**Fig. 4.** Role of SCF and APC/C ligases in the response to DNA damage. Upon DNA damage, SCF and APC/C activate S and G2 checkpoints to halt cell cycle progression. Chk1 (which is activated by ATR) promotes the SCF<sup>βTrCP</sup>-dependent of Cdc25A to inhibit CDK activity, while dephosphorylation of Cdh1 by Cdc14B allows for Plk1 degradation and subsequent stabilization of Claspin and Wee1. See main text for more details.

subsequently degraded in an SCF<sup>βTrCP</sup> dependent manner [72,73]. As a consequence, activation of Chk1 declines, and the cell recovers from the checkpoint. Moreover, SCF<sup>βTrCP</sup> also regulates translation during the silencing of the G<sub>2</sub> DNA damage checkpoint. In response to genotoxic stress, AMPK (adenosine monophosphate-activated protein kinase) phosphorylates and activates eEF2K (eukaryotic Elongation Factor 2 Kinase), which, in turn, inhibits elongation by phosphorylating and inhibiting eEF2, a protein that mediates movement of the ribosome along mRNA [74]. During checkpoint silencing, SCF<sup>βTrCP</sup> targets eEF2K for degradation to enable rapid resumption of translation elongation [74].

### 3.3. G<sub>2</sub> phase and G<sub>2</sub>/M transition

After successful and accurate completion of DNA replication, production of deoxyribonucleotides (dNTPs) must be avoided. To this end, RRM2 (ribonucleotide reductase family member 2), which converts ribonucleotides to dNTPs for DNA synthesis, is phosphorylated on threonine 33 by CDKs and ubiquitinated by the SCF<sup>CyclinF</sup> complex, reducing the availability of dNTPs (Fig. 3A) [75]. In contrast, cyclin F is downregulated in response to DNA damage in an ATR-dependent manner to stabilize RRM2 (Fig. 4). By this means, production of dNTPs is enhanced to allow for efficient DNA repair [75].

Beyond DNA replication, SCF<sup>CyclinF</sup> also regulates centrosome duplication. Cyclin F and CP110, a protein essential for centrosome duplication, physically associate on the centrioles during G<sub>2</sub> [76]. By targeting CP110 for proteasomal degradation (Fig. 3A), SCF<sup>CyclinF</sup> assures that centrosomes are replicated only once during the cell cycle, preventing centrosome overduplication, which could lead to multipolar or asymmetric mitotic spindles and subsequent chromosome aberrations [76].

With progression through G<sub>2</sub> phase, the cell prepares for mitotic entry, which is orchestrated by cyclin B-Cdk1. Therefore, cyclin B-Cdk1 is a strictly regulated kinase complex. In late G<sub>2</sub> phase, SCF<sup>βTrCP</sup> promotes activation of Cdk1 by mediating the degradation of Wee1, a Cdk1 inhibitory kinase (Fig. 3B). A prerequisite for SCF<sup>βTrCP</sup> binding to Wee1 is phosphorylation of serine 53 and serine 123 by Plk1 and Cdk1, respectively [77]. Thus, a positive feedback loop between Cdk1 activation and Wee1 degradation assures rapid activation of Cdk1 upon mitotic entry (Fig. 3B). The activity of Plk1

itself is also tightly controlled through the synergistic actions of Bora and Aurora A kinase. Accumulation of Bora during G<sub>2</sub> leads to Aurora A-mediated activation of Plk1, which contributes to Wee1 degradation and subsequent activation of Cdk1 (Fig. 3B) [78].


To preclude premature mitotic entry, early activation of cyclin B-Cdk1 must be prevented. To this end, SCF<sup>NIPa</sup> targets nuclear cyclin B1 during S and G<sub>2</sub> (Fig. 3A) [79]. However, in late G<sub>2</sub>, NIPa is phosphorylated at serine 395 by cyclin B-Cdk1, after initial phosphorylation on serine 354 and serine 359 by Erk2 (Fig. 3B) [80,81]. Phosphorylation inhibits assembly of a functional SCF<sup>NIPa</sup> complex, allowing nuclear accumulation of cyclin B1 and subsequent mitotic entry. In this context, cyclin B1 regulates its own abundance through a positive feedback loop (Fig. 3B).

## 4. Mitosis

### 4.1. Prometaphase and spindle assembly checkpoint

Cyclin B-Cdk1 orchestrates essential steps of early mitosis, including the assembly of the mitotic spindle, breakdown of the nuclear envelope, cessation of gene transcription, and condensation of chromosomes. These processes drive the cell through prophase until metaphase.

Starting from metaphase, the APC/C becomes a key player in promoting mitotic progression and, eventually, mitotic exit. Thus, inhibitory mechanisms that keep the APC/C in check throughout S- and G<sub>2</sub> phases must be removed. In this regard, SCF<sup>βTrCP</sup> mediates ubiquitylation and degradation of the APC/C inhibitor Emi1 following phosphorylation by cyclin B-Cdk1 and Plk1 in early mitosis (Fig. 5A) [82,83]. However, until all kinetochores are correctly attached to spindle fibers and the chromosomes are properly aligned on the metaphase plate, the Spindle Assembly Checkpoint (SAC) inhibits APC/C activation, blocking the onset of anaphase [84]. In the presence of unattached kinetochores, the mitotic checkpoint proteins, BubR1, Bub3, and Mad2, bind to Cdc20 to form the Mitotic Checkpoint Complex (MCC), which sequesters Cdc20 and directly inhibits activation of the APC/C<sup>Cdc20</sup> (Fig. 5A) [84–86]. Another model suggests that Cdc20 is targeted for degradation by APC/C in an MCC-dependent manner during SAC activation [87,88]. Furthermore, inhibitory phosphorylation of Cdc20 by protein kinases, such as Cdk1 (Fig. 5A), MAPK, or Bub1, contributes to Cdc20 inhibition during SAC activation [89–91]. Interestingly, the transcription of Mad2



**Fig. 5.** SCF and APC/C mediated degradation processes during mitosis. A. In early mitosis, SCF<sup>βTrCP</sup> ubiquitylates Emi1 to eliminate the repression of APC/C<sup>Cdh1</sup> (which remains inactive because of Cdh1 phosphorylation by CDKs) and, possibly, APC/C<sup>Cdc20</sup>. However, during activation of the Spindle Assembly Checkpoint (SAC), APC/C<sup>Cdc20</sup> activity is repressed, with cyclin B-Cdk1 contributing to its inhibition. B. After fulfillment of the SAC, APC/C<sup>Cdc20</sup> inhibition is relieved, allowing it to target mitotic cyclins and Securin for degradation, the latter leading to Separase activation and sister chromatid separation. With mitotic progression, APC/C<sup>Cdc20</sup> activity decreases via several mechanisms, and APC/C<sup>Cdh1</sup> assembles to target cell cycle proteins for degradation. See main text for more details.

is also regulated by SCF<sup>βTrCP</sup>. During late G<sub>2</sub>, SCF<sup>βTrCP</sup> targets REST (Repressor-Element-1-Silencing Transcription factor) for degradation, allowing transcriptional derepression of Mad2 (Fig. 5A) [92].

Despite the many mechanisms keeping APC/C<sup>Cdc20</sup> in check, its activity is not completely inhibited; a minor fraction remains active even during SAC activation. This subpopulation sustains cyclin B-Cdk1 activity during prometaphase by targeting p21 for degradation (Fig. 5A) [93]. Although p21 is degraded via SCF<sup>Skp2</sup> at the G<sub>1</sub>/S transition, it reaccumulates during G<sub>2</sub> and interacts with cyclin B-Cdk1 before it is ubiquitylated by APC/C<sup>Cdc20</sup>. Interestingly, APC/C<sup>Cdc20</sup> and APC/C<sup>Cdh1</sup> have opposing effects on p21 levels. While p21 is degraded via APC/C<sup>Cdc20</sup> in early mitosis, APC/C<sup>Cdh1</sup> indirectly stabilizes p21 in G<sub>1</sub> phase by targeting Skp2 and Cdc20 for degradation [27,94,95]. In addition, APC/C<sup>Cdc20</sup> targets two other substrates for degradation in early mitosis, cyclin A and Nek2A, even when the spindle checkpoint is active. How the checkpoint-resistant activity of the APC/C<sup>Cdc20</sup> is regulated remains unclear. Potential mechanisms include Cdc20-independent binding of substrates to APC10 or an increase in substrate affinity for Cdc20, which allows substrates to outcompete SAC proteins [96–98].

#### 4.2. Metaphase to anaphase transition and mitotic exit

Upon proper attachment of all spindle microtubules to chromosome kinetochores, the SAC is rapidly inactivated, and APC/C<sup>Cdc20</sup> is released from inhibition via an uncertain mechanism (Fig. 5B).

Checkpoint inactivation depends on Cdc20-dependent ubiquitylation. According to some, this leads to dissociation of the MCC from APC/C<sup>Cdc20</sup>, but others suggest that it causes degradation of Cdc20 [99,100]. An additional, ATP-dependent step is also required for both dissociation of the MCC from APC/C and disassembly of the free MCC [101,102]. This step has been shown to involve β-γ bond cleavage of ATP, suggesting the presence of an additional ubiquitin independent process, which may be kinase- or chaperone-related [102]. The understanding of SAC inactivation is further complicated by the fact that Cdc20 proteolysis is also stimulated by Mad2 and BubR1 to keep levels of Cdc20 low in order to avoid premature activation of APC/C and maintain the checkpoint [87,88]. Eventually, after kinetochore attachment, activated APC/C<sup>Cdc20</sup> initiates anaphase by targeting Securin for degradation (Fig. 5B) [20,103,104]. During metaphase, sister chromatids are held together by Cohesin. Securin is a chaperone that binds and inhibits Separase, an enzyme capable of cleaving Cohesin. With Securin degradation, Separase is released, and centromeric Cohesin is cleaved, which leads to sister chromatid segregation (Fig. 5B) [105]. Moreover, Shugoshin, a protector of centromeric sister chromatid cohesion, is targeted for degradation by APC/C<sup>Cdh1</sup>; however, this step is not indispensable for chromatid segregation and seems more important during meiosis [106,107].

Both mitotic cyclins, cyclin A and cyclin B, are targeted for degradation by the APC/C (Fig. 5B) [108]. Whereas cyclin A degradation starts in prometaphase, cyclin B ubiquitylation starts at the metaphase/anaphase

transition, mediated first by APC/C<sup>Cdc20</sup> and later by APC/C<sup>Cdh1</sup> [109–111]. Notably, APC/C<sup>Cdc20</sup> has different effects on Cdk1 activity depending on the mitotic phase. While it augments cyclin B-Cdk1 activity during prometaphase via degradation of p21 (Fig. 5A), it suppresses cyclin B-Cdk1 via degradation of cyclin B in late mitosis (Fig. 5B) [93]. Mitotic cyclin-CDK activity inhibits separase, so the degradation of cyclin A and cyclin B by APC/C<sup>Cdc20</sup> is directly involved in sister chromatid separation (Fig. 5B) [20]. Furthermore, drastically reduced CDK activity in anaphase allows for disassembly of the mitotic spindle, chromosome decondensation, cytokinesis, and reconstitution of the nuclear envelope [20].

Although APC/C is the most prominent E3 ubiquitin ligase orchestrating the metaphase to anaphase transition, SCF ubiquitin ligases are also involved in the regulation of anaphase onset. For instance, Bora, which works together with Plk1 and Aurora A to regulate microtubule polymerization, spindle stability, and tension between kinetochores, is targeted for degradation by SCF<sup>F<sup>β</sup>TrCP</sup> at the metaphase to anaphase transition (Fig. 5A) [112,113]. Interestingly, a prerequisite for Bora ubiquitylation is phosphorylation by Plk1, which itself is activated in a Bora-dependent manner at the G<sub>2</sub>/M transition [78,112]. Timely activation and degradation of Bora assure regulated cell cycle progression, as inhibition of Bora degradation delays anaphase onset and knockdown of Bora activates the spindle checkpoint [112,113].

In late mitosis, Aurora A and Plk1 are eventually targeted for degradation via the APC/C<sup>Cdh1</sup> to promote mitotic exit (Fig. 5B) [33,34,36]. Another important substrate of APC/C in late mitosis is Geminin (Fig. 5B), which inhibits DNA re-replication by precluding formation of the pre-replication complex (pre-RC) after S-phase, rendering replication possible during the subsequent cell cycle [114].

Overall, the APC/C is the central E3 ligase that coordinates mitotic progression, mitotic exit, and the subsequent establishment of a G<sub>1</sub> state. While anaphase onset is induced by APC/C<sup>Cdc20</sup>, both APC/C<sup>Cdc20</sup> and APC/C<sup>Cdh1</sup> orchestrate mitotic exit, and APC/C<sup>Cdh1</sup> contributes to establishment and maintenance of the G<sub>1</sub> phase of the next cell cycle. This distinct activation profile is the result of different regulatory steps. As mentioned earlier, Cdh1 is kept inactivated via phosphorylation by CDKs during S, G<sub>2</sub> and part of mitosis [115]. Degradation of cyclin B reduces CDK activity at the metaphase to anaphase transition, so the levels of dephosphorylated Cdh1 rise. Dephosphorylated Cdh1 binds APC/C and negatively regulates APC/C<sup>Cdc20</sup> by targeting Cdc20 for degradation (Fig. 5B) [94,95]. Moreover, APC/C dependent autoubiquitylation of Cdc20 in anaphase has been proposed to further contribute to the inactivation of APC/C<sup>Cdc20</sup> (Fig. 5B) [99,100]. The oscillation between Cdc20 and Cdh1 expression partly explains how the APC/C times the degradation of its various substrates. Depending on their destruction box and their affinity for Cdc20 or Cdh1, APC/C substrates will be targeted earlier or later during mitosis or G<sub>1</sub> for proteasomal degradation [116,117].

## 5. Other CRLs involved in cell cycle control

Other CRLs are also implicated in cell cycle control and the DNA damage response. CRL3 complexes appear to control mitosis via proteolytic and non-proteolytic mechanisms. The best understood cell cycle-related CRL3 substrate is Aurora B. K63-type ubiquitylation of Aurora B targets it to mitotic chromosomes to control their alignment [118]. The CRL4 ligases also control features of the cell cycle. In addition to Cul4 and Rbx1, CRL4 ligases contain the adaptor protein DDB1 (DNA Damage Binding protein 1), which recruits members of the DCAF (DDB1–Cul4 Associated Factors) family to dictate the specificity of substrate degradation [119–121]. CRL4<sup>Cdt2</sup> is a prominent member of this family implicated in DNA replication and the DNA damage response. Notably, CRL4<sup>Cdt2</sup> interacts with different UBCs (UBCH8, UBE2G1, and UBE2G2) to mediate ubiquitylation of its various substrates [122]. An important target of CRL4<sup>Cdt2</sup> is Cdt1, a replication licensing factor and pre-replication complex (pre-RC) component that

is tightly regulated to ensure that DNA replication only occurs once per cell cycle. After S phase, Cdt1 activity is attenuated both by Geminin binding and by proteasomal degradation. Importantly, CRL4<sup>Cdt2</sup>-mediated degradation of Cdt1 is initiated shortly after S phase onset [123–126]. This event requires Cdt1 binding to PCNA via a PIP (PCNA interacting protein)-box motif present in Cdt1. Upon DNA damage, Cdt1 is degraded in the same way to preclude DNA replication [127]. (Interestingly, Cdt1 is also targeted for degradation by SCF<sup>Skp2</sup> in response to CDK-dependent phosphorylation [123,128].)

An additional target of CRL4<sup>Cdt2</sup> is p21, which is degraded both in response to UV-induced DNA damage and during the unperturbed cell cycle. In the later context, CRL4<sup>Cdt2</sup> collaborates with SCF<sup>Skp2</sup> to mediate timely degradation of p21. Like Cdt1, p21 degradation by CRL4<sup>Cdt2</sup> requires interaction with PCNA via a PIP-box [129]. CRL4<sup>Cdt2</sup> is further involved in cell cycle control via regulation of histone methylation by ubiquitylation of Set8. Set8 mono-methylates Lys 20 of histone H4 (H4K20me1) during G<sub>2</sub> to promote chromatin compaction. During S phase and upon DNA damage, Set8 is degraded by CRL4<sup>Cdt2</sup> in a PCNA-dependent manner [130–132]. Perturbation of this mechanism leads to accumulation of H4K20me1, premature chromatin compaction, and activation of the G<sub>2</sub>/M checkpoint [130,131].

Interestingly, Cdt2 levels are regulated via SCF<sup>Fbxo11</sup> during the cell cycle [133,134] and inhibition of Cdt2 degradation delays cell cycle exit. In contrast to most other SCF substrates, whose phosphorylation promotes binding to the F-box protein, CDK-dependent phosphorylation of the Cdt2 degron inhibits binding to and degradation via Fbxo11 [134]. Thus, crosstalk between an SCF ligase, CDKs, and a CRL4 complex controls the abundance of Cdt2 to regulate the timing of cell cycle exit.

## 6. Clinical implications

### 6.1. SCF complexes as targets in the therapy of cancer and other diseases

The SCF complexes and the APC/C are key players in cell cycle control and the DNA damage response. As such, they govern cell proliferation and genome stability, which, when deregulated, contribute to tumorigenesis. Among the SCF ligases controlling cell growth and proliferation (discussed above), this situation has been described in particular for SCF<sup>Skp2</sup>, SCF<sup>F<sup>β</sup>TrCP</sup>, SCF<sup>Fbxw7</sup>, and SCF<sup>Fbxo9</sup>. The recent therapeutic success of the proteasome inhibitor bortezomib in multiple myeloma and certain types of B-cell Non-Hodgkin's lymphoma has expedited the efforts to develop inhibitors of ubiquitin ligases or its components [135,136]. While bortezomib exhibits a certain degree of selectivity towards cancer cells and has a noticeable therapeutic index, it non-specifically blocks the ubiquitin–proteasome-dependent degradation of all cellular proteins, resulting in considerable side effects and the development of resistance. Therapies directed against individual E3 ligases or ligase families known to be deregulated in human cancers may thus prove more efficacious. To qualify as a target for therapeutic inhibition, a ligase should be an oncoprotein (based on the activity of its substrates – e.g., tumor suppressors) and deregulated in tumors.

#### 6.1.1. Skp2

Skp2 is an oncoprotein by virtue of its function in degrading negative cell cycle regulators, including p27, p21, p130, and p57 [137,138]. Skp2 is overexpressed in various malignancies, such as gastric cancer [139,140], colon cancer [141], and breast cancer [142], with an associated decrease in p27 levels and indicating a poor prognosis. The oncogenic function of Skp2 is further underscored by mouse genetic studies. Targeted expression of Skp2 in the T-lymphoid lineage cooperates with activated N-Ras to induce T-cell lymphomas [143], while tissue specific expression of Skp2 in the prostate induces hyperplasia and low grade carcinoma [144]. In addition, a critical role for Skp2-dependent degradation of p27 in colon adenoma–carcinoma development was demonstrated using a Skp2 knock-in model [145]. Similarly, in Skp2-null cells, aberrant

oncogenic signaling or inactivation of tumor suppressor genes trigger a potent, tumor-suppressive senescence response, although Skp2 inactivation alone does not induce senescence [146]. Finally, RNAi-mediated knockdown of Skp2 inhibits the growth of tumor cells, including cell lines derived from glioblastoma [147], melanoma [148], and oral cancer [149]. Specific inhibition of Skp2 would thus be expected to be effective in the treatment of cancers with activated Skp2 signaling.

Interestingly, Skp2 also interacts with a number of proteins produced by pathogenic viruses, such as X-protein (HBV), EBNA3C (EBV), and E7 (HPV16/18) [150], suggesting that Skp2 inhibitors may be also utilized in certain viral infections.

### 6.1.2. $\beta$ TrCP

Given the diversity of its substrates (reviewed in [151]),  $\beta$ TrCP might be expected to exert both oncogenic and tumor suppressor activities. However, in certain tissues,  $\beta$ TrCP is clearly an oncoprotein based on its ubiquitylation activity against tumor suppressors, such as I $\kappa$ B, a negative regulator of NF $\kappa$ B [152], PDCD4, an inhibitor of eIF4A-mediated protein translation [39], and BimEL, a potent proapoptotic protein [40]. Indeed,  $\beta$ TrCP is overexpressed in breast cancer [153], and forced overexpression of  $\beta$ TrCP induces transformation in breast epithelium [154]. Likewise, upregulation of  $\beta$ TrCP has been shown in colon cancer, where it correlates with elevated NF $\kappa$ B activity and poor prognosis [155] and in pancreatic cancer cells, where it impacts chemoresistance [156]. Further evidence demonstrating the oncogenic potential of  $\beta$ TrCP stems from transgenic mouse models, in which tumor formation was observed in the mammary gland, liver, and kidney when  $\beta$ TrCP expression was directed to these tissues [154,157]. Interestingly, no obvious phenotype was observed upon targeted expression of  $\beta$ TrCP in lymphoid organs, suggesting that  $\beta$ TrCP-dependent tumorigenesis is tissue specific [154]. Therefore, disrupting the interaction of  $\beta$ TrCP and its tumor suppressive substrates may be an attractive therapeutic strategy in defined tumor entities.

Similarly to Skp2,  $\beta$ TrCP interacts with a number of proteins produced by pathogenic viruses, such as Tax (HTLV1). Moreover, the HIV-1 protein Vpu targets SCF $^{\beta$ TrCP to CD4 [158]. This finding, together with the evidence that other HIV-1 proteins (i.e. Vif and Vpr) bind CRLs to eliminate cellular proteins with antiviral activity (e.g., APOBEC3), indicates that SCF $^{\beta$ TrCP and other CRLs are involved in the HIV life cycle and represent potential targets in the fight against this virus.

### 6.1.3. Fbxo9 and Fbxw7

The rationale for targeting the ubiquitin–proteasome system is particularly evident in multiple myeloma, due to the high efficacy of proteasome inhibitors in this disease. Indeed, Bortezomib, a reversible proteasome inhibitor, has been approved as the first line treatment for multiple myeloma. In addition, Carfilzomib, a second generation irreversible proteasome inhibitor has recently been approved by the Food and Drug Administration (FDA) for the treatment of patients with relapsed and refractory multiple myeloma, and demonstrates increased potency and an improved therapeutic index [159,160]. Two ubiquitin ligases of the SCF family have recently been reported to contribute to the pathogenesis of multiple myeloma, SCF $^{\text{Fbxo9}}$  and SCF $^{\text{Fbxw7}\alpha}$ .

As outlined above, Fbxo9 targets Tel2/Tti1 proteins in a CK2-dependent manner to adjust mTOR signaling to the availability of growth factors. Significantly, Fbxo9 is overexpressed in multiple myeloma and drives constitutive activation of the PI3K/mTORC2/Akt pathway to promote survival [38]. In multiple myelomas with elevated Fbxo9 expression, inhibition of Fbxo9 represents an attractive approach to inhibit Akt signaling, one of the most important mediators of cell survival in this disease [159]. Inhibition of CK2, the kinase that promotes Fbxo9-mediated ubiquitylation of Tel/Tti1, may also be an interesting therapeutic approach in multiple myeloma, particularly since CK2 inhibitors are readily available [161].

Although Fbxw7 behaves as a tumor suppressor due to its ubiquitylation activity against mitogenic substrates (e.g., Notch, c-Myc, cyclin E), Fbxw7 functions as a pro-survival gene in multiple myeloma by constitutively targeting the NF $\kappa$ B inhibitor p100 in a GSK3-dependent manner [162]. While a number of cancers, including T-ALL, breast cancers, and gastric adenocarcinoma often carry mutations in the Fbxw7 gene, leading to an accumulation of mitogenic substrates [163–165], these mutations do not occur in B-cell malignancies like multiple myeloma [166]. Fbxw7 and GSK3 may thus serve as promising targets for the treatment of multiple myelomas with constitutive activation of the NF $\kappa$ B pathway.

In a short term setting, acute delivery of an Fbxw7 inhibitor could also be effective in tumors where this F-box protein plays a tumor suppressive role by inducing cancer stem cells to proliferate and, therefore, become susceptible to traditional chemotherapies. However, this hypothesis remains to be proved experimentally.

### 6.1.4. Other F-box proteins

Fbxo11 mutations are present in human cancers, such as diffuse large B cell lymphomas (DLBCLs), colon, lung, ovary, and head and neck tumors [167–171] (and Staudt L., personal communication). At least in DLBCLs, these mutations inhibit SCF $^{\text{Fbxo11}}$ . The presence of inactivating mutations suggests that Fbxo11 may function as a tumor suppressor, whose loss of function contributes to the pathogenesis of DLBCL (via BCL6 accumulation) and other cancers (through the stabilization of unidentified oncogenic substrates). Furthermore, inactivating mutations have been reported for Fbxo4 in esophageal carcinoma, and these mutations are associated with an increase in cyclin D1 levels. The mutations occur in the N-terminal regulatory regions of Fbxo4 and disrupt dimerization and activation [172]. Moreover, there is evidence for the loss of  $\alpha$ /B crystallin in breast cancer and a corresponding increase in cyclin D1 levels.  $\alpha$ /B crystallin is the substrate adaptor for cyclin D1 together with Fbxo4 [63,173]. These data suggest a tumor suppressor role for the SCF $^{\text{Fbxo4-}\alpha/\text{B crystallin}}$  ligase in human tumors based on its ability to destabilize cyclin D1. Finally, cyclin F has been reported to be downregulated in hepatocellular carcinoma (HCC). In this context, low cyclin F expression was an independent poor prognostic marker for overall survival and correlated with tumor size and clinical stage [174]. Given the role of cyclin F in regulating centrosome homeostasis and the balanced abundance of dNTPs, these findings hint at a function for cyclin F as a tumor suppressor in HCC by maintaining genomic stability [75,76].

## 6.2. Strategies to target CRLs

In theory, targets for inhibiting substrate degradation by the ubiquitin proteasome system can involve any of the three enzymes involved in ubiquitin transfer, including the E1 activating, the E2 conjugating, and the E3 ligase enzymes. These approaches would be expected to inhibit most or certain ubiquitin ligase families, when targeting E1 or E2 enzymes, respectively, and become highly specific when targeting single E3 ligases.

### 6.2.1. Targeting E1 ubiquitin activating enzymes

E1s (of which there are two in mammals) catalyze the initial step in ubiquitin conjugation, which involves an ATP-dependent covalent attachment of ubiquitin to its active cysteine site [4]. Efforts have been undertaken to target E1. These studies have yielded candidate compounds like PYR-41, an irreversible ubiquitin E1 inhibitor, but further investigations will be necessary to assess if they are clinical candidates [175]. Additional efforts have been directed to the E1s for ubiquitin like proteins (such as NEDD8, SUMO, etc.). Recently, MLN4924, an adenosine sulfamate analog, was reported to inhibit the E1 responsible for NEDDylation, the covalent addition of NEDD8 to substrates [176]. The cullin family of proteins are the most important substrates for neddylation, and the activity of SCF and other CRLs



requires cullin neddylation [12,177,178]. In contrast to an inhibitor for the ubiquitin E1s, a small molecule inhibitor of the NEDD8 E1 would only inhibit CRLs. Currently, MLN4924 is in phase I/II clinical trials for the treatment of multiple myeloma and non-Hodgkins lymphoma, and the preclinical data suggest high efficacy in AML [179,180].

#### 6.2.2. Targeting E2 ubiquitin conjugating enzymes

Together with the E3 ligase, E2s mediate the transfer of ubiquitin to the target protein and govern the type and extent of ubiquitin linkage [181]. The human genome encodes at least 38 E2 enzymes, and Cdc34 (aka Ubc3) appears to be the major E2 for SCF ligases in promoting K48-linked polyubiquitylation and proteasomal degradation [12]. Recently, CC0651, a small molecule allosteric inhibitor of Cdc34 was reported [182]. CC0651 inhibits p27 ubiquitylation and degradation, and it demonstrates convincing specificity for human Cdc34, as evidenced by a lack of reactivity against any other E2/E3 pairs tested [182]. While current data on other E2 inhibitors is limited, the development of CC0651 indicates the feasibility of developing highly selective inhibitors of E2s and encourages efforts to target other E2s in a similar manner.

#### 6.2.3. Targeting E3 ubiquitin ligases

E3 ligases determine which target protein becomes ubiquitylated. Inhibition of a single E3 ligase would be the most selective targeting approach, since it would affect a limited number of proteins, potentially translating into a better therapeutic ratio and fewer side effects. Most E3 ligases do not feature a canonical active site, and instead, the active site is a protein–protein interaction with a substrate. Inhibition of these protein–protein interactions is typically considered more difficult than inhibition of a catalytic site. However, this task appears increasingly feasible with advances in the structural understanding of these interactions [183]. MDM2 is an extensively studied ubiquitin E3 ligase with strong clinical relevance by virtue of its ability to regulate the abundance of the tumor suppressor p53. Previously, the Nutlin class of imidazoline chemotypes was identified, and they specifically disrupt the protein–protein binding interface between MDM2 and p53, stabilizing p53 [184]. Nutlin-3a is the most promising candidate and has favorable preclinical characteristics in terms of pharmacological properties and toxicity [185,186]. Currently, Nutlin-3a is being investigated in early phase clinical trials for several solid and hematological tumor entities, setting a precedent for the specific inhibition of a ligase–substrate pair as a promising clinical approach.

In the case of SCF complexes, inhibition of either the F-box protein–substrate interface or the recruitment of the F-box protein to the SCF core are attractive strategies to selectively inhibit individual ubiquitylation events. In this regard, the small molecule inhibitor CpdA has previously been found to prevent the binding of Skp2 to the SCF ligase complex, inducing G1/S arrest and apoptosis by stabilizing p27, p21, and other Skp2 target proteins. In addition, CpdA sensitized multiple myeloma cells to cytostatic agents and bortezomib, and it was active against both myeloid and lymphoblastoid leukemia blasts [187].

Independent studies have recently demonstrated the feasibility of direct inhibition of the interfaces between the F-box protein and either the substrate or SCF core. SCF-I2 is an allosteric inhibitor of substrate recognition by the yeast F-box protein Cdc4 and inhibits binding and ubiquitylation of substrates. SCF-I2 inserts itself between the  $\beta$ -strands of the WD-40 propeller domain of Cdc4, impairing recognition of Cdc4-specific phosphodegrons [188]. In addition, a recent study screened for inhibitors that selectively target the p27-binding interface formed by Skp2–Cks1. This approach yielded four compounds that stabilize the expression of p27 in different human cancer cell lines and induce cell cycle arrest in G1 [189]. Finally, a chemical genetics screen for enhancers of rapamycin identified SMER3, an inhibitor of the SCF<sup>Met30</sup> ligase. SMER3 specifically inhibits

binding of the F-box protein subunit Met30 to the SCF core, stabilizing substrates such as Met4 [190].

Another approach to prevent substrate degradation at the E3 ligase level is to inhibit the kinase that phosphorylates a substrate to mark it for degradation. This approach particularly applies to SCF ubiquitin ligases, as it is well established that substrate phosphorylation is frequently a prerequisite for F-box protein binding [13,151,191]. Indeed, kinases like Plk1, which control the degradation of tumor suppressors, are potential targets, and inhibitors have already entered early clinical trials [192]. Notably, this approach could quickly impact the clinic, as inhibitors to numerous kinases are readily available, but broader application will require further investigations.

Together, SCF ligases are promising therapeutic target structures, whose pharmacological modulation would offer highly specific therapeutic approaches to a wide variety of malignancies. The success of this undertaking will largely depend on the pharmacological progress in developing specific inhibitors of F-box proteins and a better mechanistic understanding of the many uncharacterized F-box proteins, both with regard to their substrates and the posttranslational modifications that prime them for SCF-dependent ubiquitylation. Finally, it will be critical to further define tumor entities in which distinct SCF ligases are deregulated. This data will help define distinct F-box proteins as biomarkers and allow the selection of patient subpopulations that would profit from a targeted approach against SCF ligases.

#### Acknowledgements

We thank Jeffrey Skaar for critically reading the manuscript. We apologize to colleagues whose work could not be mentioned owing to space limitations. Work in the Bassermann laboratory is supported by grants from the German Research Foundation (Emmy Noether Program, BA 2851/3-1) and the German Cancer Aid (#109543), and in the Pagano laboratory by grants from the NIH (R01-GM057587 R37-CA076584, and R21-CA161108). MP is an investigator with the Howard Hughes Medical Institute.

#### References

- [1] A.W. Murray, Recycling the cell cycle: cyclins revisited, *Cell* 116 (2004) 221–234.
- [2] M. Glotzer, A.W. Murray, M.W. Kirschner, Cyclin is degraded by the ubiquitin pathway, *Nature* 349 (1991) 132–138.
- [3] J.R. Skaar, M. Pagano, Control of cell growth by the SCF and APC/C ubiquitin ligases, *Curr. Opin. Cell Biol.* 21 (2009) 816–824.
- [4] A. Hershko, A. Ciechanover, The ubiquitin system, *Annu. Rev. Biochem.* 67 (1998) 425–479.
- [5] M.J. Clague, S. Urbe, Ubiquitin: same molecule, different degradation pathways, *Cell* 143 (2010) 682–685.
- [6] C. Behrends, J.W. Harper, Constructing and decoding unconventional ubiquitin chains, *Nat. Struct. Mol. Biol.* 18 (2011) 520–528.
- [7] P. Xu, D.M. Duong, N.T. Seyfried, D. Cheng, Y. Xie, J. Robert, J. Rush, M. Hochstrasser, D. Finley, J. Peng, Quantitative proteomics reveals the function of unconventional ubiquitin chains in proteasomal degradation, *Cell* 137 (2009) 133–145.
- [8] A.D. Jacobson, N.Y. Zhang, P. Xu, K.J. Han, S. Noone, J. Peng, C.W. Liu, The lysine 48 and lysine 63 ubiquitin conjugates are processed differently by the 26S proteasome, *J. Biol. Chem.* 284 (2009) 35485–35494.
- [9] M.J. Clague, J.M. Coulson, S. Urbe, Cellular functions of the DUBs, *J. Cell Sci.* 125 (2012) 277–286.
- [10] W. Li, M.H. Bengtson, A. Ulbrich, A. Matsuda, V.A. Reddy, A. Orth, S.K. Chanda, S. Batalov, C.A. Joazeiro, Genome-wide and functional annotation of human E3 ubiquitin ligases identifies MULAN, a mitochondrial E3 that regulates the organelle's dynamics and signaling, *PLoS One* 3 (2008) e1487.
- [11] M.B. Metzger, V.A. Hristova, A.M. Weissman, HECT and RING finger families of E3 ubiquitin ligases at a glance, *J. Cell Sci.* 125 (2012) 531–537.
- [12] M.D. Petroski, R.J. Deshaies, Function and regulation of cullin-RING ubiquitin ligases, *Nat. Rev. Mol. Cell Biol.* 6 (2005) 9–20.
- [13] T. Cardozo, M. Pagano, The SCF ubiquitin ligase: insights into a molecular machine, *Nat. Rev. Mol. Cell Biol.* 5 (2004) 739–751.
- [14] N. Zheng, B.A. Schulman, L. Song, J.J. Miller, P.D. Jeffrey, P. Wang, C. Chu, D.M. Koepp, S.J. Elledge, M. Pagano, R.C. Conaway, J.W. Conaway, J.W. Harper, N.P. Pavletich, Structure of the Cul1–Rbx1–Skp1–F boxSkp2 SCF ubiquitin ligase complex, *Nature* 416 (2002) 703–709.

- [15] C. Bai, P. Sen, K. Hofmann, L. Ma, M. Goebel, J.W. Harper, S.J. Elledge, SKP1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box, *Cell* 86 (1996) 263–274.
- [16] C. Cenciarelli, D.S. Chaiur, D. Guardavaccaro, W. Parks, M. Vidal, M. Pagano, Identification of a family of human F-box proteins, *Curr. Biol.* 9 (1999) 1177–1179.
- [17] J. Jin, T. Cardozo, R.C. Lovering, S.J. Elledge, M. Pagano, J.W. Harper, Systematic analysis and nomenclature of mammalian F-box proteins, *Genes Dev.* 18 (2004) 2573–2580.
- [18] J.S. Silverman, J.R. Skaar, M. Pagano, SCF ubiquitin ligases in the maintenance of genome stability, *Trends Biochem. Sci.* 37 (2012) 66–73.
- [19] R. van Leuken, L. Clijsters, R. Wolthuis, To cell cycle, swing the APC/C, *Biochim. Biophys. Acta* 1786 (2008) 49–59.
- [20] C. Acquaviva, J. Pines, The anaphase-promoting complex/cyclosome: APC/C, *J. Cell Sci.* 119 (2006) 2401–2404.
- [21] M. Li, P. Zhang, The function of APC/CCdh1 in cell cycle and beyond, *Cell Div.* 4 (2009) 2.
- [22] K.I. Nakayama, K. Nakayama, Regulation of the cell cycle by SCF-type ubiquitin ligases, *Semin. Cell Dev. Biol.* 16 (2005) 323–333.
- [23] N. Mailand, J.F. Diffley, CDKs promote DNA replication origin licensing in human cells by protecting Cdc6 from APC/C-dependent proteolysis, *Cell* 122 (2005) 915–926.
- [24] M. Li, Y.H. Shin, L. Hou, X. Huang, Z. Wei, E. Klann, P. Zhang, The adaptor protein of the anaphase promoting complex Cdh1 is essential in maintaining replicative lifespan and in learning and memory, *Nat. Cell Biol.* 10 (2008) 1083–1089.
- [25] C. Albanese, J. Johnson, G. Watanabe, N. Eklund, D. Vu, A. Arnold, R.G. Pestell, Transforming p21ras mutants and c-Ets-2 activate the cyclin D1 promoter through distinguishable regions, *J. Biol. Chem.* 270 (1995) 23589–23597.
- [26] T. Bashir, M. Pagano, Don't skip the G1 phase: how APC/CCdh1 keeps SCF/SKP2 in check, *Cell Cycle* 3 (2004) 850–852.
- [27] T. Bashir, N.V. Dorrello, V. Amador, D. Guardavaccaro, M. Pagano, Control of the SCF(Skp2-Cks1) ubiquitin ligase by the APC/C(Cdh1) ubiquitin ligase, *Nature* 428 (2004) 190–193.
- [28] W. Wei, N.G. Ayad, Y. Wan, G.J. Zhang, M.W. Kirschner, W.G. Kaelin Jr., Degradation of the SCF component Skp2 in cell-cycle phase G1 by the anaphase-promoting complex, *Nature* 428 (2004) 194–198.
- [29] M. Donzelli, M. Squatrito, D. Ganoth, A. Hershko, M. Pagano, G.F. Draetta, Dual mode of degradation of Cdc25 A phosphatase, *EMBO J.* 21 (2002) 4875–4884.
- [30] J.R. Skaar, M. Pagano, Cdh1: a master G0/G1 regulator, *Nat. Cell Biol.* 10 (2008) 755–757.
- [31] R.W. King, J.M. Peters, S. Tugendreich, M. Rolfe, P. Hieter, M.W. Kirschner, A 20S complex containing CDC27 and CDC16 catalyzes the mitosis-specific conjugation of ubiquitin to cyclin B, *Cell* 81 (1995) 279–288.
- [32] S. Geley, E. Kramer, C. Geffers, J. Gannon, J.M. Peters, T. Hunt, Anaphase-promoting complex/cyclosome-dependent proteolysis of human cyclin A starts at the beginning of mitosis and is not subject to the spindle assembly checkpoint, *J. Cell Biol.* 153 (2001) 137–148.
- [33] K. Honda, H. Mihara, Y. Kato, A. Yamaguchi, H. Tanaka, H. Yasuda, K. Furukawa, T. Urano, Degradation of human Aurora2 protein kinase by the anaphase-promoting complex-ubiquitin-proteasome pathway, *Oncogene* 19 (2000) 2812–2819.
- [34] A. Castro, Y. Arlot-Bonnemains, S. Vigneron, J.C. Labbe, C. Prigent, T. Lorca, APC/fizzy-related targets aurora-A kinase for proteolysis, *EMBO Rep.* 3 (2002) 457–462.
- [35] S. Stewart, G. Fang, Destruction box-dependent degradation of aurora B is mediated by the anaphase-promoting complex/cyclosome and Cdh1, *Cancer Res.* 65 (2005) 8730–8735.
- [36] C. Lindon, J. Pines, Ordered proteolysis in anaphase inactivates Plk1 to contribute to proper mitotic exit in human cells, *J. Cell Biol.* 164 (2004) 233–241.
- [37] I. Garcia-Higuera, E. Manchado, P. Dubus, M. Canamero, J. Mendez, S. Moreno, M. Malumbres, Genomic stability and tumour suppression by the APC/C cofactor Cdh1, *Nat. Cell Biol.* 10 (2008) 802–811.
- [38] V. Fernandez-Saiz, B.S. Targosz, S. Lemeer, R. Eichner, C. Langer, L. Bullinger, C. Reiter, J. Slotta-Huspenina, S. Schroeder, A.M. Knorn, J. Kurutz, C. Peschel, M. Pagano, B. Kuster, F. Bassermann, SCF(Fbxo9) and CK2 direct the cellular response to growth factor withdrawal via Tel2/Tti1 degradation and promote survival in multiple myeloma, *Nat. Cell Biol.* 15 (2013) 72–81.
- [39] N.V. Dorrello, A. Peschiaroli, D. Guardavaccaro, N.H. Colburn, N.E. Sherman, M. Pagano, S6K1- and betaTRCP-mediated degradation of PDCD4 promotes protein translation and cell growth, *Science* 314 (2006) 467–471.
- [40] E. Dehan, F. Bassermann, D. Guardavaccaro, G. Vasiliou-Shamis, M. Cohen, K.N. Lowes, M. Dustin, D.C. Huang, J. Taunton, M. Pagano, betaTRCP- and Rsk1/2-mediated degradation of BimEL inhibits apoptosis, *Mol. Cell* 33 (2009) 109–116.
- [41] S. Duan, J.R. Skaar, S. Kuchay, A. Toschi, N. Kanarek, Y. Ben-Neriah, M. Pagano, mTOR generates an auto-amplification loop by triggering the betaTRCP- and CK1alpha-dependent degradation of DEPTOR, *Mol. Cell* 44 (2011) 317–324.
- [42] D. Gao, H. Inuzuka, M.K. Tan, H. Fukushima, J.W. Locasale, P. Liu, L. Wan, B. Zhai, Y.R. Chin, S. Shaik, C.A. Lyssiotis, S.P. Gygi, A. Toker, L.C. Cantley, J.M. Asara, J.W. Harper, W. Wei, mTOR drives its own activation via SCF(betaTRCP)-dependent degradation of the mTOR inhibitor DEPTOR, *Mol. Cell* 44 (2011) 290–303.
- [43] M. Rape, M.W. Kirschner, Autonomous regulation of the anaphase-promoting complex couples mitosis to S-phase entry, *Nature* 432 (2004) 588–595.
- [44] T. Listovsky, Y.S. Oren, Y. Yudkovsky, H.M. Mahbubani, A.M. Weiss, M. Lebendiker, M. Brandeis, Mammalian Cdh1/Fzr mediates its own degradation, *EMBO J.* 23 (2004) 1619–1626.
- [45] C. Lukas, C.S. Sorensen, E. Kramer, E. Santoni-Rugiu, C. Lindenege, J.M. Peters, J. Bartek, J. Lukas, Accumulation of cyclin B1 requires E2F and cyclin-A-dependent rearrangement of the anaphase-promoting complex, *Nature* 401 (1999) 815–818.
- [46] R. Benmaamar, M. Pagano, Involvement of the SCF complex in the control of Cdh1 degradation in S-phase, *Cell Cycle* 4 (2005) 1230–1232.
- [47] J.J. Miller, M.K. Summers, D.V. Hansen, M.V. Nachury, N.L. Lehman, A. Loktev, P.K. Jackson, Emi1 stably binds and inhibits the anaphase-promoting complex/cyclosome as a pseudosubstrate inhibitor, *Genes Dev.* 20 (2006) 2410–2420.
- [48] J.Y. Hsu, J.D. Reimann, C.S. Sorensen, J. Lukas, P.K. Jackson, E2F-dependent accumulation of hEmi1 regulates S phase entry by inhibiting APC(Cdh1), *Nat. Cell Biol.* 4 (2002) 358–366.
- [49] Y. Geng, E.N. Eaton, M. Picon, J.M. Roberts, A.S. Lundberg, A. Gifford, C. Sardet, R.A. Weinberg, Regulation of cyclin E transcription by E2Fs and retinoblastoma protein, *Oncogene* 12 (1996) 1173–1180.
- [50] A. Schulze, K. Zerfass, D. Spitkovsky, S. Middendorp, J. Berges, K. Helin, P. Jansen-Durr, B. Henglein, Cell cycle regulation of the cyclin A gene promoter is mediated by a variant E2F site, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 11264–11268.
- [51] L.M. Tsvetkov, K.H. Yeh, S.J. Lee, H. Sun, H. Zhang, p27(Kip1) ubiquitination and degradation is regulated by the SCF(Skp2) complex through phosphorylated Thr187 in p27, *Curr. Biol.* 9 (1999) 661–664.
- [52] A.C. Carrano, E. Eytan, A. Hershko, M. Pagano, SKP2 is required for ubiquitin-mediated degradation of the CDK inhibitor p27, *Nat. Cell Biol.* 1 (1999) 193–199.
- [53] M. Pagano, S.W. Tam, A.M. Theodoras, P. Beer-Romero, G. Del Sal, V. Chau, P.R. Yew, G.F. Draetta, M. Rolfe, Role of the ubiquitin-proteasome pathway in regulating abundance of the cyclin-dependent kinase inhibitor p27, *Science* 269 (1995) 682–685.
- [54] H. Sutterluty, E. Chatelain, A. Marti, C. Wirbelauer, M. Senften, U. Muller, W. Krek, p45SKP2 promotes p27Kip1 degradation and induces S phase in quiescent cells, *Nat. Cell Biol.* 1 (1999) 207–214.
- [55] R.J. Sheaff, M. Groudine, M. Gordon, J.M. Roberts, B.E. Clurman, Cyclin E-CDK2 is a regulator of p27Kip1, *Genes Dev.* 11 (1997) 1464–1478.
- [56] J. Vlach, S. Hennecke, B. Amati, Phosphorylation-dependent degradation of the cyclin-dependent kinase inhibitor p27, *EMBO J.* 16 (1997) 5334–5344.
- [57] A. Montagnoli, F. Fiore, E. Eytan, A.C. Carrano, G.F. Draetta, A. Hershko, M. Pagano, Ubiquitination of p27 is regulated by Cdk-dependent phosphorylation and trimeric complex formation, *Genes Dev.* 13 (1999) 1181–1189.
- [58] D. Ganoth, G. Bornstein, T.K. Ko, B. Larsen, M. Tyers, M. Pagano, A. Hershko, The cell-cycle regulatory protein Cks1 is required for SCF(Skp2)-mediated ubiquitinylation of p27, *Nat. Cell Biol.* 3 (2001) 321–324.
- [59] C. Spruck, H. Strohmaier, M. Watson, A.P. Smith, A. Ryan, T.W. Krek, S.I. Reed, A CDK-independent function of mammalian Cks1: targeting of SCF(Skp2) to the CDK inhibitor p27Kip1, *Mol. Cell* 7 (2001) 639–650.
- [60] G. Bornstein, J. Bloom, D. Sirty-Shevah, K. Nakayama, M. Pagano, A. Hershko, Role of the SCFSkp2 ubiquitin ligase in the degradation of p21Cip1 in S phase, *J. Biol. Chem.* 278 (2003) 25752–25757.
- [61] T. Kamura, T. Hara, S. Kotoshiba, M. Yada, N. Ishida, H. Imaki, S. Hatakeyama, K. Nakayama, K.I. Nakayama, Degradation of p57Kip2 mediated by SCFSkp2-dependent ubiquitylation, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 10231–10236.
- [62] D.D. Hershko, Oncogenic properties and prognostic implications of the ubiquitin ligase Skp2 in cancer, *Cancer* 112 (2008) 1415–1424.
- [63] D.I. Lin, O. Barbash, K.G. Kumar, J.D. Weber, J.W. Harper, A.J. Klein-Szanto, A. Rustgi, S.Y. Fuchs, J.A. Diehl, Phosphorylation-dependent ubiquitination of cyclin D1 by the SCF(FBX4-alphaB crystallin) complex, *Mol. Cell* 24 (2006) 355–366.
- [64] B.E. Clurman, R.J. Sheaff, K. Thress, M. Groudine, J.M. Roberts, Turnover of cyclin E by the ubiquitin-proteasome pathway is regulated by cdk2 binding and cyclin phosphorylation, *Genes Dev.* 10 (1996) 1979–1990.
- [65] D.M. Koepf, L.K. Schaefer, X. Ye, K. Keyomarsi, C. Chu, J.W. Harper, S.J. Elledge, Phosphorylation-dependent ubiquitination of cyclin E by the SCFbeta7 ubiquitin ligase, *Science* 294 (2001) 173–177.
- [66] S. Matsuoka, M. Huang, S.J. Elledge, Linkage of ATM to cell cycle regulation by the Chk2 protein kinase, *Science* 282 (1998) 1893–1897.
- [67] Y. Zeng, K.C. Forbes, Z. Wu, S. Moreno, H. Piwnicka-Worms, T. Enoch, Replication checkpoint requires phosphorylation of the phosphatase Cdc25 by Cds1 or Chk1, *Nature* 395 (1998) 507–510.
- [68] L. Busino, M. Donzelli, M. Chiesa, D. Guardavaccaro, D. Ganoth, N.V. Dorrello, A. Hershko, M. Pagano, G.F. Draetta, Degradation of Cdc25A by beta-TRCP during S phase and in response to DNA damage, *Nature* 426 (2003) 87–91.
- [69] V.A. Smits, R. Klompaker, L. Arnaud, G. Rijkssen, E.A. Nigg, R.H. Medema, Polo-like kinase-1 is a target of the DNA damage checkpoint, *Nat. Cell Biol.* 2 (2000) 672–676.
- [70] F. Bassermann, D. Frescas, D. Guardavaccaro, L. Busino, A. Peschiaroli, M. Pagano, The Cdc14B-Cdh1-Plk1 axis controls the G2 DNA-damage-response checkpoint, *Cell* 134 (2008) 256–267.
- [71] F. Bassermann, M. Pagano, Dissecting the role of ubiquitylation in the DNA damage response checkpoint in G2, *Cell Death Differ.* 17 (2010) 78–85.
- [72] A. Peschiaroli, N.V. Dorrello, D. Guardavaccaro, M. Venere, T. Halazonetis, N.E. Sherman, M. Pagano, SCFBetaTRCP-mediated degradation of Claspin regulates recovery from the DNA replication checkpoint response, *Mol. Cell* 23 (2006) 319–329.
- [73] N. Mailand, S. Bekker-Jensen, J. Bartek, J. Lukas, Destruction of Claspin by SCFBetaTRCP restrains Chk1 activation and facilitates recovery from genotoxic stress, *Mol. Cell* 23 (2006) 307–318.
- [74] F. Kruiswijk, L. Yuniati, R. Magliozzi, T.Y. Low, R. Lim, R. Bolder, S. Mohammed, C.G. Proud, A.J. Heck, M. Pagano, D. Guardavaccaro, Coupled activation and degradation of eEF2K regulates protein synthesis in response to genotoxic stress, *Sci. Signal.* 5 (2012) ra40.
- [75] V. D'Angiolella, V. Donato, F.M. Forrester, Y.T. Jeong, C. Pellacani, Y. Kudo, A. Saraf, L. Flores, M.P. Washburn, M. Pagano, Cyclin F-mediated degradation of ribonucleotide reductase M2 controls genome integrity and DNA repair, *Cell* 149 (2012) 1023–1034.

- [76] V. D'Angioletta, V. Donato, S. Vijayakumar, A. Saraf, L. Florens, M.P. Washburn, B. Dynlacht, M. Pagano, SCF (Cyclin F) controls centrosome homeostasis and mitotic fidelity through CP110 degradation, *Nature* 466 (2010) 138–142.
- [77] N. Watanabe, H. Arai, Y. Nishihara, M. Taniguchi, T. Hunter, H. Osada, M-phase kinases induce phospho-dependent ubiquitination of somatic Wee1 by SCFbeta-TrCP, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 4419–4424.
- [78] A. Seki, J.A. Coppinger, C.Y. Jang, J.R. Yates, G. Fang, Bora and the kinase Aurora cooperatively activate the kinase Plk1 and control mitotic entry, *Science* 320 (2008) 1655–1658.
- [79] F. Bassermann, C. von Klitzing, S. Munch, R.Y. Bai, H. Kawaguchi, S.W. Morris, C. Peschel, J. Duyster, NIPA defines an SCF-type mammalian E3 ligase that regulates mitotic entry, *Cell* 122 (2005) 45–57.
- [80] F. Bassermann, C. von Klitzing, A.L. Illert, S. Munch, S.W. Morris, M. Pagano, C. Peschel, J. Duyster, Multisite phosphorylation of nuclear interaction partner of ALK (NIPA) at G2/M involves cyclin B1/Cdk1, *J. Biol. Chem.* 282 (2007) 15965–15972.
- [81] A.L. Illert, M. Zech, C. Moll, C. Albers, S. Kreutmair, C. Peschel, F. Bassermann, J. Duyster, Extracellular signal-regulated kinase 2 (ERK2) mediates phosphorylation and inactivation of nuclear interaction partner of anaplastic lymphoma kinase (NIPA) at G2/M, *J. Biol. Chem.* 287 (2012) 37997–38005.
- [82] F. Margottin-Goguet, J.Y. Hsu, A. Loktev, H.M. Hsieh, J.D. Reimann, P.K. Jackson, Prophase destruction of Emi1 by the SCF(betaTrCP/Slmb) ubiquitin ligase activates the anaphase promoting complex to allow progression beyond prometaphase, *Dev. Cell* 4 (2003) 813–826.
- [83] D. Guardavaccaro, Y. Kudo, J. Boulaire, M. Barchi, L. Busino, M. Donzelli, F. Margottin-Goguet, P.K. Jackson, L. Yamasaki, M. Pagano, Control of meiotic and mitotic progression by the F box protein beta-Trcp1 in vivo, *Dev. Cell* 4 (2003) 799–812.
- [84] G. Fang, H. Yu, M.W. Kirschner, The checkpoint protein MAD2 and the mitotic regulator CDC20 form a ternary complex with the anaphase-promoting complex to control anaphase initiation, *Genes Dev.* 12 (1998) 1871–1883.
- [85] V. Sudakin, G.K. Chan, T.J. Yen, Checkpoint inhibition of the APC/C in HeLa cells is mediated by a complex of BUBR1, BUB3, CDC20, and MAD2, *J. Cell Biol.* 154 (2001) 925–936.
- [86] I. Braunstien, S. Miniowitz, Y. Moshe, A. Hershko, Inhibitory factors associated with anaphase-promoting complex/cyloosome in mitotic checkpoint, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 4870–4875.
- [87] J. Nilsson, M. Yekezare, J. Minshall, J. Pines, The APC/C maintains the spindle assembly checkpoint by targeting Cdc20 for destruction, *Nat. Cell Biol.* 10 (2008) 1411–1420.
- [88] S. Ge, J.R. Skaar, M. Pagano, APC/C- and Mad2-mediated degradation of Cdc20 during spindle checkpoint activation, *Cell Cycle* 8 (2009) 167–171.
- [89] V. D'Angioletta, C. Mari, D. Nocera, L. Rametti, D. Grieco, The spindle checkpoint requires cyclin-dependent kinase activity, *Genes Dev.* 17 (2003) 2520–2525.
- [90] E. Chung, R.H. Chen, Phosphorylation of Cdc20 is required for its inhibition by the spindle checkpoint, *Nat. Cell Biol.* 5 (2003) 748–753.
- [91] Z. Tang, H. Shu, D. Oncel, S. Chen, H. Yu, Phosphorylation of Cdc20 by Bub1 provides a catalytic mechanism for APC/C inhibition by the spindle checkpoint, *Mol. Cell* 16 (2004) 387–397.
- [92] D. Guardavaccaro, D. Frescas, N.V. Dorrello, A. Peschiaroli, A.S. Multani, T. Cardozo, A. Lasorella, A. Iavarone, S. Chang, E. Hernandez, M. Pagano, Control of chromosome stability by the beta-TrCP-REST-Mad2 axis, *Nature* 452 (2008) 365–369.
- [93] V. Amador, S. Ge, P.G. Santamaria, D. Guardavaccaro, M. Pagano, APC/C(Cdc20) controls the ubiquitin-mediated degradation of p21 in prometaphase, *Mol. Cell* 27 (2007) 462–473.
- [94] C.M. Pflieger, M.W. Kirschner, The KEN box: an APC recognition signal distinct from the D box targeted by Cdh1, *Genes Dev.* 14 (2000) 655–665.
- [95] A. Reis, M. Levasseur, H.Y. Chang, D.J. Elliott, K.T. Jones, The CRY box: a second APCcdh1-dependent degron in mammalian cdc20, *EMBO Rep.* 7 (2006) 1040–1045.
- [96] B. Di Fiore, J. Pines, How cyclin A destruction escapes the spindle assembly checkpoint, *J. Cell Biol.* 190 (2010) 501–509.
- [97] M.J. Hayes, Y. Kimata, S.L. Wattam, C. Lindon, G. Mao, H. Yamano, A.M. Fry, Early mitotic degradation of Nek2A depends on Cdc20-independent interaction with the APC/C, *Nat. Cell Biol.* 8 (2006) 607–614.
- [98] W. van Zon, R.M. Wolthuis, Cyclin A and Nek2A: APC/C-Cdc20 substrates invisible to the mitotic spindle checkpoint, *Biochem. Soc. Trans.* 38 (2010) 72–77.
- [99] G. Varetti, C. Guida, S. Santaguida, E. Chiroli, A. Musacchio, Homeostatic control of mitotic arrest, *Mol. Cell* 44 (2011) 710–720.
- [100] I.T. Foe, S.A. Foster, S.K. Cheung, S.Z. DeLuca, D.O. Morgan, D.P. Toczycki, Ubiquitination of Cdc20 by the APC occurs through an intramolecular mechanism, *Curr. Biol.* 21 (2011) 1870–1877.
- [101] S.K. Reddy, M. Rape, W.A. Margansky, M.W. Kirschner, Ubiquitination by the anaphase-promoting complex drives spindle checkpoint inactivation, *Nature* 446 (2007) 921–925.
- [102] S. Miniowitz-Shemtov, A. Teichner, D. Sitry-Shevah, A. Hershko, ATP is required for the release of the anaphase-promoting complex/cyloosome from inhibition by the mitotic checkpoint, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 5351–5356.
- [103] H. Zou, T.J. McGarry, T. Bernal, M.W. Kirschner, Identification of a vertebrate sister-chromatid separation inhibitor involved in transformation and tumorigenesis, *Science* 285 (1999) 418–422.
- [104] J.M. Peters, The anaphase promoting complex/cyloosome: a machine designed to destroy, *Nat. Rev. Mol. Cell Biol.* 7 (2006) 644–656.
- [105] S. Hauf, I.C. Waizenegger, J.M. Peters, Cohesin cleavage by separase required for anaphase and cytokinesis in human cells, *Science* 293 (2001) 1320–1323.
- [106] Z. Karamysheva, L.A. Diaz-Martinez, S.E. Crow, B. Li, H. Yu, Multiple anaphase-promoting complex/cyloosome degrons mediate the degradation of human Sgo1, *J. Biol. Chem.* 284 (2009) 1772–1780.
- [107] T.S. Kitajima, S.A. Kawashima, Y. Watanabe, The conserved kinetochore protein shugoshin protects centromeric cohesion during meiosis, *Nature* 427 (2004) 510–517.
- [108] V. Sudakin, D. Ganoth, A. Dahan, H. Heller, J. Hershko, F.C. Luca, J.V. Ruderman, A. Hershko, The cyloosome, a large complex containing cyclin-selective ubiquitin ligase activity, targets cyclins for destruction at the end of mitosis, *Mol. Biol. Cell* 6 (1995) 185–197.
- [109] J.W. Raff, K. Jeffers, J.Y. Huang, The roles of Fzy/Cdc20 and Fzr/Cdh1 in regulating the destruction of cyclin B in space and time, *J. Cell Biol.* 157 (2002) 1139–1149.
- [110] N. den Elzen, J. Pines, Cyclin A is destroyed in prometaphase and can delay chromosome alignment and anaphase, *J. Cell Biol.* 153 (2001) 121–136.
- [111] P. Clute, J. Pines, Temporal and spatial control of cyclin B1 destruction in metaphase, *Nat. Cell Biol.* 1 (1999) 82–87.
- [112] A. Seki, J.A. Coppinger, H. Du, C.Y. Jang, J.R. Yates III, G. Fang, Plk1- and beta-TrCP-dependent degradation of Bora controls mitotic progression, *J. Cell Biol.* 181 (2008) 65–78.
- [113] E.H. Chan, A. Santamaria, H.H. Sillje, E.A. Nigg, Plk1 regulates mitotic Aurora A function through betaTrCP-dependent degradation of hBora, *Chromosoma* 117 (2008) 457–469.
- [114] T.J. McGarry, M.W. Kirschner, Geminin, an inhibitor of DNA replication, is degraded during mitosis, *Cell* 93 (1998) 1043–1053.
- [115] E.R. Kramer, N. Scheuringer, A.V. Podtelejnikov, M. Mann, J.M. Peters, Mitotic regulation of the APC activator proteins CDC20 and CDH1, *Mol. Biol. Cell* 11 (2000) 1555–1569.
- [116] A. Zur, M. Brandeis, Timing of APC/C substrate degradation is determined by fzy/fzr specificity of destruction boxes, *EMBO J.* 21 (2002) 4500–4510.
- [117] M. Rape, S.K. Reddy, M.W. Kirschner, The processivity of multiubiquitination by the APC determines the order of substrate degradation, *Cell* 124 (2006) 89–103.
- [118] I. Sumara, S. Maerki, M. Peter, E3 ubiquitin ligases and mitosis: embracing the complexity, *Trends Cell Biol.* 18 (2008) 84–94.
- [119] S. Angers, T. Li, X. Yi, M.J. MacCoss, R.T. Moon, N. Zheng, Molecular architecture and assembly of the DDB1-CUL4A ubiquitin ligase machinery, *Nature* 443 (2006) 590–593.
- [120] Y.J. He, C.M. McCall, J. Hu, Y. Zeng, Y. Xiong, DDB1 functions as a linker to recruit receptor WD40 proteins to CUL4-ROC1 ubiquitin ligases, *Genes Dev.* 20 (2006) 2949–2954.
- [121] J. Lee, P. Zhou, DCAFs, the missing link of the CUL4-DDB1 ubiquitin ligase, *Mol. Cell* 26 (2007) 775–780.
- [122] E. Shibata, T. Abbas, X. Huang, J.A. Wohlschlegel, A. Dutta, Selective ubiquitylation of p21 and Cdt1 by UBC9 and UBE2G ubiquitin-conjugating enzymes via the CRL4Cdt2 ubiquitin ligase complex, *Mol. Cell Biol.* 31 (2011) 3136–3145.
- [123] H. Nishitani, N. Sugimoto, V. Roukos, Y. Nakanishi, M. Saijo, C. Obuse, T. Tsurimoto, K.I. Nakayama, K. Nakayama, M. Fujita, Z. Lygerou, T. Nishimoto, Two E3 ubiquitin ligases, SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis, *EMBO J.* 25 (2006) 1126–1136.
- [124] E.E. Arias, J.C. Walter, PCNA functions as a molecular platform to trigger Cdt1 destruction and prevent re-replication, *Nat. Cell Biol.* 8 (2006) 84–90.
- [125] J. Jin, E.E. Arias, J. Chen, J.W. Harper, J.C. Walter, A family of diverse Cul4-Ddb1-interacting proteins includes Cdt2, which is required for S phase destruction of the replication factor Cdt1, *Mol. Cell* 23 (2006) 709–721.
- [126] W. Zhong, H. Feng, F.E. Santiago, E.T. Kipreos, CUL-4 ubiquitin ligase maintains genome stability by restraining DNA-replication licensing, *Nature* 423 (2003) 885–889.
- [127] J. Hu, C.M. McCall, T. Ohta, Y. Xiong, Targeted ubiquitination of CDT1 by the DDB1-CUL4A-ROC1 ligase in response to DNA damage, *Nat. Cell Biol.* 6 (2004) 1003–1009.
- [128] X. Li, Q. Zhao, R. Liao, P. Sun, X. Wu, The SCF(Skp2) ubiquitin ligase complex interacts with the human replication licensing factor Cdt1 and regulates Cdt1 degradation, *J. Biol. Chem.* 278 (2003) 30854–30858.
- [129] T. Abbas, U. Sivaprasad, K. Terai, V. Amador, M. Pagano, A. Dutta, PCNA-dependent regulation of p21 ubiquitylation and degradation via the CRL4Cdt2 ubiquitin ligase complex, *Genes Dev.* 22 (2008) 2496–2506.
- [130] T. Abbas, E. Shibata, J. Park, S. Jha, N. Karnani, A. Dutta, CRL4(Cdt2) regulates cell proliferation and histone gene expression by targeting PR-Set7/Set8 for degradation, *Mol. Cell* 40 (2010) 9–21.
- [131] R.C. Centore, C.G. Havens, A.L. Manning, J.M. Li, R.L. Flynn, A. Tse, J. Jin, N.J. Dyson, J.C. Walter, L. Zou, CRL4(Cdt2)-mediated destruction of the histone methyltransferase Set8 prevents premature chromatin compaction in S phase, *Mol. Cell* 40 (2010) 22–33.
- [132] S. Jorgensen, M. Eskildsen, K. Fugger, L. Hansen, M.S. Larsen, A.N. Kousholt, R.G. Syljuuasen, M.B. Trelle, O.N. Jensen, K. Helin, C.S. Sorensen, SET8 is degraded via PCNA-coupled CRL4(CDT2) ubiquitylation in S phase and after UV irradiation, *J. Cell Biol.* 192 (2011) 43–54.
- [133] T. Abbas, A. Mueller, E. Shibata, M. Keaton, M. Rossi, A. Dutta, CRL1-FBXO11 promotes Cdt2 ubiquitylation and degradation and regulates Pr-Set7/Set8-mediated cellular migration, *Mol. Cell* (in press), <http://dx.doi.org/10.1016/j.molcel.2013.02.003> (Epub 2013 Mar 7).
- [134] M. Rossi, S. Duan, Y.T. Jeong, M. Horn, A. Saraf, L. Florens, M.P. Washburn, A. Antebi, M. Pagano, Regulation of the CRL4Cdt2 ubiquitin ligase and cell cycle exit by the SCFFbxo11 ubiquitin ligase, *Mol. Cell* (in press), <http://dx.doi.org/10.1016/j.molcel.2013.02.004> (Epub 2013 Mar 7).
- [135] S.V. Rajkumar, P.G. Richardson, T. Hideshima, K.C. Anderson, Proteasome inhibition as a novel therapeutic target in human cancer, *J. Clin. Oncol.* 23 (2005) 630–639.

- [136] P. Moreau, P.G. Richardson, M. Cavo, R.Z. Orlowski, J.F. San Miguel, A. Palumbo, J.L. Housseau, Proteasome inhibitors in multiple myeloma: 10 years later, *Blood* 120 (2012) 947–959.
- [137] K.I. Nakayama, K. Nakayama, Ubiquitin ligases: cell-cycle control and cancer, *Nat. Rev. Cancer* 6 (2006) 369–381.
- [138] J.R. Skaar, V. D'Angiolella, J.K. Pagan, M. Pagano, SnapShot: F box proteins II, *Cell* 137 (2009) 1358, (1358 e1351).
- [139] T.A. Masuda, H. Inoue, H. Sonoda, S. Mine, Y. Yoshikawa, K. Nakayama, K. Nakayama, M. Mori, Clinical and biological significance of S-phase kinase-associated protein 2 (Skp2) gene expression in gastric carcinoma: modulation of malignant phenotype by Skp2 overexpression, possibly via p27 proteolysis, *Cancer Res.* 62 (2002) 3819–3825.
- [140] T.A. Masuda, H. Inoue, K. Nishida, H. Sonoda, Y. Yoshikawa, Y. Kakeji, T. Utsunomiya, M. Mori, Cyclin-dependent kinase 1 gene expression is associated with poor prognosis in gastric carcinoma, *Clin. Cancer Res.* 9 (2003) 5693–5698.
- [141] M. Shapira, O. Ben-Izhak, S. Linn, B. Futerman, I. Minkov, D.D. Hershko, The prognostic impact of the ubiquitin ligase subunits Skp2 and Cks1 in colorectal carcinoma, *Cancer* 103 (2005) 1336–1346.
- [142] S. Signorelli, L. Di Marcotullio, A. Richardson, S. Ramaswamy, B. Isaac, M. Rue, F. Monti, M. Loda, M. Pagano, Oncogenic role of the ubiquitin ligase subunit Skp2 in human breast cancer, *J. Clin. Invest.* 110 (2002) 633–641.
- [143] E. Latres, R. Chiarle, B.A. Schulman, N.P. Pavletich, A. Pellicer, G. Inghirami, M. Pagano, Role of the F-box protein Skp2 in lymphomagenesis, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 2515–2520.
- [144] E.H. Shim, L. Johnson, H.L. Noh, Y.J. Kim, H. Sun, C. Zeiss, H. Zhang, Expression of the F-box protein SKP2 induces hyperplasia, dysplasia, and low-grade carcinoma in the mouse prostate, *Cancer Res.* 63 (2003) 1583–1588.
- [145] I. Timmerbeul, C.M. Garrett-Engel, U. Kossatz, X. Chen, E. Firpo, V. Grunwald, K. Kamino, L. Wilkens, U. Lehmann, J. Buer, R. Geffers, S. Kubicka, M.P. Manns, P.L. Porter, J.M. Roberts, N.P. Malek, Testing the importance of p27 degradation by the SCFskp2 pathway in murine models of lung and colon cancer, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 14009–14014.
- [146] H.K. Lin, Z. Chen, G. Wang, C. Nardella, S.W. Lee, C.H. Chan, W.L. Yang, J. Wang, A. Egia, K.I. Nakayama, C. Cordon-Cardo, J. Teruya-Feldstein, P.P. Pandolfi, Skp2 targeting suppresses tumorigenesis by Arf-p53-independent cellular senescence, *Nature* 464 (2010) 374–379.
- [147] S.H. Lee, F. McCormick, Downregulation of Skp2 and p27/Kip1 synergistically induces apoptosis in T98G glioblastoma cells, *J. Mol. Med.* 83 (2005) 296–307.
- [148] Y. Katagiri, Y. Hozumi, S. Kondo, Knockdown of Skp2 by siRNA inhibits melanoma cell growth in vitro and in vivo, *J. Dermatol. Sci.* 42 (2006) 215–224.
- [149] Y. Kudo, S. Kitajima, I. Ogawa, M. Kitagawa, M. Miyachi, T. Takata, Small interfering RNA targeting of S phase kinase-interacting protein 2 inhibits cell growth of oral cancer cells by inhibiting p27 degradation, *Mol. Cancer Ther.* 4 (2005) 471–476.
- [150] M. Barry, K. Fruh, S.T.K.E. Sci, Viral modulators of cullin RING ubiquitin ligases: culling the host defense, *Sci. STKE* (2006) pe21.
- [151] D. Frescas, M. Pagano, Deregulated proteolysis by the F-box proteins SKP2 and beta-TRCP: tipping the scales of cancer, *Nat. Rev. Cancer* 8 (2008) 438–449.
- [152] J.T. Winston, P. Strack, P. Beer-Romero, C.Y. Chu, S.J. Elledge, J.W. Harper, The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha ubiquitination in vitro, *Genes Dev.* 13 (1999) 270–283.
- [153] W. Tang, Y. Li, D. Yu, A. Thomas-Tikhonenko, V.S. Spiegelman, S.Y. Fuchs, Targeting beta-transducin repeat-containing protein E3 ubiquitin ligase augments the effects of antitumor drugs on breast cancer cells, *Cancer Res.* 65 (2005) 1904–1908.
- [154] Y. Kudo, D. Guardavaccaro, P.G. Santamaria, R. Koyama-Nasu, E. Latres, R. Bronson, L. Yamasaki, M. Pagano, Role of F-box protein betaTRCP1 in mammary gland development and tumorigenesis, *Mol. Cell. Biol.* 24 (2004) 8184–8194.
- [155] A. Ougolkov, B. Zhang, K. Yamashita, V. Bilim, M. Mai, S.Y. Fuchs, T. Minamoto, Associations among beta-TRCP, an E3 ubiquitin ligase receptor, beta-catenin, and NF-kappaB in colorectal cancer, *J. Natl. Cancer Inst.* 96 (2004) 1161–1170.
- [156] S. Muerkoster, A. Arlt, B. Sipos, M. Witt, M. Grossmann, G. Kloppel, H. Kalthoff, U.R. Folsch, H. Schafer, Increased expression of the E3-ubiquitin ligase receptor subunit betaTRCP1 relates to constitutive nuclear factor-kappaB activation and chemoresistance in pancreatic carcinoma cells, *Cancer Res.* 65 (2005) 1316–1324.
- [157] N. Beldaidouni, M. Peuchmaur, C. Perret, A. Florentin, R. Benarous, C. Besnard-Guerin, Overexpression of human beta TRCP1 deleted of its F box induces tumorigenesis in transgenic mice, *Oncogene* 24 (2005) 2271–2276.
- [158] C. Besnard-Guerin, N. Beldaidouni, I. Lassot, E. Segeral, A. Jobart, C. Marchal, R. Benarous, HIV-1 Vpu sequesters beta-transducin repeat-containing protein (betaTRCP) in the cytoplasm and provokes the accumulation of beta-catenin and other SCFbetaTRCP substrates, *J. Biol. Chem.* 279 (2004) 788–795.
- [159] A. Palumbo, K. Anderson, Multiple myeloma, *N. Engl. J. Med.* 364 (2011) 1046–1060.
- [160] A.M. Ruschak, M. Slassi, L.E. Kay, A.D. Schimmer, Novel proteasome inhibitors to overcome bortezomib resistance, *J. Natl. Cancer Inst.* 103 (2011) 1007–1017.
- [161] R. Battistutta, Protein kinase CK2 in health and disease: structural bases of protein kinase CK2 inhibition, *Cell Mol. Life Sci.* 66 (2009) 1868–1889.
- [162] L. Busino, S.E. Millman, L. Scotto, C.A. Kyratsous, V. Basur, O. O'Connor, A. Hoffmann, K.S. Elenitoba-Johnson, M. Pagano, Fbxw7alpha- and GSK3-mediated degradation of p100 is a pro-survival mechanism in multiple myeloma, *Nat. Cell Biol.* 14 (2012) 375–385.
- [163] B.J. Thompson, S. Buonamici, M.L. Sulis, T. Palomero, T. Vilimas, G. Basso, A. Ferrando, I. Aifantis, The SCFFBW7 ubiquitin ligase complex as a tumor suppressor in T cell leukemia, *J. Exp. Med.* 204 (2007) 1825–1835.
- [164] H. Rajagopalan, P.V. Jallepalli, C. Rago, V.E. Velculescu, K.W. Kinzler, B. Vogelstein, C. Lengauer, Inactivation of hCDC4 can cause chromosomal instability, *Nature* 428 (2004) 77–81.
- [165] J. O'Neil, J. Grim, P. Strack, S. Rao, D. Tibbitts, C. Winter, J. Hardwick, M. Welcher, J.P. Meijerink, R. Pieters, G. Draetta, R. Sears, B.E. Clunman, A.T. Look, FBW7 mutations in leukemic cells mediate NOTCH pathway activation and resistance to gamma-secretase inhibitors, *J. Exp. Med.* 204 (2007) 1813–1824.
- [166] M.A. Chapman, M.S. Lawrence, J.J. Keats, K. Cibulskis, C. Sougnez, A.C. Schinzel, C.L. Harview, J.P. Brunet, G.J. Ahmann, M. Adli, K.C. Anderson, K.G. Ardlie, D. Auclair, A. Baker, P.L. Bergsagel, B.E. Bernstein, Y. Drier, R. Fonseca, S.L. Gabriel, C.C. Hofmeister, S. Jagannath, A.J. Jakubowski, A. Krishnan, J. Levy, T. Liefeld, S. Lonial, S. Mahan, B. Mfuko, S. Monti, L.M. Perkins, R. Onofrio, T.J. Pugh, S.V. Rajkumar, A.H. Ramos, D.S. Siegel, A. Sivachenko, A.K. Stewart, S. Trudel, R. Vij, D. Voet, W. Winckler, T. Zimmerman, J. Carpten, J. Trent, W.C. Hahn, L.A. Garraway, M. Meyerson, E.S. Lander, G. Getz, T.R. Golub, Initial genome sequencing and analysis of multiple myeloma, *Nature* 471 (2011) 467–472.
- [167] S. Duan, L. Cermak, J.K. Pagan, M. Rossi, C. Martinengo, P.F. di Celle, B. Chapuy, M. Shipp, R. Chiarle, M. Pagano, FBXO11 targets BCL6 for degradation and is inactivated in diffuse large B-cell lymphomas, *Nature* 481 (2012) 90–93.
- [168] J.G. Lohr, P. Stojanov, M.S. Lawrence, D. Auclair, B. Chapuy, C. Sougnez, P. Cruz-Gordillo, B. Knoechel, Y.W. Asmann, S.L. Slager, A.J. Novak, A. Dogan, S.M. Ansell, B.K. Link, L. Zou, J. Gould, G. Saksena, N. Stransky, C. Rangel-Escareno, J.C. Fernandez-Lopez, A. Hidalgo-Miranda, J. Melendez-Zajgla, E. Hernandez-Lemus, A. Schwarz-Cruz y Celis, I. Imaz-Rosshandler, A.I. Ojesina, J. Jung, C.S. Pedamallu, E.S. Lander, T.M. Habermann, J.R. Cerhan, M.A. Shipp, G. Getz, T.R. Golub, Discovery and prioritization of somatic mutations in diffuse large B-cell lymphoma (DLBCL) by whole-exome sequencing, *Proc. Natl. Acad. Sci. U. S. A.* (2012) 3879–3884.
- [169] Integrated genomic analyses of ovarian carcinoma, *Nature* 474 (2011) 609–615.
- [170] N. Stransky, A.M. Egloff, A.D. Tward, A.D. Kostic, K. Cibulskis, A. Sivachenko, G.V. Kryukov, M.S. Lawrence, C. Sougnez, A. McKenna, E. Shefler, A.H. Ramos, P. Stojanov, S.L. Carter, D. Voet, M.L. Cortes, D. Auclair, M.F. Berger, G. Saksena, C. Guiducci, R.C. Onofrio, M. Parkin, M. Romkes, J.L. Weissfeld, R.R. Seethala, L. Wang, C. Rangel-Escareno, J.C. Fernandez-Lopez, A. Hidalgo-Miranda, J. Melendez-Zajgla, W. Winckler, K. Ardlie, S.B. Gabriel, M. Meyerson, E.S. Lander, G. Getz, T.R. Golub, L.A. Garraway, J.R. Grandis, The mutational landscape of head and neck squamous cell carcinoma, *Science* 333 (2011) 1157–1160.
- [171] Z. Kan, B.S. Jaiswal, J. Stinson, V. Janakiraman, D. Bhatt, H.M. Stern, P. Yue, P.M. Havery, R. Bourgon, J. Zheng, M. Moorhead, S. Chaudhuri, L.P. Tomsho, B.A. Peters, K. Pujara, S. Cordes, D.P. Davis, V.E. Carlton, W. Yuan, L. Li, W. Wang, C. Eigenbrot, J.S. Kaminker, D.A. Eberhard, P. Waring, S.C. Schuster, Z. Modrusan, Z. Zhang, D. Stokoe, F.J. de Sauvage, M. Faham, S. Seshagiri, Diverse somatic mutation patterns and pathway alterations in human cancers, *Nature* 466 (2010) 869–873.
- [172] O. Barbash, P. Zamfirova, D.I. Lin, X. Chen, K. Yang, H. Nakagawa, F. Lu, A.K. Rustgi, J.A. Diehl, Mutations in Fbx4 inhibit dimerization of the SCF(Fbx4) ligase and contribute to cyclin D1 overexpression in human cancer, *Cancer Cell* 14 (2008) 68–78.
- [173] O. Barbash, J.A. Diehl, SCF(Fbx4)/alphaB-crystallin E3 ligase: when one is not enough, *Cell Cycle* 7 (2008) 2983–2986.
- [174] J. Fu, H. Qiu, M. Cai, Y. Pan, Y. Cao, L. Liu, J. Yun, C.Z. Zhang, Low cyclin F expression in hepatocellular carcinoma associates with poor differentiation and unfavorable prognosis, *Cancer Sci.* (2013), <http://dx.doi.org/10.1111/cas.12100>, (Epub 2013 Feb 17).
- [175] Y. Yang, J. Kitagaki, R.M. Dai, Y.C. Tsai, K.L. Lorick, R.L. Ludwig, S.A. Pierre, J.P. Jensen, I.V. Davydov, P. Oberoi, C.C. Li, J.H. Kerten, J.A. Beutler, K.H. Vousden, A.M. Weissman, Inhibitors of ubiquitin-activating enzyme (E1), a new class of potential cancer therapeutics, *Cancer Res.* 67 (2007) 9472–9481.
- [176] T.A. Soucy, P.G. Smith, M.A. Milhollen, A.J. Berger, J.M. Gavin, S. Adhikari, J.E. Brownell, K.E. Burke, D.P. Cardin, S. Critchley, C.A. Cullis, A. Doucette, J.J. Garsney, J.L. Gaulin, R.E. Gershman, A.R. Lublinsky, A. McDonald, H. Mizutani, U. Narayanan, E.J. Olhava, S. Peluso, M. Rezaei, M.D. Sintchak, T. Talreja, M.P. Thomas, T. Traore, S. Vyskocil, G.S. Weatherhead, J. Yu, J. Zhang, L.R. Dick, C.F. Claiborne, M. Rolfe, J.B. Bolen, S.P. Langston, An inhibitor of NEDD8-activating enzyme as a new approach to treat cancer, *Nature* 458 (2009) 732–736.
- [177] S.J. Goldenberg, T.C. Cascio, S.D. Shumway, K.C. Garbutt, J. Liu, Y. Xiong, N. Zheng, Structure of the Cnd1-Cul1-Roc1 complex reveals regulatory mechanisms for the assembly of the multisubunit cullin-dependent ubiquitin ligases, *Cell* 119 (2004) 517–528.
- [178] D.M. Duda, L.A. Borg, D.C. Scott, H.W. Hunt, M. Hammel, B.A. Schulman, Structural insights into NEDD8 activation of cullin-RING ligases: conformational control of conjugation, *Cell* 134 (2008) 995–1006.
- [179] T.A. Soucy, P.G. Smith, M. Rolfe, Targeting NEDD8-activated cullin-RING ligases for the treatment of cancer, *Clin. Cancer Res.* 15 (2009) 3912–3916.
- [180] R.T. Swords, K.R. Kelly, P.G. Smith, J.J. Garsney, D. Mahalingam, E. Medina, K. Oberheu, S. Padmanabhan, M. O'Dwyer, S.T. Nawrocki, F.J. Giles, J.S. Carew, Inhibition of NEDD8-activating enzyme: a novel approach for the treatment of acute myeloid leukemia, *Blood* 115 (2010) 3796–3800.
- [181] Y. Ye, M. Rape, Building ubiquitin chains: E2 enzymes at work, *Nat. Rev. Mol. Cell Biol.* 10 (2009) 755–764.
- [182] D.F. Ceccarelli, X. Tang, B. Pelletier, S. Orlicky, W. Xie, V. Plantevin, D. Neculai, Y.C. Chou, A. Ogunjimi, A. Al-Hakim, X. Varelis, J. Koszela, G.A. Wasney, M. Vedadi, S. Dhe-Paganon, S. Cox, S. Xu, A. Lopez-Girona, F. Mercurio, J. Wrana, D. Durocher, S. Meloche, D.R. Webb, M. Tyers, F. Sicheri, An allosteric inhibitor of the human Cdc34 ubiquitin-conjugating enzyme, *Cell* 145 (2011) 1075–1087.

- [183] D. Gonzalez-Ruiz, H. Gohlke, Targeting protein–protein interactions with small molecules: challenges and perspectives for computational binding epitope detection and ligand finding, *Curr. Med. Chem.* 13 (2006) 2607–2625.
- [184] L.T. Vassilev, B.T. Vu, B. Graves, D. Carvajal, F. Podlaski, Z. Filipovic, N. Kong, U. Kammlott, C. Lukacs, C. Klein, N. Fotouhi, E.A. Liu, In vivo activation of the p53 pathway by small-molecule antagonists of MDM2, *Science* 303 (2004) 844–848.
- [185] K.G. Wiman, Strategies for therapeutic targeting of the p53 pathway in cancer, *Cell Death Differ.* 13 (2006) 921–926.
- [186] S. Shangary, D. Qin, D. McEachern, M. Liu, R.S. Miller, S. Qiu, Z. Nikolovska-Coleska, K. Ding, G. Wang, J. Chen, D. Bernard, J. Zhang, Y. Lu, Q. Gu, R.B. Shah, K.J. Pienta, X. Ling, S. Kang, M. Guo, Y. Sun, D. Yang, S. Wang, Temporal activation of p53 by a specific MDM2 inhibitor is selectively toxic to tumors and leads to complete tumor growth inhibition, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 3933–3938.
- [187] Q. Chen, W. Xie, D.J. Kuhn, P.M. Voorhees, A. Lopez-Girona, D. Mendy, L.G. Corral, V.P. Krenitsky, W. Xu, L. Moutouh-de Parseval, D.R. Webb, F. Mercurio, K.I. Nakayama, K. Nakayama, R.Z. Orlowski, Targeting the p27 E3 ligase SCF(Skp2) results in p27- and Skp2-mediated cell-cycle arrest and activation of autophagy, *Blood* 111 (2008) 4690–4699.
- [188] S. Orlicky, X. Tang, V. Neduva, N. Elowe, E.D. Brown, F. Sicheri, M. Tyers, An allosteric inhibitor of substrate recognition by the SCF(Cdc4) ubiquitin ligase, *Nat. Biotechnol.* 28 (2010) 733–737.
- [189] L. Wu, A.V. Grigoryan, Y. Li, B. Hao, M. Pagano, T. Cardozo, Specific Small Molecule Inhibitors of Skp2-Mediated p27 Degradation, *Chem. Biol.* 19 (2012) 1515–1524.
- [190] M. Aghajan, N. Jonai, K. Flick, F. Fu, M. Luo, X. Cai, I. Ouni, N. Pierce, X. Tang, B. Lomenick, R. Damoiseaux, R. Hao, P.M. Del Moral, R. Verma, Y. Li, C. Li, K.N. Houk, M.E. Jung, N. Zheng, L. Huang, R.J. Deshaies, P. Kaiser, J. Huang, Chemical genetics screen for enhancers of rapamycin identifies a specific inhibitor of an SCF family E3 ubiquitin ligase, *Nat. Biotechnol.* 28 (2010) 738–742.
- [191] R.J. Deshaies, C.A. Joazeiro, RING domain E3 ubiquitin ligases, *Annu. Rev. Biochem.* 78 (2009) 399–434.
- [192] W.W. Ma, W.A. Messersmith, G.K. Dy, C.D. Weekes, A. Whitworth, C. Ren, M. Maniar, F. Wilhelm, S.G. Eckhardt, A.A. Adjei, A. Jimeno, Phase I study of Rigosertib, an inhibitor of the phosphatidylinositol 3-kinase and Polo-like kinase 1 pathways, combined with gemcitabine in patients with solid tumors and pancreatic cancer, *Clin. Cancer Res.* 18 (2012) 2048–2055.