


# Cell Reports

## The DNA Sensor AIM2 Maintains Intestinal Homeostasis via Regulation of Epithelial Antimicrobial Host Defense

### Graphical Abstract



### Authors

Shuiqing Hu, Lan Peng, Youn-Tae Kwak, ..., James S. Malter, Lora V. Hooper, Md. Hasan Zaki

### Correspondence

hasan.zaki@utsouthwestern.edu

### In Brief

AIM2 is a cytosolic DNA sensor. Hu et al. demonstrate that intestinal microbial DNA activates the AIM2 inflammasome. AIM2 activation leads to the production of IL-1 $\beta$  and IL-18, which participate in the regulation of intestinal microbiota such as *Escherichia coli* via induction of the antimicrobial peptides in intestinal epithelial cells.

### Highlights

- AIM2 senses intestinal microbial DNA and activates the inflammasome
- The AIM2 inflammasome protects mice from experimental colitis
- The AIM2 inflammasome controls the growth of commensal *Escherichia coli*
- The inflammasome triggers antimicrobial peptide production by epithelial cells



# The DNA Sensor AIM2 Maintains Intestinal Homeostasis via Regulation of Epithelial Antimicrobial Host Defense

Shuiqing Hu,<sup>1</sup> Lan Peng,<sup>1</sup> Youn-Tae Kwak,<sup>1</sup> Erin McElvania Tekippe,<sup>1,2</sup> Chandrashekhar Pasare,<sup>3</sup> James S. Malter,<sup>1</sup> Lora V. Hooper,<sup>3,4</sup> and Md. Hasan Zaki<sup>1,\*</sup>

<sup>1</sup>Department of Pathology, UT Southwestern Medical Center, Dallas, TX 75390, USA

<sup>2</sup>Children's Medical Center, Dallas, TX 75390, USA

<sup>3</sup>Department of Immunology, UT Southwestern Medical Center, Dallas, TX 75390, USA

<sup>4</sup>The Howard Hughes Medical Institute, UT Southwestern Medical Center, Dallas, TX 75390, USA

\*Correspondence: [hasan.zaki@utsouthwestern.edu](mailto:hasan.zaki@utsouthwestern.edu)

<http://dx.doi.org/10.1016/j.celrep.2015.10.040>

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## SUMMARY

Microbial pattern molecules in the intestine play immunoregulatory roles via diverse pattern recognition receptors. However, the role of the cytosolic DNA sensor AIM2 in the maintenance of intestinal homeostasis is unknown. Here, we show that *Aim2*<sup>-/-</sup> mice are highly susceptible to dextran sodium sulfate-induced colitis that is associated with microbial dysbiosis as represented by higher colonic burden of commensal *Escherichia coli*. Colonization of germ-free mice with *Aim2*<sup>-/-</sup> mouse microbiota leads to higher colitis susceptibility. In-depth investigation of AIM2-mediated host defense responses reveals that caspase-1 activation and IL-1 $\beta$  and IL-18 production are compromised in *Aim2*<sup>-/-</sup> mouse colons, consistent with defective inflammasome function. Moreover, IL-18 infusion reduces *E. coli* burden as well as colitis susceptibility in *Aim2*<sup>-/-</sup> mice. Altered microbiota in inflammasome-defective mice correlate with reduced expression of several antimicrobial peptides in intestinal epithelial cells. Together, these findings implicate DNA sensing by AIM2 as a regulatory mechanism for maintaining intestinal homeostasis.

## INTRODUCTION

The intestinal mucosal immune system of mammals evolved to coexist with densely populated microorganisms that reside in the intestinal lumen. The central physiological process for homeostatic immune response in the gut is the recognition of pathogen-associated molecular patterns (PAMPs) by host pattern recognition receptors (PRRs). There are several evolutionary conserved PRRs including Toll-like receptors (TLRs), NOD-like receptors (NLRs), RIG-I-like receptors (RLRs), C-type lectin-like receptors (CLRs), and HIN-200 family receptors. Defects in pathogen sensing systems lead to dysregulated immune responses in the intestine, resulting in the induction of intestinal inflammatory

disorders such as inflammatory bowel disease (IBD) and colorectal cancer (Cho, 2008). Clinical and experimental studies have shown that dysfunctions in several TLRs (TLR2, TLR4, and TLR9) and NLRs (NOD2, NLRP3, NLRP6, and NLRP12) are associated with the pathogenesis of intestinal inflammation and tumorigenesis (Chen et al., 2008; Elinav et al., 2011; Hugot et al., 2001; Katakura et al., 2005; Maeda et al., 2005; Rakoff-Nahoum et al., 2004; Vijay-Kumar et al., 2007; Villani et al., 2009; Zaki et al., 2010a, 2011b). However, the precise mechanisms of the regulation of intestinal homeostasis by these pathogen sensors are not clearly defined, and the functions of the majority of PRRs in intestinal physiology have yet to be explored.

Absent in melanoma 2 (AIM2), a member of interferon-inducible gene HIN-200 family, has been implicated as a cytosolic sensor of DNA (Bürckstümmer et al., 2009; Fernandes-Alnemri et al., 2009; Hornung et al., 2009). Recent studies have underscored the importance of AIM2-mediated sensing of microbial DNA in host defense responses against bacterial and viral infections caused by *Listeria monocytogenes*, *Francisella tularensis*, *Streptococcus pneumoniae*, and murine cytomegalovirus (Fang et al., 2014; Fernandes-Alnemri et al., 2010; Jones et al., 2010; Kim et al., 2010; Park et al., 2014; Rathinam et al., 2010). Structurally, AIM2 contains an N-terminal pyrin domain (PYD) and a C-terminal oligonucleotide binding HIN domain. When cytosolic dsDNA binds to its HIN domain, AIM2 recruits apoptosis speck-like protein containing a caspase recruitment domain (ASC) and caspase-1 to form a molecular platform called the inflammasome. In addition to AIM2, at least three NLR family members—NLRP1, NLRP3, and NLRC4—are known to activate the inflammasome in biological systems (Lamkanfi and Dixit, 2009). Inflammasome-mediated activation of caspase-1 is required to cleave pro-IL-1 $\beta$  and pro-IL-18 into their active forms. Mounting evidence points to the critical roles caspase-1 and other inflammatory caspases play in regulating intestinal inflammation and tumorigenesis (Allen et al., 2010; Dupaul-Chicoine et al., 2010; Elinav et al., 2011; Salcedo et al., 2010; Zaki et al., 2010a). We and others previously demonstrated that the NLRP3 inflammasome regulates intestinal inflammation and tumorigenesis in mice (Allen et al., 2010; Hirota et al., 2011; Zaki et al., 2010a, 2010b). Recent studies showed that the

inflammasome in the gut can be activated by other PRRs, such as NLRP1 and NLRP6, which also contribute to intestinal homeostasis (Elinav et al., 2011; Williams et al., 2015). However, until now, DNA-dependent activation of the AIM2 inflammasome and its physiological relevance to intestinal homeostasis in the gut has not been reported.

Here, we investigated the physiological function of AIM2 in the gut. Our results show that *Aim2*<sup>-/-</sup> mice are highly susceptible to DSS-induced colitis. AIM2-mediated protection against colonic inflammation was associated with its function as an activator of the inflammasome, since caspase-1 activation and IL-18 production were remarkably attenuated in the absence of AIM2. Interestingly, the colons of *Aim2*<sup>-/-</sup> and other inflammasome-defective mice, e.g., *Nlrp3*<sup>-/-</sup>, *caspase-1*<sup>-/-</sup> (*caspase1/11*<sup>-/-</sup>), and *Il18*<sup>-/-</sup>, contain significantly higher loads of *Escherichia coli*, an indication of dysbiosis in their gut. Transfer of gut microbiota from *Aim2*<sup>-/-</sup> to wild-type (WT) mice confers higher colitis susceptibility in the WT mice, suggesting that the AIM2 inflammasome plays a critical role in the regulation of intestinal inflammation via shaping the intestinal microbial ecology. Further investigation revealed that the AIM2 inflammasome regulates the expression of antimicrobial peptides (AMPs), including C-type lectins (Reg3β and Reg3γ), calprotectin (S100A8 and S100A9), and lipocalin 2 (Lcn2) by intestinal epithelial cells. Overall, our study elucidates a major inflammasome activation pathway in the intestine that involves the DNA sensor AIM2 and offers further insights into the mechanism of inflammasome-mediated protection against intestinal inflammation.

## RESULTS

### AIM2 Protects Mice from DSS-Induced Colitis

Previous studies have shown that *Aim2* is highly expressed in the large intestine (Cridland et al., 2012). Our real-time PCR analyses confirm that *Aim2* is ubiquitously expressed in the colon (Figures S1A and S1B). To investigate the physiological function of AIM2 in the colon, we induced colitis in *Aim2*<sup>-/-</sup> mice (Rathinam et al., 2010) by feeding them 3% DSS-containing water for 5 days. Colitis susceptibility in *Aim2*<sup>-/-</sup> and WT mice was evaluated daily by monitoring body weight and measuring clinical features such as diarrhea and occult stool bleeding. *Aim2*<sup>-/-</sup> mice lost significantly more body weight and exhibited increased diarrhea and rectal bleeding compared to WT mice (Figures 1A–1C). Mice were sacrificed at day 8 and examined macroscopically and microscopically for inflammatory changes. *Aim2*<sup>-/-</sup> mouse colons were significantly shorter compared to WT mice (Figure 1D). This was further supported by histological analysis of H&E-stained whole colon tissue sections (Figure 1E). Much of the distal colon in *Aim2*<sup>-/-</sup> mice showed complete crypt destruction, ulceration, and submucosal edema, which were far less severe in WT mice (Figures 1E and 1F). Histopathological analyses demonstrated significantly higher scores for inflammation, ulceration, and crypt distortion in the middle and distal part of the *Aim2*<sup>-/-</sup> mouse colon at day 8 post DSS administration (Figures 1F and 1G). Notably, *Aim2*<sup>-/-</sup> mice exhibited normal colonic histopathology prior to DSS administration (Figure 1E), and the expression of *Aim2* was not altered during colitis (Figures S1C–S1E). Taken together, AIM2 is a critical regulator of intestinal

homeostasis, and *Aim2*-deficiency leads to higher colitis susceptibility and severity in mice.


### Colonic Injury Leads to Hyper-Inflammation in *Aim2*<sup>-/-</sup> Mice

Although cytokine production in the intestine is essential for repair and healing of the injured epithelium, overwhelming production of cytokines and chemokines trigger pathologic mucosal tissue injury and inflammatory disorders (Zaki et al., 2011a). Therefore, the types and levels of different mediators determine the fate of colonic inflammation and damage. To understand whether AIM2 signaling regulates cytokine production, we measured the expression of IL-1β, IL-1α, IL-6, KC, TNFα, MIP2, CCL2, and CXCL10 in colon tissues collected from wild-type and *Aim2*<sup>-/-</sup> mice over the course of colitis (days 0, 3, 5, and 8). Although the initial (day 0) expression of proinflammatory cytokines and chemokines were comparable, IL-1α, IL-1β, IL-6, KC, and MIP2 were significantly increased in *Aim2*<sup>-/-</sup> mice at day 5 and day 8 (Figure 2A). In addition, CXCL10 and CCL2 were also overexpressed in *Aim2*<sup>-/-</sup> mouse colons at day 8 (Figure 2A). Elevated KC and IL-6 levels in *Aim2*<sup>-/-</sup> mouse colon homogenates were confirmed by ELISA at day 5 and day 8 of colitis (Figure 2B). However, type I interferons were not regulated by AIM2 (Figure S2A). MIP2, KC, and CCL2 are involved in the recruitment and migration of macrophages, neutrophils, and lymphocytes. Consistent with the accumulation of these chemokines in *Aim2*<sup>-/-</sup> mouse colons, we observed greater macrophage infiltration as measured by immunohistochemistry (Figure 2C). Furthermore, the inflammatory signaling pathway ERK showed elevated activation in *Aim2*<sup>-/-</sup> mouse colons collected at day 5 and day 8 post DSS as compared to those in WT mice (Figure 2D). However, we did not observe differences in epithelial cell death between WT and *Aim2*<sup>-/-</sup> mice during colitis as measured by TUNEL assay (Figures S2B and S2C).

### AIM2 Regulates Intestinal Microbiota

Colitis is initiated by the invasion of gut commensal flora into the lamina propria, driving the activation of immune cells. To investigate the possibility that increased inflammation in *Aim2*<sup>-/-</sup> mice is due to increased penetration of the intestinal commensal flora into colonic mucosa during DSS-colitis, we measured the number of culturable bacteria in the colon tissue at day 8 after DSS administration. Colon homogenates were cultured onto brain heart infusion (BHI) and blood agar, under aerobic and anaerobic conditions, respectively. There were significantly greater aerobic and anaerobic/facultative anaerobic bacteria present in the *Aim2*<sup>-/-</sup> mouse colon tissue compared to WT mice (Figure 3A). Culture of fecal homogenates revealed a higher bacterial burden in the colonic lumen of *Aim2*<sup>-/-</sup> mice compared to those of WT mice (Figure 3B). Notably, no bacteria were detected in WT or *Aim2*<sup>-/-</sup> mouse colon tissues (data not shown), and there were no significant differences in total stool bacterial burden prior to colitis induction (Figures S3A and S3B).

These above findings led us to hypothesize that increased colitis susceptibility in *Aim2*<sup>-/-</sup> mice is associated with dysregulated host defense response against pre-existing commensal bacteria. Thus, we treated a cohort of WT and *Aim2*<sup>-/-</sup> mice




### Figure 1. *Aim2*<sup>-/-</sup> Mice Are Susceptible to DSS-Induced Colitis

(A–E) WT (n = 17) and *Aim2*<sup>-/-</sup> (n = 12) mice were fed with 3% DSS for 5 days, followed by regular drinking water for 3 days. Body weight (A), stool consistency (B), and rectal bleeding (C) were scored daily. (D) Mice were sacrificed on day 8 to measure colon length. (E) Colon tissues collected from healthy mice (day 0) or at day 8 were stained with H&E. Scale bars, 100  $\mu$ m.

(F and G) Semiquantitative scoring of histopathology (n = 8–10). Data represent mean  $\pm$  SEM of a representative experiment; \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001. See also Figure S1.

with a cocktail of broad-spectrum antibiotics containing vancomycin, neomycin, ampicillin, and metronidazole for 4 weeks as described (Rakoff-Nahoum et al., 2004). Antibiotic-treated or untreated WT and *Aim2*<sup>-/-</sup> mice were then fed with 3% DSS for 5 days. Interestingly, antibiotic-treated WT and *Aim2*<sup>-/-</sup> mice lost minimum body weight as compared to untreated mice (Figure 3C). Although untreated *Aim2*<sup>-/-</sup> mice lost significantly greater body weight, exhibited shorter colons, and suffered

from severe colonic injury and inflammation as compared to those of WT mice, antibiotic treatment completely attenuated the differences between *Aim2*<sup>-/-</sup> and WT mice at day 8 after DSS (Figures 3C–3E). Consistently, the production of proinflammatory cytokines IL-6, KC, MIP2, and CCL2 was reduced to baseline in antibiotic-treated WT and *Aim2*<sup>-/-</sup> mouse colons (Figure S3C). These findings support the idea that increased colitis in *Aim2*<sup>-/-</sup> mice is due to bacterial overgrowth in the colon during acute colitis.



**Figure 2. AIM2-Deficiency Leads to Increased Inflammation in the Colon**

WT and *Aim2*<sup>-/-</sup> mice were fed with 3% DSS in drinking water for 5 days, followed by regular drinking water for 3 days.

(A) Real-time PCR analyses of *IL-1 $\alpha$* , *IL-1 $\beta$* , *IL-6*, *TNF- $\alpha$* , *MIP2*, *Cxcl-10*, and *Ccl2* expression in colons collected at days 0, 3, 5, and 8 after DSS.


(B) Colon homogenates were analyzed for the level of KC and IL-6 by ELISA. Data represent mean  $\pm$  SEM; n = 7–10/group; \*p < 0.05, \*\*p < 0.01.

(C) Colon tissues collected at day 8 was fixed in formalin and immunostained for the macrophage marker F4/80. Brown colors indicate F4/80-positive cells. Scale bars, 100  $\mu$ m.

(D) Whole colon homogenates were analyzed for the activation of ERK by western blotting.

See also Figure S2 and Table S1.





**Figure 3. AIM2 Protects Host against Colitis via Regulation of Colonic Microbiota**

WT and *Aim2*<sup>-/-</sup> mice were fed with 3% DSS in drinking water for 5 days.

(A) Colon homogenates collected at day 8 after DSS were plated on BHI and Blood agar and incubated in aerobic (BHI agar) or anaerobic (Blood agar) conditions at 37°C.

(B) Feces collected from WT and *Aim2*<sup>-/-</sup> mice at day 8 after DSS administration were diluted in PBS and plated on BHI and Blood agar. After overnight culture at 37°C, CFU was counted.

(C–E) WT and *Aim2*<sup>-/-</sup> mice were treated with a combination of antibiotics or left untreated for 4 weeks and then administered with 3% DSS for 5 days. (C) Body weight changes were monitored daily. (D) Mice were sacrificed at day 8 and colon lengths were measured. (E) Colons collected at day 8 after DSS administration

(legend continued on next page)

The composition of bacterial species present in the gut lumen regulates downstream immune responses and affects colitis severity. Accumulating evidence suggests that the presence of certain bacterial species in the gut, such as *E. coli*, *Bacteroides*, *Clostridia*, and *Firmicutes*, augments experimental colitis in mice (Frank et al., 2007; Li et al., 2015; Martinez-Medina and Garcia-Gil, 2014; Walters et al., 2014). A previous study showed that colitis susceptibility in *Nlrp6*<sup>-/-</sup>, *ASC*<sup>-/-</sup>, and *Il18*<sup>-/-</sup> mice was associated with increased number of *Prevotellaceae* and TM7 (Elinav et al., 2011). To examine the possibility that disease severity in *Aim2*<sup>-/-</sup> mice was linked to the colonic load of *Prevotellaceae* and TM7, we measured the relative abundance of 16S rDNA specific to these bacteria by real-time PCR. Interestingly, there was no significant difference in *Prevotellaceae* or TM7 between *Aim2*<sup>-/-</sup> and WT mice (Figure 3F). In addition, we also measured 16S rDNA levels for other known mouse gut commensal and colitogenic bacteria such as *Bacteroides*, *E. coli*, Enterobacteriaceae, *Lactobacillus*, mouse intestinal *Bacteroides* (MIB), *Eubacterium rectale* (EREC), segmented filamentous bacteria (SFB), *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Clostridium*, *Prophyromonas*, and *Bifidobacteria*. With the notable exception of *E. coli* and Enterobacteriaceae, none of those commensal bacteria were significantly different between the two groups of mice (Figures 3F and 3G). However, *E. coli*- and Enterobacteriaceae-specific 16S rDNA was several thousand folds higher in *Aim2*<sup>-/-</sup> mouse feces compared to those in WT mice (Figure 3G). The overgrowth of *E. coli* was confirmed by colony forming units (CFU) assay of fecal homogenates on MacConkey agar, a selective media for Enterobacteriaceae. While *E. coli* was barely detectable in WT mouse stool, *Aim2*<sup>-/-</sup> mice harbored 10<sup>6</sup> CFU of *E. coli* per gram of feces (Figure 3H). DSS administration further enhanced *E. coli* burden in *Aim2*<sup>-/-</sup> mice (Figures 3I and 3J). MacConkey agar colonies were unambiguously identified as *E. coli* by MALDI-TOF (data not shown). Other known colitogenic bacteria such as *Bacteroides*, SFB, *Clostridium*, *Bifidobacterium*, *Prevotellaceae*, and TM7 were not significantly different, even after colitis induction, in either set of mice (Figure S3D). However, increased growth of *E. coli* reflects an alteration of microbial composition in the intestine of *Aim2*<sup>-/-</sup> mice.

### Dysbiosis Regulates Colitis Susceptibility in *Aim2*-Deficient Mice

We postulated that dysbiosis, as evidenced by increased growth of *E. coli* in *Aim2*<sup>-/-</sup> mouse colons, exacerbates colitis development. To verify this hypothesis, we generated *Aim2* heterozygous mice by breeding of *Aim2*<sup>-/-</sup> female and WT male mice. Littermate WT (WT-lit) and *Aim2*<sup>-/-</sup> (*Aim2*<sup>-/-</sup>-lit) mice obtained from breeding of *Aim2* heterozygous mice were then adminis-

tered with 3% DSS for 5 days. WT-lit and *Aim2*<sup>-/-</sup>-lit mice generated in this way should harbor intestinal flora originating from *Aim2*<sup>-/-</sup> mice. Interestingly, WT-lit mice developed pronounced colitis that was phenotypically very similar to the colitis in *Aim2*<sup>-/-</sup> mice as evidenced by body weight loss, clinical scores for diarrhea and rectal bleeding, and shortening of the colon (Figures 4A–4C). Histopathological examination revealed that inflammation, ulceration, edema, and crypt destruction in WT-lit mice were morphologically indistinguishable from that seen in *Aim2*<sup>-/-</sup>-lit mice (Figure 4D). In agreement with comparable colitis susceptibility in littermate mice, an equal burden of *E. coli* was detected in WT-lit and *Aim2*<sup>-/-</sup>-lit mouse colons (Figure 4E), and similar bacterial growth was observed between WT-lit and *Aim2*<sup>-/-</sup>-lit mouse feces (Figures S4A and S4B).

The above results suggested that colitis susceptibility of *Aim2*<sup>-/-</sup> mice might be transferable. To verify this, we co-housed WT and *Aim2*<sup>-/-</sup> mice for 4 weeks. Although adult WT mice were resistant to the maternal transfer of *E. coli* and failed to develop severe colitis (Figures S4C–S4E), colitis susceptibility of co-housed *Aim2*<sup>-/-</sup> mice was more mild than *Aim2*<sup>-/-</sup> mice housed without WT mice (Figures S4D and S4E). This finding suggests that unknown microflora transferred from WT mice to co-housed *Aim2*<sup>-/-</sup> mice plays a protective role against colitis, and exacerbated colitis in *Aim2*<sup>-/-</sup> mice is partly driven by the lack of protective microbiota. To further confirm that altered microbiota is responsible for increased colitis susceptibility in *Aim2*<sup>-/-</sup> mice, we colonized germ-free WT (GF) mice with microbiota present in conventionally raised WT or *Aim2*<sup>-/-</sup> mice. Two weeks after colonization, *E. coli* burden in GF mice colonized with fecal microbiota either from WT (GF-WT) or *Aim2*<sup>-/-</sup> (GF-*Aim2*<sup>-/-</sup>) mice was measured. While GF-*Aim2*<sup>-/-</sup> mice harbored huge *E. coli* burdens, no *E. coli* was detected in GF-WT mice (Figure 4F). Colitis was then induced in these two groups of mice by feeding them with 2.5% DSS for 5 days. We observed that GF-*Aim2*<sup>-/-</sup> mice lost more than 25% of their body weight at day 7 after DSS, while GF-WT mice were relatively resistant to colitis as evaluated by body weight loss, clinical scores, and colon length (Figures 4G–4I). Consistently, there was significantly increased inflammation, production of proinflammatory cytokines, and bacterial growth in GF-*Aim2*<sup>-/-</sup> mouse colons (Figures 4J, 4K, S4F, and S4G). These results shed light on the importance of DNA sensor AIM2 in the regulation of commensal *E. coli* and other uncultured microorganisms that potentially regulate intestinal homeostasis and colitis pathogenesis.

### AIM2 Regulates the Expression of Antimicrobial Peptides in the Colon


Intestinal epithelial cells produce antimicrobial effectors that play a central role in the shaping of the gut microbial community and

were examined histologically. Scale bars, 100  $\mu$ m. Data represent mean  $\pm$  SEM; n = 7–10 and representative of three independent experiments. \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001.

(F and G) DNA was isolated from feces of healthy WT and *Aim2*<sup>-/-</sup> mice. Bacterial 16S rRNA gene sequences were quantitatively measured by real-time PCR. Data represent mean  $\pm$  SEM (n = 8–10/group); \*\*\*p < 0.001.

(H and I) Stools collected from WT and *Aim2*<sup>-/-</sup> mice before or 8 days after DSS administration were diluted in PBS and plated on MacConkey agar plate. Data represent mean  $\pm$  SEM; \*\*\*p < 0.001.

(J) 16S rRNA gene sequences of *E. coli* and Enterobacteriaceae in DNA isolated from stool at day 8 post DSS were quantitatively measured by real-time PCR. See also Figure S3 and Table S2.



**Figure 4. Colitis Susceptibility in *Aim2*<sup>-/-</sup> Mice Is Contributed by Altered Microbiota**

(A–E) Littermate WT (WT-lit; n = 9) and *Aim2*<sup>-/-</sup> (*Aim2*<sup>-/-</sup>-lit; n = 7), non-littermate WT (n = 7) and *Aim2*<sup>-/-</sup> (n = 7) mice were fed with 3% DSS for 5 days, followed by regular drinking water for 3 days. Body weight (A) and stool consistency (B) were scored daily. Mice were sacrificed on day 8 to measure colon length (C) and examined for histopathological changes in colon tissue by H&E staining (D). (E) Fecal pellets collected from healthy littermate WT and *Aim2*<sup>-/-</sup> mice were diluted in PBS and cultured on MacConkey agar plate. Cfu of *E. coli* was counted.

(F–K) Germ-free (GF) mice were co-housed with either conventionally raised WT (GF-WT) or *Aim2*<sup>-/-</sup> mice (GF-*Aim2*<sup>-/-</sup>) for 3 days. (F) Two weeks after colonization, feces were homogenized in PBS and plated on MacConkey agar plate to count *E. coli*. (G–J) Microbiota-colonized GF mice (n = 4/group) were fed with

(legend continued on next page)



protecting mucosal tissues from colonization and invasion of commensal microorganisms (Gallo and Hooper, 2012). To understand why *Aim2*<sup>-/-</sup> mouse colons harbor altered microbiota, we examined whether AIM2 regulates goblet cell function by using periodic acid-Schiff (PAS) staining (Figure 5A). We found the number of PAS-positive cells was comparable between WT and *Aim2*<sup>-/-</sup> mouse colons collected at day 5 following DSS-induced colitis (Figure 5B). We also did not see any difference in the levels of mucins (*Muc1*, *Muc2*, *Muc3*, and *Muc4*) present in WT and *Aim2*<sup>-/-</sup> mouse colons at baseline or following colitis induction (Figure 5C). Autophagy in intestinal epithelial cells plays an important role in regulating gut microbiota, but the conversion of LC3-I into LC3-II in WT and *Aim2*<sup>-/-</sup> was comparable both pre- and post-induction of colitis (Figures S5A and S5B). Intestinal epithelial cells also produce several AMPs, which either directly exert cytotoxicity or inhibit bacterial growth by limiting trace elements (Gallo and Hooper, 2012). We measured the expression of AMPs, including mouse  $\beta$ -defensin 2 (mBD2), Reg3 $\gamma$  and Reg3 $\beta$ , S100A8 and S100A9, and Lcn2 in WT and *Aim2*<sup>-/-</sup> mouse colons at day 0 and day 3 following colitis induction. Interestingly, while the expression of mBD2 and Lcn2 in healthy (day 0) WT and *Aim2*<sup>-/-</sup> mice was comparable, the level of Reg3 $\gamma$ , Reg3 $\beta$ , S100A8, and S100A9 was significantly reduced in *Aim2*<sup>-/-</sup> mouse colons as compared to those of WT mice (Figure 5D). The expression of Reg3 $\beta$ , S100A8, and S100A9 remained low in the early phases of colitis (day 3) (Figure 5D). These results suggest that AIM2 may regulate the bacterial burden of *E. coli* and Enterobacteriaceae via the production of AMPs. However, expression of Reg3 $\gamma$ , Reg3 $\beta$ , S100A8, S100A9, and Lcn2 were upregulated in *Aim2*<sup>-/-</sup> mice during acute colitis (day 8) (Figure S5C). Higher expression of these AMPs in *Aim2*<sup>-/-</sup> mice during colitis may be associated with higher bacterial burden since their expression in *Aim2*<sup>-/-</sup> and WT mice was reduced to the basal level upon treatment with antibiotics (Figure S5C).

### AIM2 Is a Major Inflammasome Activator in the Colon


Previous studies have shown that the NLRP3 inflammasome plays a protective role against experimental colitis (Allen et al., 2010; Zaki et al., 2010a). Notably, caspase-1 activation in *Nlrp3*<sup>-/-</sup> mice is only partially reduced but not completely abrogated during DSS-colitis (Figure S6A), suggesting that other mechanisms of caspase-1 activation are active in the gut. We hypothesized that the dsDNA from colonizing bacteria contributes to intestinal homeostasis via activation of the AIM2 inflammasome. To test this hypothesis, we measured caspase-1 activation in colon tissues collected from WT and *Aim2*<sup>-/-</sup> mice at different days during colitis. Interestingly, caspase-1 activation in *Aim2*<sup>-/-</sup> mouse colons at days 0, 3, and 5 after DSS administration was significantly reduced as compared to WT mouse colons (Figures 6A–6D). However, a few *Aim2*<sup>-/-</sup> mice showed near normal caspase-1 activation in the colon, which may reflect the action of the NLRP3, NLRP6, or other inflammasomes. We consistently observed

significantly reduced levels of IL-18, despite enhanced expression of *IL-18* mRNA in inflamed colons of *Aim2*<sup>-/-</sup> mice at days 3 and 5 after DSS administration (Figures 6E and S6B). Similarly, IL-1 $\beta$  production was suppressed by the absence of AIM2 (Figure 6E). Reduced activation of IL-1 $\beta$  and IL-18 in *Aim2*<sup>-/-</sup> mouse colons at healthy and colitic conditions was also seen in ex vivo colon organ culture (Figure S6C). In agreement with the function of IL-18 in epithelial cell proliferation (Dupaul-Chicoine et al., 2010; Zaki et al., 2010a), we observed significantly reduced proliferation of intestinal epithelial cells in *Aim2*<sup>-/-</sup> mice during acute colitis (day 5) but not at homeostasis (day 0) or delayed stage of colitis (day 8) (Figures S6D and S6E). These data support the hypothesis that bacterial dsDNA activates an AIM2-dependent inflammasome in the colon. To confirm that bacterial dsDNA present in the colonic lumen can activate the AIM2 inflammasome, we isolated DNA from mouse stool and transfected it into bone marrow-derived macrophages (BMDM). While fecal DNA activated caspase-1 in WT BMDM, cleaved caspase-1 was not detected in *Aim2*<sup>-/-</sup> BMDM (Figure 6F). However, activation of the NLRP3 inflammasome induced by LPS plus ATP or Nigericin was completely normal in *Aim2*<sup>-/-</sup> BMDM (Figure S6F). To understand which cellular compartments are involved in AIM2-dependent inflammasome activation, we isolated colonic epithelial cells and lamina propria mononuclear cells (LPMN) from WT and *Aim2*<sup>-/-</sup> mice at day 5 post DSS. While caspase-1 activation was significantly higher in WT epithelial cells compared to that of *Aim2*<sup>-/-</sup> mice (Figure 6G), it was barely detectable in LPMNs of either background of mice (data not shown). The notion that the epithelial compartment is the major site for the activation of the AIM2 inflammasome is further supported by bone marrow chimera studies. Chimeric mice having AIM2 in epithelial compartment, irrespective of the presence of AIM2 in myeloid cells (WT > WT and *Aim2*<sup>-/-</sup> > WT), developed less severe colitis as compared to mice deficient in AIM2 in the epithelium (*Aim2*<sup>-/-</sup> > *Aim2*<sup>-/-</sup> and WT > *Aim2*<sup>-/-</sup>) (Figures 6H and 6I). The observed defect in inflammasome activation in *Aim2*<sup>-/-</sup> mice was not dependent on altered gut microbiota since inflammasome activation remained defective in *Aim2*<sup>-/-</sup> mice co-housed with WT mice (Figure S6G).

### The AIM2 Inflammasome Regulates Gut Microbiota and Colitis Susceptibility via Production of AMPs

Next, we sought to determine whether AIM2 controls microbial growth via the inflammasome. Therefore, we measured *E. coli* burden in healthy WT, *Aim2*<sup>-/-</sup>, *Nlrp3*<sup>-/-</sup>, *caspase-1*<sup>-/-</sup>, *Il1 $\beta$* <sup>-/-</sup> and *Il18*<sup>-/-</sup> mice. The *E. coli* burden was significantly higher than WT in all inflammasome-defective mice (Figures 7A and S7A); however, *E. coli* burden in *Il1 $\beta$* <sup>-/-</sup> mice was significantly less than other inflammasome-defective mice, suggesting that IL-18 plays a major role in maintaining intestinal microbial homeostasis. Dysregulated microbiota represented by increased growth of *E. coli* in *caspase-1*<sup>-/-</sup> and *Il18*<sup>-/-</sup> mice was also associated with increased colitis susceptibility as measured by

3% DSS for 5 days. Body weight changes (G) and clinical scores for diarrhea (H) were monitored daily. (I) Colon lengths were measured at day 7 after DSS administration. (J) Colons were fixed in 10% formalin and processed for H&E staining. (K) A part of the colon was analyzed for IL-6 and KC production by ELISA. Data represent mean  $\pm$  SEM. \* $p < 0.05$ ; \*\* $p < 0.01$ . Scale bars, 100  $\mu$ m. See also Figure S4 and Table S1.



**Figure 5. AIM2 Regulates Production of Antimicrobial Peptides**

WT and *Aim2*<sup>-/-</sup> mice were fed with 3% DSS in drinking water for 5 days.

(A) Colons collected at day 5 after DSS were fixed in 10% formalin and cryosections were stained with PAS. Scale bars, 100 μm.

(B) PAS-positive cells per crypt were counted.

(C) Colons were collected at day 0 (n = 6/group) and day 5 (n = 6/group) after DSS administration. RNA was isolated and expression of Mucins (*Muc1*, *Muc2*, *Muc3*, *Muc4*) was analyzed by qPCR.


(D) Expression of antimicrobial peptides *mBD2*, *Reg3β*, *Reg3γ*, *S100a8*, *S100a9*, and *Lcn2* in colon tissues collected at day 0 and day 3 after DSS-administration was measured by real-time PCR. Data represent mean ± SEM; n = 8–10/group; \*p < 0.05, \*\*p < 0.01.

See also [Figure S5](#) and [Table S1](#).

body weight change, colon shortening, and clinical scores for diarrhea ([Figures 7B](#) and [S7B–S7D](#)).

To understand whether dysbiosis in *caspase-1*<sup>-/-</sup> mice is also due to the defects in AMP production, we measured the expression of *mBD2*, *Reg3γ*, *Reg3β*, *S100A8*, *S100A9*, and *Lcn2* at

homeostasis (day 0) or day 3 after DSS administration. Interestingly, similar to *Aim2*<sup>-/-</sup> mice ([Figure 5D](#)), the expression of *Reg3γ*, *Reg3β*, *S100A8*, and *S100A9*, was significantly reduced in *caspase-1*<sup>-/-</sup> mouse colons as compared to those in WT mice ([Figure 7C](#)). However, the level of *mBD2* and *Lcn2* was



**Figure 6. AIM2 Activates the Inflammasome in the Colon during Colitis**

WT and *Aim2*<sup>-/-</sup> mice were fed with 3% DSS in drinking water for 5 days.

(A–D) Colons collected at days 0, 3, 5, and 8 after DSS administration were homogenized and lysates were analyzed for caspase-1 activation by western blotting. Cleaved caspase-1 (p20) is indicated by arrow.

(E) Colon homogenates were assayed for IL-1β and IL-18 ELISA. Data represent mean ± SEM; n = 7–10/group; \*p < 0.05, \*\*p < 0.01.

(F) BMDM collected from WT or *Aim2*<sup>-/-</sup> mice were stimulated with 500 ng/ml LPS followed by transfected with either stool DNA or poly dA:dT (AIM2 ligand) for 4 hr. Cell lysates were analyzed for caspase-1 activation by western blotting.

(G) Epithelial cells were isolated from WT and *Aim2*<sup>-/-</sup> mice at day 5 after DSS administration. Cell lysates were analyzed for caspase-1 activation by western blotting.

(legend continued on next page)

not different between *caspase-1*<sup>-/-</sup> and WT mouse colons (Figure 7C). This observation is further supported by in vitro studies showing that Reg3 $\gamma$ , Reg3 $\beta$ , S100A9, and Lcn2 were induced by both IL-1 $\beta$  and IL-18, and S100A8 was induced by IL-18 in colonic epithelial cells (Figures 7D, S7E, and S7F). Similarly, intestinal organoids expressed Reg3 $\gamma$ , Reg3 $\beta$ , S100A9, and Lcn2 in response to IL-1 $\beta$  and IL-18 (Figure S7G). Culture supernatants of IL-1 $\beta$ - and IL-18-stimulated colon organ culture effectively killed *E. coli* (Figure S7H). In vivo infusion of IL-18 led to a dramatic reduction of colonic *E. coli* growth in *Aim2*<sup>-/-</sup> mice and protected them against DSS-induced colitis (Figures 7E and 7F). IL-1 $\beta$  infusion also demonstrated a suppressive effect on *E. coli* growth, but its effect was much less than that of IL-18 (Figures 7E and 7F). These results suggest that IL-18 produced by the AIM2 inflammasome binds to the IL-18R on epithelial and immune cells (Figures S7I and S7J), leading to the induction of antimicrobial peptides that play a central role in regulating the intestinal microbial ecology and inflammation (Figure 7G).

## DISCUSSION

The surveillance of PAMPs by diverse PRRs expressed in intestinal epithelial cells, as well as immune cells, triggers a homeostatic immune response that allows for the peaceful coexistence of the intestinal mucosal tissue and gut microbiota. Microbial DNA is known to be sensed by TLR9, AIM2, and cGAS, leading to the activation of NF- $\kappa$ B, inflammasome, and interferon signaling pathways, respectively (Hornung et al., 2009; Roberts et al., 2009; Sun et al., 2013; Tsujimura et al., 2004). TLR9-mediated recognition of DNA in the intestine exerts a protective effect against experimental colitis (Katakura et al., 2005; Rachmilewitz et al., 2002). Although the physiological relevance of DNA sensing by cGAS during intestinal inflammation has not been investigated to date, type-I interferon signaling has been shown to be critical in regulating Paneth and T cell functions (Lee et al., 2012; Tschurtschenthaler et al., 2014). However, little is known about the role of AIM2 as a DNA sensor in intestinal homeostasis. This study demonstrates the involvement of AIM2 in DNA-dependent activation of the intestinal mucosal immune system, which critically provides protection against intestinal inflammation.

Pathogenesis of inflammatory bowel diseases (IBD) such as Crohn's disease (CD) and ulcerative colitis involves both host genetic susceptibility and altered gut microbiota. Several bacterial species, including *Escherichia coli*, *Bacteroides*, *Clostridia*, *Bifidobacteria*, and *Firmicutes* have been identified in clinical samples from IBD patients, suggesting a role for these bacteria in IBD pathogenesis (Frank et al., 2007; Li et al., 2015; Martinez-Medina and Garcia-Gil, 2014; Walters et al., 2014). Therefore, how host immune system regulates the growth of these pathobionts is a long sought after goal in IBD research. A major limitation of studies investigating the regulation of gut microbiota in experimental models is the diversity of microflora in animals from different laboratories. This may explain why we found no


remarkable difference in *Prevotellaceae* and TM7 in inflammasome-deficient mice, unlike what had been previously described (Elinav et al., 2011). WT and all mutant mice included in this study were obtained from Jackson Laboratory and they were housed in the same animal facility at UT Southwestern. Thus, altered microbiota as represented by increased growth of *E. coli* in *Aim2*<sup>-/-</sup>, *Nlrp3*<sup>-/-</sup>, *caspase-1*<sup>-/-</sup>, and *Il18*<sup>-/-</sup> mice is less likely due to the difference in source, location, and other environmental factors. Notably, we did not see similar level of *E. coli* as of *Aim2*<sup>-/-</sup> and *caspase-1*<sup>-/-</sup> mice in *Nod2*<sup>-/-</sup> and *Nlrp12*<sup>-/-</sup> mice, which are also susceptible to experimental colitis and housed in the same animal facility (data not shown). However, this study warrants that microbiota-related disease phenotypes in experimental studies should be re-evaluated in different laboratory settings.

Increased growth of *E. coli* in *Aim2*<sup>-/-</sup> and other inflammasome-deficient mice is of great clinical significance. A growing number of clinical studies demonstrate that *E. coli* resides in the colonic and celiac mucosa of CD patients (Burke and Axon, 1988; Darfeuille-Michaud et al., 2004; Martinez-Medina and Garcia-Gil, 2014; Tabaqchali et al., 1978). Our understanding of the genetic link between increased *E. coli* colonization and CD susceptibility is poor. More than 100 loci associated with CD susceptibility have been identified including *NOD2*, *ATG16L*, *IRGM*, *IL-23R*, *IL-10*, and *IL-18RAP* (Cleynen et al., 2013). However, a common mechanism of controlling *E. coli* growth by these functionally diverse genes is less clear. Perhaps defects in innate and adaptive immune system lead to an altered host defense in the gut allowing increased growth of *E. coli* and other Enterobacteriaceae. Supporting this view, increased growth of Enterobacteriaceae was noted in *T-bet*<sup>-/-</sup>*Rag2*<sup>-/-</sup> mice, a mice model for spontaneous colitis (Garrett et al., 2010). An imbalance in the host immune response during inflammation has also been shown to boost *E. coli* growth in the colon (Winter et al., 2013). Our study demonstrates that the AIM2 inflammasome is a part of intestinal innate host defense against an overgrowth of commensal *E. coli*. Notably, the colitis phenotype in *Aim2*<sup>-/-</sup> mice is not solely dependent on *E. coli* since co-housed *Aim2*<sup>-/-</sup> mice developed attenuated colitis despite higher *E. coli* burden, suggesting that other unidentified bacteria that are transferred from WT to *Aim2*<sup>-/-</sup> mice during co-housing may have protected *Aim2*<sup>-/-</sup> mice against exacerbated colitis.

Although the inflammasome has emerged as a critical regulator of intestinal homeostasis, our knowledge of how it regulates the intestinal microbial community is limited. This study reveals a pathway for the regulation of gut microbiota by inflammasome-mediated production of AMPs, including Reg3 $\beta$ , Reg3 $\gamma$ , S100A8, S100A9, and Lcn2. AMPs produced by intestinal epithelial cells play critical roles in regulating the growth of the intestinal microbial community (Lai and Gallo, 2009; Ostaff et al., 2013). Previous studies have elucidated a role for Reg3 $\beta$  in host defense against Gram-negative bacteria (Stelter et al., 2011; van Ampting et al., 2012). S100A8 and S100A9 act as

---

(H and I) Lethally irradiated WT and *Aim2*<sup>-/-</sup> mice were injected with either WT or *Aim2*<sup>-/-</sup> bone marrow cells and divided into four groups: WT > WT (n = 8), WT > *Aim2*<sup>-/-</sup> (n = 6), *Aim2*<sup>-/-</sup> > WT (6), and *Aim2*<sup>-/-</sup> > *Aim2*<sup>-/-</sup> (n = 6). Seven weeks after reconstitution, chimeric mice were fed with 3% DSS for 5 days. Body weight changes were monitored daily (H) and colon length was measured at day 7 (I). Data represent mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.01. See also Figure S6.



**Figure 7. The Inflammasome Regulates Intestinal Microbiota via Production of AMPs**

(A) Feces collected from healthy WT, *Aim2*<sup>-/-</sup>, *Nlrp3*<sup>-/-</sup>, *Il1β*<sup>-/-</sup>, *caspase-1*<sup>-/-</sup>, and *Il18*<sup>-/-</sup> mice were diluted in PBS and plated on MacConkey agar. Colonies were counted after 24 hr culture at 37°C.

(legend continued on next page)



metal chelators, thus they limit the growth of pathogens by reducing the availability of essential trace elements such as  $Zn^{2+}$  and  $Mn^{2+}$  (Gallo and Hooper, 2012). Similarly, *Lcn2* was shown to inhibit bacterial growth, including *E. coli*, by sequestering iron (Berger et al., 2006; Flo et al., 2004). Although the level of *Lcn2* in healthy *Aim2*<sup>-/-</sup> and *caspase-1*<sup>-/-</sup> mouse colons were seen comparable to WT mice, its expression was strikingly induced by cytokines IL-1 $\beta$  and IL-18, suggesting that *Lcn2* may be induced by the inflammasome and contribute to antimicrobial host defense in the gut. Since *Lcn2* is also induced by MyD88-dependent pathways (Layoun et al., 2012), it is possible that an increase in the *E. coli* burden in inflammasome-defective mice may induce *Lcn2* in the gut through activation of the TLR-MyD88 pathways, making the level of *Lcn2* indistinguishable in comparison to WT mice. Taken together, these data suggest that inflammasome-associated cytokines regulate the growth *E. coli* and other gut microbiota through induction of multiple AMPs.

Our study shows that AIM2 is expressed in different cell population including myeloid, lymphocyte, and epithelial cells in the colon. However, how microbial DNA is delivered in the host cells is unknown. Since the AIM2 inflammasome is activated by diverse bacterial infections, it might be possible that gut bacteria are taken up by intestinal epithelial and myeloid cells leading to the release of bacterial DNA into the cytosol. A possibility of natural transfection of microbial DNA into intestinal immune and epithelial cell should be investigated in future studies.

In summary, this study underscores the critical role the DNA sensor AIM2 plays in regulating intestinal microbial community and inflammation. We show that AIM2 activates the inflammasome via sensing bacterial DNA in the gut. Although a protective role for the inflammasome in intestinal inflammatory disorders has been known for last five years, the underlying mechanism of inflammasome-mediated protection remains elusive. Therefore, the findings of this study, showing regulation of gut microbiota by the inflammasome via production of antimicrobial effector molecules, offer an insight into the pathogenesis of inflammatory disorders of the gut. Moreover, the AIM2 inflammasome may serve as a therapeutic target for the regulation of intestinal microbiota and inflammation.

## EXPERIMENTAL PROCEDURES

### Mice

*Aim2*<sup>-/-</sup>, *Nlrp3*<sup>-/-</sup>, *Il18*<sup>-/-</sup>, *Caspase-1*<sup>-/-</sup> (*Caspase-1/11*<sup>-/-</sup>), and WT (C57Bl6/J) mice were purchased from Jackson Laboratory. *Il1 $\beta$* <sup>-/-</sup> mice (Shor-

nick et al., 1996) were kindly shared by Dr. Chandrashekar Pasare at UT Southwestern. All mice are maintained in a specific pathogen-free (SPF) facility. All studies were approved by the Institutional Animal Care and Use Committee of UT Southwestern Medical Center and were conducted in accordance with the IACUC guidelines and NIH Guide for the Care and Use of Laboratory Animals. All experiments were conducted with sex and age-matched mice and both male and female mice were included.

### Colitis

Mice were treated with 3% (w/v) DSS (molecular mass 36–40 kDa; TdB Consultancy) dissolved in sterile, distilled water ad libitum for the experimental days 1–5 followed by normal drinking water until the end of the experiment. Body weight, stool consistency, and the presence of occult blood were determined daily up to day 8. Scoring system for diarrhea and rectal bleeding is described in the Supplemental Experimental Procedures.

### Gene Expression

For gene expression analysis, tissues were homogenized in Trizol reagent (Invitrogen). RNA was extracted according to the manufacturer's instructions and reverse transcribed into cDNA. qPCR assays were performed with an CFX Connect real-time PCR detection system (Bio-Rad) using iTaq Universal SYBR Green Supermix (Bio-Rad). Expression of GAPDH was used as an endogenous control. Primers used for real-time PCR are listed in Table S1.

### Western Blot Analyses

For western blot analysis, total proteins from tissues or cells were homogenized in RIPA lysis buffer containing protease and phosphatase inhibitor (Roche), resolved by SDS-PAGE, and transferred onto a PVDF membrane. Membranes were immunoblotted with rabbit antibodies against ERK (4695, Cell Signaling), Phospho-ERK (4370, Cell Signaling) and LC3A (4599, Cell Signaling), caspase-1 (AG-20B-0042, Adipogen), and AIM2 (13095, Cell Signaling).  $\beta$ -actin or tubulin was used as a loading control.

### Bacterial Genomic DNA Isolation and Microbiota Analysis

Frozen fecal samples were processed for DNA isolation using QIAamp Fast DNA Stool Mini Kit (QIAGEN). To analyze the relative number of specific intestinal bacterial groups, extracted bacterial genomic DNA was amplified for 16S rDNA with iTaq Universal SYBR Green Supermix (Bio-Rad). Specific 16S rDNA primer sequences for real-time PCR are listed in Table S2.

### Statistical Analysis

Data are represented as mean  $\pm$  SEM. Data were analyzed by Prism5 (GraphPad Software) and statistical significance was determined by two-tailed Student's t test.  $p < 0.05$  was considered statistically significant.

## SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, seven figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2015.10.040>.

(B) Colitis was induced in WT, *Aim2*<sup>-/-</sup>, *caspase-1*<sup>-/-</sup>, and *Il18*<sup>-/-</sup> ( $n = 8-10$ ) mice with 3% DSS. Body weights were monitored daily. Data represent mean  $\pm$  SEM of a representative experiment; \*\* $p < 0.01$ , \*\*\* $p < 0.001$ .

(C) Expression of *mBD2*, *Reg3 $\beta$* , *Reg3 $\gamma$* , *S100a8*, *S100a9*, and *Lcn2* in colons collected at days 0 and 3 was measured by real-time PCR. Data represent mean  $\pm$  SEM;  $n = 7-10$ /group; \* $p < 0.05$ , \*\* $p < 0.01$ .

(D) Colonic crypts were cultured and treated with IL-1 $\beta$  or IL-18 for 4 hr. Expression of *mBD2*, *Reg3 $\beta$* , *Reg3 $\gamma$* , *S100a8*, *S100a9*, and *Lcn2* was measured by RT-PCR. Data represent mean  $\pm$  SD of triplicate wells; \* $p < 0.05$ , \*\* $p < 0.01$ .

(E and F) *Aim2*<sup>-/-</sup> mice were treated with either IL-18 (0.5  $\mu$ g/mouse) or IL-1 $\beta$  (0.3  $\mu$ g/mouse) for 5 consecutive days. Feces collected at days 0, 3, and 6 after cytokine infusion were cultured on MacConkey agar to count *E. coli* burden. (F) WT ( $n = 5$ ), *Aim2*<sup>-/-</sup> ( $n = 3$ ), *Aim2*<sup>-/-</sup> (IL-1 $\beta$ ) ( $n = 5$ ), and *Aim2*<sup>-/-</sup> (IL-18) ( $n = 5$ ) mice were fed with 3% DSS for 5 days. IL-1 $\beta$ - and IL-18-treated mice further received IL-1 $\beta$  (0.3  $\mu$ g/mouse) and IL-18 (0.5  $\mu$ g/mouse) respectively at days 1, 3, and 5 following DSS administration. Body weight changes were monitored daily. Data represent mean  $\pm$  SEM. \* $p < 0.05$ , \*\* $p < 0.01$ .

(G) The proposed pathway for the AIM2 inflammasome-mediated regulation of intestinal inflammation and microbiota.

See also Figure S7 and Table S1.

## AUTHOR CONTRIBUTIONS

S.H., Y.-T.K., and M.H.Z. designed and performed experiments and analyzed data. L.P. performed histopathological examination of colon tissues. E.M.T. identified bacterial species isolated from feces by MALDI-TOF system. J.S.M., C.P., and L.V.H. provided reagents and helped editing the manuscript. M.H.Z. conceived this study and wrote the manuscript.

## ACKNOWLEDGMENTS

We thank John Shelton and Angie Mobley for their assistance in histopathology and flow cytometry respectively. This work was supported by Crohn's and Colitis Foundation Career Development Award (3711), a New Investigator Award from the American Cancer Society (IRG-02-196-10), and the Harold C. Simmons Comprehensive Cancer Center (National Cancer Institute/NIH, P30 CA142543), and UT Southwestern startup funding given to M.H.Z.

Received: April 3, 2015

Revised: June 29, 2015

Accepted: October 14, 2015

Published: November 19, 2015

## REFERENCES

- Allen, I.C., TeKippe, E.M., Woodford, R.M., Uronis, J.M., Holl, E.K., Rogers, A.B., Herfarth, H.H., Jobin, C., and Ting, J.P. (2010). The NLRP3 inflammasome functions as a negative regulator of tumorigenesis during colitis-associated cancer. *J. Exp. Med.* *207*, 1045–1056.
- Berger, T., Togawa, A., Duncan, G.S., Elia, A.J., You-Ten, A., Wakeham, A., Fong, H.E., Cheung, C.C., and Mak, T.W. (2006). Lipocalin 2-deficient mice exhibit increased sensitivity to *Escherichia coli* infection but not to ischemia-reperfusion injury. *Proc. Natl. Acad. Sci. USA* *103*, 1834–1839.
- Bürkstümmer, T., Baumann, C., Blüml, S., Dixit, E., Dürnberger, G., Jahn, H., Planyavsky, M., Bilban, M., Colinge, J., Bennett, K.L., and Superti-Furga, G. (2009). An orthogonal proteomic-genomic screen identifies AIM2 as a cytoplasmic DNA sensor for the inflammasome. *Nat. Immunol.* *10*, 266–272.
- Burke, D.A., and Axon, A.T. (1988). Adhesive *Escherichia coli* in inflammatory bowel disease and infective diarrhoea. *BMJ* *297*, 102–104.
- Chen, G.Y., Shaw, M.H., Redondo, G., and Núñez, G. (2008). The innate immune receptor Nod1 protects the intestine from inflammation-induced tumorigenesis. *Cancer Res.* *68*, 10060–10067.
- Cho, J.H. (2008). The genetics and immunopathogenesis of inflammatory bowel disease. *Nat. Rev. Immunol.* *8*, 458–466.
- Cleynen, I., González, J.R., Figueroa, C., Franke, A., McGovern, D., Bortlik, M., Crusius, B.J., Vecchi, M., Artieda, M., Szczypiorska, M., et al. (2013). Genetic factors conferring an increased susceptibility to develop Crohn's disease also influence disease phenotype: results from the IBDchip European Project. *Gut* *62*, 1556–1565.
- Cridland, J.A., Curley, E.Z., Wykes, M.N., Schroder, K., Sweet, M.J., Roberts, T.L., Ragan, M.A., Kassahn, K.S., and Stacey, K.J. (2012). The mammalian PYHIN gene family: phylogeny, evolution and expression. *BMC Evol. Biol.* *12*, 140.
- Darfeuille-Michaud, A., Boudeau, J., Bulois, P., Neut, C., Glasser, A.L., Barnich, N., Bringer, M.A., Swidsinski, A., Beaugerie, L., and Colombel, J.F. (2004). High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology* *127*, 412–421.
- Dupaul-Chicoine, J., Yeretssian, G., Doiron, K., Bergstrom, K.S., McIntire, C.R., LeBlanc, P.M., Meunier, C., Turbide, C., Gros, P., Beauchemin, N., et al. (2010). Control of intestinal homeostasis, colitis, and colitis-associated colorectal cancer by the inflammatory caspases. *Immunity* *32*, 367–378.
- Elinav, E., Strowig, T., Kau, A.L., Henao-Mejia, J., Thaiss, C.A., Booth, C.J., Peaper, D.R., Bertin, J., Eisenbarth, S.C., Gordon, J.I., and Flavell, R.A. (2011). NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis. *Cell* *145*, 745–757.
- Fang, R., Hara, H., Sakai, S., Hernandez-Cuellar, E., Mitsuyama, M., Kawamura, I., and Tsuchiya, K. (2014). Type I interferon signaling regulates activation of the absent in melanoma 2 inflammasome during *Streptococcus pneumoniae* infection. *Infect. Immun.* *82*, 2310–2317.
- Fernandes-Alnemri, T., Yu, J.W., Datta, P., Wu, J., and Alnemri, E.S. (2009). AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature* *458*, 509–513.
- Fernandes-Alnemri, T., Yu, J.W., Juliana, C., Solorzano, L., Kang, S., Wu, J., Datta, P., McCormick, M., Huang, L., McDermott, E., et al. (2010). The AIM2 inflammasome is critical for innate immunity to *Francisella tularensis*. *Nat. Immunol.* *11*, 385–393.
- Flo, T.H., Smith, K.D., Sato, S., Rodriguez, D.J., Holmes, M.A., Strong, R.K., Akira, S., and Aderem, A. (2004). Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* *432*, 917–921.
- Frank, D.N., St Amand, A.L., Feldman, R.A., Boedeker, E.C., Harpaz, N., and Pace, N.R. (2007). Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc. Natl. Acad. Sci. USA* *104*, 13780–13785.
- Gallo, R.L., and Hooper, L.V. (2012). Epithelial antimicrobial defence of the skin and intestine. *Nat. Rev. Immunol.* *12*, 503–516.
- Garrett, W.S., Gallini, C.A., Yatsunenko, T., Michaud, M., DuBois, A., Delaney, M.L., Punit, S., Karlsson, M., Bry, L., Glickman, J.N., et al. (2010). Enterobacteriaceae act in concert with the gut microbiota to induce spontaneous and maternally transmitted colitis. *Cell Host Microbe* *8*, 292–300.
- Hirota, S.A., Ng, J., Lueng, A., Khajah, M., Parhar, K., Li, Y., Lam, V., Potentier, M.S., Ng, K., Bawa, M., et al. (2011). NLRP3 inflammasome plays a key role in the regulation of intestinal homeostasis. *Inflamm. Bowel Dis.* *17*, 1359–1372.
- Hornung, V., Ablasser, A., Charrel-Dennis, M., Bauernfeind, F., Horvath, G., Caffrey, D.R., Latz, E., and Fitzgerald, K.A. (2009). AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature* *458*, 514–518.
- Hugot, J.P., Chamaillard, M., Zouali, H., Lesage, S., Cézard, J.P., Belaiche, J., Almer, S., Tysk, C., O'Morain, C.A., Gassull, M., et al. (2001). Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* *411*, 599–603.
- Jones, J.W., Kayagaki, N., Broz, P., Henry, T., Newton, K., O'Rourke, K., Chan, S., Dong, J., Qu, Y., Roose-Girma, M., et al. (2010). Absent in melanoma 2 is required for innate immune recognition of *Francisella tularensis*. *Proc. Natl. Acad. Sci. USA* *107*, 9771–9776.
- Katakura, K., Lee, J., Rachmilewitz, D., Li, G., Eckmann, L., and Raz, E. (2005). Toll-like receptor 9-induced type I IFN protects mice from experimental colitis. *J. Clin. Invest.* *115*, 695–702.
- Kim, S., Bauernfeind, F., Ablasser, A., Hartmann, G., Fitzgerald, K.A., Latz, E., and Hornung, V. (2010). *Listeria monocytogenes* is sensed by the NLRP3 and AIM2 inflammasome. *Eur. J. Immunol.* *40*, 1545–1551.
- Lai, Y., and Gallo, R.L. (2009). AMPed up immunity: how antimicrobial peptides have multiple roles in immune defense. *Trends Immunol.* *30*, 131–141.
- Lamkanfi, M., and Dixit, V.M. (2009). Inflammasomes: guardians of cytosolic sanctity. *Immunol. Rev.* *227*, 95–105.
- Layoun, A., Huang, H., Calvé, A., and Santos, M.M. (2012). Toll-like receptor signal adaptor protein MyD88 is required for sustained endotoxin-induced acute hyperferremic response in mice. *Am. J. Pathol.* *180*, 2340–2350.
- Lee, S.E., Li, X., Kim, J.C., Lee, J., González-Navajas, J.M., Hong, S.H., Park, I.K., Rhee, J.H., and Raz, E. (2012). Type I interferons maintain Foxp3 expression and T-regulatory cell functions under inflammatory conditions in mice. *Gastroenterology* *143*, 145–154.
- Li, J., Butcher, J., Mack, D., and Stintzi, A. (2015). Functional impacts of the intestinal microbiome in the pathogenesis of inflammatory bowel disease. *Inflamm. Bowel Dis.* *21*, 139–153.
- Maeda, S., Hsu, L.C., Liu, H., Bankston, L.A., Iimura, M., Kagnoff, M.F., Eckmann, L., and Karin, M. (2005). Nod2 mutation in Crohn's disease potentiates NF-kappaB activity and IL-1beta processing. *Science* *307*, 734–738.

- Martinez-Medina, M., and Garcia-Gil, L.J. (2014). *Escherichia coli* in chronic inflammatory bowel diseases: An update on adherent invasive *Escherichia coli* pathogenicity. *World J. Gastrointest. Pathophysiol.* *5*, 213–227.
- Ostaf, M.J., Stange, E.F., and Wehkamp, J. (2013). Antimicrobial peptides and gut microbiota in homeostasis and pathology. *EMBO Mol. Med.* *5*, 1465–1483.
- Park, E., Na, H.S., Song, Y.R., Shin, S.Y., Kim, Y.M., and Chung, J. (2014). Activation of NLRP3 and AIM2 inflammasomes by *Porphyromonas gingivalis* infection. *Infect. Immun.* *82*, 112–123.
- Rachmilewitz, D., Karmeli, F., Takabayashi, K., Hayashi, T., Leider-Trejo, L., Lee, J., Leoni, L.M., and Raz, E. (2002). Immunostimulatory DNA ameliorates experimental and spontaneous murine colitis. *Gastroenterology* *122*, 1428–1441.
- Rakoff-Nahoum, S., Paglino, J., Eslami-Varzaneh, F., Edberg, S., and Medzhitov, R. (2004). Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell* *118*, 229–241.
- Rathinam, V.A., Jiang, Z., Waggoner, S.N., Sharma, S., Cole, L.E., Waggoner, L., Vanaja, S.K., Monks, B.G., Ganesan, S., Latz, E., et al. (2010). The AIM2 inflammasome is essential for host defense against cytosolic bacteria and DNA viruses. *Nat. Immunol.* *11*, 395–402.
- Roberts, T.L., Idris, A., Dunn, J.A., Kelly, G.M., Burnton, C.M., Hodgson, S., Hardy, L.L., Garceau, V., Sweet, M.J., Ross, I.L., et al. (2009). HIN-200 proteins regulate caspase activation in response to foreign cytoplasmic DNA. *Science* *323*, 1057–1060.
- Salcedo, R., Worschech, A., Cardone, M., Jones, Y., Gyulai, Z., Dai, R.M., Wang, E., Ma, W., Haines, D., O’Hugin, C., et al. (2010). MyD88-mediated signaling prevents development of adenocarcinomas of the colon: role of interleukin 18. *J. Exp. Med.* *207*, 1625–1636.
- Shornick, L.P., De Togni, P., Mariathasan, S., Goellner, J., Strauss-Schoenberger, J., Karr, R.W., Ferguson, T.A., and Chaplin, D.D. (1996). Mice deficient in IL-1 $\beta$  manifest impaired contact hypersensitivity to trinitrochlorobenzene. *J. Exp. Med.* *183*, 1427–1436.
- Stelter, C., Käppeli, R., König, C., Krah, A., Hardt, W.D., Stecher, B., and Bumann, D. (2011). Salmonella-induced mucosal lectin RegIII $\beta$  kills competing gut microbiota. *PLoS ONE* *6*, e20749.
- Sun, L., Wu, J., Du, F., Chen, X., and Chen, Z.J. (2013). Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* *339*, 786–791.
- Tabaqchali, S., O’Donoghue, D.P., and Bettelheim, K.A. (1978). *Escherichia coli* antibodies in patients with inflammatory bowel disease. *Gut* *19*, 108–113.
- Tschurtschenthaler, M., Wang, J., Fricke, C., Fritz, T.M., Niederreiter, L., Adolph, T.E., Sarcevic, E., Künzel, S., Offner, F.A., Kalinke, U., et al. (2014). Type I interferon signalling in the intestinal epithelium affects Paneth cells, microbial ecology and epithelial regeneration. *Gut* *63*, 1921–1931.
- Tsujimura, H., Tamura, T., Kong, H.J., Nishiyama, A., Ishii, K.J., Klinman, D.M., and Ozato, K. (2004). Toll-like receptor 9 signaling activates NF- $\kappa$ B through IFN regulatory factor-8/IFN consensus sequence binding protein in dendritic cells. *J. Immunol.* *172*, 6820–6827.
- van Ampting, M.T., Loonen, L.M., Schonewille, A.J., Konings, I., Vink, C., Iovanna, J., Chamailard, M., Dekker, J., van der Meer, R., Wells, J.M., and Bo-vee-Oudenhoven, I.M. (2012). Intestinally secreted C-type lectin Reg3b attenuates salmonellosis but not listeriosis in mice. *Infect. Immun.* *80*, 1115–1120.
- Vijay-Kumar, M., Sanders, C.J., Taylor, R.T., Kumar, A., Aitken, J.D., Sitaraman, S.V., Neish, A.S., Uematsu, S., Akira, S., Williams, I.R., and Gewirtz, A.T. (2007). Deletion of TLR5 results in spontaneous colitis in mice. *J. Clin. Invest.* *117*, 3909–3921.
- Villani, A.C., Lemire, M., Fortin, G., Louis, E., Silverberg, M.S., Collette, C., Baba, N., Libioulle, C., Belaiche, J., Bitton, A., et al. (2009). Common variants in the NLRP3 region contribute to Crohn’s disease susceptibility. *Nat. Genet.* *41*, 71–76.
- Walters, W.A., Xu, Z., and Knight, R. (2014). Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS Lett.* *588*, 4223–4233.
- Williams, T.M., Leeth, R.A., Rothschild, D.E., Coutermarsh-Ott, S.L., McDaniel, D.K., Simmons, A.E., Heid, B., Cecere, T.E., and Allen, I.C. (2015). The NLRP1 inflammasome attenuates colitis and colitis-associated tumorigenesis. *J. Immunol.* *194*, 3369–3380.
- Winter, S.E., Winter, M.G., Xavier, M.N., Thiennimitr, P., Poon, V., Keestra, A.M., Laughlin, R.C., Gomez, G., Wu, J., Lawhon, S.D., et al. (2013). Host-derived nitrate boosts growth of *E. coli* in the inflamed gut. *Science* *339*, 708–711.
- Zaki, M.H., Boyd, K.L., Vogel, P., Kastan, M.B., Lamkanfi, M., and Kanneganti, T.D. (2010a). The NLRP3 inflammasome protects against loss of epithelial integrity and mortality during experimental colitis. *Immunity* *32*, 379–391.
- Zaki, M.H., Vogel, P., Body-Malapel, M., Lamkanfi, M., and Kanneganti, T.D. (2010b). IL-18 production downstream of the Nlrp3 inflammasome confers protection against colorectal tumor formation. *J. Immunol.* *185*, 4912–4920.
- Zaki, M.H., Lamkanfi, M., and Kanneganti, T.D. (2011a). The Nlrp3 inflammasome: contributions to intestinal homeostasis. *Trends Immunol.* *32*, 171–179.
- Zaki, M.H., Vogel, P., Malireddi, R.K., Body-Malapel, M., Anand, P.K., Bertin, J., Green, D.R., Lamkanfi, M., and Kanneganti, T.D. (2011b). The NOD-like receptor NLRP12 attenuates colon inflammation and tumorigenesis. *Cancer Cell* *20*, 649–660.