



Cortical aPKC kinase activity distinguishes neural stem cells from progenitor cells by ensuring asymmetric segregation of Numb

Jill M. Haenfler^a, Chaoyuan Kuang^{a,b}, Cheng-Yu Lee^{a,c,d,e,*}

^a Program in Cellular and Molecular Biology, University of Michigan, Ann Arbor, MI 48109, USA

^b Medical Scientist Training Program, University of Michigan, Ann Arbor, MI 48109, USA

^c Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, MI 48109, USA

^d Division of Molecular Medicine and Genetics, Department of Internal Medicine, University of Michigan, Ann Arbor, MI 48109, USA

^e Center for Stem Cell Biology, Life Sciences Institute, University of Michigan, Ann Arbor, MI 48109, USA

ARTICLE INFO

Article history:

Received for publication 16 December 2011

Revised 16 February 2012

Accepted 20 February 2012

Available online 25 February 2012

Keywords:

Asymmetric division

Cell polarity

Intermediate neural progenitor

Neuroblast

Lethal (2) giant larvae

Numb

ABSTRACT

During asymmetric stem cell division, polarization of the cell cortex targets fate determinants unequally into the sibling daughters, leading to regeneration of a stem cell and production of a progenitor cell with restricted developmental potential. In mitotic neural stem cells (neuroblasts) in fly larval brains, the antagonistic interaction between the polarity proteins Lethal (2) giant larvae (*Lgl*) and atypical Protein Kinase C (aPKC) ensures self-renewal of a daughter neuroblast and generation of a progenitor cell by regulating asymmetric segregation of fate determinants. In the absence of *lgl* function, elevated cortical aPKC kinase activity perturbs unequal partitioning of the fate determinants including Numb and induces supernumerary neuroblasts in larval brains. However, whether increased aPKC function triggers formation of excess neuroblasts by inactivating Numb remains controversial. To investigate how increased cortical aPKC function induces formation of excess neuroblasts, we analyzed the fate of cells in neuroblast lineage clones in *lgl* mutant brains. Surprisingly, our analyses revealed that neuroblasts in *lgl* mutant brains undergo asymmetric division to produce progenitor cells, which then revert back into neuroblasts. In *lgl* mutant brains, Numb remained localized in the cortex of mitotic neuroblasts and failed to segregate exclusively into the progenitor cell following completion of asymmetric division. These results led us to propose that elevated aPKC function in the cortex of mitotic neuroblasts reduces the function of Numb in the future progenitor cells. We identified that the acyl-CoA binding domain containing 3 protein (ACBD3) binding region is essential for asymmetric segregation of Numb in mitotic neuroblasts and suppression of the supernumerary neuroblast phenotype induced by increased aPKC function. The ACBD3 binding region of Numb harbors two aPKC phosphorylation sites, serines 48 and 52. Surprisingly, while the phosphorylation status at these two sites directly impinged on asymmetric segregation of Numb in mitotic neuroblasts, both the phosphomimetic and non-phosphorylatable forms of Numb suppressed formation of excess neuroblasts triggered by increased cortical aPKC function. Thus, we propose that precise regulation of cortical aPKC kinase activity distinguishes the sibling cell identity in part by ensuring asymmetric partitioning of Numb into the future progenitor cell where Numb maintains restricted potential independently of regulation by aPKC.

© 2012 Elsevier Inc. All rights reserved.

Introduction

During asymmetric stem cell divisions, polarization of the cell cortex allows unequal partitioning of the cell fate determinants that instruct the daughter progeny to either self-renew as a stem cell or adopt the progenitor cell identity (Neumüller and Knoblich, 2009; Prehoda, 2009). Progenitor cells possess restricted developmental potential and undergo limited rounds of cell division that give rise to differentiated progeny. Mis-regulation of cortical polarity in

asymmetrically dividing stem cells can impinge on the accumulation and/or function of fate determinants in the intended recipient cell. Such defects can lead to generation of progenitor cells that possess stem cell-like properties, perturbing homeostasis and contributing to tumor initiation. Thus, insight into the mechanisms that distinguish sibling cell identity during normal tissue development will likely improve our understanding of aberrant processes from congenital birth defects to tumorigenesis.

In fly larval brains, two classes of neuroblast lineages can be unambiguously identified based on expression of the cell fate markers and properties of their daughter progeny (Chia et al., 2008; Doe, 2008; Egger et al., 2008; Knoblich, 2008, 2010; Weng and Lee, 2011) (Fig. S1A). A type I neuroblast expresses Deadpan (Dpn) and

* Corresponding author. Fax: +1 734 615 5520.

E-mail address: leecheng@umich.edu (C.-Y. Lee).

Asense (Ase) and divides asymmetrically to self-renew a neuroblast and to generate a progenitor cell called a ganglion mother cell (GMC). In contrast, a type II neuroblast (Dpn^+Ase^-) divides asymmetrically to self-renew and to generate an immature intermediate neural progenitor cell (INP) that lacks the expression of *Dpn* and *Ase* and is transiently arrested in the cell cycle while acquiring INP identity (Bello et al., 2008; Boone and Doe, 2008; Bowman et al., 2008). Following maturation, an INP (Dpn^+Ase^+) undergoes limited rounds of asymmetric divisions to regenerate and to produce GMCs. A key functional property that distinguishes these two neuroblast lineages rests on their dependence on *Notch* signaling for maintenance of their identity (Bowman et al., 2008; Song and Lu, 2011; Weng et al., 2011). While dispensable for maintenance of a type I neuroblast, *Notch* signaling is crucial in maintaining type II neuroblasts (Figs. S1B–E).

The mutually antagonistic interaction between *Lgl* and aPKC in mitotic neuroblasts ensures that *Numb* segregates exclusively into the cortex of the presumptive progenitor cell where *Numb* functions to specify progenitor cell identity (Lee et al., 2006; Rolls et al., 2003; Wang et al., 2006). In *lgl* mutant brains, increased cortical aPKC function disrupts asymmetric segregation of *Numb* in mitotic neuroblasts and triggers formation of supernumerary neuroblasts. Consistent with *Numb* acting as a conserved inhibitor of *Notch* signaling, neuroblasts lacking *numb* function or expressing constitutively active *Notch* generate supernumerary neuroblasts at the expense of progenitor cells (Bowman et al., 2008; Frise et al., 1996; Guo et al., 1996; Lee et al., 2006; Rhyu et al., 1994; San-Juán and Baonza, 2011; Wang et al., 2006; Zhong et al., 1997). Thus, elevated cortical aPKC kinase activity induces supernumerary neuroblast formation likely by attenuating *Numb*-dependent regulation of *Notch* signaling. The fly *Numb* protein contains five evolutionarily conserved aPKC phosphorylation sites, and the non-phosphorylatable form of the *Numb* transgenic protein at these sites (*Numb*^{5A}) fails to segregate asymmetrically in mitotic sensory organ precursor cells (Dho et al., 2006; Nishimura and Kaibuchi, 2007; Smith et al., 2007). aPKC can indeed directly phosphorylate *Numb* through these sites and render *Numb* non-functional (Dho et al., 2006; Nishimura and Kaibuchi, 2007; Smith et al., 2007; Wirtz-Peitz et al., 2008). Together, these results led to the hypothesis that increased cortical aPKC kinase activity induces supernumerary neuroblasts by perturbing the localization and the function of *Numb*. Thus far, evidence supporting this proposed mechanism appears largely correlative. First, direct evidence linking aPKC kinase activity to the de-localization of *Numb* from the cortex of mitotic neuroblasts is absent. Second, whether phosphorylation by aPKC indeed renders *Numb* inactive in progenitor cells has never been tested. Finally, type II neuroblasts require *Notch* signaling for maintenance of their identity; therefore, over-expression of *Numb* or *Numb*^{5A} most likely induces supernumerary type II neuroblasts in *lgl* mutant brains to undergo premature differentiation rather than restoring proper specification of INP identity (Wirtz-Peitz et al., 2008) (Figs. S1B–G). As such, whether increased cortical aPKC kinase activity induces supernumerary neuroblasts by impinging on the localization and the function of *Numb* remains an open question.

In this study, we show that despite failing to segregate *Numb* asymmetrically, neuroblasts in *lgl* mutant brains reproducibly undergo asymmetric division to generate progenitor cells. This result suggests that increased cortical aPKC kinase activity impinged on the segregation but not the function of *Numb*. Surprisingly, the non-phosphorylatable *Numb*^{5A} at the five conserved aPKC phosphorylation sites exclusively partitioned in mitotic neuroblasts, indicating that *Numb* contains additional aPKC phosphorylation sites required for asymmetric segregation. Indeed, the two aPKC phosphorylation sites, serines 48 and 52, in the ACBD3 binding region played a pivotal role in asymmetrically segregating *Numb* into the cortex of the future progenitor cell. Most unexpectedly, *Numb* suppressed supernumerary neuroblasts induced by increased cortical aPKC function regardless

of the phosphorylation status at serines 48 and 52. Thus, we propose that the antagonistic interaction between *Lgl* and aPKC ensures that sufficient *Numb* reaches the future progenitor cells where *Numb* maintains their limited potential irrespective of its phosphorylation by aPKC.

Materials and methods

Fly stocks and transgenes

The novel *lgl* mutants were generated by EMS mutagenesis following standard procedures. The *numb* deletion constructs were generated by site-directed mutagenesis of the *numb* cDNA, sequenced, and cloned in the *pUAST-HA* vector for germline transformation. The *UAS-numb*^{S2A} and *UAS-numb*^{S2D} flies were generated using the *pUAST-attB-HA* vector for insertion into an identical docking site in the fly genome via the ϕ C31 integrase-mediated transgenesis (Bischof and Basler, 2008). *Erm-lacZ* flies were generated by cloning the *R9D11* enhancer element upstream of a minimal promoter and the *lacZ* gene followed by ϕ C31 integrase-mediated transgenesis. *Drosophila* cultures were maintained at 25 °C under standard conditions. Other mutant and transgenic flies used in the study include *lgl*³³⁴ (Peng et al., 2000), *aPKC*^{O6403} (Rolls et al., 2003), *UAS-aPKC*^{caax} (Sotillos et al., 2004), *Erm-GAL4* (Pfeiffer et al., 2008), *Wor-GAL4*, *Ase-GAL4* (Zhu et al., 2006), *UAS-numb* (Knoblich et al., 1997), *UAS-numb*^{5A} (Smith et al., 2007), *UAS-numb*^{ΔN} (Knoblich et al., 1997), *UAS-Notch*_{intra} (Chung and Struhl, 2001), and *aph-1*⁵⁰⁷² (Weng et al., 2011). The *UAS-Notch*_{RNAi} and *UAS-spd*_{RNAi} lines were obtained from the Vienna *Drosophila* Resource Center. *Oregon R*, *Sca-GAL4*, *UAS-dcr2*, *aph-1*^{D35}, *hs-flp*, *Act-FRT-Stop-FRT-GAL4*, *UAS-GFP*, *UAS-flp*, *Act-FRT-Stop-FRT-lacZ*, and *tubGal80*^{ts} flies were obtained from the Bloomington *Drosophila* Stock Center.

Immunofluorescent staining and antibodies

Antibody staining was performed as previously described. Antibodies used in this study include rat anti-*Dpn* (1:1), rabbit anti-*Ase* (1:400), mouse anti-*Pros* (1:100), sheep anti-*Lgl* (1:1000, S. Goode), guinea pig anti-*Numb* (1:2500, J. Skeath), mouse anti-*Dlg* (1:50, DSHB), mouse anti- β -gal (1:100, Sigma), chicken anti-GFP (1:2000, Aves Labs), rat anti- α -tubulin (1:100, Serotec), rabbit anti-aPKC (1:1000, Sigma), rabbit anti-phospho-HistoneH3 (1:1000, Upstate), mouse anti-HA (1:1000, Covance), and mouse anti-c-Myc (1:50, DSHB). Secondary antibodies were from Invitrogen and Jackson ImmunoResearch (details are available upon request). Fluorescent conjugates of phalloidin (Invitrogen), which stain F-actin, were used to mark the cell cortex. All images are single confocal sections acquired on a Leica SP5 scanning confocal microscope.

Lineage clone induction

For neuroblast clone induction, wild-type or *lgl* mutant larvae containing *hs-flp* were heat-shocked as follows to induce recombination and marking by the *Act-FRT-STOP-FRT-GAL4* driving *UAS-GFP* expression. After hatching, larvae were cultured for 24 h at 25 °C, subjected to a 1 hour heat-shock at 37 °C, and returned to 25 °C for 24 or 48 h as indicated. For INP clone induction, wild-type or *lgl* mutant larvae containing *erm-GAL4*, *tubGal80*^{ts}, and *UAS-flp* were cultured at 33 °C for 72 h to induce recombination and marking by the *Act-FRT-STOP-FRT-lacZ* reporter. For clones over-expressing *Notch*_{intra}, larvae containing *hs-flp* were cultured at 25 °C and heat-shocked at 24 h after hatching for 2 h at 37 °C to induce *Act-FRT-STOP-FRT-GAL4* recombination and expression of *UAS-Notch*_{intra} and *UAS-GFP*. Larval brains were then dissected and stained as described previously.

Over-expression experiments

For expression of transgenes using *wor-GAL4* alone or in the *lgl* rescue experiments using *ase-GAL4*, larvae were cultured at 32.5 °C for 72 h. For *aPKC^{caax}* overexpression studies, larvae were cultured at 31 °C for 72 h and at 33 °C for 96 h when using *Erm-GAL4*. Larval brains were then dissected and stained as described previously.

Results

Supernumerary neuroblasts in *lgl* mutant brains most likely arise from progenitor cells

Increased cortical aPKC kinase activity phosphorylates Numb, an evolutionarily conserved protein instrumental for specification of the daughter sibling cell fate during asymmetric cell division, possibly rendering it inactive in *lgl* mutant brains (Guo et al., 1996; Nishimura

and Kaibuchi, 2007; Rhyu et al., 1994; Smith et al., 2007; Wirtz-Peitz et al., 2008; Zhong et al., 1997). In addition, analyses of various *lgl* mutant alleles showed that both type I and II neuroblasts become aberrantly expanded in *lgl* mutant brains (Figs. 1E, J and S2). These data led us to hypothesize that supernumerary neuroblasts in *lgl* mutant brains arise from symmetric neuroblast divisions. We tested this hypothesis by assessing the identity of cells in the lineage clones derived from single type I neuroblasts in wild-type or *lgl* mutant brains at 24 or 48 h after clone induction. In wild-type brains, we detected a single neuroblast (Dpn⁺Ase⁻) per type I neuroblast clone, and the neuroblast was always surrounded by GMCs (Dpn⁻Ase⁺) and their daughter progeny (Dpn⁻Ase⁻) (Figs. 1A, C, E, K and S1A). Unexpectedly, all 24-hour type I neuroblast clones in *lgl* mutant brains also contained a single neuroblast surrounded by GMCs and their daughter progeny (Figs. 1B, E and K). However, 25% of the 48-hour type I neuroblast clones in *lgl* mutant brains contained more than one neuroblast per clone, and we frequently observed supernumerary neuroblasts formed basally in the clones (Figs. 1D, E and K; n=91). Thus, we




Fig. 1. *lgl* is required for the maintenance, but not specification, of progenitor cells. (A–B) At 24 h after clone induction, type I neuroblast clones in both wild-type and *lgl^{334/3644}* mutant brains contained a single neuroblast surrounded by GMCs and neurons (n = 30 and 28 clones, respectively). (C–D) At 48 h after clone induction, type I neuroblast clones in wild-type brains still contained a single neuroblast surrounded by GMCs and neurons; however, type I neuroblast clones in *lgl^{334/3644}* mutant brains contained a parental neuroblast immediately adjacent to GMCs and neurons with supernumerary neuroblasts located in the basal portion of the clone (n = 11 and 91 clones, respectively). (E) Quantification of the number of neuroblasts per type I neuroblast clone is shown for wild-type and *lgl* mutant clones at 24 (open) and 48 (filled) hours after clone induction. (F–G) At 24 h after clone induction, type II neuroblast clones in both wild-type and *lgl^{334/3644}* mutant brains contained a single neuroblast surrounded by immature INPs, INPs, GMCs and neurons (n = 15 and 25 clones, respectively). (H–I) At 48 h after clone induction, type II neuroblast clones in wild-type brains still contained a single neuroblast surrounded by immature INPs, INPs, GMCs and neurons; however, type II neuroblast clones in *lgl^{334/3644}* mutant brains contained a parental neuroblast isolated from the supernumerary neuroblasts by many immature INPs, INPs, GMCs and neurons (n = 8 and 144 clones, respectively). (J) Quantification of the number of neuroblasts per type II neuroblast clone is shown for wild-type and *lgl* mutant clones at 24 (open) and 48 (filled) hours after clone induction. (K) The table shows the frequency of clones containing supernumerary neuroblasts. Brains were stained with the indicated markers. Single neuroblast clones marked by GFP are circled by the dotted line. Arrows indicate the neuroblasts (white, Type I; yellow, Type II). Single confocal planes of the same clone are shown at 0 mm and +7.5 μm (C–D and H–I). All scale bars are 10 μm.

propose that in *lgl* mutant brains, type I neuroblasts divide asymmetrically to self-renew and to generate GMCs, which revert back into type I neuroblasts.

The 24-hour or 48-hour type II neuroblast lineage clones in wild-type brains contain a single neuroblast (Dpn^+Ase^-) per clone, and the neuroblast was always directly surrounded by immature INPs (Dpn^-Ase^- or Dpn^-Ase^+) while INPs (Dpn^+Ase^+), GMCs and their daughter progeny were typically one or more cells away (Figs. 1F, H, J–K and S1A). Surprisingly, all 24-hour type II neuroblast clones in *lgl* mutant brains also contained a single neuroblast surrounded by immature INPs, INPs and their daughter progeny (Figs. 1G, J and K). Most importantly, 47% of the 48-hour clones in *lgl* mutant brains contained more than one neuroblast per clone, and we reproducibly observed supernumerary neuroblasts formed basally in the clones (Figs. 1I, J and K; $n = 144$). These results led us to conclude that in *lgl* mutant brains, type II neuroblasts also divide asymmetrically to self-renew and to produce immature INPs that mature into INPs, but INPs revert back into type II neuroblasts. Based on these data, we propose that Lgl functions to maintain restricted potential in progenitor cells including INPs and GMCs.

Increased cortical aPKC kinase activity triggers reversion of INPs back into neuroblasts

If Lgl indeed functions to maintain restricted potential in progenitor cells, we predict that the genetic clones derived from INPs in *lgl* mutant brains should contain supernumerary type II neuroblasts. In order to induce the INP lineage clones, we first examined whether INPs in *lgl* mutant brains show expression of the INP-specific *earmuff-lacZ* reporter transgene (Weng et al., 2010) (this study). *earmuff-LacZ* was detected in small Dpn^+Ase^+ cells surrounding type II neuroblasts but was undetectable in both type I and II neuroblasts in wild-type and *lgl* mutant brains (Figs. 2A–B). Thus, the INP-specific enhancer element in the *earmuff* gene remains active in *lgl* mutant brains, allowing us to

induce lineage clones derived from INPs in wild-type or *lgl* mutant brains by expressing flipase driven by the *earmuff-Gal4* (Weng et al., 2010). All INP clones in wild-type brains ($n = 31$) contained only progeny that lack Dpn and Ase expression but never type II neuroblasts (Fig. 2C). In contrast, *lgl* mutant brains (86%, $n = 21$) contained INP clones with one or more type II neuroblasts (Fig. 2D). These aberrant neuroblasts can indeed undergo asymmetric division to self-renew and to produce progenitor cells as indicated by the presence of immature INPs within the clones. Thus, INPs can indeed revert back into type II neuroblasts in *lgl* mutant brains.

Since Lgl functions with aPKC in mitotic neuroblasts, we examined if reversion of INPs back into type II neuroblasts in *lgl* mutant brains occurs due to increased cortical aPKC kinase activity. We first tested if reduced function of aPKC can suppress supernumerary type II neuroblasts and INPs in *lgl* mutant brains. While a wild-type brain lobe contained 8 type II neuroblasts and 58 ± 8 INPs ($Dpn^+Ase^+earmuff-LacZ^+$), an *lgl* mutant brain lobe possessed 36 ± 9 type II neuroblasts and 131 ± 25 INPs (Fig. S3B). Consistent with our hypothesis, an *lgl* mutant brain lobe heterozygous for *aPKC* contained 13 ± 5 type II neuroblasts and 66 ± 19 INPs (Fig. S3B). We next directly assessed if unrestrained cortical aPKC kinase activity is sufficient to trigger reversion of INPs back into type II neuroblasts. Indeed, INPs ectopically expressing constitutively membrane localized aPKC^{caax} under the control of *earmuff-Gal4* generated supernumerary type II neuroblasts (Figs. 2E–F and S3A). Thus, precise regulation of aPKC kinase activity plays a critical role in maintaining restricted potential in INPs.

Numb requires the ACBD3 binding region for its localization and function in neuroblasts

Numb, which localized in the basal cortex of mitotic INPs and type I neuroblasts, is an excellent candidate for acting downstream of Lgl




Fig. 2. *lgl* mutant and *aPKC^{caax}* overexpressing INPs revert back to type II neuroblasts. (A–B) Wild-type and *lgl*^{334/3644} mutant brains expressed *erm-lacZ* specifically in INPs, but not in neuroblasts. (C–D) While an INP-derived clone in a wild-type brain only contained neurons, an INP-derived clone in an *lgl*^{334/3644} mutant brain contained multiple supernumerary type II neuroblasts ($n = 31$ and 21 brains, respectively). (E–F) Overexpression of *aPKC^{caax}* driven by *Erm-GAL4* leads to supernumerary type II neuroblasts in comparison to a wild-type brain ($n = 8$ and 9 brains, respectively). Brains were stained with the indicated markers. Clones marked by β-galactosidase are circled by the dotted line. Arrows indicate the neuroblasts (white, Type I; yellow, Type II) and yellow arrowheads indicate the INPs. All scale bars are 10 μm.




Fig. 3. The localization and function of Numb in neuroblasts requires the ACBD3 binding region. (A–B) Numb localized to the basal cortex of a wild-type INP, but distributed uniformly throughout the cortex of an *lgl*^{334/3644} mutant INP (n = 8 and 9, respectively). (C–H) Wild-type neuroblasts showed asymmetric localization of Numb at metaphase and telophase, while *lgl*^{334/3644} mutant neuroblasts showed uniform cortical Numb localization at metaphase and basal enrichment of Numb at telophase. Additionally, the heterozygous mutant *aPKC*^{06403/+} restored the asymmetric localization of Numb in *lgl*^{334/3644} mutant neuroblasts at metaphase and telophase (n = 30, 23, 46, 13, 9 and 9, respectively). (I) The diagram shows an illustration of the Numb protein as well as the deletion and mutant constructs used. (J–N) Similar to the full-length Numb transgenic protein, Numb^{SA}, Numb^{ΔN}, and Numb^{ΔPTB} localized exclusively in the basal cortex of telophase neuroblasts; however, Numb^{ΔAB} did not localize exclusively to the basal cortex (tag indicates the myc epitope tag in J–L and the HA epitope tag in M–N) (n = 21, 10, 15, 57, and 27, respectively). (O–R) Supernumerary expression of either full-length Numb or Numb^{ΔC} using *Ase-GAL4* suppressed the supernumerary type I neuroblasts in *lgl*^{334/3644} mutant brains, but expression of Numb^{ΔAB} or Numb^{ΔPTB} failed to rescue the supernumerary neuroblast phenotype (n = 7, 10, 6, and 5, respectively). (S) Quantification of the number of type I neuroblasts per brain lobe is shown for expression of each transgenic Numb protein by *Ase-GAL4* in *lgl*^{334/3644} mutant brains. (T–V) Expression of Numb^{ΔC}, but not Numb^{ΔAB} or Numb^{ΔPTB}, suppressed the supernumerary neuroblasts induced by expression of *aPKC*^{caax} driven by *Wor-GAL4* (n = 6 per genotype). Brains were stained with the indicated markers. Dotted lines mark the location of the apical cortex. White arrows indicate the type I neuroblasts. Scale bars are 5 μm (A–B), 10 μm (C–N), and 25 μm (O–V).

to maintain restricted potential in progenitor cells (Figs. 3A, C–D). In *lgl* mutant brains, Numb localized uniformly in the cortex of metaphase INPs (100%, n = 9) and type I neuroblasts (100%, n = 46) and became enriched in the cortex of the future GMC in telophase neuroblasts (76%, n = 13) (Figs. 3B, E and F). Furthermore, heterozygosity of *aPKC* restored asymmetric localization and segregation of Numb in mitotic INPs and type I neuroblasts in *lgl* mutant brains (Figs. 3G–H;

metaphase = 78%, n = 9; telophase = 89%, n = 9). Moreover, neuroblasts ectopically expressing *aPKC*^{caax} showed uniform cortical localization of Numb in both metaphase and telophase (Fig. S4) (Wang et al., 2006). These data led us to conclude that increased cortical *aPKC* kinase activity perturbs asymmetric localization of Numb in the cortex of mitotic neuroblasts and likely reduces the function of Numb in the cortex of the future progenitor cell in *lgl* mutant brains.

We tested if aPKC regulates asymmetric localization of Numb in mitotic neuroblasts via the five conserved aPKC phosphorylation sites proposed by a previous study (Wirtz-Peitz et al., 2008). Surprisingly, ectopic expression of the non-phosphorylatable Numb^{S2A} transgenic protein at these sites in the presence of the endogenous Numb segregated exclusively into the cortex of the future progenitor cell in telophase neuroblasts (Fig. 3J; 100%, n = 21). This result strongly suggested that aPKC regulates Numb via alternative phosphorylation sites prompting us to first identify the domain(s) required for asymmetrically localizing Numb in mitotic brain neuroblasts. We ectopically expressed the *UAS-numb* transgenes that encode various truncated forms of Numb in type I neuroblasts in the presence of endogenous Numb and examined their localization pattern (Fig. 3I). Identical to the full-length Numb transgenic protein, Numb^{ΔN}, Numb^{ΔPTB} and Numb^{ΔC} segregated exclusively into the cortex of the future GMC in the telophase neuroblasts (Figs. 3K–L and N; data not presented; 100%, n = 10, 15, and 27, respectively). In contrast, the Numb^{ΔAB} transgenic protein failed to segregate exclusively into the cortex of the future GMC in the telophase neuroblasts (Fig. 3M; 61%, n = 57). Thus, we conclude that the ACBD3 binding region is necessary for asymmetric segregation of Numb.

We next ectopically expressed this series of the *UAS-numb* transgenes in type I neuroblasts, where *Notch* signaling is dispensable for maintenance of their identity, to determine which domains mediate the function of Numb in suppressing reversion of progenitor cells in *lgl* mutant brains. While ectopic expression of Numb, Numb^{ΔN} and Numb^{ΔC} efficiently suppressed supernumerary neuroblasts in *lgl* mutant brains, expression of Numb^{ΔPTB} did not have any effects on the supernumerary neuroblast phenotype (Figs. 3O and Q–S). The PTB domain mediates Numb binding to the Notch receptor protein and is essential for Numb suppression of *Notch* signaling (Frise et al., 1996; Yaich et al., 1998). Additionally, ectopic expression of Numb in type I neuroblasts of wild-type brains using *ase-GAL4* had no effect on neuroblast number (data not presented). These data strongly suggest that aberrant activation of *Notch* signaling leads to supernumerary type I neuroblasts in *lgl* mutant brains. Most importantly, the Numb^{ΔAB} transgenic protein also failed to suppress supernumerary neuroblasts in *lgl* mutant brains (Figs. 3P and S). We independently

tested whether the ACBD3 binding region is indeed necessary for suppressing supernumerary neuroblasts induced by unrestrained cortical aPKC kinase activity. While ectopic expression of Numb or Numb^{ΔC} efficiently suppressed massive supernumerary neuroblasts induced by aPKC^{caax}, expression of Numb^{ΔAB} or Numb^{ΔPTB} did not have any effects on the supernumerary neuroblast phenotype (Figs. 3T–V; data not presented). Thus, Numb requires the ACBD3 binding region to suppress reversion of progenitor cells in *lgl* mutant brains. Together, we conclude that the ACBD3 binding region is necessary for the localization and the function of Numb during asymmetric division of brain neuroblasts.

Serines 48 and 52 are required for asymmetric localization of Numb but likely dispensable for regulating specification of progenitor cells

Our data showed that the ACBD3 binding region is indispensable for the localization and the function of Numb during neuroblast asymmetric division; therefore, we investigated if aPKC might regulate Numb through the phosphorylation sites in this domain. The ACBD3 binding region of Numb harbors two aPKC phosphorylation sites, serines 48 and 52 (Nishimura and Kaibuchi, 2007; Smith et al., 2007). If aPKC indeed regulates the localization of Numb through the ACBD3 binding region, the phosphorylation status at serines 48 and 52 should directly impinge on the distribution of the Numb protein in mitotic neuroblasts. Consistently, the non-phosphorylatable Numb^{S2A} transgenic protein localized throughout the cortex of the telophase neuroblasts (Figs. 3I and 4A; 100%, n = 17). In contrast, the phosphomimetic Numb^{S2D} transgenic protein became basally enriched in the telophase neuroblasts (Figs. 3I and 4B; 74%, n = 19). Together, these data strongly suggest that aPKC excludes Numb from the apical cortex of mitotic neuroblasts by phosphorylating serines 48 and 52.

We next tested if increased cortical aPKC kinase activity inactivates the function of Numb by phosphorylating serines 48 and 52 in the ACBD3 binding region during neuroblast asymmetric division. We ectopically expressed Numb^{S2A} or Numb^{S2D} in type I neuroblasts in *lgl* mutant brains. Surprisingly, either Numb^{S2A} or Numb^{S2D} efficiently suppressed supernumerary type I neuroblasts in *lgl* mutant




Fig. 4. Phosphorylation of Numb at serines 48 and 52 regulates its localization, but not function. (A–B) Numb^{S2A} localized uniformly throughout the entire cortex of the telophase neuroblast, while Numb^{S2D} was enriched in the basal cortex (n = 17 and 19, respectively). (C–D) Expression of Numb^{S2A} or Numb^{S2D} driven by *Ase-GAL4* suppressed the supernumerary neuroblasts in *lgl^{334/3644}* mutant brains (n = 8 and 9, respectively). (E) Quantification of the number of type I neuroblasts per brain lobe is shown for expression of the transgenic Numb proteins by *Ase-GAL4* in *lgl^{334/3644}* mutant brains. (F–I) Similar to full-length Numb, expression of Numb^{S2A} or Numb^{S2D} suppressed the supernumerary neuroblasts induced by expression of aPKC^{caax} driven by *Wor-GAL4*, while expression of Numb^{ΔAB} did not have an effect on the supernumerary neuroblast phenotype (n = 8 per genotype). Brains were stained with the indicated markers. White arrows indicate the type I neuroblasts. Scale bars are 10 μm (A–B) and 25 μm (C–I).




Fig. 5. The ACBD3 binding region of Numb specifically mediates the inhibition of *Notch* signaling in larval neuroblasts. (A) Quantification of type II neuroblasts per brain lobe is shown for *Wor-GAL4* driving expression of *Numb*^{ΔAB}, *Numb*^{S2A}, or *Numb*^{S2D} (n = 5 per genotype). (B–F) Compared to wild-type flies, overexpression of *Numb*^{ΔAB}, *Numb*^{S2A}, *Numb*^{S2D}, or *Numb*^{ΔC} driven by *Sca-GAL4* leads to a loss of scutellar bristles (n = 259, 187, 83, and 34, respectively). (G) Quantification of the number of bristles per scutellum is shown for *Sca-GAL4* driving expression of *Numb*^{ΔAB}, *Numb*^{S2A}, or *Numb*^{S2D}.

brains (Figs. 4C–E). In addition, while *Numb*^{ΔAB} failed to suppress supernumerary neuroblasts induced by *aPKC*^{caax} in larval brains, *Numb*^{S2A} or *Numb*^{S2D} completely suppressed the supernumerary neuroblast phenotype in the same genetic background (Figs. 4F–I). Thus, the phosphorylation status of serine 48 and 52 has no effects on the ability of the Numb transgenic protein to restore restricted potential in progenitor cells in *lgl* mutant brains. We propose that serines 48 and 52 play a critical role in asymmetric localization of Numb but are likely dispensable for regulation of progenitor cell potential.

The ACBD3 binding region mediates Numb-dependent suppression of Notch signaling specifically in brain neuroblasts

The ACBD3 binding region is necessary for the function of the mouse Numb protein, but how this domain mediates the function of the fly Numb protein has never been investigated (Zhou et al., 2007). The ACBD3 binding region is necessary for Numb to suppress supernumerary type I neuroblasts in larval brains lacking *lgl* function or ectopically expressing *aPKC*^{caax}, phenotypes that required

activation of *Notch* signaling (Figs. 3 and S5). Thus, we hypothesize that the ACBD3 binding region mediates Numb suppression of *Notch* signaling. We tested this hypothesis by ectopically expressing the *UAS-numb*^{ΔAB} transgene under the control of a pan-neuroblast *Wor-Gal4* driver in the larval brain. While increased function of *numb* or decreased function of *Notch* led to premature differentiation of type II neuroblasts, expression of *Numb*^{ΔAB} did not have any effects on maintenance of the type II neuroblast identity (Figs. S1C–E, G and 5A). Importantly, expression of *Numb*^{S2A} or *Numb*^{S2D} led to complete loss of type II neuroblasts prematurely in larval brains, indicating that the phosphorylation status at serines 48 and 52 does not affect the function of Numb to antagonize *Notch* signaling (Fig. 5A).

Asymmetric divisions of sensory organ precursors give rise to the bristles on the scutellum of the adult fly and are highly sensitive to changes in *Notch* signaling (Frise et al., 1996; Knoblich et al., 1997; Rhyu et al., 1994; Yaich et al., 1998). Similar to over-expression of *Numb*^{ΔC}, unexpectedly, ectopic expression of *Numb*^{ΔAB}, *Numb*^{S2A} or *Numb*^{S2D} all led to cell fate transformation in the sensory organ precursor lineage and resulted in decreased bristles on the scutellum (Figs. 5B–G). This result indicates that the ACBD3 binding region is dispensable for Numb-mediated suppression of *Notch* signaling in




Fig. 6. Diagram of proposed model. In wild-type neuroblasts, the mutual antagonism between *Lgl* and *aPKC* ensures asymmetric segregation of Numb into the progenitor cell, where Numb antagonizes *Notch* signaling via its PTB and ACBD3 protein binding domains. In the absence of *lgl*, increased cortical *aPKC* kinase activity redistributes Numb in the cortex of the dividing neuroblast, potentially leading to insufficient Numb to inhibit *Notch* in the progenitor cell. This increases the *Notch* activity in the progenitor cells and drives the progenitor cell to revert back into a neuroblast. Furthermore, expression of the phosphomimetic *Numb*^{S2D} rescues the *lgl* mutant supernumerary neuroblast phenotype by restoring Numb and the inhibition of *Notch* activity in the progenitor cell.

sensory organ precursor cells in the peripheral nervous system. Together, we conclude that the ACBD3 binding region specifically mediates the function of Numb in suppressing *Notch* signaling in the brain regardless of the phosphorylation by aPKC.

Discussion

The antagonistic interaction between the polarity proteins Lgl and aPKC provides an evolutionarily conserved mechanism for regulating the cell fate determinants inherited into the daughter siblings during asymmetric cell division (Beatty et al., 2010; Betschinger et al., 2003; Hooge et al., 2010; Lee et al., 2006). However, regulation of the localization and the function of fate determinants by the polarity proteins can be uncoupled in a context-dependent manner. Our study led us to conclude that the antagonistic interaction between Lgl and aPKC maintains limited potential in progenitor cells at least in part by ensuring asymmetric partitioning of Numb into the future progenitor cells, where Numb acts irrespective of its phosphorylation by aPKC (Fig. 6). We showed that neuroblasts in *lgl* mutant brains undergo asymmetric division to self-renew and to generate progenitor cells, which can produce post-mitotic progeny but can also revert back into neuroblasts (Figs. 1 and 2). This indicates that although progenitor cells initially establish the proper identity these cells fail to maintain their limited potential. Additionally, Numb remained localized in the cortex of telophase neuroblasts lacking *lgl* function or ectopically expressing aPKC^{caax} (Fig. 3F and data not presented). The phosphomimetic Numb^{2D} transgenic protein at serines 48 and 52, which are essential for asymmetric cortical localization of Numb, also remained localized in the cell cortex of telophase neuroblasts (Fig. 4B). Finally, increased function of Numb efficiently suppressed supernumerary neuroblasts in *lgl* mutant brains irrespective of the phosphorylation status at serines 48 and 52 (Figs. 4C–D). Thus, we propose that Lgl antagonizes aPKC to ensure a necessary threshold of Numb in the progenitor cells where Numb maintains limited potential regardless of phosphorylation by aPKC (Fig. 6).

Lgl maintains limited potential in progenitor cells

How Lgl suppresses formation of supernumerary neuroblasts in larval brains has remained a mystery largely due to the existence of a phenomenon called “telophase rescue” (Albertson and Doe, 2003; Cai et al., 2001). In *lgl* mutant brains, the basal proteins including Miranda and Numb fail to localize to the basal crescent in metaphase neuroblasts but by and large re-localize asymmetrically in telophase neuroblasts (Lee et al., 2006; Rolls et al., 2003; Wirtz-Peitz et al., 2008). Furthermore, Miranda and Numb appear to localize independently of each other in mitotic neuroblasts (Lu et al., 1998; Shen et al., 1997). Thus, the transcription factor Dpn whose expression and localization pattern is not impinged upon by defective cortical cell polarity provides an excellent cell identity marker to investigate the cellular origin of supernumerary neuroblasts in *lgl* mutant brains (Fig. S1A; Komori and Lee, unpublished). Surprisingly, only the 48-hour, but not the 24-hour, type I and II neuroblast lineage clones in *lgl* mutant brains contained supernumerary neuroblasts, which frequently localized basally from the parental neuroblasts (Fig. 1). The most recently born daughter always remains immediately adjacent to the parental neuroblasts while the earlier born progeny gradually becomes displaced away from the parental neuroblasts (Bayraktar et al., 2010; Bowman et al., 2008; Weng et al., 2010). Thus, supernumerary neuroblasts in *lgl* mutant brains most likely originated from the progenitor cells rather than symmetric neuroblast division. We propose that in the type I neuroblast lineage Lgl prevents aPKC kinase activity in the basal cortex to ensure that GMCs maintain limited potential and generate only post-mitotic progeny. In the type II neuroblast lineage, Lgl prevents aPKC kinase activity in the basal cortex to ensure that after

maturation the INP can maintain limited potential and generate only GMCs during limited rounds of asymmetric divisions.

Although these results do not exclude the possibility that GMCs in the type II neuroblast lineages in *lgl* mutant brains can revert back into neuroblasts, we believe that reacquisition of the type II neuroblast fate by GMCs might be less likely. First, the basal protein Prospero plays a critical role in regulating the function of GMCs, and mosaic clones derived from *prospero* mutant INPs contained massive supernumerary INPs but never supernumerary type II neuroblasts (Hirata et al., 1995; Knoblich et al., 1995; Spana and Doe, 1995; Weng et al., 2010). Second, mosaic clones derived from *numb* mutant INPs contained supernumerary INPs but never supernumerary type II neuroblasts (Komori and Lee, unpublished). Thus, blocking differentiation in GMCs allows them to retain the identity of their immediate parental cell type, which is INP in the type II neuroblast lineage. Until the enhancer elements that exhibit GMC-specific expression become available, we cannot conclusively rule out the possibility that GMCs can re-acquire the type II neuroblast fate in *lgl* mutant brains.

aPKC regulates asymmetric localization but not the function of Numb in neuroblasts

A previous study strongly suggested that increased cortical aPKC kinase activity induces supernumerary neuroblasts in *lgl* mutant brains by phosphorylating Numb, therefore, displacing it from the neuroblast cortex and inactivating its function (Wirtz-Peitz et al., 2008). This proposed mechanism was in part based on studies in vertebrates showing that phosphorylation by aPKC perturbs cortical localization and the function of Numb (Nishimura and Kaibuchi, 2007; Smith et al., 2007). Inconsistent with this proposed mechanism, we reproducibly detected disruption in asymmetric distribution of Numb in the cortex of mitotic neuroblasts in *lgl* mutant brains (Figs. 3E–H). Failure to displace Numb from the cortex of mitotic neuroblasts in *lgl* mutant brains was unlikely due to insufficient aPKC kinase activity as Numb remained localized uniformly in the cortex of neuroblasts over-expressing aPKC^{caax} (Fig. S4). Additionally, the phosphomimetic Numb^{2D} transgenic protein at serines 48 and 52, two residues required for asymmetric segregation of Numb in mitotic neuroblasts, remained localized in the neuroblast cortex (Fig. 4B). Thus, increased cortical aPKC kinase activity most likely disperses Numb in the cortex of mitotic neuroblasts and reduces accumulation of Numb in the cortex of the future progenitor cell in *lgl* mutant brains. Most importantly, over-expression of Numb^{2D} suppressed supernumerary neuroblasts in *lgl* mutant brains as efficiently as Numb^{2A}, strongly suggesting that phosphorylation by aPKC does not inactivate the function of Numb (Fig. 4). Together, these data led us to propose that increased cortical aPKC kinase activity induces supernumerary neuroblasts in *lgl* mutant brains by reducing accumulation of Numb rather than inactivation of Numb in the progenitor cells (Fig. 6).

Studies in vertebrates identified five conserved aPKC phosphorylation sites in the fly Numb, and the non-phosphorylatable Numb^{5A} transgenic protein at these sites localized uniformly cortical in mitotic sensory organ precursors (Nishimura and Kaibuchi, 2007; Smith et al., 2007). Surprisingly, Numb^{5A} localized asymmetrically in mitotic brain neuroblasts, indicating that these sites are dispensable for exclusion of Numb from the apical cortex by aPKC (Fig. 3J). Thus, many important questions regarding the significance of these five conserved aPKC phosphorylation sites on the localization and the function of Numb remain to be tested. For example, does the phosphomimetic Numb^{5D} transgenic protein indeed fail to localize to the cell cortex of mitotic sensory organ precursors? Furthermore, is Numb^{5D} indeed non-functional? Does over-expression of Numb^{5D} have any effects on cell fate determination in the sensory organ precursor cell lineage? Are the non-conserved aPKC phosphorylation sites dispensable for the function of Numb in the sensory organ

precursor cell lineage? Most importantly, the extent to which these conserved and non-conserved aPKC phosphorylation sites might impinge on other biological processes regulated by Numb requires additional direct and rigorous assessment.

The ACBD3 binding region mediates Numb suppression of Notch signaling specifically in the brain

Numb is a highly conserved protein and exerts its antagonistic effect on *Notch* signaling via the PTB domain, which mediates direct binding to the Notch receptor protein (Frise et al., 1996; Knoblich et al., 1997; Yaich et al., 1998; Zhong et al., 1997). A previous study identified that the ACBD3 binding region is required for asymmetric localization of the Numb protein and Numb-dependent suppression of *Notch* signaling (Zhou et al., 2007). Our study extended this result and showed that the ACBD3 binding region mediates asymmetric cortical localization of the fly Numb protein in mitotic neuroblasts through an aPKC-regulated mechanism (Fig. 3). Interestingly, the ACBD3 binding region regulates tissue-specific suppression of *Notch* signaling by Numb despite the presence of the PTB domain (Fig. 5). Since the ACBD3 binding region appears to mediate direct protein-protein interactions, we propose that this domain serves as a platform in which tissue-specific regulators can exert precise control of the Numb function in antagonizing *Notch* signaling. Identification and functional characterization of proteins that interact with Numb through the ACBD3 binding region will provide novel mechanistic insight into how the evolutionarily conserved Numb-dependent suppression of *Notch* signaling can be precisely regulated in a tissue-specific manner.

Supplementary materials related to this article can be found online at doi:10.1016/j.ydbio.2012.02.027.

Acknowledgments

We thank Drs. C. Doe, S. Goode, J. Knoblich, F. Schweisguth, J. Skeath and G. Struhl for fly stocks and antibody reagents. We thank the Bloomington *Drosophila* Stock Center and Vienna *Drosophila* RNAi Center for fly stocks and the Developmental Studies Hybridoma Bank for antibodies. We thank the anonymous reviewer and members of the Lee lab for reading the manuscript and providing critical comments. J.M.H. was supported by the NIH Cellular and Molecular Biology Training Grant T32-GM007315. C.K. was also supported by the NIH MSTP Training Grant T32-GM07863. C.-Y.L. was supported by University of Michigan start-up, the Burroughs Wellcome Fund Career Award in the Biomedical Sciences (1006160.01), a Sontag Foundation Distinguished Scientist Award and an NIH grant R01-GM092818.

References

Albertson, R., Doe, C.Q., 2003. Dlg, Scrib and Lgl regulate neuroblast cell size and mitotic spindle asymmetry. *Nat. Cell Biol.* 5, 166–170.

Bayraktar, O.A., Boone, J.Q., Drummond, M.L., Doe, C.Q., 2010. *Drosophila* type II neuroblast lineages keep Prospero levels low to generate large clones that contribute to the adult brain central complex. *Neural Dev.* 26 Oct 1.

Beatty, A., Morton, D., Kempthues, K., 2010. The *C. elegans* homolog of *Drosophila* Lethal giant larvae functions redundantly with PAR-2 to maintain polarity in the early embryo. *Development* 137, 3995–4004.

Bello, B.C., Izergina, N., Caussinus, E., Reichert, H., 2008. Amplification of neural stem cell proliferation by intermediate progenitor cells in *Drosophila* brain development. *Neural Dev.* 3.

Betschinger, J., Mechtler, K., Knoblich, J.A., 2003. The Par complex directs asymmetric cell division by phosphorylating the cytoskeletal protein Lgl. *Nature* 422, 326–330.

Bischof, J., Basler, K., 2008. Recombinases and their use in gene activation, gene inactivation, and transgenesis. *Methods Mol. Biol.* 420, 175–195.

Boone, J.Q., Doe, C.Q., 2008. Identification of *Drosophila* type II neuroblast lineages containing transit amplifying ganglion mother cells. *Dev. Neurobiol.* 68, 1185–1195.

Bowman, S.K., Rolland, V., Betschinger, J., Kinsey, K.A., Emery, G., Knoblich, J.A., 2008. The tumor suppressors Brat and Numb regulate transit-amplifying neuroblast lineages in *Drosophila*. *Dev. Cell* 14, 535–546.

Cai, Y., Chia, W., Yang, X., 2001. A family of snail-related zinc finger proteins regulates two distinct and parallel mechanisms that mediate *Drosophila* neuroblast asymmetric divisions. *EMBO J.* 20, 1704–1714.

Chia, W., Somers, W.G., Wang, H., 2008. *Drosophila* neuroblast asymmetric divisions: cell cycle regulators, asymmetric protein localization, and tumorigenesis. *J. Cell Biol.* 180, 267–272.

Chung, H.M., Struhl, G., 2001. Nicastrin is required for Presenilin-mediated transmembrane cleavage in *Drosophila*. *Nat. Cell Biol.* 3, 1129–1132.

Dho, S.E., Trejo, J., Siderovski, D.P., McClade, C.J., 2006. Dynamic regulation of mammalian numb by G protein-coupled receptors and protein kinase C activation: structural determinants of numb association with the cortical membrane. *Mol. Biol. Cell* 17, 4142–4155.

Doe, C.Q., 2008. Neural stem cells: balancing self-renewal with differentiation. *Development* 135, 1575–1587.

egger, B., Chell, J.M., Brand, A.H., 2008. Insights into neural stem cell biology from flies. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 39–56.

Frise, E., Knoblich, J.A., Younger-Shepherd, S., Jan, L.Y., Jan, Y.N., 1996. The *Drosophila* Numb protein inhibits signaling of the Notch receptor during cell-cell interaction in sensory organ lineage. *Proc. Natl. Acad. Sci. U. S. A.* 93, 11925–11932.

Guo, M., Jan, L.Y., Jan, Y.N., 1996. Control of daughter cell fates during asymmetric division: interaction of Numb and Notch. *Neuron* 17, 27–41.

Hirata, J., Nakagoshi, H., Nabeshima, Y., Matsuzaki, F., 1995. Asymmetric segregation of the homeodomain protein Prospero during *Drosophila* development. *Nature* 377, 627–630.

Hoege, C., Constantinescu, A.T., Schwager, A., Goehring, N.W., Kumar, P., Hyman, A.A., 2010. LGL can partition the cortex of one-cell *Caenorhabditis elegans* embryos into two domains. *Curr. Biol.* 20, 1296–1303.

Knoblich, J.A., 2008. Mechanisms of asymmetric stem cell division. *Cell* 132, 583–597.

Knoblich, J.A., 2010. Asymmetric cell division: recent developments and their implications for tumour biology. *Nat. Rev. Mol. Cell Biol.* 11, 849–860.

Knoblich, J.A., Jan, L.Y., Jan, Y.N., 1995. Asymmetric segregation of Numb and Prospero during cell division. *Nature* 377, 624–627.

Knoblich, J.A., Jan, L.Y., Jan, Y.N., 1997. The N terminus of the *Drosophila* Numb protein directs membrane association and actin-dependent asymmetric localization. *Proc. Natl. Acad. Sci. U. S. A.* 94, 13005–13010.

Lee, C.Y., Robinson, K.J., Doe, C.Q., 2006. Lgl, Pins and aPKC regulate neuroblast self-renewal versus differentiation. *Nature* 439, 594–598.

Lu, B., Rothenberg, M., Jan, L.Y., Jan, Y.N., 1998. Partner of Numb colocalizes with Numb during mitosis and directs Numb asymmetric localization in *Drosophila* neural and muscle progenitors. *Cell* 95, 225–235.

Neumüller, R.A., Knoblich, J.A., 2009. Dividing cellular asymmetry: asymmetric cell division and its implications for stem cells and cancer. *Genes Dev.* 23, 2675–2699.

Nishimura, T., Kaibuchi, K., 2007. Numb controls integrin endocytosis for directional cell migration with aPKC and PAR-3. *Dev. Cell* 13, 15–28.

Peng, C.Y., Manning, L., Albertson, R., Doe, C.Q., 2000. The tumour-suppressor genes *Igl* and *dlg* regulate basal protein targeting in *Drosophila* neuroblasts. *Nature* 408, 596–600.

Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.T., Misra, S., Murphy, C., Scully, A., Carlson, J.W., Wan, K.H., Lavery, T.R., Mungall, C., Svirskas, R., Kadonaga, J.T., Doe, C.Q., Eisen, M.B., Celniker, S.E., Rubin, G.M., 2008. Tools for neuroanatomy and neurogenetics in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* 105, 9715–9720.

Prehoda, K.E., 2009. Polarization of *Drosophila* neuroblasts during asymmetric division. *Cold Spring Harb. Perspect. Biol.* 1, a001388 Aug.

Rhyu, M.S., Jan, L.Y., Jan, Y.N., 1994. Asymmetric distribution of numb protein during division of the sensory organ precursor cell confers distinct fates to daughter cells. *Cell* 76, 477–491.

Rolls, M.M., Albertson, R., Shih, H.P., Lee, C.Y., Doe, C.Q., 2003. *Drosophila* aPKC regulates cell polarity and cell proliferation in neuroblasts and epithelia. *J. Cell Biol.* 163, 1089–1098.

San-Juan, B.P., Baonza, A., 2011. The bHLH factor deadpan is a direct target of Notch signaling and regulates neuroblast self-renewal in *Drosophila*. *Dev. Biol.* 352, 70–82.

Shen, C.P., Jan, L.Y., Jan, Y.N., 1997. Miranda is required for the asymmetric localization of Prospero during mitosis in *Drosophila*. *Cell* 90, 449–458.

Smith, C.A., Lau, K.M., Rahmani, Z., Dho, S.E., Brothers, G., She, Y.M., Berry, D.M., Bonnell, E., Thibault, P., Schweisguth, F., Le Borgne, R., McClade, C.J., 2007. aPKC-mediated phosphorylation regulates asymmetric membrane localization of the cell fate determinant Numb. *EMBO J.* 26, 468–480.

Song, Y., Lu, B., 2011. Regulation of cell growth by Notch signaling and its differential requirement in normal vs. tumor-forming stem cells in *Drosophila*. *Genes Dev.* 25, 2644–2658.

Sotillos, S., Díaz-Meco, M.T., Caminero, E., Moscat, J., Campuzano, S., 2004. DaPKC-dependent phosphorylation of Crumbs is required for epithelial cell polarity in *Drosophila*. *J. Cell Biol.* 166, 549–557.

Spana, E.P., Doe, C.Q., 1995. The prospero transcription factor is asymmetrically localized to the cell cortex during neuroblast mitosis in *Drosophila*. *Development* 121, 3187–3195.

Wang, H., Somers, G.W., Bashirullah, A., Heberlein, U., Yu, F., Chia, W., 2006. Aurora-A acts as a tumor suppressor and regulates self-renewal of *Drosophila* neuroblasts. *Genes Dev.* 20, 3453–3463.

Weng, M., Golden, K.L., Lee, C.Y., 2010. dFzefz/Earmuff maintains the restricted developmental potential of intermediate neural progenitors in *Drosophila*. *Dev. Cell* 18, 126–135.

- Weng, M., Haenfler, J.M., Lee, C.Y., 2011. Changes in Notch signaling coordinates maintenance and differentiation of the *Drosophila* larval optic lobe neuroepithelia. *Dev. Neurobiol.* doi:10.1002/dneu.20995.
- Weng, M., Lee, C.Y., 2011. Keeping neural progenitor cells on a short leash during *Drosophila* neurogenesis. *Curr. Opin. Neurobiol.* 21, 36–42.
- Wirtz-Peitz, F., Nishimura, T., Knoblich, J.A., 2008. Linking cell cycle to asymmetric division: Aurora-A phosphorylates the Par complex to regulate Numb localization. *Cell* 135, 161–173.
- Yaich, L., Ooi, J., Park, M., Borg, J.P., Landry, C., Bodmer, R., Margolis, B., 1998. Functional analysis of the Numb phosphotyrosine-binding domain using site-directed mutagenesis. *J. Biol. Chem.* 273, 10381–10388.
- Zhong, W., Jiang, M.M., Weinmaster, G., Jan, L.Y., Jan, Y.N., 1997. Differential expression of mammalian Numb, Numbl like and Notch1 suggests distinct roles during mouse cortical neurogenesis. *Development* 124, 1887–1897.
- Zhou, Y., Atkins, J.B., Rompani, S.B., Bancescu, D.L., Petersen, P.H., Tang, H., Zou, K., Stewart, S.B., Zhong, W., 2007. The mammalian Golgi regulates numb signaling in asymmetric cell division by releasing ACBD3 during mitosis. *Cell* 129, 163–178.
- Zhu, S., Lin, S., Kao, C.F., Awasaki, T., Chiang, A.S., Lee, T., 2006. Gradients of the *Drosophila* Chinmo BTB-zinc finger protein govern neuronal temporal identity. *Cell* 127 (2), 409–422.