



ELSEVIER

journal homepage: www.elsevier.com/locate/febsopenbio

Diffuse binding of Zn²⁺ to the denatured ensemble of Cu/Zn superoxide dismutase 1



Scarlett Szpryngiel, Mikael Oliveberg, Lena Mäler*

Department of Biochemistry and Biophysics, The Arrhenius Laboratory, Stockholm University, 10691 Stockholm, Sweden

ARTICLE INFO

Article history:

Received 22 October 2014

Revised 30 December 2014

Accepted 30 December 2014

Keywords:

NMR

SOD1

Folding

Metal-binding

ABSTRACT

The stability and structural properties of the metalloprotein superoxide dismutase 1 (SOD1) are found to depend critically on metal ions. Native SOD1 monomers coordinate one structural Zn²⁺ and one redox-active Cu^{2+/1+} to the active site. To do this, the Zn²⁺ ions need to interact with the SOD1 protein on the denatured side of the folding barrier, prior to the formation of the folding nucleus. In this study, we have examined at residue level the nature of this early Zn²⁺ binding by NMR studies on the urea denatured-state of SOD1. Nearly complete backbone chemical shift assignments were obtained in 9 M urea at physiological pH, conditions at which NMR studies are scarce. Our results demonstrate that SOD1 is predominantly unstructured under these conditions. Chemical-shift changes upon Zn²⁺ titration show that denatured SOD1 retains a significant affinity to Zn²⁺ ions, even in 9 M urea. However, the Zn²⁺ interactions are not limited to the native metal-binding ligands in the two binding sites, but are seen for all His residues. Moreover, the native Cu^{2+/1+} ligand H46 seems not to bind as well as the other His residues, while the nearby non-native H43 does bind, indicating that the binding geometry is relaxed. The result suggests that the Zn²⁺-binding observed to catalyze folding of SOD1 in physiological buffer is initiated by diffuse, non-specific coordination to the coil, which subsequently funnels by ligand exchange into the native coordination geometry of the folded monomer. Altogether, this diffuse binding is a result with fundamental implications for folding of metalloproteins in general.

© 2015 The Authors. Published by Elsevier B.V. on behalf of the Federation of European Biochemical Societies. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Superoxide dismutases are ubiquitous enzymes that neutralize reactive oxygen species in the cells. Human Cu/Zn superoxide dismutase (SOD1) is a 153 amino acid residue homo-dimeric protein with two cofactors per subunit; one copper and one zinc (Fig. 1). The copper ion is, in its oxidized form, bound by four histidines (H46, H48, H63 and H120). All these histidines are essential for enzyme activity [1]. The zinc ion is bound by three histidines (H63, H71 and H80) and one aspartate (D83), all part of what is commonly called “the zinc binding loop” (loop IV, residues 48–84). This loop and its zinc ion stabilize the protein structure,

including both metal centers [2–5] but Cu²⁺ is necessary for catalysis and native structure (Fig. 1). Both metals, as well as many other divalent cations, are able to bind to both metal sites in natively folded SOD1 *in vitro* [1].

Zn²⁺ binds to both the zinc and copper sites, and this observation has been used to elucidate how metals affect folding of SOD1 [5–7]. These studies showed that Zn²⁺ catalyses protein folding by binding transiently to the SOD1 Cu^{2+/1+} ligands. Once folding is completed, the zinc ion is transferred to the thermodynamically favorable position in the Zn²⁺ site [7]. A question then arises concerning the nature of this Zn²⁺ coordination. There are many potential competing ligands (e.g. His residues, charged residues) in the protein and furthermore there is no tertiary structural context that would favor the native ligands. To investigate the nature of the early Zn²⁺ binding we have in this work focused on the metal-binding properties of unfolded SOD1. We have examined possible structures that may arise due to specific metal interaction of the urea-denatured state of the monomeric SOD1, F50E/G51E, C6A/C111A variant (SOD1^{G111/50/51}). The first two substitutions hinder dimer formation [8], whereas the latter two abolish erroneous

Abbreviations: SOD1, superoxide dismutase 1; HSQC, heteronuclear single-quantum coherence; TOCSY, total correlation spectroscopy; NOESY, nuclear Overhauser enhancement spectroscopy; IDP, intrinsically disordered protein

Database: The backbone chemical shift assignments for urea-denatured human SOD1 have been deposited with the Biological Magnetic Resonance Bank (BMRB) under the accession code 18968.

* Corresponding author. Tel.: +46 8 162448.

E-mail address: lena.maler@dbb.su.se (L. Mäler).

<http://dx.doi.org/10.1016/j.fob.2014.12.003>

2211-5463/© 2015 The Authors. Published by Elsevier B.V. on behalf of the Federation of European Biochemical Societies. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

disulfide bond formation [9], which provides a way to study the folding and metal-binding properties of the monomeric protein. Our aim with the study was to examine the early folding events as introduced by Zn^{2+} coordination. For doing this, we need relatively high Zn^{2+} concentrations in order to capture what may be relevant Zn^{2+} -interactions for the unfolded state even at physiological conditions. Urea-denatured SOD1 may be a simple model for the unfolded state, but nevertheless presents a way to capture possible ways for the protein to interact with metal ions. In this way we are able to capture the effect of metalation at the very start of the folding reaction, prior to the formation of a folding nucleus. We have furthermore characterized the residual structure in the denatured protein and examined the effect of urea at several urea concentrations. The results presented here demonstrate that Zn^{2+} -binding to unfolded SOD1 occurs through diffuse and non-native coordination. This suggests that the ligands are shuffled towards the native coordination during the course of the folding reaction providing a favorable funnel-like restriction of the conformational entropy of the coil.

2. Results

2.1. SOD1 is unstructured in 9 M urea

The overall appearance of the ^{15}N -HSQC is indicative of a protein with no or very little secondary structure (Fig. 2). We have assigned the resonances of backbone atoms (N^{H} , C^{α} , C' and H^{α}) in apo-SOD1^{6/111/50/51} in 9 M urea and pH 6.3 to a completeness of 97%. The *N*-terminal residues A1 and T2 lack assignment since there were no detectable peaks in any of the spectra that could be assigned to them. C' of H63, F64 and P66 were not assigned whereas the rest of the assignment for the backbone is complete.

To evaluate the extent of remaining structure of urea-denatured SOD1^{6/111/50/51}, we compared the backbone carbon atom chemical shifts to those previously measured for short peptides [10] using the chemical shift index method [11]. This simple analysis verified that only little structure existed in the protein (Fig. 3). Most of the very small deviations in the chemical shifts, which were only observed for C^{α} , were slightly positive, corresponding to α -helical structure. The analysis of $\Delta\delta(\text{C}')$ supported this observation, indicating a random variation along the sequence. Since the deviations were so small (on average less than 0.4 ppm) and did not indicate β -sheet structure we concluded that neither the $\Delta\delta(\text{C}^{\alpha})$ nor the $\Delta\delta(\text{C}')$ analyses support the presence of any significant degree of

structure in SOD1^{6/111/50/51}. Additionally, and as expected for random coil conformations, no long-range NOEs were found. This further underlines that there are no secondary structure elements in SOD1 under the conditions studied. Therefore, we conclude that there is no residual β -sheet structure in the protein.

Since we assigned urea-denatured SOD1, we also compared the chemical shifts with data obtained for denatured GGXGG peptides by Schwarzingger et al. [12,13], which were measured at much lower pH (pH 2.3) than in the present study. No significant differences between this analysis and the one using the dataset by Wishart et al. were observed, except that all aspartic and glutamic acids in the amino acid sequence of SOD1 showed extreme values (data not shown), as expected for pH-dependent residues. Also, $\Delta\delta(\text{C}^{\alpha})$ for N65 displayed a large deviation from what is expected for an unfolded protein, most likely due to the proline situated subsequent to it [10,13]. This effect was also seen for all glycines followed by prolines (G12, G27, G61 and G73). Similar deviations in the $\Delta\delta(\text{C}^{\alpha})$ for proline-neighboring residues was seen when using the Wishart dataset but was compensated for, as described [10].

To determine the effect of the denaturant on the protein structure the urea concentration was lowered in steps of 0.5 M. Fig. 4 shows the effect on the chemical shifts of H^{N} and N^{H} when diluting the urea from 9 to 4 M. The changes ($\Delta_{\text{avg}}\delta$) were generally very small, on the order of 0.01–0.08 ppm. It is noteworthy that the region that displayed the largest shifts when lowering the urea concentration comprises amino acid residues H63–H71, which in turn is part of the zinc-binding loop. Especially residues S68 and R69 clearly shifted towards the expected chemical shifts in the natively folded protein, as seen in a previous assignment of apo-SOD1 [14]. Two other regions displayed small but consistent chemical shift differences that may be significant, I17–K23 and R143–I149. These regions correspond to β -strands 2 and 8 in the folded protein.

2.2. Zn^{2+} binds to all His residues in SOD1

To monitor Zn^{2+} -binding of urea-denatured SOD1^{6/111/50/51} we performed a series of ^{15}N -HSQC experiments (Fig. 5A). The protein was exposed to 9 M urea and increasing Zn^{2+} concentrations and the resulting chemical shift changes, $\Delta_{\text{avg}}\delta$ were plotted as a function of relative Zn^{2+} concentration (Fig. 5B). When zinc ions were added at lower ratio than 1:1 molar concentration there were no noticeable changes in the spectra ($\Delta_{\text{avg}}\delta < 0.01$ ppm, Fig. 5B). At equivalent concentrations of Zn^{2+} and protein (1:1 Zn^{2+} :SOD1)




Fig. 1. Metal binding sites in folded, dimeric, holo-SOD1. The zinc ion (gray) is bound by three histidines (cyan) and one aspartate (red). The copper ion (orange) is bound by four histidines. H63 is bridging both metals in the oxidized form. Loop IV (the so-called zinc-binding loop) is depicted in pink, loop VII (the “electrostatic loop”) in blue. The figure was made in VMD with the coordinates taken from the PDB (accession code 2GBT, www.pdb.org).




Fig. 2. A ^{15}N -HSQC indicating the backbone assignment of SOD1^{6/111/50/51} in 9 M urea. Basically all residues have typical random coil chemical shifts, giving the spectrum the typical clustering appearance one normally see for unfolded proteins without residual elements.

the peaks of all eight His residues shifted slightly to higher values for H^{N} and lower for N^{H} ($\Delta_{\text{avg}}\delta \sim 0.01$ ppm for all His residues, Fig. 5B), and this became even more pronounced at a 2:1 Zn^{2+} :SOD1 ratio. No consistent line-broadening for the shifting His peaks and no additional peaks were observed. We therefore conclude that the interaction is most likely characterized by a fast exchange event. At higher concentrations, certain differences between the His residues could be discerned. The chemical shifts of H46 were significantly less affected than what was seen for the other residues; the maximum change was around 0.06 ppm, which can be compared to H71 for which the maximum was 0.16 ppm (Fig. 5B). Fig. 6A shows the chemical shift changes at a Zn^{2+} :SOD1 ratio of 12:1, and from this plot it is evident that all His peaks but one, His46, has shifted their position significantly. At this concentration, also peaks neighboring the His residues (i.e. V47, E49, F64 and E121) display shifts that are larger than what is observed for the bulk of the protein (Fig. 6), supporting

the observation that the effect is specific for His residues. At a very high concentration of Zn^{2+} , corresponding to more than a 30:1 ratio, the resonances of most peaks shifted in a more uniform way and the specificity of the interaction was less pronounced. To ascertain that this effect at a high zinc concentration was purely ionic, measurement with a “charge equivalent” of NaCl corresponding to a 30:1 Zn^{2+} :SOD1 ratio was performed. No shifts were, however, observed in the spectrum by adding NaCl. Hence, a high ZnSO_4 concentration induces a general effect in the protein that was not observed using NaCl. Adding NaCl also demonstrated that the effect observed for the His residues is specific for Zn^{2+} and not a consequence of altering the sample conditions.

Next, the patterns of the shifts were analyzed (Fig. 5B). All of the His resonances were initially shifted in a linear manner with increasing concentrations of Zn^{2+} . However, at the same ratio where we also observed more general shifts in the protein, there was a change in this linear pattern for several residues (H43, H46, H48 and H80), and the H^{N} resonances were now shifted to lower values, indicating a possibility for one or several intermediate species along the titration series. This effect continued until a concentration around 60:1 Zn^{2+} :SOD1. Here saturation was achieved, with no further changes. None of the other residues, including the native Zn^{2+} -ligand D83 display large shifts even at these high Zn^{2+} concentrations (data not shown). In conclusion, we find that all His residues display chemical shift changes, but that for H46 the change is much smaller than for the others, indicating that this residue is less involved in metal coordination.

3. Discussion

NMR studies of intrinsically disordered proteins (IDPs) have provided a wealth of information about how to best describe the properties of them [15–19]. It has been established that there often is residual structure present and that there is variation in the dynamic behavior along the backbone [20,21]. There are, however, not that many examples of NMR studies of proteins in high concentrations of urea at physiological pH. Suggestions have been made that residual structure remains also in high amounts of urea, but these studies were conducted at low pH [22]. There have been arguments both for and against residual structure in urea-unfolded proteins, as reviewed by McCarney et al. [23]. The urea-unfolded barstar was demonstrated to contain little native structure [24], while studies in lower amounts of urea revealed residual helical structure [25]. For the tRNA methyltransferase YibK it was on the other hand suggested that residual structure remained in 8 M urea at pH 7.5, although the protein was mostly described as random coil [26]. Therefore, to understand the molecular properties related to folding of proteins, studies of unfolded states are necessary. Here we report on the backbone assignment of monomeric urea-denatured SOD1^{6/111/50/51}. Our analysis of the backbone chemical shifts shows that SOD1 in 9 M urea is unstructured, and remains largely unstructured to a urea concentration of less than 4 M. At low urea concentration evidence for slight structure induction in two of the β -strands (strands 2 and 8) and in the zinc-loop is observed. This is in agreement with an earlier study by Assfalg et al. that demonstrated that monomeric SOD1 is largely unstructured in guanidinium chloride (GdmCl) [27].

Although we needed high concentrations of Zn^{2+} to map out the binding sites in the urea-denatured state of SOD1, the results demonstrate the possible interaction sites also in unfolded SOD1 under more physiological conditions. Theoretically, the insertion of metal ions and the folding of a polypeptide chain may take place in any order. Either the metal ion is bound to the completely or partially folded structure, or to the unfolded polypeptide chain. In the case of active (holo) azurin, a blue-copper protein, the complete folding




Fig. 3. Secondary chemical shifts for SOD1^{6/111/50/51} in 9 M urea. The position of the secondary structure elements as seen in the X-ray structure of monomeric SOD1 (accession code 3HFF, www.pdb.org) are shown under the sequence. In panel A C_α is shown, in panel B C^β secondary shifts according to Zhang et al. [8].

pathway has been shown to be 4000 times faster if copper is allowed to bind before the rest of the polypeptide folds [28,29]. A few years later it was shown that the metal ions could bind in several ways, but with one major population [30]. Here we have shown that in unfolded SOD1 most of the His residues, and not only the ligands in the two metal-binding sites, are sensitive to addition of Zn^{2+} ions. We also noted that high concentrations of Zn^{2+} were needed to induce significant changes in the NMR spectrum of urea-denatured SOD1. The relatively high concentration needed may indicate that all of the His residues are transiently involved in binding of several ions, but it should also be noted that urea is known to chelate Zn^{2+} , and that the effective concentration present for interaction with the His residues may be much lower [31]. Under the conditions used in this study, with high Zn^{2+} concentrations, the effects that may very well be common to most proteins that are abundant in His residues, and not specific for SOD1. Another general observation is that the Zn^{2+} -binding appears to proceed via one or several intermediate states. This is evidenced by the non-linear change in both H^N and N^H chemical shifts with increasing concentration. This has recently been explored in detail in a study of the ultrafast folding of the minipro-

tein Trp-cage [32]. Our data, however, point toward a more distinct change in the chemical shift pattern. The data indicates an onset of a different Zn^{2+} interaction at very high ion concentrations rather than an exchange between Zn^{2+} -free unstructured SOD1, fully Zn^{2+} -bound, and one or several intermediate states.

Zn^{2+} seems to coordinate, to some extent, to all the His residues in the unfolded protein, including the two non-native metal ligands H110 and H43. Our data reveal that there is no selectivity for any of the two native metal centers (Fig. 6). We can also conclude that only His residues, and no other common ligands such as negatively charged amino acid residues, bind the cationic Zn^{2+} ions in urea-denatured SOD1. Even D83, the only non-His metal binding ligand in the native protein, is largely unaffected by Zn^{2+} , and no large shifts are observed (Fig. 6). This finding indicates a His-specific binding and not only an ionic effect. H120, one of the copper ligands, is affected to the greatest extent by the presence of zinc ions. We also note that binding to the non-native copper ligand H43 seems to be preferred over H46. This implies that it may not be favorable for Zn^{2+} to bind to both H46 and H48 simultaneously in unfolded SOD1, since these residues may be situated too close in the sequence. Therefore it appears that in the unfolded




Fig. 4. The weighted average difference in H^N and N^H chemical shifts when lowering the urea concentration from 9 to 4 M. The region H63–H71 (yellow), part of the zinc-binding loop (pink), is most affected. Loop VII is marked in blue. The positions of the secondary structure elements as seen in the X-ray structure of monomeric SOD1 (accession code 3HFF, www.pdb.org) are shown under the sequence.

protein, zinc coordination to H43 and H48 is more favorable than to H46 and H48, i.e. in an extended structure this relaxes the binding geometry of the protein [33,34]. Together these results demonstrate that the initial binding of zinc to the protein occurs via a diffuse coordination, which in turn suggests that the ligands have to be rearranged during the course of the folding process to reach the native metal-bound state. This is in agreement with a study of SOD1 in GdmCl, which demonstrated structural rearrangements of the both native metal-binding sites [27]. The study in GdmCl further showed that the presence of metals induced a somewhat more compact protein, indicating (transient) metal interactions.

A recent study demonstrated that Zn^{2+} appears to be coordinated to the copper site in the protein in the early folding process, and that the zinc site forms late in the folding process [7]. Contrary to these results, it has been argued that the Zn^{2+} site is preformed in the presence of Zn^{2+} , and that this stabilizes the folding nucleus [35]. Here we see that the diffuse binding of zinc, where the metal is coordinated to residues in both metal-binding sites, provides a means for SOD1 to explore several structural states at the very start of the folding reaction. Clearly, early zinc binding does not induce a pre-organized metal-binding site. This is consistent with the result that the zinc-binding site forms late in the folding process.

The region known as the zinc-binding loop in the folded protein (loop IV) appears to be the one that undergoes the most significant changes when slowly lowering the urea concentration. Small, but significant changes in chemical shifts for e.g. residues S68 and R69 in the zinc-binding loop are observed (Fig. 6), indicating that this region changes its structural characteristics. This implies that some sort of rearrangement of the loop takes place in the early stages of folding. NMR relaxation data for the apo protein has shown that loop IV is highly dynamic [36], and it has been demonstrated that the structural integrity of the protein relies on metal coordination [5]. The present finding indicating that the zinc-binding loop undergoes dynamic rearrangements, together with the unspecific metal-binding in the unfolded protein provides an indication that transient structural rearrangements in the zinc-binding loop may be important in the folding of human SOD1. For the protein to fold, several structural states are explored, which may provide a means for the protein to initiate folding. Finally, most proteins that contain His residues will most likely coordinate to

Zn^{2+} in a diffuse and non-specific way, providing a means for metal-bound proteins to explore several conformational states. The diffuse metal binding in the early folding pathway observed here may well be a general mechanism by how metalloproteins initiate the folding process.

4. Materials and methods

4.1. Materials

EDTA was purchased from Merck (Whitehouse station, U.S.A.), guanidine hydrochloride (ultrapure) from AppliChem (Darmstadt, Germany) and bis-tris hydrochloride from Sigma (St. Louis, U.S.A.). Urea (ultra pure) was bought from MP Biomedicals (Illkirch, France). ^{15}N - NH_4Cl , ^{13}C -glucose and D_2O was purchased from Cambridge Isotope Laboratories (Andover, MA, USA). $ZnSO_4$ was from Fluka (Buchs, Switzerland). Spectra/Por dialysis membranes were from Spectrum Laboratories (Rancho Dominguez, U.S.A.)

4.2. Expression and purification

The construct used, SOD1^{6/111/50/51}, was over-expressed in *Escherichia coli* grown in M9 minimal medium, containing ^{15}N -labeled ammonium chloride and/or ^{13}C -labeled D -glucose as sole nitrogen and carbon sources. The expression and purification of SOD1 were conducted as described in [37], but with no $CuSO_4$ addition upon induction.

4.3. Metal depletion and unfolding of SOD1

Apo-SOD1 was prepared by dialysis against 500 mL 50 mM EDTA and 4 M guanidine hydrochloride, buffered by 10 mM bis-Tris (pH 6.3). The dialysis membranes had a pore size of 6–8000 MWCO and the dialysis continued in this solution for at least four hours in room temperature with stirring. The tubing and its content was then moved into a solution containing 250 mL 10 M urea and 10 mM bis-Tris (pH 6.3) and dialyzed again for 2×2 h in room temperature. This was done to keep the protein completely unfolded, remove EDTA and reduce the salt content in the sample. Addition of 10% D_2O (for the field frequency lock stabilization in




Fig. 6. Zn²⁺-induced changes in SOD1^{G111/S50/S51} chemical shifts. Correlation between the chemical shift changes in SOD1 due to the addition of 3 mM Zn²⁺–260 μM SOD1 for H^N and N^H (A). The Zn²⁺:SOD1 ratios were obtained by taking into account the (small) differences in sample volumes as a consequence of adding Zn²⁺. The Zn²⁺-induced weighted average difference in ¹H^N and ¹⁵N chemical shifts as a function of amino acid sequence (B).

Acknowledgements

This work was supported by grants from the Swedish Research Council (grant number 621-2011-5964). We thank Dr. Helena Kovacs for assistance with some of the NMR experiments. SS, MO and LM conceived and designed the project, SS acquired and analyzed the data, SS MO and LM interpreted the data, and SS and LM wrote the paper.

References

- [1] Ferraroni, M., Rypniewski, W., Wilson, K., Viezzoli, M., Banci, L., Bertini, I. and Mangani, S. (1999) The crystal structure of the monomeric human SOD mutant F50E/G51E/E133Q at atomic resolution. The enzyme mechanism revisited. *J. Mol. Biol.* 288, 413–426.
- [2] Lynch, S.M., Boswell, S.A. and Colón, W. (2004) Kinetic stability of Cu/Zn superoxide dismutase is dependent on its metal ligands: implications for ALS. *Biochemistry* 43, 16525–16531.
- [3] Roberts, B.R., Tainer, J.A., Getzoff, E.D., Malencik, D.A., Anderson, S.R., Bomben, V.C., Meyers, K.R., Karplus, A. and Beckman, J.S. (2007) Structural characterization of zinc-deficient human superoxide dismutase and implications for ALS. *J. Mol. Biol.* 373, 877–890.
- [4] Mulligan, V.K., Kerman, A., Ho, S. and Chakrabarty, A. (2008) Denaturational stress induces formation of zinc-deficient monomers of Cu, Zn superoxide dismutase: implications for pathogenesis in amyotrophic lateral sclerosis. *J. Mol. Biol.* 383, 424–436.
- [5] Nordlund, A., Leinartaite, L., Saraboji, K., Aisenbrey, C., Gröbner, G., Zetterström, P., Danielsson, J., Logan, D.T. and Oliveberg, M. (2009) Functional features cause misfolding of the ALS-provoking enzyme SOD1. *Proc. Natl. Acad. Sci. U.S.A.* 106, 9667–9672.
- [6] Estévez, A.G., Crow, J.P., Sampson, J.B., Reiter, C., Zhuang, Y., Richardson, G.J., Tarpey, M.M., Barbeito, L. and Beckman, J.S. (1999) Induction of nitric oxide-dependent apoptosis in motor neurons by zinc-deficient superoxide dismutase. *Science* 286, 2498–2500.
- [7] Leinartaite, L., Saraboji, K., Nordlund, A., Logan, D.T. and Oliveberg, M. (2010) Folding catalysis by transient coordination of Zn²⁺ to the Cu ligands of the ALS-associated enzyme Cu/Zn superoxide dismutase 1. *J. Am. Chem. Soc.* 132, 13495–13504.
- [8] Bertini, I., Piccioli, M., Viezzoli, M.S., Chiu, C.Y. and Mullenbach, G.T. (1994) A spectroscopic characterization of a monomeric analog of copper, zinc superoxide dismutase. *Eur. Biophys. J.* 23, 167–176.
- [9] Lepock, J., Frey, H.E. and Hallewell, R. (1990) Contribution of conformational stability and reversibility of unfolding to the increased thermostability of human and bovine superoxide dismutase mutated at free cysteines. *J. Biol. Chem.* 265, 21612–21618.
- [10] Wishart, D.S., Bigam, C.G., Holm, A., Hodges, R.S. and Sykes, B.D. (1995) ¹H, ¹³C and ¹⁵N random coil NMR chemical shifts of the common amino acids. I. Investigations of nearest-neighbor effects. *J. Biomol. NMR* 5, 67–81.
- [11] Wishart, D.S. and Sykes, B.D. (1994) The ¹³C chemical-shift index: a simple method for the identification of protein secondary structure using ¹³C chemical-shift data. *J. Biomol. NMR* 4, 171–180.

- [12] Schwarzwinger, S., Kroon, G.J.A., Foss, T.R., Chung, J., Wright, P.E. and Dyson, H.J. (2001) Sequence-dependent correction of random coil NMR chemical shifts. *J. Am. Chem. Soc.* 123, 2970–2978.
- [13] Schwarzwinger, S., Kroon, G.J.A., Foss, T.R., Wright, P.E. and Dyson, H.J. (2000) Random coil chemical shifts in acidic 8 M urea: implementation of random coil shift data in NMR view. *J. Biomol. NMR* 18, 43–48.
- [14] Teilum, K., Smith, M.H., Schulz, E., Christensen, L.C., Solomentsev, G., Oliveberg, M. and Akke, M. (2009) Transient structural distortion of metal-free Cu/Zn superoxide dismutase triggers aberrant oligomerization. *Proc. Natl. Acad. Sci. U.S.A.* 106, 18273–18278.
- [15] Tompa, P. (2002) Intrinsically unstructured proteins. *Trends Biochem. Sci.* 27, 527–533.
- [16] Dyson, H.J. and Wright, P.E. (2005) Intrinsically unstructured proteins and their functions. *Nat. Rev. Mol. Cell Biol.* 6, 197–208.
- [17] Tompa, P. (2011) Unstructural biology coming of age. *Curr. Opin. Struct. Biol.* 21, 419–425.
- [18] Uversky, V.N. (2011) Intrinsically disordered proteins from A to Z. *Int. J. Biochem. Cell Biol.* 43, 1090–1103.
- [19] Kosol, S., Contreras-Martos, S., Cedeno, C. and Tompa, P. (2013) Structural characterization of intrinsically disordered proteins by NMR spectroscopy. *Molecules* 18, 10802–10828.
- [20] Marsh, J.A., Singh, V.K., Jia, Z. and Forman-Kay, J.D. (2006) Sensitivity of secondary structure propensities to sequence differences between α - and γ -synuclein: Implications for fibrillation. *Protein Sci.* 15, 2795–2804.
- [21] Camilloni, C., De Simone, A., Vranken, W.F. and Vendruscolo, M. (2012) Determination of secondary structure populations in disordered states of proteins using nuclear magnetic resonance chemical shifts. *Biochemistry* 51, 2224–2231.
- [22] Neri, D., Billeter, M., Wider, G. and Wüthrich, K. (1992) NMR determination of residual structure in a urea-denatured protein, the 434-repressor. *Science* 257, 1559–1563.
- [23] McCarney, E.R., Kohn, J.E. and Plaxco, K.W. (2005) Is there or is not there? The case for (and against) residual structure in chemically denatured proteins. *Crit. Rev. Biochem. Mol. Biol.* 40, 181–189.
- [24] Bhavesh, N.S., Juneja, J., Udgaonkar, J.B. and Hosur, R.V. (2004) Native and nonnative conformational preferences in the urea-unfolded state of barstar. *Protein Sci.* 13, 3085–3091.
- [25] Wong, K., Freund, S. and Fersht, A.R. (1996) Cold denaturation of barstar: ^1H , ^{15}N and ^{13}C NMR assignment and characterisation of residual structure. *J. Mol. Biol.* 259, 805–818.
- [26] Hsieh, S.M., Mallam, A.L., Jackson, S.E. and Hsu, S.D. (2013) Backbone NMR assignments of a topologically knotted protein in urea-denatured state. *Biomol. NMR Assign.* 8, 283–285.
- [27] Assfalg, M., Banci, L., Bertini, I., Turano, P. and Vasos, P.R. (2003) Superoxide dismutase folding/unfolding pathway: role of the metal ions in modulating structural and dynamical features. *J. Mol. Biol.* 330, 145–158.
- [28] Pozdnyakova, I. and Wittung-Stafshede, P. (2001) Copper binding before polypeptide folding speeds up formation of active (holo) *Pseudomonas aeruginosa* azurin. *Biochemistry* 40, 13728–13733.
- [29] Pozdnyakova, I. and Wittung-Stafshede, P. (2001) Biological relevance of metal binding before protein folding. *J. Am. Chem. Soc.* 123, 10135–10136.
- [30] Pozdnyakova, I. and Wittung-Stafshede, P. (2003) Approaching the speed limit for Greek Key β -barrel formation: transition-state movement tunes folding rate of zinc-substituted azurin. *Biochim. Biophys. Acta* 1651, 1–4.
- [31] Ozutsumi, K., Taguchi, Y. and Kawashima, T. (1995) Thermodynamics of formation of urea complexes with manganese (II), nickel (II) and zinc (II) ions in *N,N*-dimethylformamide. *Talanta* 42, 535–541.
- [32] Rovó, P., Stráner, P., Láng, A., Bartha, I., Huszár, K., Nyitray, L. and Perczel, A. (2013) Structural insights into the Trp-cage folding intermediate formation. *Chem. Eur. J.* 19, 2628–2640.
- [33] Krantz, B.A. and Sosnick, T.R. (2001) Engineered metal binding sites map the heterogeneous folding landscape of a coiled coil. *Nat. Struct. Mol. Biol.* 8, 1042–1047.
- [34] Bosco, G.L., Baxa, M. and Sosnick, T.R. (2009) Metal binding kinetics of bi-histidine sites used in ψ analysis: evidence of high-energy protein folding intermediates. *Biochemistry* 48, 2950–2959.
- [35] Kayatekin, C., Zitzewitz, J.A. and Matthews, C.R. (2008) Zinc binding modulates the entire folding free energy surface of human Cu, Zn superoxide dismutase. *J. Mol. Biol.* 384, 540–555.
- [36] Banci, L., Bertini, I., Cramaro, F., Del Conte, R. and Viezzoli, M.S. (2003) Solution structure of Apo Cu, Zn superoxide dismutase: role of metal ions in protein folding. *Biochemistry* 42, 9543–9553.
- [37] Lindberg, M.J., Tibell, L. and Oliveberg, M. (2002) Common denominator of Cu/Zn superoxide dismutase mutants associated with amyotrophic lateral sclerosis: decreased stability of the apo state. *Proc. Natl. Acad. Sci. U.S.A.* 99, 16607–16612.
- [38] Goddard, T. and Kneller, D. (2004) SPARKY 3, University of California, San Francisco.
- [39] Zhang, H., Neal, S. and Wishart, D.S. (2003) RefDB: a database of uniformly referenced protein chemical shifts. *J. Biomol. NMR* 25, 173–195.
- [40] Tamiola, K., Acar, B. and Mulder, F.A.A. (2010) Sequence-specific random coil chemical shifts of intrinsically disordered proteins. *J. Am. Chem. Soc.* 132, 18000–18003.
- [41] Grzesiek, S., Bax, A., Clore, G.M., Gronenborn, A.M., Hu, J.S., Kaufman, J., Palmer, I., Stahl, S.J. and Wingfield, P.T. (1996) The solution structure of HIV-1 Nef reveals an unexpected fold and permits delineation of the binding surface for the SH3 domain of Hck tyrosine protein kinase. *Nat. Struct. Biol.* 3, 340–345.
- [42] Garrett, D.S., Seok, Y.J., Peterkofsky, A., Clore, G.M. and Gronenborn, A.M. (1997) Identification by NMR of the binding surface for the histidine-containing phosphocarrier protein HPr on the *N*-terminal domain of enzyme I of the *Escherichia coli* phosphotransferase system. *Biochemistry* 36, 4393–4398.