



Bull. Sci. math. 134 (2010) 12–36

 BULLETIN DES
 SCIENCES
 MATHÉMATIQUES

www.elsevier.com/locate/bulsci

Sobolev embeddings for Riesz potential spaces of variable exponents near 1 and Sobolev's exponent

Yoshihiro Mizuta ^a, Takao Ohno ^{b,*}, Tetsu Shimomura ^c

^a *Department of Mathematics, Faculty of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima 739-8521, Japan*

^b *General Arts, Hiroshima National College of Maritime Technology, Higashino Oosakikamijima Toyotagun 725-0231, Japan*

^c *Department of Mathematics, Graduate School of Education, Hiroshima University, Higashi-Hiroshima 739-8524, Japan*

Received 1 April 2009

Available online 11 September 2009

Abstract

Our aim in this paper is to deal with Sobolev embeddings for Riesz potentials of order α for functions f satisfying the Orlicz type condition

$$\int |f(y)|^{p(y)} (\log(c + |f(y)|))^{q(y)} dy < \infty,$$

where $p(\cdot)$ and $q(\cdot)$ are variable exponents satisfying the log-Hölder conditions.

© 2009 Elsevier Masson SAS. All rights reserved.

MSC: 46E30

Keywords: Sobolev embeddings; Riesz potential; Variable exponent

1. Introduction

The Sobolev space is a useful tool of the study for the existence and regularity of solutions of partial differential equations. The famous Sobolev's inequality says that the Riesz potential $I_\alpha f$

* Corresponding author.

E-mail addresses: yomizuta@hiroshima-u.ac.jp (Y. Mizuta), ohno@hiroshima-cmt.ac.jp (T. Ohno), tshimo@hiroshima-u.ac.jp (T. Shimomura).

of order α with $f \in L^p(\mathbf{R}^n)$ belongs to $L^{p^*}(\mathbf{R}^n)$ when $1 < p < \infty$ and $1/p^* = 1/p - \alpha/n > 0$. If f satisfies the Orlicz condition

$$\int |f(y)|^p (\log(e + |f(y)|))^q dy < \infty,$$

then it is known (see e.g. Cianchi [1]) that

$$\int (|I_\alpha f(x)| (\log(e + |f(y)|))^{q/p})^{p^*} dx < \infty.$$

When $p = 1$, the situation is a little different (see O’Neil [13]).

In the present paper, we aim to establish Sobolev’s inequality for Riesz potentials of functions in the Orlicz spaces of variable exponent. In recent years, the generalized Lebesgue spaces have attracted more and more attention, in connection with the study of elasticity, fluid mechanics and differential equations with $p(\cdot)$ -growth; see for example Orlicz [14], Kováčik–Rákosník [9], Edmunds–Rákosník [4] and Růžička [15]. In the limiting case we are also concerned with exponential integrabilities of Trudinger type and continuity for Riesz potentials.

For $0 < \alpha < n$, we define the Riesz potential of order α for a locally integrable function f on \mathbf{R}^n by

$$I_\alpha f(x) = \int_{\mathbf{R}^n} |x - y|^{\alpha-n} f(y) dy.$$

Here it is natural to assume that

$$\int_{\mathbf{R}^n} (1 + |y|)^{\alpha-n} |f(y)| dy < \infty$$

(see [10, Theorem 1.1, Chapter 2]).

In this paper, following Cruz-Uribe and Fiorenza [2], we consider variable exponents $p(\cdot)$ and $q(\cdot)$ such that

$$|p(x) - p(y)| \leq \frac{a}{\log(e + 1/|x - y|)} \quad \text{for all } x, y \in \mathbf{R}^n, \tag{1.1}$$

$$|q(x) - q(y)| \leq \frac{b}{\log(e + \log(e + 1/|x - y|))} \quad \text{for all } x, y \in \mathbf{R}^n, \tag{1.2}$$

$$1 \leq p^- \equiv \inf_{x \in \mathbf{R}^n} p(x) \leq \sup_{x \in \mathbf{R}^n} p(x) \equiv p^+ < \infty \tag{1.3}$$

and

$$-\infty < q^- \equiv \inf_{x \in \mathbf{R}^n} q(x) \leq \sup_{x \in \mathbf{R}^n} q(x) \equiv q^+ < \infty \tag{1.4}$$

for $a, b > 0$. Moreover, suppose there exist $\varepsilon_0 > 0$ and $C > 0$ such that

$$s^{p(x)-1} (\log(e + s))^{q(x)-\varepsilon_0} \leq Ct^{p(x)-1} (\log(e + t))^{q(x)-\varepsilon_0} \tag{1.5}$$

whenever $0 < s < t$ and $x \in \mathbf{R}^n$. This is true if

$$K(p(x) - 1) + q(x) \geq \varepsilon_0$$

for some positive constant K .

Let G be a bounded Borel set in \mathbf{R}^n . Let $\Phi(x, t)$ be a nonnegative function on $G \times \mathbf{R}$ such that

- (1) $\Phi(\cdot, t)$ is measurable for each t .
- (2) $\Phi(x, \cdot)$ is continuous and convex for each fixed $x \in G$.

Define the norm by

$$\|f\|_{\Phi(\cdot, \cdot)(G)} = \inf \left\{ \lambda > 0: \int_G \Phi(x, |f(x)/\lambda|) dx \leq 1 \right\}$$

and denote by $\Phi(\cdot, \cdot)(G)$ the space of all measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} < \infty$.

If $p(x) < n/\alpha$, then we set

$$1/p^*(x) = 1/p(x) - \alpha/n.$$

Define

$$\begin{aligned} \Phi(x, t) &= t^{p(x)} (\log(c + t))^{q(x)}, \\ \Psi(x, t) &= \{t (\log(c + t))^{q(x)/p(x)}\}^{p^*(x)} \end{aligned}$$

and

$$\tilde{\Psi}(x, t) = \{t (\log(c + t))^{q(x)/p(x) - p(x)/p^*(x)}\}^{p^*(x)},$$

where $c \geq e$ is chosen so that $\Phi(x, \cdot)$, $\Psi(x, \cdot)$, $\tilde{\Psi}(x, \cdot)$ are all convex on $[0, \infty)$.

For $0 < \delta < n/\alpha$, divide G into four sets:

$$\begin{aligned} G_1 &= \{x \in G: 1 \leq p(x) < 1 + \delta\}, \\ G_2 &= \{x \in G: 1 + \delta \leq p(x) < n/\alpha\}, \\ G_3 &= \{x \in G: p(x) \geq n/\alpha \text{ and } q(x) < p(x) - 1\}, \\ G_4 &= \{x \in G: p(x) \geq n/\alpha \text{ and } q(x) \geq p(x) - 1\}. \end{aligned}$$

Denote by χ_E the characteristic function of a measurable set E .

Our main result is the following, which is an extension of Futamura–Mizuta [5], Futamura–Mizuta–Shimomura [6] and Harjulehto–Hästö [7].

Theorem 1.1. *Let $p(\cdot)$ and $q(\cdot)$ be as above. Then there exist constants $c_1, c_2, c_3, c_4 > 0$ such that*

$$\begin{aligned} &\int_G \left\{ \tilde{\Psi}(x, I_\alpha f_1(x)) \chi_{G_1}(x) + \Psi(x, c_1^{-1} \gamma_1(x)^{-1} I_\alpha f_2(x)) \chi_{G_2}(x) \right. \\ &\quad + \exp\left(\frac{I_\alpha f_3(x)^{p(x)/(p(x)-q(x)-1)}}{(c_2 \gamma_3(x))^{p(x)/(p(x)-q(x)-1)}}\right) \chi_{G_3}(x) \\ &\quad \left. + \exp\left(\exp\left(\frac{I_\alpha f_4(x)^{p(x)/(p(x)-1)}}{c_3^{p(x)/(p(x)-1)}}\right)\right) \chi_{G_4}(x) \right\} dx \leq c_4 \end{aligned}$$

for all nonnegative measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$, where $f_j = f \chi_{G_j}$ ($j = 1, 2, 3, 4$),

$$\gamma_1(x) = p^*(x)^{(q(x)+p(x)-1)/p(x)} (\log p^*(x))^{q(x)/p(x)}$$

and

$$\gamma_3(x) = \gamma_2(x)^{-(p(x)-1)/p(x)} (\log(1/\gamma_2(x)))^{q(x)/p(x)}$$

with $\gamma_2(x) = \min\{p(x) - q(x) - 1, 1/2\}$.

The proof is given by discussing the cases (i) $1 \leq p(x) \leq p^+ < n/\alpha$ (Section 2), (ii) $1 < p^- \leq p(x) < n/\alpha$ (Section 3), (iii) $p^- \geq n/\alpha, q(x) < p(x) - 1$ (Section 4) and (iv) $p^- \geq n/\alpha, q(x) \geq p(x) - 1$ (Section 5), separately.

In their paper [7], Harjulehto–Hästö gave an integrability result of Sobolev functions by diving the domain of integration into countably many measurable sets.

With the aid of O’Neil [13], one sees that if $f \in L^1(G)$, then

$$\int_G |I_\alpha f(x)|^{p^*} (\log(e + |I_\alpha f(x)|))^{-\beta} dx < \infty$$

when $\beta > 1$; this is not true when $\beta = 1$. Theorem 1.1 extends his result to the valuable exponent case. In case $p > 1$, the maximal function is a crucial tool by Hedberg’s trick (see Hedberg [8]). In case $p^- = 1$, our strategy is to give an estimate of $I_\alpha f$ by use of a logarithmic type potential

$$\int_{\{y \in G: |x-y|^{-\varepsilon} < f(y)\}} (\log(c + |x - y|^{-1}))^{\varepsilon-1} |x - y|^{-n} (\log(c + f(y)))^{-\varepsilon} g(y) dy,$$

which plays a role of maximal functions, where $g(y) = \Phi(y, f(y))$. Thus our proof is quite different from that of O’Neil [13].

In Section 7, we are concerned with continuity of Riesz potentials when $p(x) \geq n/\alpha$ and $q(x) > p(x) - 1$ for $x \in \mathbf{R}^n$.

We define the logarithmic potential for a locally integrable function f on \mathbf{R}^n by

$$I_n f(x) = \int_{\mathbf{R}^n} (\log^+(1/|x - y|)) f(y) dy,$$

where $\log^+ r = \max\{0, \log r\}$. Here it is natural to assume that

$$\int_{\mathbf{R}^n} (\log(e + |y|)) |f(y)| dy < \infty$$

(see [10, Theorem 1.1, Chapter 2]).

Finally, in Section 8, we show the exponential integrability of logarithmic potentials $I_n f$.

2. Variable exponents near 1

Throughout this paper, let C denote various positive constants independent of the variables in question and $C(a, b, \dots)$ be a constant that depends on a, b, \dots .

Let us begin with Sobolev’s inequality for Riesz potentials of functions in $\Phi(\cdot, \cdot)(G)$ when $1 \leq p(x) \leq p^+ < n/\alpha$, which gives an extension of O’Neil [13].

Theorem 2.1. *Let $p(\cdot)$ and $q(\cdot)$ be as in the Introduction. Suppose*

$$1 \leq p(x) \leq p^+ < n/\alpha$$

for $x \in \mathbf{R}^n$. Then there exists a constant $c_1 > 0$ such that

$$\|I_\alpha f\|_{\tilde{\Psi}(\cdot, \cdot)(G)} \leq c_1 \|f\|_{\Phi(\cdot, \cdot)(G)}$$

for all $f \in \Phi(\cdot, \cdot)(G)$.

When $p(x) = p = 1$ and $q(x) = \varepsilon_0 > 0$ for $x \in \mathbf{R}^n$, Theorem 2.1 says that

$$\left(\int_G |I_\alpha f(x)|^{p^*} (\log(e + |I_\alpha f(x)|))^{\varepsilon_0 p^* - 1} dx \right)^{1/p^*} \leq C$$

for all measurable functions f satisfying

$$\int_G |f(y)| (\log(e + |f(y)|))^{\varepsilon_0} dy \leq 1,$$

which is a consequence of O’Neil [13].

The case $p^- > 1$ is treated in the next section, by use of maximal functions. However the maximal operator fails to be bounded in $L^{p(\cdot)}(G)$ when $p^- = 1$. To show Theorem 2.1, we introduce the logarithmic type potential

$$J \equiv \int_{\{y \in G: |x-y|^{-\varepsilon} < f(y)\}} \rho_{\varepsilon-1}(|x-y|) (\log(c + f(y)))^{-\varepsilon} g(y) dy,$$

which plays a role of maximal functions; here $g(y) = \Phi(y, |f(y)|)$ and $\rho_{\varepsilon-1}(r) = r^{-n} (\log(c + 1/r))^{\varepsilon-1}$ for $0 < \varepsilon < \varepsilon_0/2$ with ε_0 in (1.5).

We use the notation $B(x, r)$ to denote the open ball centered at $x \in \mathbf{R}^n$ of radius $r > 0$.

Lemma 2.2. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 2.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. If*

$$\delta = J^{-1/n} (\log(c + J))^{-p(x)/n},$$

then

$$\begin{aligned} I &\equiv \int_{\{y \in G \cap B(x, \delta): |x-y|^{-\varepsilon} < f(y)\}} |x-y|^{\alpha-n} (\log(c + 1/|x-y|))^{-\varepsilon} (\log(c + f(y)))^\varepsilon f(y) dy \\ &\leq C \{ J^{1/p^*(x)} (\log(c + J))^{p(x)/p^*(x) - q(x)/p(x)} + 1 \}. \end{aligned}$$

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. First consider the case when $J \geq 1$. We have by (1.5) for $k > 0$

$$\begin{aligned} I &\leq k (\log(c + k))^\varepsilon \int_{\{y \in G \cap B(x, \delta): |x-y|^{-\varepsilon} < f(y)\}} |x-y|^{\alpha-n} (\log(c + 1/|x-y|))^{-\varepsilon} dy \\ &\quad + \int_{\{y \in B(x, \delta): |x-y|^{-\varepsilon} < f(y)\}} |x-y|^{\alpha-n} (\log(c + 1/|x-y|))^{-\varepsilon} (\log(c + f(y)))^\varepsilon f(y) \\ &\quad \times C \left(\frac{f(y)}{k} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + k)} \right)^{q(y)-2\varepsilon} dy \end{aligned}$$

$$\begin{aligned} &\leq C \left\{ k(\log(c+k))^\varepsilon \delta^\alpha (\log(c+1/\delta))^{-\varepsilon} \right. \\ &\quad \left. + \delta^\alpha (\log(c+1/\delta))^{1-2\varepsilon} \int_{\{y \in B(x,\delta): |x-y|^{-\varepsilon} < f(y)\}} \rho_{\varepsilon-1}(|x-y|) (\log(c+f(y)))^{-\varepsilon} g(y) \right. \\ &\quad \left. \times \left(\frac{1}{k}\right)^{p(y)-1} \left(\frac{1}{\log(c+k)}\right)^{q(y)-2\varepsilon} dy \right\}. \end{aligned}$$

Set

$$k = J^{1/p(x)} (\log(c+J))^{-q(x)/p(x)}.$$

Since

$$\frac{\log J}{\log(1/\delta)} = \frac{\log J}{\log(J^{1/n}(\log(c+J))^{p(x)/n})} \leq C,$$

we see that if $y \in B(x, \delta)$, then

$$J^{-p(y)} = J^{-p(y)+p(x)} J^{-p(x)} \leq J^{a/\log(1/\delta)} J^{-p(x)} \leq C J^{-p(x)}$$

and similarly

$$(\log(c+J))^{p(y)} \leq (\log(c+J))^{a/\log(1/\delta)} (\log(c+J))^{p(x)} \leq C (\log(c+J))^{p(x)},$$

so that

$$k^{-p(y)} \leq C J^{-1} (\log(c+J))^{q(x)}$$

and

$$(\log(c+k))^{-q(y)} \leq C (\log(c+J))^{-q(x)}.$$

Consequently it follows that

$$\begin{aligned} I &\leq C \left\{ J^{1/p(x)} (\log(c+J))^{-q(x)/p(x)} (\log(c+J))^\varepsilon \delta^\alpha (\log(c+1/\delta))^{-\varepsilon} \right. \\ &\quad \left. + \delta^\alpha (\log(c+1/\delta))^{1-2\varepsilon} J^{1/p(x)} (\log(c+J))^{-q(x)/p(x)+2\varepsilon} \right\} \\ &\leq C J^{1/p(x)-\alpha/n} (\log(c+J))^{-q(x)/p(x)-\alpha p(x)/n+1}. \end{aligned}$$

In the case $J \leq 1$, we set $k = 1$. The above considerations gives $I \leq C$. Now the result follows. \square

Lemma 2.3. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 2.1. Suppose $p^+ < n/\alpha$. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{G \setminus B(x,\delta)} |x-y|^{\alpha-n} f(y) dy \leq C \delta^{\alpha-n/p(x)} (\log(c+1/\delta))^{-q(x)/p(x)}$$

for all $x \in G$ and $\delta > 0$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. For $y \in G \setminus B(x, \delta)$, set

$$N(x, y) = |x-y|^{-n/p(x)} (\log(c+|x-y|^{-1}))^{-q(x)/p(x)}.$$

By conditions (1.1), (1.2) and (1.5), we see that

$$\begin{aligned}
 & \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\
 & \leq \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} N(x, y) dy \\
 & \quad + C \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) \left(\frac{f(y)}{N(x, y)} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + N(x, y))} \right)^{q(y)} dy \\
 & \leq C \left\{ \delta^{\alpha-n/p(x)} (\log(c + 1/\delta))^{-q(x)/p(x)} \right. \\
 & \quad \left. + \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n/p(x)} (\log(c + 1/|x - y|))^{-q(x)/p(x)} g(y) dy \right\} \\
 & \leq C \delta^{\alpha-n/p(x)} (\log(c + 1/\delta))^{-q(x)/p(x)} \left(1 + \int_{G \setminus B(x, \delta)} g(y) dy \right) \\
 & \leq C \delta^{\alpha-n/p(x)} (\log(c + 1/\delta))^{-q(x)/p(x)},
 \end{aligned}$$

where $g(y) = f(y)^{p(y)} (\log(c + f(y)))^{q(y)}$, as required. \square

We denote by $|E|$ the volume of E .

Proof of Theorem 2.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. For

$$\delta = J^{-1/n} (\log(c + J))^{-p(x)/n},$$

write

$$\begin{aligned}
 I_\alpha f(x) &= \int_{G \cap B(x, \delta)} |x - y|^{\alpha-n} f(y) dy + \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\
 &= I_1 + I_2.
 \end{aligned}$$

For $0 < \varepsilon < \min\{\varepsilon_0/2, \alpha\}$, set

$$J = \int_{\{y \in G: |x-y|^{-\varepsilon} < f(y)\}} \rho_{\varepsilon-1}(|x - y|) (\log(c + f(y)))^{-\varepsilon} g(y) dy.$$

In view of Lemma 2.2, we find

$$\begin{aligned}
 I_1 &\leq \int_{G \cap B(x, \delta)} |x - y|^{\alpha-n-\varepsilon} dy \\
 &\quad + \int_{\{y \in B(x, \delta): |x-y|^{-\varepsilon} < f(y)\}} |x - y|^{\alpha-n} \frac{(\log(c + f(y)))^\varepsilon}{(\log(c + |x - y|^{-\varepsilon}))^\varepsilon} f(y) dy \\
 &\leq C \left\{ 1 + J^{1/p^*(x)} (\log(c + J))^{p(x)/p^*(x)-q(x)/p(x)} \right\}.
 \end{aligned}$$

Moreover, Lemma 2.3 yields

$$I_2 \leq C \delta^{\alpha-n/p(x)} (\log(c + 1/\delta))^{-q(x)/p(x)},$$

so that

$$\begin{aligned} I_\alpha f(x) &\leq C \left\{ 1 + J^{1/p^*(x)} (\log(c + J))^{p(x)/p^*(x)-q(x)/p(x)} \right. \\ &\quad \left. + \delta^{\alpha-n/p(x)} (\log(c + 1/\delta))^{-q(x)/p(x)} \right\} \\ &\leq C \left\{ 1 + J^{1/p^*(x)} (\log(c + J))^{p(x)/p^*(x)-q(x)/p(x)} \right\}. \end{aligned}$$

Hence we have

$$\int_G \tilde{\Psi}(x, I_\alpha f(x)) dx \leq C \int_G (1 + J) dx.$$

By using Fubini’s theorem, we obtain

$$\begin{aligned} &\int_G \tilde{\Psi}(x, I_\alpha f(x)) dx \\ &\leq C \left\{ \int_G \left(\int_{\{y \in G: |x-y|^{-\varepsilon} < f(y)\}} \rho_{\varepsilon-1}(|x-y|) dx \right) (\log(c + f(y)))^{-\varepsilon} g(y) dy + |G| \right\} \\ &\leq C \left\{ \int_G g(y) dy + |G| \right\} \leq C, \end{aligned}$$

which completes the proof. \square

Remark 2.4. Let $p(x) = p = 1$ and

$$q(x) = \frac{1}{\log(\log(e + |x|^{-1}))}$$

for $x \in \mathbf{B} = B(0, 1)$. Here, note that $q(\cdot)$ satisfies the condition (1.2) since

$$\begin{aligned} \frac{1}{\log(\log(1/t))} - \frac{1}{\log(\log(1/s))} &= \int_0^{t-s} (\log(\log(1/(r+s))))^{-2} (\log(1/(r+s)))^{-1} \frac{dr}{r+s} \\ &\leq C \int_0^{t-s} (\log(\log(1/r)))^{-2} (\log(1/r))^{-1} \frac{dr}{r} \\ &= \frac{C}{\log(\log(1/(t-s)))} \end{aligned}$$

whenever $0 < s < t < 1/e$. Set $p^* = n/(n - \alpha)$. Then one can find $f \in \Phi(\cdot, \cdot)(\mathbf{B})$ such that

$$\int_{\mathbf{B}} |I_\alpha f(x)|^{p^*} (\log(c + |I_\alpha f(x)|))^{p^*q(x)-1} dx = \infty.$$

To show this, for $0 < \gamma < 1/p^*$, let f be a nonnegative function on \mathbf{B} such that

$$f(y) = |y|^{-n} (\log(c + 1/|y|))^{-1} (\log(c + \log(c + 1/|y|)))^{-\gamma-1}.$$

Then we have

$$\int_{\mathbf{B}} f(y)(\log(c + f(y)))^{q(y)} dy < \infty$$

and for $x \in \mathbf{B}$

$$\begin{aligned} I_\alpha f(x) &\geq \int_{B(0, |x|/2)} |x - y|^{\alpha-n} f(y) dy \\ &\geq C|x|^{\alpha-n} \int_{B(0, |x|/2)} |y|^{-n} (\log(c + 1/|y|))^{-1} (\log(c + \log(c + 1/|y|)))^{-\gamma-1} dy \\ &\geq C|x|^{\alpha-n} (\log(c + \log(c + 1/|x|)))^{-\gamma}. \end{aligned}$$

Hence it follows that

$$\begin{aligned} \int_{\mathbf{B}} |I_\alpha f(x)|^{p^*} (\log(c + |I_\alpha f(x)|))^{p^*q(x)-1} dx \\ \geq C \int_{\mathbf{B}} |x|^{-n} (\log(c + 1/|x|))^{-1} (\log(c + \log(c + 1/|x|)))^{-\gamma p^*} dx = \infty \end{aligned}$$

when $\gamma < 1/p^*$.

3. Variable exponents near Sobolev’s exponent

Set

$$\Psi(x, t) = \{t(\log(c + t))^{q(x)/p(x)}\}^{p^*(x)}$$

and denote by $\Psi(\cdot, \cdot)(G)$ the family of all measurable functions f on G such that

$$\|f\|_{\Psi(\cdot, \cdot)(G)} = \inf \left\{ \lambda > 0: \int_G \Psi(x, |f(x)/\lambda|) dx \leq 1 \right\} < \infty.$$

Theorem 3.1. *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that*

$$1 < p^- \leq p(x) < n/\alpha$$

for $x \in G$. Then there exists a constant $c_1 > 0$ such that

$$\|\gamma_1(\cdot)^{-1} I_\alpha f\|_{\Psi(\cdot, \cdot)(G)} \leq c_1 \|f\|_{\Phi(\cdot, \cdot)(G)}$$

for all $f \in \Phi(\cdot, \cdot)(G)$, where

$$\gamma_1(x) = p^*(x)^{(q(x)+p(x)-1)/p(x)} (\log p^*(x))^{q(x)/p(x)}.$$

The case $\sup_{x \in G} p(x) < n/\alpha$ was shown in [11, Theorem 2.8].

Remark 3.2. For $0 < \delta < 1$, we can find $f \in \Phi(\cdot, \cdot)(G)$ such that

$$\int_G \Psi(x, \gamma_1(\cdot)^{-\delta} I_\alpha f(x)) dx = \infty$$

with $q(x) = 0$, so that the weight $\gamma_1(\cdot)^{-1}$ in Theorem 3.1 is needed.

For this, consider

$$p(x) = \frac{n}{\alpha} - \frac{1}{\log(1/|x|)}$$

for $x \in B_0 = B(0, 1/e)$ and

$$f(y) = |y|^{-\alpha} (\log(1/|y|))^{-\beta}$$

for $y \in B_0$. If $\alpha/n < \beta < 1$, then

$$\int_{B_0} f(y)^{p(y)} dy < \infty.$$

If $x \in B_0$, then

$$\begin{aligned} I_\alpha f(x) &\geq \int_{\{y \in B_0: |y| \geq |x|/2\}} |x - y|^{\alpha-n} f(y) dy \\ &\geq C \int_{\{y \in B_0: |y| \geq |x|/2\}} |y|^{-n} (\log(1/|y|))^{-\beta} dy \geq C (\log(1/|x|))^{-\beta+1} \end{aligned}$$

when $\beta < 1$. Now take β such that $\alpha/n < \beta < 1$ and

$$-\delta(1 - \alpha/n) - \beta + 1 > 0.$$

Since $p^*(x) = (n/\alpha)^2 \log(1/|x|) - n/\alpha$, there exists a constant $c_0 > 0$ such that $\gamma_1(x) \leq C (\log(1/|x|))^{1-\alpha/n}$ and $p^*(x) > c_0 \log(1/|x|)$ for $x \in B_0$. Hence we find

$$\begin{aligned} \int_{B_0} \Psi(x, \gamma_1(\cdot)^{-\delta} I_\alpha f(x)) dx &\geq C \int_{B_0} (\log(1/|x|))^{p^*(x)(-\delta(1-\alpha/n)-\beta+1)} dx \\ &\geq C \int_{B_0} (\log(1/|x|))^{c_0(-\delta(1-\alpha/n)-\beta+1)\log(1/|x|)} dx = \infty. \end{aligned}$$

For a proof of Theorem 3.1, we prepare several results.

Lemma 3.3. *Suppose $0 < a \leq R_0$ and $0 \leq b \leq R_0$. Then there exists a constant $C(R_0) > 0$ such that*

$$\int_{\delta}^{1/2} t^{-a} (\log(1/t))^{-b} \frac{dt}{t} \leq C(R_0) a^{-b-1} \delta^{-a} (\log(1/\delta))^{-b}$$

for all $0 < \delta < 1/2$.

Proof. Note that $u_a(s) = s^{-a} (\log(1/s))^{-b}$ attains a minimum value of $e^b b^{-b} a^b$ at $s = e^{-b/a}$ for $0 < s < 1$. If $1/2 \leq e^{-b/a}$, then u_a is decreasing on $(0, 1/2]$. Hence

$$u_a(t) \leq u_a(\delta) \quad \text{for } 0 < \delta \leq t < 1/2.$$

If $e^{-b/a} < 1/2$, then u_a is decreasing on $(0, e^{-b/a}]$ and increasing on $[e^{-b/a}, 1/2]$. Hence, in the case $e^{-b/a} \leq \delta$ we have

$$u_a(t) \leq \frac{u_a(1/2)}{u_a(e^{-b/a})} u_a(\delta) = \frac{2^a (\log 2)^{-b}}{e^b b^{-b} a^b} u_a(\delta) \quad \text{for } 0 < \delta \leq t < 1/2,$$

and, in the case $0 < \delta < e^{-b/a}$ we have

$$\begin{aligned} u_a(t) &\leq u_a(\delta) \quad \text{for } 0 < \delta \leq t < e^{-b/a}, \\ u_a(t) &\leq \frac{2^a (\log 2)^{-b}}{e^b b^{-b} a^b} u_a(\delta) \quad \text{for } e^{-b/a} \leq t < 1/2. \end{aligned}$$

Therefore, we obtain

$$u_a(t) \leq C(R_0) a^{-b} u_a(\delta) \quad \text{for } 0 < \delta \leq t < 1/2, \tag{3.1}$$

so that

$$\begin{aligned} \int_{\delta}^{1/2} t^{-a} (\log(1/t))^{-b} \frac{dt}{t} &\leq C(R_0) (a/2)^{-b} u_{a/2}(\delta) \int_{\delta}^{1/2} t^{-a/2} \frac{dt}{t} \\ &\leq C(R_0) 2^{b+1} a^{-b-1} \delta^{-a} (\log(1/\delta))^{-b} \end{aligned}$$

for all $0 < \delta < 1/2$, as required. \square

Lemma 3.4. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 3.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \leq C \gamma_1(x) \delta^{\alpha-n/p(x)} (\log(1/\delta))^{-q(x)/p(x)}$$

for all $x \in G$ and $0 < \delta < 1/2$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

First note that

$$\begin{aligned} \int_{G \setminus B(x, p^*(x)^{-1/n})} |x - y|^{\alpha-n} f(y) dy &\leq p^*(x)^{1-\alpha/n} \int_G f(y) dy \\ &\leq p^*(x)^{1-\alpha/n} \int_G \{1 + g(y)\} dy \\ &\leq C p^*(x)^{1-\alpha/n} \\ &\leq C p^*(x)^{1-1/p(x)} \\ &\leq C \gamma_1(x) \end{aligned}$$

since $p^*(x)^{1/p^*(x)} \leq C$, where $g(y) = f(y)^{p(y)} (\log(c + f(y)))^{q(y)}$ as before.

Setting $\eta(x) = p^*(x)^{-1/p(x)} (\log p^*(x))^{q(x)/p(x)}$ and $N(x, y) = |x - y|^{-n/p(x)} (\log(1/|x - y|))^{-q(x)/p(x)}$, we have

$$\begin{aligned}
 & \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\
 & \leq \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} \{ \eta(x) N(x, y) \} dy \\
 & \quad + \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) \\
 & \quad \times C \left(\frac{f(y)}{\eta(x) N(x, y)} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + \eta(x) N(x, y))} \right)^{q(y)} dy.
 \end{aligned}$$

Here note that for $y \in B(x, p^*(x)^{-1/n})$

$$\{ \eta(x) N(x, y) \}^{-p(y)} \leq C \eta(x)^{-p(x)} |x - y|^n (\log(1/|x - y|))^{q(x)}$$

since

$$p^*(x)^{p(y)} \leq p^*(x)^{a/(\log p^*(x)^{1/n})} p^*(x)^{p(x)} \leq C p^*(x)^{p(x)}$$

and

$$\eta(x)^{-p(y)} \leq C \eta(x)^{-p(x)}.$$

Further, noting that

$$\log(c + st) \leq C \log(c + s) \log(c + t) \quad \text{when } s, t > 0, \tag{3.2}$$

we obtain

$$\begin{aligned}
 \{ \log(c + \eta(x) N(x, y)) \}^{-q(y)} & \leq C (\log p^*(x))^{q(y)} (\log(1/|x - y|))^{-q(y)} \\
 & \leq C (\log p^*(x))^{q(x)} (\log(1/|x - y|))^{-q(x)}
 \end{aligned}$$

for $y \in B(x, p^*(x)^{-1/n})$. It follows from Lemma 3.3 and (3.1) that

$$\begin{aligned}
 & \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\
 & \leq C \left\{ \eta(x) (n/p(x) - \alpha)^{-q(x)/p(x)-1} \delta^{\alpha-n/p(x)} (\log(1/\delta))^{-q(x)/p(x)} \right. \\
 & \quad + \eta(x)^{1-p(x)} (\log p^*(x))^{q(x)} \\
 & \quad \times \left. \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n/p(x)} (\log(1/|x - y|))^{-q(x)/p(x)} g(y) dy \right\} \\
 & \leq C \left\{ \gamma_1(x) \delta^{\alpha-n/p(x)} (\log(1/\delta))^{-q(x)/p(x)} + \eta(x)^{1-p(x)} (\log p^*(x))^{q(x)} \right. \\
 & \quad \times (n/p(x) - \alpha)^{-q(x)/p(x)} \delta^{\alpha-n/p(x)} (\log(1/\delta))^{-q(x)/p(x)} \left. \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} g(y) dy \right\}
 \end{aligned}$$

$$\begin{aligned} &\leq C\gamma_1(x)\delta^{\alpha-n/p(x)}(\log(1/\delta))^{-q(x)/p(x)}\left(1 + \int_{B(x, p^*(x)^{-1/n}) \setminus B(x, \delta)} g(y) dy\right) \\ &\leq C\gamma_1(x)\delta^{\alpha-n/p(x)}(\log(1/\delta))^{-q(x)/p(x)}, \end{aligned}$$

which proves the lemma. \square

For a locally integrable function f on G , we set

$$f_B = \frac{1}{|B|} \int_{B \cap G} f(y) dy.$$

We consider the maximal function Mf defined by

$$Mf(x) = \sup_B |f|_B,$$

where the supremum is taken over all balls $B = B(x, r)$. Diening [3] was the first who proved the local boundedness of maximal functions in the Lebesgue spaces of variable exponents satisfying the log-Hölder condition. In our case, we need the following result (see also D. Cruz-Urbe and A. Fiorenza [2]):

Proposition 3.5. (See [11, Theorem 2.7].) *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that*

$$p^- > 1.$$

Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then

$$\int_G \Phi(x, Mf(x)) dx \leq C.$$

Proof of Theorem 3.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Write

$$f = f\chi_{\{y \in G: f(y) < 1\}} + f\chi_{\{y \in G: f(y) \geq 1\}} = f_1 + f_2.$$

Then $I_\alpha f_1(x) \leq C$.

We have by Lemma 3.4

$$\begin{aligned} I_\alpha f(x) &= \int_{B(x, \delta)} |x - y|^{\alpha-n} f(y) dy + \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\ &\leq C \left\{ \delta^\alpha Mf(x) + \gamma_1(x) \delta^{\alpha-n/p(x)} (\log(1/\delta))^{-q(x)/p(x)} \right\} \end{aligned}$$

for $0 < \delta < 1/2$. Here, considering

$$\delta = C(\gamma_1(x)^{-1} Mf(x))^{-p(x)/n} (\log(\gamma_1(x)^{-1} Mf(x)))^{-q(x)/n}$$

when $\gamma_1(x)^{-1} Mf(x) \geq 1$, we find

$$I_\alpha f(x) \leq C\gamma_1(x)^{\alpha p(x)/n} Mf(x)^{1-\alpha p(x)/n} (\log(c + \gamma_1(x)^{-1} Mf(x)))^{-\alpha q(x)/n}.$$

If $\gamma_1(x)^{-1}Mf(x) < 1$, then

$$I_\alpha f(x) \leq C \{Mf(x) + \gamma_1(x)\} \leq C\gamma_1(x).$$

Hence it follows that

$$\begin{aligned} \gamma_1(x)^{-1}I_\alpha f(x) &\leq C \{(\gamma_1(x)^{-1}Mf(x))^{1-\alpha p(x)/n} (\log(c + \gamma_1(x)^{-1}Mf(x)))^{-\alpha q(x)/n} + 1\} \\ &\leq C \{Mf(x)^{p(x)/p^*(x)} (\log(c + Mf(x)))^{-\alpha q(x)/n} + 1\} \end{aligned}$$

since $\gamma_1(x)^{-1/p^*(x)} \leq C$, so that

$$\{c_1^{-1}\gamma_1(x)^{-1}I_\alpha f(x) (\log(c + \gamma_1(x)^{-1}I_\alpha f(x)))^{q(x)/p(x)}\}^{p^*(x)} \leq C \{\Phi(x, Mf(x)) + 1\}.$$

By Proposition 3.5, we have

$$\int_G \Psi(x, c_1^{-1}\gamma_1(x)^{-1}I_\alpha f(x)) dx \leq C \int_G \{\Phi(x, Mf(x)) + 1\} dx \leq C,$$

which completes the proof. \square

4. Trudinger’s exponential integrability

This section is concerned with the exponential integrability of Trudinger’s type.

Theorem 4.1. *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that*

$$p(x) \geq n/\alpha \quad \text{and} \quad q(x) < p(x) - 1$$

for $x \in G$. Then there exist constants $c_1, c_2 > 0$ such that

$$\int_G \exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-q(x)-1)}}{(c_1\gamma_3(x))^{p(x)/(p(x)-q(x)-1)}}\right) dx \leq c_2$$

for all nonnegative measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$, where

$$\gamma_3(x) = \gamma_2(x)^{-(p(x)-1)/p(x)} (\log(1/\gamma_2(x)))^{q(x)/p(x)}$$

with $\gamma_2(x) = \min\{p(x) - q(x) - 1, 1/2\}$.

Corollary 4.2. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 4.1. Then there exists a constant $c_3 > 0$ such that*

$$\int_G \left\{ \exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-q(x)-1)}}{(c_3\gamma_3(x))^{p(x)/(p(x)-q(x)-1)}}\right) - 1 \right\} dx \leq 1$$

for all nonnegative measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

Remark 4.3. For $0 < \delta < 1$, we can find $f \in \Phi(\cdot, \cdot)(G)$ such that

$$\int_{B_0} \exp((\gamma_3(x)^{-\delta} I_\alpha f(x))^{p(x)/(p(x)-q(x)-1)}) dx = \infty.$$

For this, consider

$$p(x) = \frac{n}{\alpha} + \frac{1}{\log(1/|x|)}$$

for $x \in B_0 = B(0, 1/4)$ and

$$f(y) = |y|^{-\alpha} (\log(1/|y|))^{-1} (\log \log(1/|y|))^{-\beta}$$

for $y \in B_0$. If $q(x) = p(x) - 1 - 1/\log \log(1/|x|)$ and $\beta > \alpha/n$, then

$$\int_{B_0} f(y)^{p(y)} (\log f(y))^{q(y)} dy < \infty.$$

If $x \in B_0$, then, as in Remark 3.2, we find

$$I_\alpha f(x) \geq C (\log \log(1/|x|))^{-\beta+1}$$

when $\beta < 1$. Now take β and $\varepsilon > 0$ such that $\alpha/n < \beta < 1$ and

$$-\delta(1 - \alpha/n + \varepsilon) - \beta + 1 > 0.$$

Since

$$\gamma_3(x) \leq C (\log \log(1/|x|))^{1-\alpha/n} (\log \log \log(1/|x|))^{(n-\alpha)/n} \leq C (\log \log(1/|x|))^{1-\alpha/n+\varepsilon}$$

and

$$p(x)/(p(x) - q(x) - 1) > (n/\alpha) \log \log(1/|x|),$$

we have

$$\begin{aligned} & \int_{B_0} \exp((\gamma_3(x)^{-\delta} I_\alpha f(x))^{p(x)/(p(x)-q(x)-1)}) dx \\ & \geq \int_{B_0} \exp((C \log \log(1/|x|))^{(-\delta(1-\alpha/n+\varepsilon)-\beta+1)(n/\alpha) \log \log(1/|x|)}) dx = \infty. \end{aligned}$$

Before proving Theorem 4.1, we prepare the following result.

Lemma 4.4. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 4.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \leq C \gamma_3(x) (\log(1/\delta))^{(p(x)-q(x)-1)/p(x)}$$

for all $x \in G$ and $0 < \delta < 1/2$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

First note that

$$\int_{G \setminus B(x, \gamma_2(x)^{1/n})} |x - y|^{\alpha-n} f(y) dy \leq C \gamma_2(x)^{(\alpha-n)/n} \leq C \gamma_2(x)^{-1+1/p(x)}.$$

Setting $\eta(x) = \gamma_2(x)^{1/p(x)}(\log(1/\gamma_2))^{q(x)/p(x)}$ and $N(x, y) = |x - y|^{-n/p(x)}(\log(1/|x - y|))^{-(q(x)+1)/p(x)}$, we have

$$\begin{aligned} & \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\ & \leq \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} \{\eta(x)N(x, y)\} dy + \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) \\ & \quad \times C \left(\frac{f(y)}{\eta(x)N(x, y)} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + \eta(x)N(x, y))} \right)^{q(y)} dy. \end{aligned}$$

If $y \in B(x, \gamma_2(x)^{1/n})$, then $\gamma_2(x)^{-p(y)} \leq C\gamma_2(x)^{-p(x)}$, so that

$$\eta(x)^{-p(y)} \leq C\eta(x)^{-p(x)}.$$

Hence

$$\{\eta(x)N(x, y)\}^{-p(y)} \leq C\eta(x)^{-p(x)}|x - y|^n (\log(1/|x - y|))^{q(x)+1}$$

and by inequality (3.2)

$$\{\log(c + \eta(x)N(x, y))\}^{-q(y)} \leq C(\log(1/\gamma_2(x)))^{q(x)}(\log(1/|x - y|))^{-q(x)}$$

for $y \in B(x, \gamma_2(x)^{1/n})$. Consequently it follows from Lemma 3.3 that

$$\begin{aligned} & \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\ & \leq C \left\{ \eta(x)\gamma_2(x)^{-1}(\log(1/\delta))^{(p(x)-q(x)-1)/p(x)} + \eta(x)^{-p(x)+1}(\log(1/\gamma_2(x)))^{q(x)} \right. \\ & \quad \left. \times \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} |x - y|^{\alpha-n/p(x)}(\log(1/|x - y|))^{(p(x)-q(x)-1)/p(x)} g(y) dy \right\} \\ & \leq C\gamma_3(x)(\log(1/\delta))^{(p(x)-q(x)-1)/p(x)} \left(1 + \int_{B(x, \gamma_2(x)^{1/n}) \setminus B(x, \delta)} g(y) dy \right) \\ & \leq C\gamma_3(x)(\log(1/\delta))^{(p(x)-q(x)-1)/p(x)}, \end{aligned}$$

where $g(y) = f(y)^{p(y)}(\log(c + f(y)))^{q(y)}$ as before.

Thus we have proved that

$$\begin{aligned} \int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy & \leq C \{ \gamma_2(x)^{-1+1/p(x)} + \gamma_3(x)(\log(1/\delta))^{(p(x)-q(x)-1)/p(x)} \} \\ & \leq C\gamma_3(x)(\log(1/\delta))^{(p(x)-q(x)-1)/p(x)} \end{aligned}$$

for $0 < \delta < 1/2$, which gives the lemma. \square

Proof of Theorem 4.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then Lemma 4.4 gives

$$I_\alpha f(x) \leq C \{ \delta^\alpha Mf(x) + \gamma_3(x) (\log(1/\delta))^{(p(x)-q(x)-1)/p(x)} \}$$

for $0 < \delta < 1/2$. Here, considering

$$\delta = C (\gamma_3(x)^{-1} Mf(x))^{-1/\alpha} (\log(\gamma_3(x)^{-1} Mf(x)))^{(p(x)-q(x)-1)/(\alpha p(x))}$$

when $\gamma_3(x)^{-1} Mf(x) \geq 1$, we find

$$\begin{aligned} I_\alpha f(x) &\leq C \{ \gamma_3(x) (\log(c + \gamma_3(x)^{-1} Mf(x)))^{(p(x)-q(x)-1)/p(x)} + \gamma_3(x) \} \\ &\leq C \gamma_3(x) (\log(c + Mf(x)))^{(p(x)-q(x)-1)/p(x)}. \end{aligned}$$

Hence it follows that

$$c_1^{-1} \gamma_3(x)^{-1} I_\alpha f(x) \leq (\log(c + Mf(x)))^{(p(x)-q(x)-1)/p(x)},$$

so that

$$\exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-q(x)-1)}}{(c_1 \gamma_3(x))^{p(x)/(p(x)-q(x)-1)}}\right) \leq c + Mf(x) \leq C \{ \Phi(x, Mf(x)) + 1 \}.$$

By Proposition 3.5, we have

$$\int_G \exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-q(x)-1)}}{(c_1 \gamma_3(x))^{p(x)/(p(x)-q(x)-1)}}\right) dx \leq C \left(\int_G \Phi(x, Mf(x)) dx + 1 \right) \leq C,$$

as required. \square

5. Trudinger’s double exponential integrability

This section is devoted to the study of the double exponential integrability.

Theorem 5.1. *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that*

$$p(x) \geq n/\alpha \quad \text{and} \quad q(x) \geq p(x) - 1$$

for $x \in \mathbf{R}^n$. Then there exist constants $c_1, c_2 > 0$ such that

$$\int_G \exp\left(\exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-1)}}{c_1^{p(x)/(p(x)-1)}}\right)\right) dx \leq c_2$$

for all nonnegative measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

Lemma 5.2. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 5.1 and let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{G \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \leq C (\log(\log(1/\delta)))^{(p(x)-1)/p(x)}$$

for all $x \in G$ and $0 < \delta < 1/2$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

First note that

$$\int_{G \setminus B(x, 1/4)} |x - y|^{\alpha-n} f(y) dy \leq C.$$

Next, setting $N(x, y) = |x - y|^{-n/p(x)} (\log(1/|x - y|))^{-1} (\log(\log(1/|x - y|)))^{-1/p(x)}$, we have

$$\begin{aligned} & \int_{B(x, 1/4) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) dy \\ & \leq \int_{B(x, 1/4) \setminus B(x, \delta)} |x - y|^{\alpha-n} N(x, y) dy \\ & \quad + \int_{B(x, 1/4) \setminus B(x, \delta)} |x - y|^{\alpha-n} f(y) \left(\frac{f(y)}{N(x, y)} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + N(x, y))} \right)^{q(y)} dy \\ & \leq C \left\{ (\log(\log(1/\delta)))^{(p(x)-1)/p(x)} + \int_{B(x, 1/4) \setminus B(x, \delta)} |x - y|^{\alpha-n/p(x)} \right. \\ & \quad \left. \times (\log(1/|x - y|))^{p(x)-q(x)-1} (\log(\log(1/|x - y|)))^{(p(x)-1)/p(x)} g(y) dy \right\} \\ & \leq C (\log(\log(1/\delta)))^{(p(x)-1)/p(x)}, \end{aligned}$$

where $g(y) = f(y)^{p(y)} (\log(c + f(y)))^{q(y)}$, as required. \square

Proof of Theorem 5.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then Lemma 5.2 gives

$$I_\alpha f(x) \leq C \delta^\alpha Mf(x) + C (\log(\log(1/\delta)))^{(p(x)-1)/p(x)}$$

for $0 < \delta < 1/4$. Here, considering

$$\delta = C Mf(x)^{-1/\alpha} (\log(\log(Mf(x))))^{(p(x)-1)/(ap(x))}$$

when $Mf(x) \geq e^2$, we find

$$I_\alpha f(x) \leq C (\log(c + \log(c + Mf(x))))^{(p(x)-1)/p(x)}.$$

Hence it follows that

$$c_1^{-1} I_\alpha f(x) \leq (\log(\log(c + Mf(x))))^{(p(x)-1)/p(x)},$$

so that

$$\exp\left(\exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-1)}}{c_1^{p(x)/(p(x)-1)}}\right)\right) \leq C \{\Phi(x, Mf(x)) + 1\}.$$

Now Proposition 3.5 yields

$$\int_G \exp\left(\exp\left(\frac{I_\alpha f(x)^{p(x)/(p(x)-1)}}{c_1^{p(x)/(p(x)-1)}}\right)\right) dx \leq C \left(\int_G \Phi(x, Mf(x)) dx + 1\right) \leq c_2,$$

as required. \square

6. Proof of Theorem 1.1

Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Let $p_1(x) = \min\{p(x), 1 + \delta\}$ for $0 < \delta < n/\alpha$. Then Theorem 2.1 yields

$$\int_{G_1} \tilde{\Psi}(x, I_\alpha f_1(x)) dx \leq C.$$

Letting $p_2(x) = \max\{p(x), 1 + \delta\}$, we see by Theorem 3.1 that

$$\int_{G_2} \Psi(x, c_1^{-1} \gamma_1(x)^{-1} I_\alpha f_2(x)) dx \leq C,$$

where $c_1 > 0$ is as in Theorem 3.1. Hence, in view of Theorems 4.1 and 5.1, Theorem 1.1 is proved. \square

7. Continuity of Riesz potentials

In this section we are concerned with continuity properties of Riesz potentials.

Theorem 7.1. *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that*

$$p(x) \geq n/\alpha \quad \text{and} \quad q(x) > p(x) - 1$$

for $x \in \mathbf{R}^n$. If f is a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$, then $I_\alpha f(x)$ is continuous and

$$|I_\alpha f(z) - I_\alpha f(x)| \leq C \gamma_5(x) (\log(1/|z - x|))^{-(q(x) - p(x) + 1)/p(x)}$$

as $z \rightarrow x$ for each $x \in G$, where

$$\gamma_5(x) = \gamma_4(x)^{-(p(x) - 1)/p(x)} (\log(1/\gamma_4(x)))^{q(x)/p(x)}$$

with $\gamma_4(x) = \min\{q(x) - p(x) + 1, 1/2\}$.

Lemma 7.2. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 7.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{B(x, \delta)} |x - y|^{\alpha - n} f(y) dy \leq C \gamma_5(x, \delta) (\log(1/\delta))^{-(q(x) - p(x) + 1)/p(x)}$$

for all $x \in G$ and $0 < \delta < 1/4$, where

$$\begin{aligned} \gamma_5(x, t) &= \gamma_4(x)^{-(p(x) - 1)/p(x) - a/(p(x) \log(1/t))} \\ &\quad \times (\log(1/\gamma_4(x)))^{q(x)/p(x) - aq(x)/(p(x) \log(1/t)) + b/\log(\log(1/t))}. \end{aligned}$$

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

Setting $\eta(x) = \gamma_4(x)^{1/p(x)} (\log(1/\gamma_4(x)))^{q(x)/p(x)}$ and $N(x, y) = |x - y|^{-n/p(x)} (\log(1/|x - y|))^{-(q(x) + 1)/p(x)}$, we have

$$\begin{aligned} & \int_{B(x,\delta)} |x - y|^{\alpha-n} f(y) dy \\ & \leq \int_{B(x,\delta)} |x - y|^{\alpha-n} \eta(x) N(x, y) dy \\ & \quad + \int_{B(x,\delta)} |x - y|^{\alpha-n} f(y) \left(\frac{f(y)}{\eta(x)N(x, y)} \right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + \eta(x)N(x, y))} \right)^{q(y)} dy. \end{aligned}$$

Note that

$$\{\eta(x)N(x, y)\}^{-p(y)} \leq \eta(x)^{-p(x)-a/\log(1/\delta)} |x - y|^n (\log(1/|x - y|))^{q(x)+1}$$

and

$$\{\log(c + \eta(x)N(x, y))\}^{-q(y)} \leq C(\log(1/\gamma_4(x)))^{q(x)+b/\log(\log(1/\delta))} (\log(1/|x - y|))^{-q(x)}$$

for $y \in B(x, \delta)$. Consequently it follows that

$$\begin{aligned} & \int_{B(x,\delta)} |x - y|^{\alpha-n} f(y) dy \\ & \leq C \left\{ \eta(x)\gamma_4(x)^{-1} (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)} \right. \\ & \quad \left. + \eta(x)^{-p(x)+1-a/\log(1/\delta)} (\log(1/\gamma_4(x)))^{q(x)+b/\log(\log(1/\delta))} \right. \\ & \quad \left. \times \int_{B(x,\delta)} |x - y|^{\alpha-n/p(x)} (\log(1/|x - y|))^{-(q(x)-p(x)+1)/p(x)} g(y) dy \right\} \\ & \leq C\gamma_5(x, \delta) (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)} \left(1 + \int_{B(x,\delta)} g(y) dy \right) \\ & \leq C\gamma_5(x, \delta) (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)}, \end{aligned}$$

where $g(y) = f(y)^{p(y)} (\log(c + f(y)))^{q(y)}$, as required. \square

Lemma 7.3. *Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 7.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then*

$$\int_{G \setminus B(x,\delta)} |x - y|^{\alpha-n-1} f(y) dy \leq C\delta^{-1} (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)}$$

for all $x \in G$ and $0 < \delta < 1/2$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

First note that

$$\int_{G \setminus B(x,1/2)} |x - y|^{\alpha-n-1} f(y) dy \leq C.$$

Setting $N(x, y) = |x - y|^{-n/p(x)} (\log(1/|x - y|))^{-(q(x)+1)/p(x)}$, we have

$$\begin{aligned} & \int_{B(x,1/2)\setminus B(x,\delta)} |x - y|^{\alpha-n-1} f(y) dy \\ & \leq \int_{B(x,1/2)\setminus B(x,\delta)} |x - y|^{\alpha-n-1} N(x, y) dy \\ & \quad + \int_{B(x,1/2)\setminus B(x,\delta)} |x - y|^{\alpha-n-1} f(y) \left(\frac{f(y)}{N(x, y)}\right)^{p(y)-1} \left(\frac{\log(c + f(y))}{\log(c + N(x, y))}\right)^{q(y)} dy. \end{aligned}$$

Since

$$\{N(x, y)\}^{-p(y)} \leq C|x - y|^n (\log(1/|x - y|))^{q(x)+1}$$

and

$$\{\log(c + N(x, y))\}^{-q(y)} \leq C(\log(1/|x - y|))^{-q(x)}$$

for $y \in B(x, 1/2)$, it follows from Lemma 3.3 that

$$\begin{aligned} & \int_{B(x,1/2)\setminus B(x,\delta)} |x - y|^{\alpha-n-1} f(y) dy \\ & \leq C \left\{ \delta^{-1} (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)} \right. \\ & \quad \left. + \int_{B(x,1/2)\setminus B(x,\delta)} |x - y|^{\alpha-n/p(x)-1} (\log(1/|x - y|))^{-(q(x)-p(x)+1)/p(x)} g(y) dy \right\} \\ & \leq C \delta^{-1} (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)} \left(1 + \int_{B(x,1/2)\setminus B(x,\delta)} g(y) dy \right) \\ & \leq C \delta^{-1} (\log(1/\delta))^{-(q(x)-p(x)+1)/p(x)}, \end{aligned}$$

where $g(y) = f(y)^{p(y)} (\log(c + f(y)))^{q(y)}$, as required. \square

Proof of Theorem 7.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

Write

$$\begin{aligned} I_\alpha f(x) - I_\alpha f(z) &= \int_{B(x,2|x-z|)} |x - y|^{\alpha-n} f(y) dy - \int_{B(x,2|x-z|)} |z - y|^{\alpha-n} f(y) dy \\ & \quad + \int_{G \setminus B(x,2|x-z|)} (|x - y|^{\alpha-n} - |z - y|^{\alpha-n}) f(y) dy. \end{aligned}$$

By Lemma 7.2, we have

$$\int_{B(x,2|x-z|)} |x - y|^{\alpha-n} f(y) dy \leq C\gamma_5(x, 3|x - z|) (\log(1/|z - x|))^{-(q(x)-p(x)+1)/p(x)}$$

and

$$\begin{aligned} \int_{B(x,2|x-z|)} |z-y|^{\alpha-n} f(y) dy &\leq \int_{B(z,3|x-z|)} |z-y|^{\alpha-n} f(y) dy \\ &\leq C\gamma_5(z,3|x-z|)(\log(1/|z-x|))^{-(q(z)-p(z)+1)/p(z)} \end{aligned}$$

for $0 < |x-z| < 1/2$. On the other hand, by the mean value theorem for analysis, we have by Lemma 7.3

$$\begin{aligned} \int_{G \setminus B(x,2|x-z|)} \left| |x-y|^{\alpha-n} - |z-y|^{\alpha-n} \right| f(y) dy \\ \leq C|x-z| \int_{G \setminus B(x,2|x-z|)} |x-y|^{\alpha-n-1} f(y) dy \\ \leq C(\log(1/|z-x|))^{-(q(x)-p(x)+1)/p(x)}. \end{aligned}$$

Now we establish

$$\begin{aligned} |I_\alpha f(x) - I_\alpha f(z)| &\leq C \left\{ \gamma_5(x,3|x-z|)(\log(1/|z-x|))^{-(q(x)-p(x)+1)/p(x)} \right. \\ &\quad \left. + \gamma_5(z,3|x-z|)(\log(1/|z-x|))^{-(q(z)-p(z)+1)/p(z)} \right\} \end{aligned}$$

for $0 < |x-z| < 1/4$, which implies

$$|I_\alpha f(z) - I_\alpha f(x)| \leq C\gamma_5(x)(\log(1/|z-x|))^{-(q(x)-p(x)+1)/p(x)}$$

as $z \rightarrow x$ for all $x \in G$. \square

8. Logarithmic potentials

In this section we discuss Sobolev’s theorem for logarithmic potentials.

Theorem 8.1. *Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.2) such that $p(x) = 1$ and*

$$0 \leq q(x) < 1$$

for $x \in \mathbf{R}^n$. Then there exist constants $c_1, c_2 > 0$ such that

$$\int_G \left\{ \exp((c_1 I_n f(x))^{1/(1-q(x))}) - 1 \right\} dx \leq c_2$$

for all nonnegative measurable functions f on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$.

To show this, we estimate $I_n f$ by the logarithmic potential

$$J = \int_G \rho_{-\beta}(|x-y|)g(y) dy,$$

where $\rho_{-\beta}(r) = r^{-n}(\log(2+1/r))^{-\beta}$ with $\beta > 1$ and $g(y) = f(y)(\log(e+f(y)))^{q(y)}$.

Lemma 8.2. Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 8.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then

$$F \equiv \int_{B(x, \delta)} \rho_{-\beta}(|x - y|) f(y) dy \leq CJ \{ (\log(e + J))^{-q(x)} + (\log(e + 1/\delta))^{-q(x)} \}$$

for all $x \in G$ and $\delta > 0$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. We have for $k > 0$

$$F \leq k \int_G \rho_{-\beta}(|x - y|) dy + \int_{B(x, \delta)} \rho_{-\beta}(|x - y|) f(y) \left(\frac{\log(e + f(y))}{\log(e + k)} \right)^{q(y)} dy.$$

If $J \leq \delta^{-n}$, then we set

$$k = J (\log(e + J))^{-q(x)}.$$

Since $\delta \leq J^{-1/n}$, we see that

$$(\log(e + k))^{-q(y)} \leq C (\log(e + J))^{-q(x)}$$

for $y \in B(x, \delta)$. Consequently it follows that

$$F \leq CJ (\log(e + J))^{-q(x)}.$$

If $J > \delta^{-n}$, then we set

$$k = \delta^{-n} (\log(e + 1/\delta))^{-q(x)}$$

and obtain

$$\begin{aligned} F &\leq C \{ \delta^{-n} (\log(e + 1/\delta))^{-q(x)} + (\log(e + 1/\delta))^{-q(x)} J \} \\ &\leq C (\log(e + 1/\delta))^{-q(x)} J. \end{aligned}$$

Now the result follows. \square

Lemma 8.3. Let $p(\cdot)$ and $q(\cdot)$ be as in Theorem 8.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Then

$$\int_{G \setminus B(x, \delta)} \log^+(1/|x - y|) f(y) dy \leq C (\log(e + 1/\delta))^{-q(x)+1}$$

for all $x \in G$ and $\delta > 0$.

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. Let $0 < \gamma < n$. For $y \in G \setminus B(x, \delta)$ and $\delta > 0$, set

$$N(x, y) = |x - y|^{-\gamma}.$$

By condition (1.2), we see that

$$\begin{aligned}
 & \int_{G \setminus B(x, \delta)} \log^+(1/|x - y|) f(y) dy \\
 & \leq \int_G \log^+(1/|x - y|) N(x, y) dy \\
 & \quad + \int_{G \setminus B(x, \delta)} \log^+(1/|x - y|) f(y) \left(\frac{\log(e + f(y))}{\log(e + N(x, y))} \right)^{q(y)} dy \\
 & \leq C \left\{ 1 + \int_{G \setminus B(x, \delta)} (\log(e + 1/|x - y|))^{-q(y)+1} g(y) dy \right\} \\
 & \leq C \left\{ 1 + (\log(e + 1/\delta))^{-q(x)+1} \int_{G \setminus B(x, \delta)} g(y) dy \right\} \\
 & \leq C (\log(e + 1/\delta))^{-q(x)+1},
 \end{aligned}$$

where $g(y) = f(y)^{p(y)} (\log(e + f(y)))^{q(y)}$, as required. \square

Proof of Theorem 8.1. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} \leq 1$. For $x \in G$ and $\delta > 0$, write

$$\begin{aligned}
 I_n f(x) &= \int_{B(x, \delta)} \log^+(1/|x - y|) f(y) dy + \int_{G \setminus B(x, \delta)} \log^+(1/|x - y|) f(y) dy \\
 &= I_1 + I_2.
 \end{aligned}$$

For $\beta > 1$, we infer from Lemma 8.2 that

$$\begin{aligned}
 I_1 &\leq C \delta^n (\log(e + 1/\delta))^{1+\beta} \int_{B(x, \delta)} \rho_{-\beta}(|x - y|) f(y) dy \\
 &\leq C \delta^n (\log(e + 1/\delta))^{1+\beta} J \{ (\log(e + 1/\delta))^{-q(x)} + (\log(e + J))^{-q(x)} \}.
 \end{aligned}$$

Hence, in view of Lemma 8.3, we find

$$\begin{aligned}
 I_n f(x) &\leq C \{ \delta^n (\log(e + 1/\delta))^{1+\beta} J \{ (\log(e + 1/\delta))^{-q(x)} + (\log(e + J))^{-q(x)} \} \\
 &\quad + (\log(e + 1/\delta))^{-q(x)+1} \}.
 \end{aligned}$$

Now, considering $\delta = J^{-1/n} (\log(e + J))^{-\beta/n}$, we find

$$I_n f(x) \leq C (\log(e + J))^{-q(x)+1},$$

so that

$$\exp((c_1 I_n f(x))^{1/(1-q(x))}) \leq e + J.$$

Integrating both sides over G gives

$$\int_G \exp((c_1 I_n f(x))^{1/(1-q(x))}) dx \leq c_2,$$

which proves the required result. \square

Theorem 8.4. Let $p(\cdot)$ and $q(\cdot)$ be two variable exponents on \mathbf{R}^n satisfying (1.1)–(1.4) such that

$$p(x) > 1 \quad \text{or} \quad q(x) \geq 1$$

for all $x \in \bar{G}$. If f is a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} < \infty$, then $I_n f$ is continuous on G .

Proof. Let f be a nonnegative measurable function on G with $\|f\|_{\Phi(\cdot, \cdot)(G)} < \infty$. Then note that

$$\int_G f(y)(\log(e + f(y))) dy < \infty.$$

Hence, it follows from [10, Theorem 9.1, Section 5.9] that $I_n f$ is continuous on G . \square

In the same manner as Lemmas 8.2 and 8.3 we can show that

$$I_n f(z) - I_n f(x) = o(|z - x|^{n(1-1/p(x))} (\log(1/|z - x|))^\gamma)$$

as $z \rightarrow x$, $x \in G$ when $p(x) < n'$ and $\gamma > 2 - (q(x) + 1)/p(x)$.

For further results, we refer the reader to the paper by Ohno [12].

References

- [1] A. Cianchi, Strong and weak type inequalities for some classical operators in Orlicz spaces, J. London Math. Soc. 60 (1999) 187–202.
- [2] D. Cruz-Uribe, A. Fiorenza, $L \log L$ results for the maximal operator in variable L^p spaces, Trans. Amer. Math. Soc. 361 (2009) 2631–2647.
- [3] L. Diening, Maximal functions in generalized $L^{p(\cdot)}$ spaces, Math. Inequal. Appl. 7 (2) (2004) 245–254.
- [4] D.E. Edmunds, J. Rákosník, Sobolev embeddings with variable exponent, II, Math. Nachr. 246–247 (2002) 53–67.
- [5] T. Futamura, Y. Mizuta, Continuity properties of Riesz potentials for functions in $L^{p(\cdot)}$ of variable exponent, Math. Inequal. Appl. 8 (4) (2005) 619–631.
- [6] T. Futamura, Y. Mizuta, T. Shimomura, Sobolev embedding for variable exponent Riesz potentials on metric spaces, Ann. Acad. Sci. Fenn. Math. 31 (2006) 495–522.
- [7] P. Harjulehto, P. Hästö, Sobolev inequalities for variable exponents attaining the values 1 and n , Publ. Mat. 52 (2008) 347–363.
- [8] L.I. Hedberg, On certain convolution inequalities, Proc. Amer. Math. Soc. 36 (1972) 505–510.
- [9] O. Kováčik, J. Rákosník, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, Czechoslovak Math. J. 41 (1991) 592–618.
- [10] Y. Mizuta, Potential Theory in Euclidean Spaces, Gakkōtoshō, Tokyo, 1996.
- [11] Y. Mizuta, T. Ohno, T. Shimomura, Sobolev's inequalities and vanishing integrability for Riesz potentials of functions in $L^{p(\cdot)}(\log L)^{q(\cdot)}$, J. Math. Anal. Appl. 345 (2008) 70–85.
- [12] T. Ohno, Continuity properties for logarithmic potentials of functions in Morrey spaces of variable exponent, Hiroshima Math. J. 38 (2008) 363–383.
- [13] R. O'Neil, Fractional integration in Orlicz spaces. I, Trans. Amer. Math. Soc. 115 (1965) 300–328.
- [14] W. Orlicz, Über konjugierte Exponentenfolgen, Studia Math. 3 (1931) 200–211.
- [15] M. Růžička, Electrorheological Fluids: Modeling and Mathematical Theory, Lecture Notes in Math., vol. 1748, Springer, 2000.