



# Comparison of ouabain-sensitive and -insensitive Na/K pumps in HEK293 cells

Jens Kockskämper<sup>a</sup>, Günter Gisselmann<sup>b</sup>, Helfried Günther Glitsch<sup>a,\*</sup>

<sup>a</sup> *Arbeitsgruppe Muskelphysiologie, Ruhr-Universität, D-44780 Bochum, Germany*

<sup>b</sup> *Lehrstuhl für Zellphysiologie, Ruhr-Universität, D-44780 Bochum, Germany*

Received 19 August 1996; revised 11 December 1996; accepted 17 December 1996

## Abstract

The Na/K pump current  $I_p$  of single HEK293 cells either untransfected (endogenous  $I_p$ ) or transfected with the  $\alpha 1$  subunit of the rat Na/K pump (exogenous  $I_p$ ) was investigated in Na-containing solution by means of whole-cell recording at 30°C. The endogenous  $I_p$  was irreversibly blocked by  $10^{-4}$  M ouabain or  $2 \cdot 10^{-4}$  M dihydro-ouabain (DHO). Its density amounted to  $0.33 \text{ pA pF}^{-1}$  at 0 mV and 5.4 mM  $K_o$ . It was half maximally activated at 1.5 mM  $K_o$  and increased linearly with depolarization over the entire voltage range studied ( $-80$  to  $+60$  mV). In contrast, HEK293 cells stably transfected with cDNA for the cardiac glycoside-resistant  $\alpha 1$  subunit of the rat Na/K pump showed an  $I_p$  in the presence of  $10^{-4}$  M ouabain and  $2 \cdot 10^{-4}$  M DHO, respectively. This exogenous  $I_p$  was reversibly blocked by  $10^{-2}$  M ouabain. Half maximal activation of the exogenous  $I_p$  occurred at 1.7 mM  $K_o$ . Its amplitude increased linearly with depolarization at negative voltages but remained almost constant at positive membrane potentials. Comparison with the  $I_p$  of isolated rat cardiac ventricular myocytes strongly suggests that the exogenous  $I_p$  in HEK293 cells is generated by the  $\alpha 1$  subunit of the rat Na/K pump since it displays identical properties. Therefore, HEK293 cells represent an expression system well suited for the electrophysiological analysis of recombinant, cardiac glycoside-resistant Na/K pumps by means of whole-cell recording.

**Keywords:** Sodium pump; Whole-cell recording; HEK293 cell; Transfection; Expression system; Rat  $\alpha 1$  subunit

## 1. Introduction

In recent years HEK293 cells (human embryonic kidney cells), a mammalian cell line, have been used as a system for the heterologous expression of transporters and ion channels [1,2]. Due to their small volume, the cells are well suited for studies on endogenous and exogenous ion channels and transporters by means of whole-cell recording. In contrast to conventional voltage-clamping on larger cells (e.g.,

*Xenopus* oocytes) the patch-clamp technique [3] allows a good control of the composition of the cell interior and renders possible a change of the intracellular solution during the experiment [4]. Apart from some fundamental data (e.g. Ref. [1]), information about the electrophysiological characteristics of HEK293 cells is scanty. It is, however, clear that endogenous, voltage-activated Ca or Na channels are absent in these cells [2]. The aim of the experiments described below was two-fold. First, we studied the current produced by the endogenous Na/K pump molecules in order to contribute to the electrophysio-

\* Corresponding author: Fax: +49 234 7094129.

logical characterization of the cells. Second, we tested whether HEK293 cells could express exogenous Na/K pumps and thus could provide a system for the analysis of the Na/K pump current generated by exogenous pump molecules by means of whole-cell recording. Such a system would markedly facilitate the electrophysiological analysis of the Na/K pump structure-function relationship. Most recently, Yamamoto and co-workers [5] reported on a successful electrophysiological analysis of recombinant, mutated Na/K pumps expressed in HeLa cells. Their study concentrated on the kinetics of  $K_o^-$  and cardiac glycoside-binding to wild-type and mutated pump molecules. The experiments described below, however, characterize different aspects of endogenous and recombinant Na/K pumps in HEK293 cells. In addition the properties of the rat  $\alpha 1$  subunit in recombinant and wild-type pumps are compared. Some of the results have been published in abstract form [6].

## 2. Materials and methods

### 2.1. Cell culture

HEK293 cells were maintained in culture flasks (25 cm<sup>2</sup>) with Minimal Essential Medium (MEM; Life Technologies GmbH, Eggenstein, Germany) supplemented with 10% (v/v) fetal bovine serum (Bio Whittaker, Verviers, Belgium), 20 mM L-glutamine, 1% (v/v) MEM non-essential amino acids, 100 000 I.U./l penicillin, and 100 mg/l streptomycin (all from Life Technologies, Eggenstein, Germany) at 37°C and 5% CO<sub>2</sub>. The culture medium was replaced every 2–3 days. When the cells were grown to near confluency, they were trypsinized and seeded at a density of 4000 cm<sup>-2</sup> in a new culture flask.

### 2.2. Transfection and selection of stably transfected cells

The eucaryotic expression plasmid pR  $\alpha 1$  was used for the transfection of the HEK293 cells [7]. It is derived from the plasmid pRc/CMV (Invitrogen Corporation, San Diego, USA) and contains the cDNA of the  $\alpha 1$  subunit of the rat Na/K pump

under the control of the enhancer-promotor regions of the immediate early gene from the human cytomegalovirus. HEK293 cells were transfected with pR  $\alpha 1$  by means of the Ca phosphate coprecipitation method [8]. Shortly, two days before transfection  $\approx 1 \cdot 10^5$  HEK293 cells were seeded in a culture flask. On the day of transfection, 250  $\mu$ l 2  $\times$  HEPES-buffered saline (in mM: 280 NaCl, 2.8 Na<sub>2</sub>HPO<sub>4</sub>, 50 N-2-hydroxyethylpiperazine-N'-2-ethane-sulfonic acid (Hepes), pH 7.2 (NaOH)) were gently mixed with 250  $\mu$ l of a 250 mM CaCl<sub>2</sub> solution containing 14  $\mu$ g pR  $\alpha 1$ . Five min later, 240  $\mu$ l of the resulting solution were added dropwise to the cells in the culture flask (2.5 ml culture medium). The cells were kept in the incubator (37°C, 5% CO<sub>2</sub>) for 3 h. Afterwards they were washed twice with phosphate-buffered saline (in mM: 0.9 CaCl<sub>2</sub> · 2 H<sub>2</sub>O, 2.7 KCl, 1.5 KH<sub>2</sub>PO<sub>4</sub>, 0.5 MgCl<sub>2</sub> · 6 H<sub>2</sub>O, 106 NaCl, 6.5 Na<sub>2</sub>HPO<sub>4</sub>, pH 7.2 (NaOH)) and kept in culture medium for two days. To obtain stably transfected cells, 10<sup>-4</sup> M dihydro-ouabain (DHO) was then permanently added to the medium.

Mock transfection (plasmid pRc/CMV only) revealed that all cells died within two days in the DHO-containing culture medium. This observation excludes the possibility that HEK293 cells became ouabain-resistant by mutations of their endogenous  $\alpha 1$  subunit.

### 2.3. Preparation of single rat ventricular myocytes

Isolation of single myocytes was carried out according to Bechem and co-workers [9]. Briefly, adult female Wistar rats were anaesthetized with diethylether and killed by decapitation. The aorta was cannulated before the heart was excised, mounted on a Langendorff apparatus, and perfused at 37°C for several minutes with the following solutions (in mM): (1) Ca-free solution: 204 sucrose, 35 NaCl, 5.4 KCl, 1.0 MgCl<sub>2</sub>, 2.0 ethylene glycol-bis( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), 10 Hepes; (2) enzyme solution: 204 sucrose, 35 NaCl, 5.4 KCl, 1.0 MgCl<sub>2</sub>, 0.1 EGTA, 10 Hepes; 0.5 mg/ml bovine serum albumine (BSA; Sigma, Germany), 0.4 mg/ml collagenase B (Boehringer Mannheim, Germany), 10  $\mu$ l/ml elastase (Serva, Heidelberg, Germany), 0.3 mg/ml protease (Sigma, Germany), 0.15 mg/ml

DNase (Sigma, Germany). The pH of the oxygenated solutions was 7.4. The digested ventricles were cut into pieces and gently stirred at 20–22°C in tissue wells containing a nominally Ca-free solution with the following composition (in mM): 204 sucrose, 35 NaCl, 5.4 KCl, 1.0 MgCl<sub>2</sub>, 10 Hepes; 1.0 mg/ml BSA, 0.3 mg/ml DNase, pH 7.4. [Ca] was increased to 1.3 mM by stepwise replacement with Hanks' medium 199 (PAA, Linz, Austria) supplemented with 100.000 I.U./l penicillin, 100 mg/l streptomycin, and 250 µg/l amphotericin B (all from Sigma, Germany). Isolated myocytes were transferred to culture dishes (Ø 35 mm) and used for the electrophysiological experiments either immediately after isolation or after one day in culture (37°C, 3% CO<sub>2</sub>).

#### 2.4. Whole-cell recording

A culture dish (Ø 35 mm) with either HEK293 cells or rat ventricular myocytes was fixed to the stage of an inverted microscope (Diaphot TMD, Nikon, Tokyo, Japan). The dish was perfused with extracellular solution from a reservoir connected to the dish by plastic tubes. It was placed about 100 cm above the stage. Solution flow, driven by gravity, was 1 ml/min. The solution level in the dish was held constant by means of an outlet opposite to the inlet from the reservoir. A multibarreled pipette (inner diameter: 500 µm) was placed 100–200 µm from the cell under investigation. The cell was continuously superfused at a rate of 0.4 ml/min with one of the solutions from reservoirs connected with the pipette via plastic tubes. The reservoirs were about 30 cm above the stage. Solution changes, controlled by electromagnetic valves (The Lee Company, Westbrook, USA), had a time constant of 200–600 ms [10]. The temperature in the vicinity of the cell was 30°C when superfused with one of the test solutions. Patch pipettes were made from borosilicate glass capillaries (GC150TF-10, Clark Electromedical Instruments, Reading, UK) and back-filled with the pipette solution. They had an initial resistance between 2.5 and 7 MΩ for experiments with HEK293 cells and between 1 and 2.5 MΩ for experiments with rat ventricular myocytes. The tip potential between the patch pipette and the extracellular solution was compensated before establishing a GΩ seal (≥ 10 GΩ). Gentle suction was applied to a patch

pipette positioned at the cell membrane to obtain the whole-cell configuration of the patch-clamp technique [3]. The cells were voltage-clamped by means of an EPC-7 patch clamp amplifier (List Medical, Darmstadt, Germany). The amplifier was connected to a personal computer via 12 bit AD- and DA-converters, respectively. Currents were low-pass filtered at 200 Hz and digitized with 1 kHz (200 Hz in early experiments). The programme ISO2 (MFK, Niedernhausen, Germany) was used to generate the voltage protocols and to record the resulting currents. The cell capacitance was determined by a programme routine that applied depolarizing and hyperpolarizing voltage ramps 10 mV in amplitude starting from a holding potential of 0 mV. HEK293 cells displayed membrane capacitances between 20 and 85 pF, whereas rat ventricular myocytes showed values between 60 and 160 pF.

#### 2.5. Solutions for whole-cell recording

The extracellular solution for whole-cell recording contained (in mM): 144 NaCl, 0–10.8 KCl, 5 NiCl<sub>2</sub>, 2 BaCl<sub>2</sub>, 1.8 CaCl<sub>2</sub>, 0.5 MgCl<sub>2</sub>, 10 Hepes, pH 7.4 (NaOH). Barium and nickel served to suppress potassium and calcium conductances, respectively, as well as the sarcolemmal Na/Ca exchanger. DHO (Roth, Karlsruhe, Germany) and ouabain (Fluka Chemie AG, Buchs, Switzerland) were added from 10<sup>-2</sup> M stock solutions. The DHO stock solution was aqueous, whereas the ouabain stock solution contained 10% (v/v) ethanol. In experiments, in which [ouabain] was 10<sup>-2</sup> M, the drug was solved directly in the extracellular solution. In this case, the superfusate contained 2% (v/v) ethanol. Control experiments revealed that this [ethanol] had no influence on the membrane current under the present experimental conditions.

The pipette solution was composed of (in mM): 110 CsCl, 40 NaCl, 10 NaOH, 3 MgCl<sub>2</sub>, 6 EGTA, 16 Hepes, 10 MgATP, pH 7.4 (CsOH). Assuming a contamination of ≈ 1 · 10<sup>-6</sup> M Ca, free [Ca] and [Mg] were calculated to be 1.3 · 10<sup>-11</sup> M and 2.4 · 10<sup>-3</sup> M, respectively [11]. The solution contained cesium to block potassium conductances and a high [Na] to strongly activate the Na/K pump at internal sites.

## 2.6. Statistics

Whenever possible data are presented as mean  $\pm$  S.E.M. The S.E.M. is only shown in the figures when it exceeds the size of the symbol.  $n$  indicates the number of cells studied. Differences between data points were checked by Student's two-tailed, unpaired  $t$ -test and considered significant if  $P < 0.05$ .

## 3. Results

### 3.1. The endogenous $I_p$

The Na/K pump is an ATP-consuming enzyme, that transports 3 Na ions out of and 2 K ions into the cell per each ATP molecule split. The net movement of one positive charge out of the cell during each Na/K pump cycle thus produces an outward current. The Na/K pump current  $I_p$  depends on  $[K]_o$  and can be blocked by cardiac glycosides, specific inhibitors of the Na/K pump. Fig. 1 shows the identification of the Na/K pump current of HEK293 cells (endogenous  $I_p$ ). The holding potential  $V_C$  is 0 mV. Horizontal lines above the current trace indicate changes of the superfusion solution. In Fig. 1A, a HEK293 cell is initially superfused with extracellular solution, that contains 144 mM Na and 5.4 mM K. Switching to K-free solution (0 K) immediately shifts the holding current by about 20 pA in the inward direction. The following application of  $10^{-4}$  M ouabain results in an inward current shift of the same amplitude but with a slower time course. Fig. 1B displays a similar experiment with another HEK293 cell. At the beginning the cell is superfused with a solution containing 5.4 mM K. K-free medium instantaneously blocks an outward current of 10 pA.  $2 \cdot 10^{-4}$  M DHO more slowly blocks an outward current component of the same amplitude. As can be seen in Fig. 1A,B, under the chosen experimental conditions the cardiac glycoside-sensitive outward current is virtually identical to the current activated by  $K_o$ . This outward current is the current generated by the Na/K pump. However, in contrast to K-free medium the block of  $I_p$  by cardiac glycosides was nearly irreversible. Even after 5–10 min in cardiac glycoside free solution, there was no recovery of  $I_p$ . The  $I_p$  density at 5.4 mM  $K_o$  and 0 mV amounted to  $0.33 \pm 0.02$  pA  $\cdot$  pF $^{-1}$  ( $n = 28$ ).




Fig. 1. Identification of the endogenous  $I_p$  in HEK293 cells.  $I_p$  is identified as current inhibited by cardiac glycosides or K-free solution. In this and the following figures solution changes are indicated by horizontal lines above the current traces and the horizontal calibration mark denotes zero current level. The holding potential  $V_C$  is 0 mV. (A) Initially, the superfusate contains 5.4 mM  $K_o$ . Application of K-free solution or of a medium containing  $10^{-4}$  M ouabain (5.4 mM  $K_o$ ) results in an identical inward shift of the membrane current. Thus, the amplitudes of  $I_p$  estimated by either procedure are identical. (B) Different cell. The  $I_p$  amplitudes estimated by application of a K-free medium or a solution containing  $2 \cdot 10^{-4}$  M dihydro-ouabain (DHO; 5.4 mM  $K_o$ ) are almost the same.

The following experiments were all performed at a holding potential of 0 mV, except for the measurements of the Na/K pump current-voltage ( $I_p$ -V) relationships.

The concentration-dependent inhibition of the endogenous  $I_p$  by DHO is illustrated in Fig. 2. Fig. 2A shows the effect of  $5 \cdot 10^{-6}$  M DHO on the endogenous  $I_p$  of a HEK293 cell. At the beginning of the record the cell is superfused with a solution containing 5.4 mM K. Solution changes are indicated by the bars above the current trace. Switching to K-free solution reveals an  $I_p$  amplitude of 12 pA. A [DHO] of  $5 \cdot 10^{-6}$  M slowly shifts the holding current by 2.5 pA in the inward direction, i.e., it inhibits 21% of the initial  $I_p$ . Changing the superfusate to K-free solution during the steady-state inhibition by DHO causes a further inward shift of the current by 9.5 pA, corre-




Fig. 2. Inhibition of  $I_p$  by DHO in native HEK293 cells. (A) Original record. At the beginning the cell is superfused with a solution containing 5.4 mM  $K_o$ .  $I_p$  amplitudes are estimated by short pulses of K-free solution. Application of  $5 \cdot 10^{-6}$  M DHO (with 5.4 mM  $K_o$ ) reduces the initial  $I_p$  amplitude (12 pA) by 2.5 pA or 21%. (B) Concentration–response curve of steady-state  $I_p$  inhibition by DHO (with 5.4 mM  $K_o$ ). Complete inhibition of  $I_p$  in a K-free medium is arbitrarily set to 1. Data are shown as means  $\pm$  S.E.M. ( $n \geq 3$ ). Error bars are given only if they exceed the size of the symbols. The sigmoid curve fitted to the data represents a logarithmic version of the Hill equation. Half maximal inhibition of  $I_p$  ( $EC_{50}$ ) occurs at  $1.1 \cdot 10^{-5}$  M DHO with a Hill coefficient  $n_H$  of 1.2.

sponding to 79% of the initial  $I_p$  amplitude. The application of various  $[DHO]$  on 21 HEK293 cells resulted in the concentration–response curve presented in Fig. 2B. The relative steady-state inhibition of  $I_p$  by DHO is plotted versus the logarithm of the  $[DHO]$ . Each data point represents the mean of at least three measurements. The sigmoid curve fitted to the data obeys a logarithmic version of the Hill equation. According to this fit half maximal  $I_p$  inhibition ( $EC_{50}$ ) occurs at  $1.1 \cdot 10^{-5}$  M DHO with a Hill coefficient  $n_H$  of 1.2, in line with the notion that one DHO molecule binds to one Na/K pump molecule.

We next studied the dependence of the endogenous  $I_p$  on  $[K]_o$ . For this purpose solutions containing various  $[K]$  were applied to the cells. An original record of such an experiment is shown in Fig. 3A. A

HEK293 cell is superfused with solutions containing 0, 1.35, 2.7, or 5.4 mM K, as indicated by the lines above the current trace. The membrane current is shifted in the outward direction with increasingly higher  $[K]_o$ . The difference in holding current between K-containing and K-free solution represents the  $I_p$  amplitude at the respective  $[K]_o$ .  $I_p$  amplitudes were normalized to the amplitude at 5.4 mM  $K_o$  and plotted versus  $[K]_o$ . The result is shown in Fig. 3B. Each data point is the mean of 2–12 measurements. The curve fitted to the data points represents the Hill equation:

$$I_p = I_{p(max)} \cdot [K]_o^{n_H} / (K_{0.5}^{n_H} + [K]_o^{n_H}). \quad (1)$$

The maximal Na/K pump current ( $I_{p(max)}$ ) is 1.21, half-maximal activation of  $I_p$  occurs at 1.5 mM  $K_o$  ( $K_{0.5}$ ), and the Hill coefficient  $n_H$  amounts to 1.2.




Fig. 3. Activation of the endogenous  $I_p$  by  $K_o$ . (A) Original record. Increasing external K concentrations (horizontal bars) evoke an increasing outward component of membrane current (bottom trace). Note that the  $K_o$ -activated outward current tends to saturate at high  $[K]_o$ . (B) Normalized concentration–response curve of  $I_p$  activation by  $K_o$ .  $I_p$  amplitudes were normalized to the amplitude at 5.4 mM  $K_o$ , which was arbitrarily set to 1. Numbers in parentheses indicate the number of cells studied. The curve fitted to the data obeys the Hill equation (Eq. (1)):  $I_{p(max)} = 1.21$ ;  $K_{0.5} = 1.5$  mM  $K_o$ ;  $n_H = 1.2$ .

It is well known that  $I_p$  is voltage-dependent (review: Ref. [12]). Fig. 4 illustrates the voltage-dependence of the endogenous  $I_p$ , which was determined by means of the protocol depicted in Fig. 4A. The upper trace in Fig. 4A represents the voltage proto-




Fig. 4.  $I_p$ - $V$  curve of native HEK293 cells. (A) Estimation of  $I_p$  at various membrane potentials (top line). The superfusate contains 5.4 mM  $K_o$ .  $I_p$  is measured as current reactivated by  $K_o$  after a short pulse of a K-free medium, indicated by the short lines above the current trace (bottom). During the application of K-free solution the membrane current is shifted in the inward direction at each potential tested. (B) Enlarged current traces from (A), ordered by membrane potential. Bars indicating changes to K-free medium are left.  $I_p$  is estimated as current reactivated by 5.4 mM  $K_o$  after superfusion with K-free medium.  $I_p$  increases with depolarization. (C) Normalized mean  $I_p$ - $V$  relationship of native HEK293 cells.  $I_p$  amplitudes are normalized to the corresponding amplitude at zero potential. Data represent means  $\pm$  S.E.M. ( $n = 4$ –15). The inset shows the normalized  $I_p$ - $V$  curve obtained in the experiment illustrated in (A) and (B).

col, whereas the lower trace shows the resulting membrane current of a HEK293 cell. The cell was clamped to voltages between  $-80$  and  $+60$  mV in 20 mV steps. Changes in the membrane potential result in capacitative artifacts visible as rapid upward and downward deflections in the current trace, respectively. The cell is continuously superfused with a solution containing 5.4 mM K. During each clamp potential the superfusate is temporarily changed to K-free solution, as marked by the horizontal bars above the current trace. K-free solution shifts the membrane current in the inward direction at each potential tested. This is shown in further detail in Fig. 4B. Magnified current traces from Fig. 4A are presented. They were obtained during superfusion with K-free solution. For clarity bars indicating changes to K-free solution are left. Two traces at zero potential are shown, the first from the beginning and the second from the end of the record in Fig. 4A. Since the difference current at this potential remains constant, changes of the seal resistance during the record can be excluded. It is clear from Fig. 4B that the difference in membrane current between K-containing and K-free solution increases with increasing membrane potential. This difference current represents  $I_p$  because K-activated and cardiac glycoside-sensitive current are identical under the present experimental conditions (cf. Fig. 1). Control experiments (not shown) confirmed that this is not only true for 0 mV but for all potentials in the voltage range studied.  $I_p$  amplitudes were normalized to the amplitude at 0 mV and plotted versus membrane potential to yield the  $I_p$ - $V$  relationship shown in the inset in Fig. 4C. Normalized  $I_p$  amplitudes range from 0.41 at  $-80$  mV to 1.32 at  $+60$  mV. The  $I_p$ - $V$  curve is nearly linear over this voltage range. The main Fig. 4C illustrates the mean  $I_p$ - $V$  relationship for the endogenous  $I_p$ . Mean values of 4–15 measurements for each potential are presented. Normalized  $I_p$  amplitudes are  $0.55 \pm 0.03$  ( $n = 10$ ) at  $-80$  mV and  $1.38 \pm 0.05$  ( $n = 4$ ) at  $+60$  mV. This mean  $I_p$ - $V$  curve is almost linear over the entire voltage range studied.

### 3.2. The exogenous $I_p$

After the characterization of the cardiac glycoside-sensitive endogenous  $I_p$ , we investigated

whether a cardiac glycoside-resistant  $I_p$  could be detected in HEK293 cells transfected with the  $\alpha 1$  subunit of the rat Na/K pump. Fig. 5 displays the membrane current of a HEK293 cell transfected with the  $\alpha 1$  subunit of the rat Na/K pump and cultured for 44 days in a medium containing  $10^{-4}$  M DHO. This [DHO] blocks the endogenous  $I_p$  (cf. Fig. 2B). The horizontal lines above the current trace indicate changes of the superfusate. At the beginning of the record the cell is superfused with a solution containing 5.4 mM  $K_o$  plus  $2 \cdot 10^{-4}$  M DHO. As can be seen from Fig. 5 a K-free solution or a solution containing 5.4 mM K plus  $10^{-2}$  M ouabain block an almost identical outward component of the membrane current. Thus, a Na/K pump current is present under conditions which completely block the endogenous  $I_p$ . The density of this Na/K pump current amounted to  $0.16 \pm 0.02$  pA  $\cdot$  pF $^{-1}$  ( $n = 11$ ). In contrast to the irreversible blockade of the endogenous  $I_p$  by ouabain or DHO the  $I_p$  inhibition by ouabain in transfected cells is reversible.  $I_p$  recovers in the presence of a solution containing  $2 \cdot 10^{-4}$  M DHO. Obviously, the transfected cells expressed Na/K pumps exhibiting a much lower ouabain sensitivity than the endogenous Na/K pumps. Assuming one-to-one binding of




Fig. 5. Blockade of the exogenous  $I_p$  by  $10^{-2}$  M ouabain is (partially) reversible. Original current record (bottom) from a HEK293 cell stably transfected with the rat  $\alpha 1$  subunit. The cell is continuously superfused with a medium containing 5.4 mM  $K_o$  plus  $2 \cdot 10^{-4}$  M DHO except for short periods of K-free or ouabain-containing ( $10^{-2}$  M) solution, as marked by the bars above the current trace.  $I_p$  is estimated either as current blocked by K-free solution or as current inhibited by  $10^{-2}$  M ouabain (5.4 mM  $K_o$ ). Both estimates are nearly identical. At the end of the record  $I_p$  amounts to  $\approx 75\%$  of the initial amplitude. The apparent increase and decrease, respectively, of the current noise in passing zero current level (dashed line) is an artifact due to the data acquisition programme.

ouabain to the Na/K pump an estimation of the apparent  $K_d$  value ( $K'_d$ ) for  $I_p$  inhibition by ouabain can be derived from the kinetics of ouabain-binding and -unbinding (for a detailed description of this procedure see Ref. [13]). Consequently,  $I_p$  inhibition and recovery were presumed to proceed monoexponentially. The time constant of  $I_p$  inhibition ( $\tau$ ) amounted to 1.3 s, whereas that for  $I_p$  recovery ( $\tau_{off}$ ) was 64.9 s. Hence, the  $K'_d$  value was calculated to be  $2.0 \cdot 10^{-4}$  M. Similar results were obtained on two other cells ( $K'_d$  values: 2.5 and  $2.7 \cdot 10^{-4}$  M, respectively). The mean  $K'_d$  for the inhibition of the exogenous  $I_p$  by ouabain was  $(2.4 \pm 0.2) \cdot 10^{-4}$  M ( $n = 3$ ). Since rat Na/K pumps are known to be ouabain-insensitive, these results strongly suggest that the transfected cells expressed a functional Na/K pump which contained the rat  $\alpha 1$  subunit and displayed, therefore, the characteristic ouabain-resistance.

The  $K_o$ -dependence of the exogenous  $I_p$  was determined in media containing different  $[K]_o$  plus either  $10^{-4}$  M ouabain or  $2 \cdot 10^{-4}$  M DHO to block the endogenous  $I_p$ . Raising  $[K]_o$  resulted in an outward shift of the holding current (not shown).  $I_p$  was estimated as the difference in the holding current measured during application of a K-free solution.  $I_p$  amplitudes were normalized to the  $I_p$  amplitude at 5.4 mM  $K_o$  and plotted versus  $[K]_o$ . Fitting with the Hill equation (Eq. (1)) yielded a maximal  $I_p$  of 1.33, a  $K_{0.5}$  of 1.7 mM  $K_o$ , and a Hill coefficient of 1. These values hardly differ from those obtained for the endogenous  $I_p$  (cf. Fig. 3B).

The voltage dependence of the exogenous  $I_p$  in transfected HEK293 cells was studied as shown in Fig. 4 for the endogenous  $I_p$ . The cells were continuously superfused with solutions containing 5.4 mM  $K_o$  plus  $2 \cdot 10^{-4}$  M DHO or  $10^{-4}$  M ouabain, respectively.  $I_p$  was estimated as  $K_o$ -activated current at membrane potentials between  $-80$  and  $+60$  mV in 20 mV steps. For this purpose a K-free solution was applied to the cell under study for a few seconds at each membrane potential. Measurements on 6 cells (summarized in Fig. 7, open circles) revealed that  $I_p$  increased with depolarization at negative voltages. However, it remained essentially constant at positive potentials in contrast to the endogenous  $I_p$  of untransfected cells which increased linearly over the entire range of membrane potentials (Fig. 4C). Normalized mean amplitudes of the exogenous  $I_p$  varied between

$0.42 \pm 0.04$  at  $-80$  mV ( $n = 2$  only) and  $1.09 \pm 0.03$  at  $+60$  mV ( $n = 4$  only).

### 3.3. The contribution of the endogenous and exogenous $I_p$ to the total $I_p$ of transfected HEK293 cells

The contribution of endogenous and exogenous Na/K pumps to the total  $I_p$  of transfected HEK293 cells was estimated in the following way. First, the transfected cells were cultured for 2–3 days in a DHO-free medium in order to abolish the inhibition of the endogenous Na/K pumps which was induced by the DHO-containing culture medium. Afterwards the total  $I_p$  of the cells was estimated as  $K_o$ -activated current. The total  $I_p$  density amounted to  $0.20 \pm 0.03$  pA · pF<sup>-1</sup> ( $n = 7$ ). The total  $I_p$  was then inhibited in two steps. Application of a superfusate containing 5.4 mM K plus  $10^{-4}$  M ouabain completely blocked the endogenous  $I_p$  of the HEK293 cells. Subsequent superfusion with a solution containing 5.4 mM K plus  $10^{-2}$  M ouabain blocked the residual  $I_p$  which was most probably generated by pump molecules containing the exogenous  $\alpha 1$  subunit of the rat Na/K pump. Fig. 6 illustrates this procedure. Fig. 6A displays an original record from a HEK293 cell. Changes of the superfusate are indicated by the horizontal lines above the current trace. At the beginning of the experiment the total  $I_p$  is estimated as the current inhibited by K-free solution. It amounts to 15 pA. The application of  $10^{-4}$  M ouabain shifts the membrane current in the inward direction by 5.2 pA or 35% of the total  $I_p$ . The residual 65% (9.8 pA) of the total  $I_p$  are then blocked by  $10^{-2}$  M ouabain. As a mean from 7 cells  $37 \pm 3\%$  of the total  $I_p$  were inhibited by  $10^{-4}$  M ouabain. Therefore, at least  $\approx 63\%$  of the total pump current were generated by Na/K pumps containing the exogenous, ouabain-resistant rat  $\alpha 1$  subunit. If  $10^{-4}$  M ouabain blocked some of these pump molecules, the contribution of the exogenous pumps to the total  $I_p$  would be even larger. For comparison, similar measurements were carried out on rat ventricular myocytes. It is known from electrophysiological and biochemical studies [14,15] that about 70–80% of the Na/K pumps in these cells contain the ouabain-resistant  $\alpha 1$  subunit. Fig. 6B shows the membrane current of a rat ventricular cell. Initially, the total  $I_p$  is estimated as current inhibited by K-free solution.  $I_p$  amounts to 60 pA.




Fig. 6. Contribution of ouabain-insensitive  $I_p$  to the total  $I_p$  of transfected HEK293 cells and native rat ventricular myocytes. (A) Contribution of the exogenous, ouabain-resistant  $I_p$  to the total  $I_p$  in a stably transfected HEK293 cell. The cell was cultured for two days in DHO-free medium in order to abolish inhibition of the endogenous Na/K pumps. Original record of membrane current (bottom trace). Solution changes are indicated by the horizontal lines above the current trace. At the beginning of the record the cell is superfused with a medium containing 5.4 mM  $K_o$ . Total  $I_p$  is estimated twice as current blocked by K-free solution. Application of  $10^{-4}$  M ouabain (with 5.4 mM  $K_o$ ) diminishes the total  $I_p$  by 35%. The residual  $I_p$  is blocked by  $10^{-2}$  M ouabain (with 5.4 mM  $K_o$ ). Thus, the exogenous, ouabain-resistant  $I_p$  contributes at least 65% to the total  $I_p$  of the cell. (B) Contribution of ouabain-insensitive Na/K pump molecules to the total  $I_p$  of a rat ventricular myocyte. Application of different media is visualized by the bars above the current trace. Initially, the superfusate contains 5.4 mM  $K_o$ . Total  $I_p$  is measured as current blocked by K-free solution. A medium containing  $10^{-4}$  M ouabain (with 5.4 mM  $K_o$ ) inhibits 17% of the total  $I_p$ . Complete block of  $I_p$  occurs following application of  $10^{-2}$  M ouabain (with 5.4 mM  $K_o$ ). The ouabain-insensitive Na/K pumps contribute at least 83% to the total  $I_p$  of the myocyte. Dashed lines in (A) and (B) are drawn to facilitate the identification of the  $I_p$  components.

$10^{-4}$  M ouabain inhibits 10 pA or 17% of the total  $I_p$ . The remaining pump current is completely blocked by  $10^{-2}$  M ouabain. This component is most probably due to the ouabain-resistant pump molecules containing the  $\alpha 1$  subunit. Corresponding experiments



on 5 myocytes revealed that  $23 \pm 4\%$  of the total  $I_p$  were blocked by  $10^{-4}$  M ouabain. Thus, 77% (at least) of the total  $I_p$  were generated by pump molecules containing the ouabain-resistant  $\alpha 1$  subunit. These results are in quantitative agreement with the reports mentioned above. It is, however, difficult to exclude that a small portion of the pumps containing the  $\alpha 1$  subunit were already blocked by  $10^{-4}$  M ouabain.

### 3.4. $I_p$ -V relationships of transfected and not-transfected HEK293 cells and of rat ventricular myocytes

Fig. 7 displays mean  $I_p$ -V relationships of normal (●;  $n = 4-15$ ) and transfected (○;  $n = 4-6$ ,  $n = 2$  only at  $-80$  mV) HEK293 cells. For comparison the  $I_p$ -V curve of rat ventricular myocytes is also presented (□;  $n = 4-10$ ). For each series of experiments the  $I_p$  amplitudes measured at the various potentials were normalized to the respective amplitude at 0 mV (cf. Fig. 4). The pump current of the native (untransfected) HEK293 cells, i.e., the endogenous  $I_p$ , increases almost linearly with depolarization




Fig. 7. Normalized mean  $I_p$ -V relationships of HEK293 cells and rat ventricular myocytes. In each series of experiments the  $I_p$  amplitudes measured at the various membrane potentials are normalized to the corresponding amplitude at zero potential, arbitrarily set to 1. The endogenous  $I_p$  (●) of native HEK293 cells ( $n = 4-15$ ) exhibits a linear voltage dependence over the entire range of membrane potentials studied. In contrast, the  $I_p$ -V curves of transfected HEK293 cells (exogenous  $I_p$ : ○;  $n = 4-6$ ;  $n = 2$  only at  $-80$  mV) and of rat ventricular myocytes (□;  $n = 4-10$ ) show nearly constant  $I_p$  amplitudes at positive voltages. Both curves are almost identical. Asterisks mark significantly different  $I_p$  amplitudes ( $P < 0.05$ ).

over the entire voltage range studied. This is in contrast to the  $I_p$ -V relationship of the transfected HEK293 cells and to the  $I_p$ -V curve of the ventricular myocytes. Both curves are nearly identical. They show a positive slope at negative membrane potentials but little variation at positive voltages. The similarity between both  $I_p$ -V curves suggests once more that the transfected HEK293 cells express Na/K pumps which contain the rat  $\alpha 1$  subunit like the majority of the rat cardiac Na/K pump molecules [16].

## 4. Discussion

In recent years, molecular biological approaches enabled physiologists to examine structure-function relationships of the Na/K pump. Most studies, however, used biochemical or conventional voltage clamp methods to characterize recombinant Na/K pumps. A cell system suitable for the investigation of recombinant pump molecules by means of the patch clamp technique would be a powerful tool for the electrophysiological analysis of structure-function relationships of the Na/K pump. Therefore, the aim of the present study was to establish such a system. We chose HEK293 cells, a human cell line widely used for the expression of recombinant proteins. These cells are well suited for whole-cell recording, because of high seal resistances of up to  $60 \text{ G}\Omega$  and because of their relatively small size. Membrane capacitance ranged between 20 and 85 pF, in good agreement with an earlier report [1]. Little is known about the Na/K pump of these cells, so we first characterized the Na/K pump current of HEK293 cells (endogenous  $I_p$ ).

### 4.1. Characteristics of the endogenous $I_p$

The endogenous  $I_p$  was determined as current blocked by the cardiac glycosides ouabain or DHO.  $I_p$  density at 0 mV amounted to  $0.33 \text{ pA} \cdot \text{pF}^{-1}$  in a solution containing 5.4 mM  $\text{K}_o$ . This value is rather small compared to excitable cells like ventricular myocytes, where the  $I_p$  density is around  $1 \text{ pA} \cdot \text{pF}^{-1}$  or beyond [17]. Half maximal steady-state  $I_p$  inhibition occurred at  $1.1 \cdot 10^{-5}$  M DHO (5.4 mM  $\text{K}_o$ ) with a Hill coefficient of 1.2, suggesting one-to-one

binding to the Na/K pump as previously demonstrated by Hermans and co-workers [17]. Similar values were obtained in guinea pig ventricular myocytes ( $1.4 \cdot 10^{-5}$  M DHO (5.4 mM  $K_o$ ); [17]) or in cardiac Purkinje cells (canine:  $0.4 \cdot 10^{-5}$  M DHO (8 mM  $K_o$ ) [18]; rabbit:  $3.5 \cdot 10^{-5}$  M DHO (10.8 mM  $K_o$ ) [13]). In human endothelial cells half maximal steady-state inhibition of  $I_p$  occurred at  $2.1 \cdot 10^{-5}$  M DHO (11 mM  $K_o$ ) [19]. In the same study, a nearly irreversible  $I_p$  inhibition by DHO was observed. This was also the case for the endogenous  $I_p$  of the HEK293 cells. Similarly, the endogenous  $I_p$  of HeLa cells, a human cell line, was blocked nearly irreversibly by ouabain, digitoxin, and digoxin [5]. The reason for this nearly irreversible  $I_p$  blockade is not known. To our knowledge such an irreversible  $I_p$  inhibition has only been observed in the human cell types mentioned above raising the possibility that this is a property of human Na/K pumps.

Under our experimental conditions  $I_p$  (i.e., the outward current blocked by cardiac glycosides) was identical to the current activated by  $K_o$ . Half maximal activation of  $I_p$  occurred at 1.5 mM  $K_o$  with a Hill coefficient of 1.2. Comparable results were obtained for rat and guinea-pig ventricular myocytes [17,20,21], for rabbit sino-atrial node cells [22], for cardiac Purkinje cells [18], for human endothelial cells [19], and for various other cell types (cf. Ref. [23]). Obviously,  $K_o$ -activation of  $I_p$  does not differ substantially between different cell types and species.

The endogenous  $I_p$  was voltage dependent.  $I_p$  increased linearly over the whole voltage range studied, i.e., between  $-80$  and  $+60$  mV. This monotonic increase argues for only one voltage-dependent partial reaction in the pump cycle under the present experimental conditions (cf. Ref. [12]). The main voltage-dependent step is most probably located in the Na-translocating branch of the pump cycle [24,25] and is favoured by more positive membrane potentials. An  $I_p$ -V relationship similar to that in HEK293 cells was found in human endothelial cells [19].  $I_p$  increased linearly between  $-150$  and  $+50$  mV in these cells. Such linear  $I_p$ -V curves are in contrast, however, to those obtained in the majority of cell types, including various cardiac cells, where  $I_p$  increases linearly with depolarization at negative membrane potentials but shows little voltage dependence at positive potentials [12,22,26–28]. The lack of volt-

age dependence at positive potentials can be explained by assuming that Na translocation is no longer rate-determining at these voltages. For the linear  $I_p$ -V curves of HEK293 and human endothelial cells it means that Na translocation must be rate-determining even at positive potentials up to  $+60$  mV.

#### 4.2. The heterologous expression of Na/K pump molecules in various cells

The heterologous expression of cardiac glycoside-resistant Na/K pumps in cells expressing glycoside-sensitive endogenous pumps is an established method for analyzing the structure–function relationship of the Na/K-ATPase. For example, HeLa cells have been widely used for the expression of glycoside-resistant, mutated  $\alpha$  subunits of the rat or sheep Na/K pump in order to study the functional differences between various  $\alpha$  subunits and the significance of distinct amino acids for cation pumping (e.g., Refs. [29–31]). These studies were mainly carried out by means of biochemical methods. Similarly, *Xenopus laevis* oocytes represent a suitable expression system for recombinant Na/K pump molecules [32,33]. The properties of the exogenous pump molecules can be measured both by biochemical and conventional electrophysiological methods. In the present study HEK293 cells were chosen for the expression of Na/K pump molecules containing the cardiac glycoside-resistant rat  $\alpha 1$  subunit of the pump.

#### 4.3. The identification of exogenous Na/K pumps in HEK293 cells

We kept the transfected HEK293 cells for several weeks in media containing  $10^{-4}$  M DHO, which completely blocked the endogenous glycoside-sensitive Na/K pumps. Under these conditions a Na/K pump current was identified (Fig. 5), suggesting the functional expression of pump molecules containing the transfected glycoside-resistant  $\alpha 1$  subunit of the rat Na/K pump. Similar to the rat cardiac Na/K-ATPase (e.g., Ref. [14]) these pumps were blocked only by the extremely high ouabain concentration of  $10^{-2}$  M (Fig. 5). The functional expression of Na/K pumps containing the rat  $\alpha 1$  subunit implies the assembly of the exogenous  $\alpha$  and the endogenous  $\beta$  subunits of the molecules. Such an assembly has been

reported before from *Xenopus* oocytes [34], COS-1 cells [35], and HeLa cells [30]. The assembly of the rat  $\alpha 1$  subunit and the endogenous  $\beta$  subunit in HEK293 cells is not unexpected since human and rat  $\alpha$  subunits are nearly homologous [33]. The total  $I_p$  density of the transfected HEK293 cells amounted to  $0.20 \text{ pA} \cdot \text{pF}^{-1}$  or 61% of the  $I_p$  density of untransfected cells. Similarly, Putnam and co-workers [36] reported that the total ATPase activity of HeLa cells transfected with the rat  $\alpha 1$  subunit was about 60% of the controls. Since exogenous and endogenous  $\alpha$  subunits compete for assembly with the available endogenous  $\beta$  subunits, increased  $I_p$  density in the transfected cells would be surprising. Furthermore, the procedure of transfection per se tends to diminish the level of expressed pump molecules [35,37]. In contrast to these findings, however, is the observation of Yamamoto and co-workers [5] that the Na/K pump current densities in native and transfected HeLa cells were almost identical. Our results illustrated in Fig. 6A revealed that at least 63% of the total  $I_p$  density in transfected HEK293 cells are generated by pump molecules containing the rat  $\alpha 1$  subunit. This is in line with the observation that 51% of the Na/K-ATPase activity of COS-1 cells transfected with the rat  $\alpha 1$  subunit are caused by pumps containing the rat subunit [35]. According to Putnam and co-workers [36] less than 50% of the total Na/K-ATPase activity in HeLa cells transfected with various rat  $\alpha$  subunits are due to molecules containing the exogenous subunit. However, in HeLa cells transfected with a mutated sheep  $\alpha 1$  subunit up to 90% of the total Na/K-ATPase activity are produced by pumps containing the sheep subunit [29]. Finally, in HeLa cells transfected with a mutated human or sheep  $\alpha 1$  subunit the whole pump current is apparently generated by recombinant pump molecules [5].

#### 4.4. Characteristics of the exogenous $I_p$

In contrast to the inhibition of  $I_p$  by ouabain in native HEK293 cells the blockade of  $I_p$  by the cardiac glycoside in transfected cells was reversible. Estimation of the apparent  $K_d$  value ( $K'_d$ ) from the kinetics of ouabain-binding and -unbinding yielded  $2.4 \cdot 10^{-4} \text{ M}$ , a concentration that completely blocks the endogenous  $I_p$ . Since both the time constant for solution exchange (200–600 ms) as well as that for  $I_p$

inhibition (1.1–1.4 s) are similar, the calculated  $K'_d$  value has to be considered a rough estimate that tends to overestimate the ‘correct’  $K'_d$ . Despite this limitation, the value obtained by our procedure agrees fairly well with the  $K'_d$  value ( $3 \cdot 10^{-4} \text{ M}$ ) reported by Berlin and co-workers [14] for the inhibition of  $I_p$  by ouabain in rat ventricular myocytes under slightly different conditions (15 mM  $\text{Na}_{\text{pip}}$ ; 15 mM  $\text{K}_o$ ). Both the reversibility and the  $K'_d$  value of the  $I_p$  inhibition by ouabain clearly demonstrate the expression of pump molecules containing the rat  $\alpha 1$  subunit in the transfected HEK293 cells. Similarly, the shape of the  $I_p$ -V curve in transfected HEK293 cells strongly suggests the functional expression of pumps with the rat subunit. Like the  $I_p$ -V relationship of rat ventricular myocytes, the  $I_p$ -V curve of transfected HEK293 cells displays little variation of  $I_p$  at positive membrane potentials, whereas the  $I_p$ -V relationship of native HEK293 cells is almost linear over the entire range of potentials tested (Fig. 7). However,  $I_p$  activation by  $\text{K}_o$  is similar in native and transfected HEK293 cells. The  $[\text{K}]_o$  required for half maximal  $I_p$  activation amounted to 1.5 mM in native cells and to 1.7 mM in transfected cells. Comparable values are reported for rat ventricular myocytes ([20]: 2.9 mM  $\text{K}_o$ ; [17]: 1 mM  $\text{K}_o$ ).

In conclusion, the shape of the  $I_p$ -V relationship and the effect of ouabain on  $I_p$ , but not the kinetics of  $I_p$  activation by  $\text{K}_o$  can be used to differentiate between endogenous Na/K pumps and exogenous pump molecules containing the rat  $\alpha 1$  subunit in HEK293 cells. The different ouabain-sensitivity and voltage dependence of the rat Na/K pump as well as the activation of the rat  $I_p$  by  $\text{K}_o$  are well conserved in the recombinant Na/K pumps in HEK293 cells, demonstrating that this cell line is suited for the expression and whole-cell patch clamp analysis of the ouabain-resistant  $\alpha 1$  subunit of the rat Na/K pump.

#### Acknowledgements

We are indebted to Professor Jerry Lingrel (Cincinnati) for a generous gift of the plasmid pR  $\alpha 1$ . The authors would like to thank Professors Barry W. Ache (Daytona) and Wolfgang Schwarz (Frankfurt) for helpful comments on an earlier draft of the manuscript.

**References**

- [1] Galli, A., DeFelice, L.J., Duke, B.-J., Moore, K.R. and Blakely, R.D. (1995) *J. Exp. Biol.* 198, 2197–2212.
- [2] Perez-Garcia, M.T., Kamp, T.J. and Marban, E. (1995) *J. Gen. Physiol.* 105, 289–306.
- [3] Hamill, O.P., Marty, A., Neher, E., Sakmann, B. and Sigworth, F.J. (1981) *Pflügers Arch.* 391, 85–100.
- [4] Soejima, M. and Noma, A. (1984) *Pflügers Arch.* 400, 424–431.
- [5] Yamamoto, S., Askew, G.R., Heiny, J., Masaki, H. and Yatani, A. (1996) *Am. J. Physiol.* 270, C457–C464.
- [6] Kockskämper, J., Gisselmann, G. and Glitsch, H.G. (1996) *Pflügers Arch.* 431, R106.
- [7] Van Huysse, J.W., Jewell, E.A. and Lingrel, J.B. (1993) *Biochemistry* 32, 819–826.
- [8] Gorman, C.M., Gies, D.R. and McRay, G. (1990) *DNA Prot. Eng. Tech.* 2, 3–10.
- [9] Bechem, M., Pott, L. and Rennebaum, H. (1983) *Eur. J. Cell Biol.* 31, 366–369.
- [10] Banach, K., Hüser, J., Lipp, P., Wellner, M.-C. and Pott, L. (1993) *J. Physiol.* 461, 263–281.
- [11] Fabiato, A. and Fabiato, F. (1979) *J. Physiol. (Paris)* 75, 463–505.
- [12] De Weer, P., Gadsby, D.C. and Rakowski, R.F. (1988) *Annu. Rev. Physiol.* 50, 225–241.
- [13] Bielen, F.V., Glitsch, H.G. and Verdonck, F. (1992) *Naunyn-Schmiedeberg's Arch. Pharmacol.* 345, 100–107.
- [14] Berlin, J.R., Fielding, A.J. and Ishizuka, N. (1992) *Ann. N.Y. Acad. Sci.* 671, 440–442.
- [15] Lucchesi, P.A. and Sweadner, K.J. (1991) *J. Biol. Chem.* 266, 9327–9331.
- [16] Sweadner, K.J. (1989) *Biochim. Biophys. Acta* 988, 185–220.
- [17] Hermans, A.N., Glitsch, H.G. and Verdonck, F. (1995) *J. Physiol.* 484, 617–628.
- [18] Cohen, I.S., Datyner, N.B., Gintant, G.A., Mulrine, N.K. and Pennefather, P. (1987) *J. Physiol.* 383, 251–267.
- [19] Oike, M., Droogmans, G., Casteels, R. and Nilius, B. (1993) *Pflügers Arch.* 424, 301–307.
- [20] Kinard, T.A., Liu, X.-Y., Liu, S. and Stimers, J.R. (1994) *Am. J. Physiol.* 266, C37–C41.
- [21] Nakao, M. and Gadsby, D.C. (1989) *J. Gen. Physiol.* 94, 539–565.
- [22] Sakai, R., Hagiwara, N., Matsuda, N., Kasanuki, H. and Hosoda, S. (1996) *J. Physiol.* 490, 51–62.
- [23] De Weer, P. (1992) in *The Kidney: Physiology and Pathophysiology* (Seldin, D.W. and Giebisch, G., eds.), pp. 93–112. Raven Press, New York.
- [24] Hilgemann, D.W. (1994) *Science* 263, 1429–1432.
- [25] Nakao, M. and Gadsby, D.C. (1986) *Nature* 323, 628–630.
- [26] Gadsby, D.C., Kimura, J. and Noma, A. (1985) *Nature* 315, 63–65.
- [27] Glitsch, H.G., Krahn, T. and Pusch, H. (1989) *Pflügers Arch.* 414, 52–58.
- [28] Stimers, J.R., Liu, S. and Kinard, T.A. (1993) *J. Membr. Biol.* 135, 39–47.
- [29] Argüello, J.M. and Lingrel, J.B. (1995) *J. Biol. Chem.* 270, 22764–22771.
- [30] Jewell, E.A. and Lingrel, J.B. (1991) *J. Biol. Chem.* 266, 16925–16930.
- [31] Jewell-Motz, E.A. and Lingrel, J.R. (1993) *Biochemistry* 32, 13523–13530.
- [32] Schwarz, W. and Gu, Q. (1988) *Biochim. Biophys. Acta* 945, 167–174.
- [33] Vasilets, L.A. and Schwarz, W. (1993) *Biochim. Biophys. Acta* 1154, 201–222.
- [34] Jaunin, P., Jaisser, F., Beggah, A.T., Takeyasu, K., Mangeat, P., Rossier, B.C., Horisberger, J.-D. and Geering, K. (1993) *J. Cell Biol.* 123, 1751–1759.
- [35] Shanbaky, N.M. and Pressley, T.A. (1995) *Biochem. Cell Biol.* 73, 261–268.
- [36] Putnam, D.S., Jewell, E.A., Lescale-Matys, L., Magyar, C.E. and McDonough, A.A. (1994) in *The Sodium Pump: Structure Mechanism, Hormonal Control, and its Role in Disease* (Bamberg, E. and Schoner, W., eds.), pp. 234–237. Steinkopff, Darmstadt.
- [37] Kaji, D.M., Bates, J., Goyzueta, J.D., Prasadan, K., Yu, H. and Kumar, S. (1996) *J. Membr. Biol.* 149, 49–55.