

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Journal of Pediatric Surgery CASE REPORTS

journal homepage: www.jpascasereports.com

A right colonic volvulus requiring extensive colectomy in an infant with trisomy 13



Kazuto Suda*, Hajime Kawakami, Takato Sasaki, Miki Ishikawa, Miki Toma, Toshihiro Yanai, Toshihiro Muraji

Department of Pediatric Surgery, Ibaraki Children's Hospital, Ibaraki, Japan

ARTICLE INFO

Article history:

Received 30 July 2015
Received in revised form
20 October 2015
Accepted 20 October 2015

Key words:

Colonic volvulus
Neurodevelopmental delay
Trisomy 13

ABSTRACT

Colonic volvulus is a rare surgical emergency condition in children. Only approximately 40 children with cecal volvulus have been reported in English literature in the past 50 years. Among these, a right colonic volvulus involving the long segment from the ileal end to the transverse colon, as in our case, is limited to a few reports. Neurodevelopmental delay and a history of chronic constipation have been reported as common associated disorders. This is the first report about a case of right colonic volvulus in an infant with trisomy 13 who required extensive colectomy during an emergency laparotomy.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Colonic volvulus is an unusual cause of intestinal obstruction, and it is responsible for approximately 3–5% of all colonic obstructions in adults. In the pediatric population, colonic volvulus has a lower incidence and is extremely rare [1–6]. However, common disorders associated with colonic volvulus include neurodevelopmental delay, a history of chronic constipation, and aerophagia [1,2,4–8]. The purpose of this report was to describe the unusual occurrence of right colonic volvulus in a patient with trisomy 13, and consider the precipitating factors.

1. Case report

A 1-year-old female infant with trisomy 13 had sudden onset of abdominal distension and continuous vomiting. On admission to our hospital for close evaluation, it was difficult to assess whether she presented with an acute abdomen because of lack of adequate communication. She presented with a high fever of 39 °C, neutrophilia, and signs of mild dehydration, with no elevation of the C-reactive protein level or white blood cell count. The initial abdominal radiograph showed a dilated left bowel (Fig. 1). After conservative treatment for a diagnosis of chronic constipation,

aerophagia, and bacterial enteritis, her condition worsened. An enhanced computed tomography (CT) scan showed a dilated bowel from the transverse to the sigmoid colon, indicating a left colonic obstruction. An internal hernia and intussusception were ruled out (Fig. 2a and b). Emergency laparotomy was required, with suspicion of transverse colonic volvulus; a contrast enema demonstrated poor flow of gastrografin into the transverse colon (Fig. 3), and blood testing suggested intestinal necrotic changes with elevated myogenic enzymes (Table 1).

During surgery, a transverse abdominal incision was made. The cecum and ascending colon were surprisingly located on the left side in the abdominal cavity, with fecal impaction throughout (Fig. 4a). Necrotic changes from the ileal end to the transverse colon, due to right colonic volvulus with a nonrotation form of malrotation, were found (Fig. 4b). The necrotic, dilated bowel was resected, and primary anastomosis was performed after removing the volvulus (Fig. 5). A midgut volvulus was ruled out because the small intestine was intact. Histopathological examination showed total necrosis of the resected bowel, with no evidence of chronic inflammation. The patient is presently doing well and is asymptomatic 4 months postoperatively. No complication was noted at this follow-up.

2. Discussion

With malformation of the mesentery said to be the primary etiology of colonic volvulus [9], the detailed factors are as follows.

* Corresponding author. Department of Pediatric Surgery, Ibaraki Children's Hospital, 3-3-1 Futabadai, Mito, Ibaraki 311-4145, Japan. Tel.: +81 29 254 1151; fax: +81 29 254 2382.

E-mail address: kdsuda@juntendo.ac.jp (K. Suda).



Fig. 1. An abdominal radiograph showing an extensively dilated left bowel; gas in the rectum is indistinct.




Fig. 3. Contrast enema showing poor flow of gastrografin into the transverse colon (arrowheads).

The possible congenital factors are a redundant and mobile colon, narrow base of the mesenteric root, elongation of the mesentery with a freely mobile bowel, and absent ligamentous attachments at the splenic or hepatic flexures, according to some reports [2,3,5,9]. In contrast, precipitating acquired factors are reported to be neurodevelopmental delay, severe chronic constipation, and aerophagia [1,2,4–8]. The coexistence of these factors causes chronic bowel distension, and inadequate treatment increases the incidence of colonic volvulus. Folaranmi et al. [1] hypothesized that these acquired factors may promote stretching of the ligaments responsible for fixation. By the time the mesentery of the colon becomes longer, a volvulus can occur.

In our case, several acquired risk factors for colonic volvulus, including neurodevelopmental delay, were also present. In addition to providing conservative treatment for a diagnosis of

chronic constipation, aerophagia, and bacterial enteritis, midgut volvulus was suspected as the cause of an acute abdomen because malrotation is presumed to be a major complication in children with trisomy 13. However, this was excluded by imaging. At one point, the influence of a nonrotation form of malrotation on causing a colonic volvulus was presumed. Nonrotation may be a risk factor of colonic volvulus because the cecum and ascending colon were shifted to the left side in the abdominal cavity in our case. In addition, the lack of broad colonic fixation to the retroperitoneum and the absence of ligamentous attachments at the hepatic and splenic flexures may be factors that influence the range of the strangulated intestinal segment. Berger et al. [10] also reported a case of volvulus of the ascending colon as an unusual complication of nonrotation of the midgut. Thus, some




Fig. 2. a and b) An enhanced computed tomography image showing a dilated transverse and descending colon, and dilated intestinal tract wall. The small intestine is not dilated.

Table 1

Pre/postoperative changes in the hematologic tests.

	WBC ($\times 10^3/\mu\text{L}$)	Hb (g/dL)	Plt ($\times 10^3/\mu\text{L}$)	CPK (IU/L)	LDH (IU/L)	CRP (mg/dL)
Preoperative	5.5	8.8	141	4949	1849	34.66
Postoperative day 5	13.3	8.7	133	133	378	2.87

WBC: white blood cells; Hb: hemoglobin; Plt: platelet; CPK: creatine phosphokinase; LDH: lactate dehydrogenase; CRP: C-reactive protein.

disorders, which are often complicated with malrotation, need to be monitored carefully, such as chromosome aberration, situs inversus, and diaphragm hernia. Although there are few reports about a correlation between trisomy 13 and colonic volvulus, surgeons should be careful not to delay diagnosis and emergency treatment for both midgut volvulus and extensive colonic volvulus in patients with neurodevelopmental delay, especially with trisomy 13.

In enhanced CT, a localized bowel loop with poor contrasting effects in parts of the colon can be a decisive finding to diagnose a colonic volvulus. Although this finding in itself will indicate the need of an emergency laparotomy, surgeons at least perform a contrast enema immediately, as mentioned below. There are two radiological features specific for colonic volvulus suggestive of the diagnosis on contrast enema. The coffee bean sign on abdominal radiography is highly suggestive of colonic volvulus, although it is present in only about 30% of cases [9]. The typical finding on contrast enema is a narrowed, twisted colon with a bird's-beak deformity [1,2,4]. However, a contrast enema is not always useful for patients with an acute abdominal emergency, as it may only delay definitive surgical treatment of the volvulus [4,9]. In our case, colonic volvulus became suspect as the diagnosis with poor flow of gastrografin into the transverse colon in a contrast enema, although the study did not demonstrate a typical bird's-beak deformity. We presume that extensive right colonic volvulus with a nonrotation form of malrotation is difficult to evaluate by imaging examination because this condition leads to complete reversal of the lateral axis of anatomical location. The dilated left bowel, seen on preoperative enhanced CT, was in fact the strangulated right colon (not the left colon), which was quite an unexpected discovery at surgery.




Fig. 4. a and b) On laparotomy, the cecum, ascending colon, and appendix (arrowheads) are surprisingly located on the left side in the abdominal cavity. At this point, necrotic changes from the ileal end to the transverse colon, due to right colonic volvulus with a nonrotation form of malrotation, can be recognized. The descending colon (dc) was intact.




Fig. 5. The descending colon (dc) and ileal end (ie) were anastomosed after removing the volvulus.

An endoscopic approach and contrast enema have been used as conservative methods to resolve colonic volvulus. Mellor et al. [3] noted that endoscopic reduction has been successful in adults, but it has not been used as a treatment of choice in children. This is due to lack of experience because of the infrequency of colonic volvulus in children. Some have reported that detorsion of the volvulus may occur during a contrast enema [2,3,7]. However, the risk of intestinal perforation will not permit reduction with a contrast enema in a situation in which necrotic change is assumed because high hydrostatic pressure would be needed to resolve a right colonic volvulus.

3. Conclusions

In summation, a rapid decision on laparotomy is necessary when acute abdomen with an extensive right colonic volvulus is assumed if patients with neurodevelopmental delay present with vague findings on physical examination. Additionally, great caution is suggested for an evolving right colonic volvulus when patients with neurodevelopmental delay, especially with trisomy 13, are suspected to be acutely symptomatic.

Conflicts of interest

None.

References

- [1] Folaranmi SE, Cho A, Tareen F, Morabito A, Rakoczy G, Cserni T. Proximal large bowel volvulus in children: 6 new cases and review of the literature. *J Pediatr Surg* 2012;47:1572–5.
- [2] Hsueh CC, Jaw TS, Lin JY, Chan HM. Splenic flexure colonic volvulus: a pediatric case report. *Kaohsiung J Med Sci* 2007;23:207–10.
- [3] Mellor MF, Drake DG. Colonic volvulus in children: value of barium enema for diagnosis and treatment in 14 children. *AJR Am J Roentgenol* 1994;162:1157–9.
- [4] Takada K, Hamada Y, Sato M, Fujii Y, Teraguchi M, Kaneko K, et al. Cecal volvulus in children with mental disability. *Pediatr Surg Int* 2007;23:1011–4.
- [5] Samuel M, Boddy SA, Nicholls E, Capps S. Large bowel volvulus in childhood. *Aust N Z J Surg* 2000;70:258–62.
- [6] Mansoor K, Al Hamidi S, Khan AM, Samujh R. Rare case of pediatric cecal volvulus. *J Indian Assoc Pediatr Surg* 2009;14:110–2.
- [7] Trillis Jr F, Gauderer MW, Ponsky JL, Morrison SC. Transverse colon volvulus in a child with pathologic aerophagia. *J Pediatr Surg* 1986;21:966–8.
- [8] Jancar J, Speller CJ. Fatal intestinal obstruction in the mentally handicapped. *J Intellect Disabil Res* 1994;38:413–22.
- [9] Mulas C, Bruna M, Garcia-Armengol J, Roig JV. Management of colonic volvulus. Experience in 75 patients. *Rev Esp Enferm Dig* 2010;102:239–48.
- [10] Berger RB, Hillemeier AC, Stahl RS, Markowitz RI. Volvulus of the ascending colon: an unusual complication of non-rotation of the midgut. *Pediatr Radiol* 1982;12:298–300.