

amyloidogenesis progresses much slower, on the days and weeks timescale. As conditions are changed from folding to misfolding, formation of the native structure slows down indicating the increase of the barrier separating the molten globule and native states. In the meantime, the native state becomes more unstable as well. Amyloid formation is only observed among solvent conditions where folding is absent.

2946-Pos Board B51

Probing Aggrecan Interactions by Atomic Force Microscopy

Preethi L. Chandran, Emiliós K. Dimitriadis, Peter J. Bassar, Ferenc Horkay.

Aggrecan, the major extracellular matrix proteoglycan in cartilage, is a highly charged bottlebrush shaped macromolecule. It consists of negatively charged glycosaminoglycan (GAG) chains attached to a protein backbone. The bottlebrush structure enables aggrecan to maintain an extended conformation responsible for the high osmotic pressure sustaining compressive loads in cartilage. Alterations in aggrecan bottlebrush structure with age and disease lead to bone deformities, dwarfism, arthritis, and other pathological conditions. In solution, aggrecan bottlebrushes show distinct osmotic pressure versus concentration regimes. They self-assemble and form large clusters. Using Atomic Force Microscopy aggrecan molecules adsorbed on controlled mica surfaces were imaged. On positively charged APS mica, the average extension and height of the bottlebrush side chains were practically unaffected by the presence of calcium ions. With increasing aggrecan concentration transition takes place from dispersed, non-interacting bottlebrushes to clusters of conforming chains. At higher concentrations aggrecan molecules form a continuous monolayer. These surface observations are consistent with aggrecan properties in solution. On negatively charged mica, aggrecan shows interesting network patterns at higher concentrations. Understanding aggrecan adsorption onto charged surfaces provides insight into its interactions with bone and implants in the biological milieu.


2947-Pos Board B52

Electrostatic Analysis of the Aggregation of TrV Viral Particles

Marcelo D. Costabel, María J. Amundarain, Fernando Zamarreño, Rubén Sánchez-Eugenia, Jon Agirre, Diego M.A. Guérin.

Triatoma virus (TrV) is a member of the insect virus family Dicistroviridae, a family similar to the vertebrate picornaviruses. TrV virions consist of a non-enveloped capsid that encloses the viral genome, a molecule of linear positive sense single-stranded RNA of about 9000 bases. The atomic structure of the capsid is an icosahedron ($T=1$, pseudo-equivalence $p=3$) of about 30 nm in diameter that encloses 60 repeats of three structural proteins VP1, VP2, and VP3 (MWs of about 39 kDa, 37 kDa, and 33 kDa; RCSB PDB code 3NAP). TrV particles stand very acidic conditions (pHs lower than 3.0) and at high concentration undergo a reversible aggregation. In this work we calculate the electrostatic energy for the interaction between two TrV particles at different solvent conditions, and we compare these results with the corresponding experimentally measured Static Light Scattering Second Virial Coefficient. The interaction mechanism does not appear to be dominated by an electrostatic effect of the charge at the capsid surface, and a dependence on the solvent ionic strength is observed. Keywords: Dicistroviridae; Triatoma virus; electrostatic energy; virus aggregation; Second Virial Coefficient.

Figure: TrV surface charge at pH 4.0.



Protein Folding & Stability III

2948-Pos Board B53

High Pressure FTIR Studies on Model α -Helical Peptides

Teraya Donaldson, Alice Smith-Gicklhorn, Sean M. Decatur.

High-pressure conditions force water into internal cavities of proteins; the presence of water in a hydrophobic interior can destabilize the tertiary structure resulting in protein unfolding. This effect has been predicted in molecular dynamics simulations for model alpha-helical peptides at high pressures, where water is forced into closer contact with backbone carbonyls (as measured by an increased in backbone hydration). According to simulations, this increase hydration depends on local sequence and the impact that specific side chains have on backbone conformation. We have investigated the effects of pressure on a series of 20-residue peptides. Model peptides based on the alanine and lysine

repeats form water soluble, stable alpha helices (1). The sequence (AAAAK)3-AAAY is a well-characterized, synthetic polypeptide, ideal for the study of helix properties(2). Using the model peptides, we experimentally confirm that the helical content is conserved under pressure. Perturbations were monitored using a probe of secondary structure, the amide I' band, with infrared (IR) spectroscopy and a diamond anvil cell. Amide I' mode shifts to a lower frequency with the increase of pressure. Local information is obtained by measuring amide I' bands in ¹³C labeled peptides, where ¹³C alanines are placed at different positions relative to the lysines within the peptide. We examined the shielding effects of lysine at the various positions and confirm that shielding of backbone carbonyls occurs.

1. Starzyk, A., Barber-Armstrong, W., Sridharan, M., and Decatur, S. M. (2005) Spectroscopic evidence for backbone desolvation of helical peptides by 2,2,2-trifluoroethanol: an isotope-edited FTIR study, *Biochemistry* 44, 369-376.

2. Paschek, D., Gnanakaran, S., and Garcia, A. E. (2005) Simulations of the pressure and temperature unfolding of an alpha-helical peptide, *Proc Natl Acad Sci U S A* 102, 6765-6770.

2949-Pos Board B54

SAXS Study of Cytochrome-C Cold Denaturation

Margaret Elmer, Christopher Asta, Katherine Butler, Apratim Dhar, Martin Gruebele, Liang Guo, Thomas Irving, Joseph Marcus, Sarah Rice, Eric Landahl.

We present a study of the cold denaturation of proteins using Small Angle X-Ray Scattering. The size and shape of equine cytochrome-c is determined at varying salt and pH conditions from -25 to 60° C and compared to the two-state Ideal Thermal Protein model of Ghosh and Dill (PNAS 2009). The incorporation of a temperature-dependent pH and solvent dielectric constant is critical to model electrostatic interactions over this broad temperature range and properly predict the observed protein stability from sequence. Under suitable conditions, the protein can be made to increase in size by nearly 9 Angstroms (over 60% of its native radius of gyration) when dropped in temperature from 0 to -25 C. Cold denaturation under these conditions is also verified by monitoring fluorescence from the native tryptophan in this protein. This allows us to compare denaturation monitored at one location inside the protein with global structural changes observed by SAXS.

2950-Pos Board B55

Apolipoprotein B Reconstruction at Single Molecular Level

Hsueh-Liang Chu, Tsai-Mu Cheng, Yu-Chuan Chang, Hung-Wei Chen, Wei-Hsien Chung, Chia-Ching Chang.

Low density lipoprotein-cholesterol (LDL-C) is a clinical significant marker of cardiovascular disease risk. Each particle of LDL contains only one protein, apolipoprotein B (apoB). In human, the liver secretes full-length apoB (apoB-100) which also serves as a ligand for receptor-mediated uptake of LDL by a variety of cell types, such as monocyte and A549. Many diseases' progression may cause by the deficiency of the LDL assembly. Therefore, it is desired to reveal the folding/assembly process of apoB. However, it is challenge to refold the apoB, because of its high insolubility in solution. In this study, the native apoB had been purified from LDL by ice cheer method, and truncated mutants of various lengths of apoB were recombinant expressed from *E. coli* system. The apoB-100 and its truncated proteins then were dissolved in the denature buffer which contained additional detergents and were refolded via an over-critical refolding process. The folding intermediates of apoB-100 and its truncated mutants could be observed by immunofluorescence microscopy at the single molecular level and the lipidation processes and secondary structures of apoBs can be also analyzed by SRCD. Moreover, the confocal microscopy showed that the refolded and native LDL could be absorbed by THP-1 and A549 cell. According to our observation the LDL assembly process can be proposed. This is the first study to refold the structural and function of apoB *in vitro*. Meanwhile, this refolded lipoprotein can be used as carrier for hydrophobic particles delivery.

2951-Pos Board B56

Folding Studies of Beta-Strand-Containing Repeat Proteins Through Naturally-Occurring and Consensus-Designed Sequences

Thuy P. Dao, Ananya Majumdar, Doug Barrick.

Repeat proteins, devoid of sequence-distant contacts observed in globular proteins, are ideal candidates for the dissection of local stability, nearest-neighbor contact parameters, cooperativity, and determination of folding energy landscapes. Recent studies on HEAT, TPR, Ankyrin, and Armadillo have contributed to the understanding of the folding of helical repeat proteins. Moreover, constructs composed of simplified "consensus" sequence

representations of helical repeat proteins have been found to be well-behaved and highly stable, and facilitate energy dissection using simple nearest-neighbor models.

In contrast, little progress has been made towards understanding the folding of β -sheet containing repeat. To determine the folding stability and cooperativity of these proteins, and to better understand the sequence determinants of structure and stability within these ubiquitous families, we have initiated studies on a series of LRR proteins of both naturally occurring and consensus-designed sequences. We find the LRR proteins PP32 and LC1 to be well-behaved, and to fold in a highly cooperative transition that is consistent with a two-state mechanism. NMR H^2H exchange shows the repeating β -strands on the concave surface of both proteins to be more protected than the rest of the molecules, and can be regarded as an exchange-resistant core, whereas the terminal caps and convex structural elements are more labile. However, truncations and sequence substitution demonstrate that the caps significantly influence stability and kinetics.

To further simplify our analysis of β -sheet containing repeat protein folding, we designed consensus LRR sequences. On their own, these constructs are unfolded and/or aggregated. By fusing these consensus sequences with naturally occurring LRR protein YopM, we have obtained solubilized, folded arrays that exhibit increased stability and drastically decreased unfolding and refolding rates with repeat number. Further studies are needed to dissect the complex folding pathways taken by these constructs.

2952-Pos Board B57

The Structural and Functional Role of the Sole Tryptophan Residue in the Human Acidic Fibroblast Growth Factor

Hannah M. Henson, Anna E. Daily, Suresh Kumar.

Department of Chemistry & Biochemistry, University of Arkansas, Fayetteville, AR.

Abstract:

Fibroblast Growth Factor-1 (FGF-1) is a 16kDa heparin binding protein, which has been associated with a variety of important functions including angiogenesis and wound repair. In order for FGF-1 to enter the cell it must interact with the FGF-1 receptor on the cell surface. One vital residue involved in the binding of FGF-1 to the receptor is tryptophan 121 (W121). This study aims to examine the role of W121 on the conformation and functionality of FGF-1. Site-directed mutagenesis will be used to incorporate mutations at position 121. The effect of these mutations will be characterized using various biophysical techniques including fluorescence, CD, ITC, and multi-dimensional NMR spectroscopy. As FGFs are involved in many crucial cellular processes, the gain from this study is expected to provide useful information on the regulation of the FGF signaling process.

2953-Pos Board B58

Folding Mechanism Revealing of PGB1 by FRET and Molecular Simulation

Chung Wei-Hsien, Yi-Chen Yeh, Nan-Yow Cheng, Chung-Yu Mou, Chia-Ching Chang.

1. Department of Biological Science and Technology, National Chiou Tung University, Hsinchu, Taiwan.

2. Institute of Physics, Academia Sinica, Taipei Taiwan.

3. Department of Physics, National Tsing Hua University, Hsinchu, Taiwan.

Immunoglobulin-binding B1 domain of streptococcal protein G (PGB1) is a small (56 residues) protein with an α -helix (from Ala23 to Asp 36) lying on top of two pairs of anti-parallel β -sheets, β -hairpin 1 (from Met1 to Ala20) and β -hairpin 2 (from Glu42 to Glu56), covering the hydrophobic core. PGB1 contains no disulfide bonds in its structure and makes it an excellent model protein for folding study. Our simulation results showed that the α -helix and β -hairpin 2 had interacted prior than β -hairpin 1 in early folding stage. Similar experimental results can be observed by monitoring the folding intermediates of PGB1, which are performed by an over-critical refolding process, by using fluorescence resonance energy transfer (FRET) technique, the technique is a precisely optical technique which can reveal the distance difference within angstrom scale. The FRET analysis of PGB1 also indicated that the distance between α -helix and β -hairpin 2 remained approximately unchanged in all folding intermediates. However, the distance between β -hairpin 1 and β -hairpin 2 decreased during the folding process. Therefore both simulation and FRET analysis were in consistency. The molecular dynamics of PGB1 during its folding process can be demonstrated.

Membrane Protein Functions

2954-Pos Board B59

Probing Mechanism for the Enhancement of Uptake of Fatty Acid into Cells by the Membrane Protein CD36

Su Xu, Michael Kirber, Nasi Huang, Kellen Brunaldi, James A. Hamilton.

CD36 is a membrane protein found in various cell types including adipocytes and endothelial cells. Physiological studies have suggested that CD36 is associated with insulin-resistant diabetes, however, the mechanism remains poorly understood. It is well recognized as a receptor for multiple ligands such as collagen and thrombospondin 1, and possibly acts as a catalyst to enhance the rate of transmembrane movement of fatty acid. To separate the two independent events transport across the plasma membrane and the subsequent intracellular metabolism, we have applied biophysical approaches and metabolic analyses to cells in vitro. Our results using cultured cells (HEK293 cells overexpressing CD36 and 3T3-L1 adipocyte cells) as well as mice adipocyte cells isolated from a CD36 null mouse showed that fatty acids diffuse through the plasma membrane rapidly with or without CD36. In HEK 298 cells, which normally synthesize triglycerides very slowly and to a limited extent, expression of CD36 enhanced the rate and extent of synthesis. Even in the presence of CD36, incorporation into triglycerides is a much slower process (min) relative to the transmembrane movement (sec), indicating that the rate-limiting step of the regulation of fatty acid uptake by CD36 is intracellular metabolism. Lastly, by PCR array analysis of 84 proteins, we have identified several enzymes involved in human fatty acid metabolism with gene expression levels altered by overexpression of CD36. Taken together, our results showed that CD36 increases fatty acid uptake by enhancing triglycerides synthesis rather than acting as a membrane transporter, but as yet by unidentified molecular mechanisms.

2955-Pos Board B60

Membrane Transport of CO₂ and H₂S: No Facilitator Required

Florian Zocher, John C. Mathai, Andreas Missner, Mark L. Zeidel,

Peter Pohl.

The observation that some membranes and epithelia have no demonstrable gas permeability suggested that membrane channels may be involved in CO₂ transport. Both aquaporins and Rhesus proteins were reported to serve as pathways for CO₂ and H₂S. In contrast, we show here that membrane lipid has such high CO₂ and H₂S permeabilities that the presence of a protein channel does not enhance the flux. Therefore we reconstituted aquaporins into lipid bilayers and used scanning microelectrodes to monitor pH in the immediate vicinity of planar lipid bilayers. The lower limits of lipid bilayer permeabilities to CO₂ and to H₂S were equal to 3.2 ± 1.6 cm/s^[1] and 0.5 ± 0.4 cm/s^[2], respectively. We also observed that the CO₂ flux through the lipid bilayer decreases several fold when the rate of CO₂ formation from HCO₃⁻ was not augmented by carbonic anhydrase (CA). Experiments with epithelial cell monolayers grown on permeable support revealed the same result. Inhibition of CA transformed these otherwise highly CO₂ permeable cell monolayers into CO₂ barriers. Finally we tested the CO₂ permeability of the epithelium of the mammalian bladder. It was impermeable to CO₂ even after uroplakin knock-out. We found that the lack of intrinsic intracellular CA activity of these epithelial cells hampers the CO₂ exchange between blood and urine.

[1] A. Missner, P. Kügler, S. M. Saparov, K. Sommer, J. C. Mathai, M. L. Zeidel, P. Pohl, J.Biol.Chem. 2008, 283 25340-25347.

[2.] J. C. Mathai, A. Missner, P. Kügler, S. M. Saparov, M. L. Zeidel, J. K. Lee, P. Pohl, Proc.Natl.Acad.Sci.U.S.A. 2009, 106 16633-16638.

2956-Pos Board B61

F65S Mutation in RhAG is Associated with Decreased Ammonia Flux Through Overhydrated Stomatocytic Erythrocytes

Sandrine Genetet, Pierre Ripoché, Julien Picot, Sylvain Bigot, Jean Delaunay, Corinne Amari-Alla, Yves Colin, Isabelle Mouro-Chanteloup.

The Overhydrated Stomatocytic (OHSt) erythrocytes of four patients, characterized by the F65S mutation (Bruce et al, Blood 2009) located in the pore of the ammonia channel Rh Associated Glycoprotein (RhAG), were studied. We have previously demonstrated that the equivalent substitution (F74L) in the non-erythroid analogue RhCG resulted in a reduction of the ammonia influx of 50% in transfected HEK293 cells.

Ghosts prepared by hypotonic lysis of OHSt and control (Ctl) erythrocytes, resealed in the presence of a pH-sensitive probe (pyranine), exhibited echinocytic morphology. Images of 1667 (Ctl) and 1998 (OHSt) echinocytes, visualized by light microscopy, allowed the determination of average ghost diameters: 5.81 ± 0.2 μ m (Ctl) and 5.83 ± 0.06 μ m (OHSt). RhAG densities, as determined by flow cytometry, were similar for Ctl and OHSt (78 000 to 80 000 copies/cell).