



REVISTA BRASILEIRA DE ANESTESIOLOGIA

Official Publication of the Brazilian Society of Anesthesiology
www.sba.com.br



ERRATUM

Association of Pain Catastrophizing with the Incidence and Severity of Acute and Persistent Perineal Pain after Natural Childbirth: Longitudinal Cohort Study

Anne Danielle Santos Soares^a, Tânia Cursino de Menezes Couceiro^{*,a,b}, Luciana Cavalcanti Lima^{c,d}, Fernanda Lobo Lago Flores^a, Eusa Maria Belarmino Alcoforado^e, Roberto de Oliveira Couceiro Filho^a

^a Instituto de Medicina Integral Professor Fernando Figueira, Recife, PE, Brazil

^b Universidade Federal de Pernambuco, Recife, PE, Brazil

^c Universidade Estadual Paulista Júlio de Mesquita Filho, São Paulo, SP, Brazil

^d Faculdade Pernambucana de Saúde, Recife, PE, Brazil

^e Ministério da Saúde, Brazil

In the original article “Association of Pain Catastrophizing with the Incidence and Severity of Acute and Persistent Perineal Pain after Natural Childbirth: Longitudinal Cohort Study” (Rev Bras Anesthesiol. 2013;63(4):317-321), on page 318, where it is read:

Table 1 Incidence and Severity of Acute and Persistent Perineal Pain.		
Incidence	Acute Perineal Pain (%)	Persistent Perineal Pain (%)
Absent		
Present		
Total		
Severity	Acute Perineal Pain (%)	Persistent Perineal Pain (%)
Mild	18 (47.3%)	6 (75%)
Moderate	19 (50%)	2 (25%)
Severe	1 (2.6%)	-
Total	38 (100%)	8 (100%)

It should read:

Table 1 Incidence and Severity of Acute and Persistent Perineal Pain.		
Incidence	Acute Perineal Pain (%)	Persistent Perineal Pain (%)
Absent	17 (31%)	47 (85.5%)
Present	38 (69%)	8 (14.5%)
Total	55 (100%)	55 (100%)
Severity	Acute Perineal Pain (%)	Persistent Perineal Pain (%)
Mild	18 (47.3%)	6 (75%)
Moderate	19 (50%)	2 (25%)
Severe	1 (2.6%)	-
Total	38 (100%)	8 (100%)

*Corresponding author.

E-mail: taniacouceiro@yahoo.com.br (T.C.M. Couceiro)