



Weak Schur numbers and the search for G.W. Walker's lost partitions

S. Eliahou^{a,b,c,*}, J.M. Marín^d, M.P. Revuelta^d, M.I. Sanz^d

^a Univ Lille Nord de France, F-59000 Lille, France

^b ULCO, LMPA J. Liouville, B.P. 699, F-62228 Calais, France

^c CNRS, FR 2956, France

^d Escuela Técnica Superior de Ingeniería de Edificación, Departamento de Matemática Aplicada I, Avenida Reina Mercedes 4, C.P. 41012 Sevilla, Spain

ARTICLE INFO

Article history:

Received 19 November 2010

Received in revised form 2 September 2011

Accepted 3 November 2011

Keywords:

Schur numbers

Sum-free sets

Weakly sum-free sets

Boolean variables

SAT problem

SAT solver

ABSTRACT

A set A of integers is *weakly sum-free* if it contains no three *distinct* elements x, y, z such that $x + y = z$. Given $k \geq 1$, let $WS(k)$ denote the largest integer n for which $\{1, \dots, n\}$ admits a partition into k weakly sum-free subsets. In 1952, G.W. Walker claimed the value $WS(5) = 196$, without proof. Here we show $WS(5) \geq 196$, by constructing a partition of $\{1, \dots, 196\}$ of the required type. It remains as an open problem to prove the equality. With an analogous construction for $k = 6$, we obtain $WS(6) \geq 572$. Our approach involves translating the construction problem into a Boolean satisfiability problem, which can then be handled by a SAT solver.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

A set A of integers is called *sum-free* if it contains no elements $x, y, z \in A$ satisfying $x + y = z$. It is called *weakly sum-free* [1] if it contains no *pairwise distinct* elements $x, y, z \in A$ satisfying $x + y = z$. Clearly, sum-free implies weakly sum-free; the converse is false, as shown by $A = \{1, 2\}$.

This paper is concerned with partitions of the set $[1, n] = \{1, 2, \dots, n\}$ into k sum-free, or k weakly sum-free parts, with k fixed and n as large as possible with respect to k .

1.1. Schur numbers

A theorem of Schur states that, given $k \geq 1$, there is indeed a largest integer n for which $[1, n]$ admits a partition into k sum-free sets [2]. This largest n is called the k -th Schur number and is denoted by $S(k)$.

For instance, one has $S(1) = 1$ and $S(2) = 4$. For $k = 2$, a partition of $[1, 4]$ into 2 sum-free sets is provided by

$$\{1, 2, 3, 4\} = \{1, 4\} \sqcup \{2, 3\},$$

and it is easy to check that there is no such partition for $[1, 5]$. Only two more exact values of $S(k)$ are known so far, namely

$$S(3) = 13, \quad S(4) = 44.$$

The currently available bounds for $S(5)$ are

$$160 \leq S(5) \leq 305, \tag{1}$$

* Corresponding author at: Univ Lille Nord de France, F-59000 Lille, France.

E-mail addresses: eliahou@lmpa.univ-littoral.fr (S. Eliahou), jmarin@us.es (J.M. Marín), pastora@us.es (M.P. Revuelta), isanz@us.es (M.I. Sanz).

and it is conjectured in [3] that the lower bound 160, settled in [4], is perhaps sharp. For the upper bound 305, see [5,6]. While the value $S(3) = 13$ can still be settled by hand, the value $S(4) = 44$ relies on an exhaustive computer search [7,8]. Also note the bounds $S(6) \geq 536$ and $S(7) \geq 1680$ established in [3]. For general $k \geq 1$, Schur proved the following lower and upper bounds [2]:

$$(3^k - 1)/2 \leq S(k) \leq \lfloor k!e \rfloor - 1.$$

1.2. Weak Schur numbers

As for partitions of $[1, n]$ into weakly sum-free sets, a result analogous to Schur's holds, due to Rado [9]. See also [10,1,11]. Let us denote by $WS(k)$ the largest integer n for which $[1, n]$ can be partitioned into k weakly sum-free parts. We call $WS(k)$ the k -th weak Schur number. Note that some authors prefer to speak of $N_k = WS(k) + 1$, the smallest integer n' for which every k -coloring of $[1, n']$ contains a monochromatic triple $\{a, b, a + b\}$ with $a \neq b$.

The current state of knowledge concerning $WS(k)$ is quite confused. The problem seems to have been first considered in [12], which is Walker's solution to Problem E 985 proposed a year earlier, in 1951, by Leo Moser¹. Subsequent mentions appear in [13,1,14,10,11,6], in the chronological order. The values $WS(1) = 2$ and $WS(2) = 8$ are easy to check. It is established in [10], by exhaustive computer search, that

$$WS(3) = 23,$$

$$WS(4) = 66,$$

$$WS(5) \geq 189.$$

However, the authors of [10] seem to have been unaware of Walker's note [12]. Indeed, that note contains amazing claims that go beyond [10]. Not only does it give the exact values for $WS(3)$ and $WS(4)$, it further claims the equality

$$WS(5) = 196.$$

Unfortunately, Walker only discusses the case $k = 3$, by giving a suitable partition of $[1, 23]$ and explaining why 23 is optimal. He gives no details for $k = 4$ and 5, not even suitable partitions which would establish $WS(4) \geq 66$ and $WS(5) \geq 196$. Nobody today seems to know how Walker managed to make these amazing claims back in 1952, when computers were not generally available. The situation is somewhat reminiscent of Fermat's claimed Last Theorem which, incidentally, was the main motivation behind Schur's discovery of his numbers $S(k)$ in [2].

Finally, let us mention the following upper bound, improving an earlier one by Irving [1] and due to Bornshtein [14]:

$$WS(k) \leq \lfloor k!ke \rfloor.$$

1.3. Comparing $S(k)$ and $WS(k)$

It is clear from the definitions that

$$S(k) \leq WS(k) \tag{2}$$

for all $k \geq 1$. Indeed, the set $[1, S(k)]$ admits a partition into k sum-free, and hence weakly sum-free, subsets. This inequality is important, as it provides a natural upper bound for the Schur numbers $S(k)$. In particular, Walker's claim on $WS(5)$ yields a major potential improvement, apparently not mentioned before, of the best known upper bound on $S(5)$:

$$S(5) \leq WS(5) \stackrel{?}{=} 196,$$

as compared to $S(5) \leq 305$ in (1). This certainly provides a strong call to definitively settle the exact value of $WS(5)$.

There is a long time interval, from [1] in 1973 to [14] in 2002, during which Walker's note seems to have fallen into oblivion. One possible reason is that yet another claim of Walker's, namely $WS(k+1) \leq 3WS(k) + 1$ for $k \geq 3$, was pointed out by Irving [1] as being incompatible with the following bound of Abbott and Hanson [15] on the ordinary Schur numbers:

$$S(k) \geq c 89^{k/4}$$

for all $k \geq 4$, where $c = 44/89$.

1.4. Links with multicolor Ramsey numbers

The classical and weak Schur numbers are related to some multicolor Ramsey numbers, as we now recall. For integers $k, m \geq 1$, denote by $R_k(m)$ the smallest integer $n \geq 1$ such that, for every k -coloring of the edges of the complete graph K_n

¹ In fact, Moser's informal challenge was about $S(3)$; Walker considered $WS(3)$ instead.

on n vertices, there is a subgraph K_m , all of whose edges are colored the same. A short argument (see e.g. [16, p. 69]) yields the bound

$$S(k) \leq R_k(3) - 2;$$

the idea is to transport a k -coloring of $[1, n - 1]$ to a k -coloring of the edges of K_n by assigning to any edge $\{x, y\}$ the color of $|x - y|$. As for weak Schur numbers, suitable adaptations of this idea yield two different bounds, namely

$$WS(k) \leq R_k(4) - 2$$

and

$$WS(k) \leq R_{2k}(3) - 2.$$

See [10, p. 2] for the first bound, and [11, p. 303] for the second one.

1.5. Contents

In this paper, we do give a proof of the inequality $WS(5) \geq 196$, by providing an actual partition of $[1, 196]$ into 5 weakly sum-free sets. Our efforts to do the same with $[1, 197]$ completely failed. It remains as a challenge to prove, by theory or by machine, that 196 is the exact value of $WS(5)$.

Our construction showing $WS(5) \geq 196$ is given in Section 2, together with reasons pointing to the probable sharpness of this bound. In Section 3, we treat the case of 6-partitions and obtain $WS(6) \geq 572$, apparently the first known realistic lower bound on $WS(6)$. Our method is described in Section 4. It involves translating the problem of constructing partitions of the desired type into a Boolean satisfiability problem, to be handled by a SAT solver. The actual computations are briefly commented in Section 5.

2. Is it true that $WS(5) = 196$?

There are no details in [12] substantiating the claim $WS(5) = 196$, not even an actual partition of $[1, 196]$ into 5 weakly sum-free subsets which would establish $WS(5) \geq 196$. Surely Walker knew such partitions, but we do not know how he proceeded, and these are probably lost forever. Here we fill this literature gap by providing one such partition, constructed using the methods of Section 4 and the SAT solver `march` [17].

As a matter of notation, we shall abbreviate runs of consecutive integers as intervals. For instance, $[8, 9] 12 [14, 17]$ stands for the set $\{8, 9, 12, 14, 15, 16, 17\}$.

Theorem 2.1. $WS(5) \geq 196$.

Proof. Consider the following partition $A_1 \sqcup A_2 \sqcup A_3 \sqcup A_4 \sqcup A_5$ of $[1, 196]$:

A_1 : 1 2 4 8 11 22 25 50 63 69 135 140 150 155 178 183 193

A_2 : 3 [5, 7] 19 21 23 [51, 53] [64, 66] [137, 139] [151, 153] [180, 182] [194, 196]

A_3 : [9, 10] [12, 18] 20 [54, 62] [141, 149] [184, 192]

A_4 : 24 [26, 49] 154 [156, 177] 179

A_5 : [67, 68] [70, 134] 136.

It is straightforward to check that each A_i is weakly sum-free. This finishes the proof. \square

Is this lower bound on $WS(5)$ sharp? If not, there would exist a partition of $[1, 197]$ into 5 weakly sum-free sets. In order to try and find one, we applied the same methods as above. But these attempts completely failed, as no conclusion of existence or non-existence was reached after several weeks of running time. This strongly supports Walker's claim. Yet it remains as an open problem to prove or disprove the inequality $WS(5) \leq 196$.

2.1. Fixing the 5th part

Here are two more computational results in support of Walker's claim. Fixing the 5th part of a tentative partition of $[1, n]$ into 5 weakly sum-free sets reduces the number of Boolean variables involved in our method from $3n$ to $2n$. The reason, made clear in Section 4, is that $\lceil \log_2(5) \rceil = 3$ whereas $\lceil \log_2(4) \rceil = 2$. This reduction allows `march` to terminate its computations and reach definitive conclusions, even for $n = 197$. The following results were obtained in this way.

The first statement concludes an attempt to improve Theorem 2.1 by constructing a partition of $[1, 197]$ into 5 weakly sum-free sets, while keeping the same 5th part A_5 . Not surprisingly, the conclusion is negative.

Computational Theorem 2.2. *There is no partition of $[1, 197]$ into 5 weakly sum-free parts with $A_5 = [67, 68] \cup [70, 134] \cup \{136\}$ as one part.*

Our second statement deals with the following question: to what extent is it possible to replace A_5 in Theorem 2.1 by a single interval? More precisely, we looked for the largest possible n for which $[1, n]$ admits a partition into 5 weakly sum-free sets B_1, \dots, B_5 such that:

- B_1, B_2, B_3, B_4 is a partition of $[1, 66]$,
- B_5 is a single interval.

Observe that the first requirement is satisfied by A_1, A_2, A_3, A_4 in Theorem 2.1, and recall that $66 = WS(4)$. Consequently, B_5 must contain 67 and cannot be strictly larger than $[67, 134]$. Thus, without loss of generality, we may and will assume $B_5 = [67, 134]$. Quite surprisingly, the largest admissible n turns out to be $n = 194$ only.

Computational Theorem 2.3. *The largest n for which $[1, n]$ admits a partition into 5 weakly sum-free parts, with $[67, 134]$ as one part, is $n = 194$.*

For the record, here is such a partition of $[1, 194]$.

B_1 : 1 2 4 8 11 22 25 50 66 138 148 153 176 181 194

B_2 : 3 [5, 7] 19 21 23 [51, 53] [63, 65] [135, 137] [149, 151] [178, 180] [191, 193]

B_3 : [9, 10] [12, 18] 20 [54, 62] [139, 147] [182, 190]

B_4 : 24 [26, 49] 152 [154, 175] 177

B_5 : [67, 134].

These two results were reached in about 17 and 18 h, respectively, on a 3.33 GHz Intel i7 processor PC with the SAT solver march.

3. A lower bound on WS(6)

We obtain here the lower bound $WS(6) \geq 572$, by exhibiting a suitable partition of $[1, 572]$ into 6 weakly sum-free subsets. Instructed by a fair amount of experimentation, we think that this bound is quite realistic, with a margin of error possibly less than 10. In the partition below, we keep the same notational convention with intervals as in the preceding section.

Theorem 3.1. $WS(6) \geq 572$.

Proof. Consider the following partition $A_1 \sqcup A_2 \sqcup A_3 \sqcup A_4 \sqcup A_5 \sqcup A_6$ of $[1, 572]$:

A_1 : 1 2 4 8 11 22 25 50 63 69 135 140 150 155 178 183 193 395 412 516 526 531 554 559 572

A_2 : 3[5, 7] 19 21 23 [51, 53] [64, 66] [137, 139] [151, 153] [180, 182] [194, 196] [396, 398] [408, 410] 435 [513, 515] [527, 529] [556, 558] [569, 571]

A_3 : [9, 10] [12, 18] 20 [54, 62] [141, 149] [184, 192] [399, 407] [437, 445] [517, 525] [560, 568]

A_4 : 24 [26, 49] 154 [156, 177] 179 411 [413, 434] 436 530 [532, 553] 555

A_5 : [67, 68] [70, 134] 136 [446, 512]

A_6 : [197, 394].

Again, it is straightforward to check that each A_i is weakly sum-free. \square

The only previously available firm lower bound on $WS(6)$ was 536, which is the lower bound for $S(6)$ given in [3]. Another lower bound could be obtained from the claimed inequality

$$WS(k) \geq 3(3^k + 2k - 1)/4 - 1,$$

which gives 554 at $k = 6$. However, this inequality does not seem to be backed up by any available proof. It is attributed to Braun in [12]. Note that it is sharp for $k = 1, 2, 3$, and gives $65 = WS(4) - 1$ at $k = 4$.

4. Reformulation as a SAT problem

Our idea for constructing the above partitions is to express the corresponding combinatorial constraints as Boolean satisfiability problems, to be then fed to a SAT solver. See [18–21] for earlier successful uses of SAT solvers in combinatorial number theory.

Recall that a logical formula over Boolean variables x_1, \dots, x_n is said to be *satisfiable* if there is an assignment of the x_i 's to True or False in such a way that the formula evaluates to True.

Let $n, k \geq 2$. Let T_1, \dots, T_r be a family of subsets of $[1, n]$. Assume that we are seeking k -colorings of $[1, n]$ for which no T_j is monochromatic. Such k -colorings correspond of course to k -partitions of $[1, n]$ for which no T_j is contained in a single part. We shall translate this existence problem into one asking whether some associated logical formula is satisfiable or not. In the applications in Section 5, the subsets T_j will be all possible triples in $[1, n]$ of the form $\{a, b, a + b\}$ with $a \neq b$.

It is sometimes convenient to express logical formulas in *conjunctive normal form*, or *CNF* for short. That is, as conjunctions

$$\bigwedge_{l=1}^t C_l \tag{3}$$

of clauses C_1, \dots, C_t , a *clause* being a disjunction of the form

$$x_{i_1} \vee \dots \vee x_{i_s} \vee \neg x_{j_1} \vee \dots \vee \neg x_{j_t}.$$

Here, as usual, the symbols \wedge , \vee and \neg denote the logical operations AND, OR and NOT, respectively. Most formulas below are in CNF.

From now on, we shall write 1 for True and 0 for False. In particular, we have $\neg 1 = 0$ and $\neg 0 = 1$.

4.1. The case $k = 2$

We start with two colors. Let x_1, \dots, x_n be n Boolean variables. There is a bijective correspondence between $\{0, 1\}$ -assignments of the x_i 's and 2-colorings of $[1, n]$. For a subset $T \subset [1, n]$, define the CNF formula

$$cl_2(T, (x_1, \dots, x_n)) = \left(\bigvee_{i \in T} x_i \right) \wedge \left(\bigvee_{i \in T} \neg x_i \right). \tag{4}$$

Lemma 4.1. *Let $T \subset [1, n]$. The 2-colorings of $[1, n]$ for which T is non-monochromatic correspond to the $\{0, 1\}$ -assignments $\epsilon_i (i = 1, \dots, n)$ for which $cl_2(T, (\epsilon_1, \dots, \epsilon_n)) = 1$.*

Proof. By construction, $cl_2(T, (\epsilon_1, \dots, \epsilon_n)) = 1$ if and only if there are indices $i, j \in T$ such that $\epsilon_i = \neg \epsilon_j = 1$, or equivalently $\epsilon_i = 1$ and $\epsilon_j = 0$; this happens if and only if T is non-monochromatic for the corresponding 2-coloring of $[1, n]$. \square

When several subsets of $[1, n]$ are required to be simultaneously non-monochromatic, it suffices to satisfy the conjunction of the corresponding formulas. This yields the following equivalence.

Proposition 4.2. *Let $n \geq 1$ and let T_1, \dots, T_r be subsets of $[1, n]$. Let x_1, \dots, x_n be Boolean variables. There exists a 2-coloring of $[1, n]$ such that no T_j is monochromatic if and only if the formula*

$$\bigwedge_{j=1}^r cl_2(T_j, (x_1, \dots, x_n))$$

is satisfiable. \square

4.2. The case $k = 2^t$

We first extend the above considerations to $k = 2^t$ for any integer $t \geq 1$. Our set of 2^t colors is taken to be the Cartesian product $\{0, 1\}^t$. Let

$$(x_{i,l})$$

$(1 \leq i \leq n, 1 \leq l \leq t)$ be a collection of nt Boolean variables. The unknown color of any $i \in [1, n]$ may and will be represented by the t -tuple

$$(x_{i,1}, \dots, x_{i,t}).$$

Let $T \subset [1, n]$. The 2^t -colorings of $[1, n]$ for which T is non-monochromatic correspond to those $\{0, 1\}$ -assignments $(\epsilon_{i,l})$ of $(x_{i,l})$ for which there are indices $l \in [1, t]$ and $i, j \in T$ such that

$$\epsilon_{i,l} \neq \epsilon_{j,l}.$$

By the case of two colors, this condition is equivalent to the Boolean one

$$cl_2(T, (\epsilon_{1,l}, \dots, \epsilon_{n,l})) = 1.$$

Since indices l where a difference occurs may be arbitrary, one needs to take the disjunction over all $l \in [1, t]$ of the above formula. Moreover, when several subsets of $[1, n]$ are involved, the conjunction of the corresponding formulas must be satisfied. This yields the following generalization of Proposition 4.2.

Proposition 4.3. Let $n, t \geq 1$ and let T_1, \dots, T_r be subsets of $[1, n]$. Let $(x_{i,l})(1 \leq i \leq n, 1 \leq l \leq t)$ be nt Boolean variables. There exists a 2^t -coloring of $[1, n]$ such that no T_j is monochromatic if and only if the formula

$$\bigwedge_{j=1}^r \left(\bigvee_{l=1}^t cl_2(T_j, (x_{1,l}, \dots, x_{n,l})) \right)$$

is satisfiable. \square

This formula is not in CNF, but this can be fixed using the distributivity of \vee over \wedge . To wit, an equivalent CNF formula is given by

$$\bigwedge_{j=1}^r \left(\bigwedge_{U \sqcup V = [1,t]} \left(\left(\bigvee_{i \in T_j} \bigvee_{u \in U} x_{i,u} \right) \vee \left(\bigvee_{i \in T_j} \bigvee_{v \in V} \neg x_{i,v} \right) \right) \right),$$

where \sqcup denotes a disjoint union.

4.3. The general case

We now treat any number $k \geq 2$ of colors. Let $t \geq 1$ be the unique integer such that

$$2^{t-1} + 1 \leq k \leq 2^t.$$

That is, $t = \lceil \log_2(k) \rceil$. Within the set $\{0, 1\}^t$ of 2^t colors, we forbid some $2^t - k$ ones. The remaining k colors then constitute our final palette of colors.

It remains to translate the requirement that some colors are forbidden into the satisfiability of appropriate Boolean formulas. For this, it suffices to consider a single forbidden color; the case of several ones follows by taking the conjunction of the corresponding formulas.

First observe that, for $x, y \in \{0, 1\}$, the condition $x \neq y$ is equivalent to the logical formula

$$(x \vee y) \wedge (\neg x \vee \neg y) = 1.$$

Now, let $\mu = (\mu_1, \dots, \mu_t) \in \{0, 1\}^t$ be a fixed color. Given t Boolean variables z_1, \dots, z_t , define

$$f_\mu(z_1, \dots, z_t) = \bigvee_{l=1}^t (z_l \vee \mu_l) \wedge (\neg z_l \vee \neg \mu_l).$$

It then follows from the above observation that, for all $\epsilon = (\epsilon_1, \dots, \epsilon_t) \in \{0, 1\}^t$, we have

$$\epsilon \neq \mu \Leftrightarrow f_\mu(\epsilon) = 1.$$

Thus, forbidding color μ may be achieved using formula f_μ . This yields the following result.

Theorem 4.4. Let n, k, t be integers with $k, n \geq 2$ and $2^{t-1} + 1 \leq k \leq 2^t$. Let T_1, \dots, T_r be subsets of $[1, n]$. The existence of k -colorings of $[1, n]$ for which no T_j is monochromatic is equivalent to the satisfiability of the formula

$$\bigwedge_{j=1}^r \left(\bigvee_{l=1}^t cl_2(T_j, (x_{1,l}, \dots, x_{n,l})) \right) \wedge \bigwedge_{i=1}^n \bigwedge_{s=1}^{2^t-k} f_{\mu_s}(x_{i,1}, \dots, x_{i,t}),$$

where $\mu_1, \dots, \mu_{2^t-k}$ is any choice of $2^t - k$ distinct elements in $\{0, 1\}^t$.

Proof. By Proposition 4.3, satisfying the left-hand subformula gives 2^t -colorings of $[1, n]$ with no T_j monochromatic. Satisfying the right-hand one guarantees that no μ_s is used in those colorings. \square

5. Applications

The above result implies the following SAT characterization of the weak Schur numbers, made explicit for completeness.

Corollary 5.1. Let n, k, t be integers, with $k, n \geq 2$ and $2^{t-1} + 1 \leq k \leq 2^t$. The following conditions are equivalent.

1. $WS(k) \geq n$.
2. The formula

$$\bigwedge_{a < b} \left(\bigvee_{l=1}^t cl_2(\{a, b, a + b\}, (x_{1,l}, \dots, x_{n,l})) \right) \wedge \bigwedge_{i=1}^n \bigwedge_{j=1}^{2^t-k} f_{\mu_j}(x_{i,1}, \dots, x_{i,t})$$

is satisfiable, where a, b run over all integers satisfying $1 \leq a < b \leq n - a - b$, and where $\mu_1, \dots, \mu_{2^t-k}$ are any $2^t - k$ elements in $\{0, 1\}^t$.

Moreover, every variable assignment for which the above formula is satisfied corresponds to an actual partition of $[1, n]$ into k weakly sum-free subsets.

Proof. This directly follows from the definition of $WS(k)$ and from Theorem 4.4, specialized to the case where the subsets T_j of $[1, n]$ are all triples $\{a, b, a + b\}$ with $a < b$. \square

This SAT reformulation, together with the SAT solver `march`, allowed us to construct the weakly sum-free partitions of Sections 2 and 3. Recall that these partitions yield the lower bounds $WS(5) \geq 196$ and $WS(6) \geq 572$, respectively. Our Computational Theorems 2.2 and 2.3 were obtained with those same tools.

However, for this attack on $WS(5)$ and $WS(6)$, the corresponding SAT problems are somewhat too large. In order to reduce the number of variables and clauses, we performed experiments with selected elements of $[1, n]$ pre-located in the same part of the tentative partitions. For instance, we looked for 5-partitions of $[1, 196]$ into weakly sum-free sets which would extend chosen 4-partitions of $[1, 66]$, where $66 = WS(4)$. This removes many variables. Without such reductions, it seems difficult for a SAT solver to construct from scratch a partition of $[1, 196]$ of the desired type, let alone to conclude $WS(5) < 197$.

The case $k = 4$, in contrast, can be fully handled. We were able to recover the known values $WS(4) = 66$ and $S(4) = 44$, the latter with a SAT characterization of $S(k)$ similar to that of $WS(k)$ above. The corresponding running times are displayed below.

| Output | Conclusion | Time in seconds |
|-------------------------------------|-----------------|-----------------|
| A suitable 4-partition of $[1, 44]$ | $S(4) \geq 44$ | 0 |
| “Unsatisfiable” | $S(4) < 45$ | 60 |
| A suitable 4-partition of $[1, 66]$ | $WS(4) \geq 66$ | 917 |
| “Unsatisfiable” | $WS(4) < 67$ | 24,450 |

We confirmed the equality $WS(4) = 66$ by running `march` on two distinct files embodying Corollary 5.1, namely `wschur4_66.txt` and `wschur4_67.txt`. These files are available at [22], for the reader wishing to reproduce these computations with any SAT solver. They contain, in standard DIMACS format, a list of CNF clauses whose satisfiability or not is equivalent to the existence or not of a suitable 4-partition of $[1, 66]$ and $[1, 67]$, respectively.

The file `wschur4_66.txt` contains 4224 clauses on 132 Boolean variables. Its first clause reads `1 2 3 67 68 69 0`, with 0 as a closing symbol, and codes for $x_1 \vee x_2 \vee x_3 \vee x_{67} \vee x_{68} \vee x_{69}$. Its next clause reads `-1 -2 -3 67 68 69 0` and codes for $\neg x_1 \vee \neg x_2 \vee \neg x_3 \vee x_{67} \vee x_{68} \vee x_{69}$. Any SAT solver running it should end up displaying an assignment of the variables that will satisfy all the clauses. As for the file `wschur4_67.txt`, which contains 4356 clauses on 134 Boolean variables, any SAT solver running it should conclude that its set of clauses is unsatisfiable.

We end with a few technical details. The version of `march` we used was `march_hi` [17], running on an Intel i7 processor PC with a CPU clock speed of 3.33 GHz and 16 GB of RAM memory. As far as we know, the multi-core architecture of the CPU is not exploited by this implementation of `march_hi`.

Acknowledgments

We thank Amine Boumaza, Jonathan Chappelon, Philippe Marion, Virginie Marion-Poty, Denis Robilliard and Dominique Verhaghe for helpful discussions and/or technical help during the preparation of this paper. We also thank both referees for their careful reading and very useful comments.

Note. While this paper was being refereed, our colleagues Amine Boumaza, Cyril Fonlupt, Virginie Marion-Poty and Denis Robilliard succeeded in improving our lower bound on $WS(6)$, from 572 to 574. They used a completely different method, namely an enhanced tabu search scheme. See their forthcoming paper, to appear in the proceedings of Artificial Evolution 2011.

Note added in proof. We have further improved the lower bound on $WS(6)$, which is now given by $WS(6) \geq 575$.

References

- [1] Robert W. Irving, An extension of Schur's theorem on sum-free partitions, *Acta Arith.* 25 (1973/74) 55–64.
- [2] Issai Schur, Über die Kongruenz $x^m + y^m \equiv z^m \pmod{p}$, *Jahresber. Deutsch. Math.-Verein.* 25 (1916) 114–117.
- [3] Harold Fredricksen, Melvin M. Sweet, Symmetric sum-free partitions and lower bounds for Schur numbers, *Electron. J. Combin.* 7 (2000) Research Paper 32, 9, electronic.
- [4] Geoffrey Exoo, A lower bound for Schur numbers and multicolor Ramsey numbers of K_3 , *Electron. J. Combin.* 1 (1994) Research Paper 8, approx. 3, electronic.
- [5] S. Radziszowski, Small Ramsey numbers, *Electron. J. Combin.* 1 (1994) Dynamic Survey DS1, Revision #6 (1999), 35, electronic.
- [6] M.I. Sanz, Números de Schur y de Rado, Ph.D Thesis, Departamento de Matemática Aplicada I, Universidad de Sevilla, 2010.
- [7] Leonard D. Baumert, Sum-free sets, *J.P.L. Res. Summary* 1 (36–10) (1961) 16–18.
- [8] Solomon W. Golomb, Leonard D. Baumert, Backtrack programming, *J. Assoc. Comput. Mach.* 12 (1965) 516–524.
- [9] R. Rado, Some solved and unsolved problems in the theory of numbers, *Math. Gaz.* 25 (1941) 72–77.
- [10] Peter F. Blanchard, Frank Harary, Rogerio Reis, Partitions into sum-free sets, *Integers* 6 (A7) (2006) 10, electronic.

- [11] A. Soifer, *The Mathematical Coloring Book. Mathematics of Coloring and the Colorful Life of its Creators*, Springer, New York, ISBN: 978-0-387-74640-1, 2009.
- [12] G.W. Walker, A problem in partitioning, *Amer. Math. Monthly* 59 (1952) 253.
- [13] W. Sierpiński, *Elementary theory of numbers*. Translated from Polish by A. Hulanicki. *Monografie Matematyczne, Tom 42 Państwowe Wydawnictwo Naukowe*, Warsaw, 1964.
- [14] Pierre Bornsztein, On an extension of a theorem of Schur, *Acta Arith.* 101 (2002) 395–399.
- [15] H.L. Abbott, D. Hanson, A problem of Schur and its generalizations, *Acta Arith.* 20 (1972) 175–187.
- [16] R.L. Graham, B.L. Rothschild, J.H. Spencer, *Ramsey theory*. Second edition. *Wiley-Interscience Series in Discrete Mathematics and Optimization*, John Wiley & Sons, Inc., New York, 1990.
- [17] <http://www.st.eui.tudelft.nl/sat/>.
- [18] Michael R. Dransfield, Lengning Liu, Victor W. Marek, Mirosław Truszczyński, Satisfiability and computing van der Waerden numbers, *Electron. J. Combin.* 11 (1) (2004) Research Paper 41, 15, electronic.
- [19] P.R. Herwig, M.J.H. Heule, P.M. van Lambalgen, H. van Maaren, A new method to construct lower bounds for van der Waerden numbers, *Electron. J. Combin.* 14 (2007) Research Paper 6, 18, electronic.
- [20] M. Kouril, J.L. Paul, The van der Waerden number $W(2, 6)$ is 1132, *Experiment. Math.* 17 (2008) 53–61.
- [21] D. Robilliard, A. Boumaza, V. Marion-Poty, Meta-heuristic search and square Erickson matrices, in: *Proceeding of the IEEE Congress on Evolutionary Computation (CEC'10)*, pp. 3237–3244, IEEE, 2010.
- [22] <http://www-imp.littoral.fr/~eliahou/schur/>.