


Contents lists available at [ScienceDirect](http://ScienceDirect)

## International Journal of Infectious Diseases

journal homepage: [www.elsevier.com/locate/ijid](http://www.elsevier.com/locate/ijid)

## Short Communication

Retrospective search for dengue vector mosquito *Aedes albopictus* in areas visited by a German traveler who contracted dengue in Japan

Mutsuo Kobayashi<sup>a,\*</sup>, Osamu Komagata<sup>a</sup>, Mayuko Yonejima<sup>b</sup>, Yoshihide Maekawa<sup>a</sup>, Kimio Hirabayashi<sup>c</sup>, Toshihiko Hayashi<sup>a</sup>, Naoko Nihei<sup>a</sup>, Masahiro Yoshida<sup>d</sup>, Yoshio Tsuda<sup>a</sup>, Kyoko Sawabe<sup>a</sup>

<sup>a</sup> National Institute of Infectious Diseases (NIID), Department of Medical Entomology, Toyama 1-23-1, Shinjuku-ku, Tokyo 162-8640, Japan

<sup>b</sup> Ritsumeikan University, Institute of Disaster Mitigation for Urban Cultural Heritage, Kyoto, Japan

<sup>c</sup> Shinshu University, Faculty of Textile Science and Technology, Ueda, Japan

<sup>d</sup> Bio-Research, Osaka, Japan

## ARTICLE INFO

## Article history:

Received 25 April 2014

Received in revised form 2 June 2014

Accepted 5 June 2014

**Corresponding Editor:** Eskild Petersen, Aarhus, Denmark

## Keywords:

German traveler

Dengue

Autochthonous infection

Japan

*Aedes albopictus*

Population density

## SUMMARY

A German traveler developed dengue fever in late August 2013, following a direct flight from Germany. Autochthonous dengue virus (DENV) infection has not been reported in Japan. To evaluate the risk of autochthonous DENV transmission in Japan, the authors performed a retrospective search of the five areas visited by the German patient to determine the population density of dengue vector mosquito, *Aedes albopictus*. The annual mean temperature of each area was higher than 12 °C, which is considered suitable for the establishment of *A. albopictus* populations. Our retrospective search revealed the population density of *A. albopictus* to be high in the urban areas of Japan.

© 2014 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/3.0/>).

## 1. Introduction

*Aedes aegypti* (Linnaeus) was detected at Tokyo Narita International Airport in 2012.<sup>1</sup> Currently, however, there are no data on *A. aegypti* infestations in urban residential areas in Japan. Dengue outbreaks occurred in Japan during World War II, at which time *Aedes albopictus* was the vector mosquito.<sup>2</sup> Of note, *A. albopictus* was reported to be the main vector in dengue outbreaks in Hawaii in 2001 and Reunion in 2012.<sup>3,4</sup> Climatological analysis using a geographic information system (GIS) showed that the current distribution of *A. albopictus* in Japan is strongly correlated with areas in which the annual mean temperature is higher than 11 °C.<sup>5</sup> Similarly, in Europe and the USA, the annual mean temperature of more than 94% of the *A. albopictus* infested areas is higher than 11 °C.<sup>6</sup> Since 2010, there has been a marked increase


in the number of imported dengue cases reported in Japan, with 249 cases reported in 2013.<sup>7</sup> More than 70% of the imported dengue cases were confirmed during the mosquito breeding season from May to October.

Coincidentally, a German tourist contracted dengue in Japan in August 2013. The patient, who arrived in Japan on a direct flight from Germany, visited five urban areas, including Hiroshima, Kyoto, Fufuki, Ueda, and Tokyo, during her 2 weeks of travel.<sup>8</sup> In this study, we determined the population density of *A. albopictus* in the areas visited by the tourist (or neighboring areas in the cases of Fufuki and Hiroshima, from where sufficient data could not be obtained), in order to evaluate the risk of autochthonous dengue infection in Japan.

## 2. Mosquito surveillance in areas visited by the German patient

We used human bait collection (HBC), a method of catching mosquitoes by sweeping for 8 min with an insect net before mosquitoes land on the exposed surface of collectors. The average

\* Corresponding author. Tel.: +81 3 5285 1111 ext. 2422; fax: +81 3 5285 1147. E-mail addresses: [mutsuo@nih.go.jp](mailto:mutsuo@nih.go.jp), [komagata@niid.go.jp](mailto:komagata@niid.go.jp) (M. Kobayashi).




**Figure 1.** Average number of *Aedes albopictus* collected using human bait collection and CDC miniature traps in areas visited by the dengue patient and surrounding areas of Japan. Gray color shows prefectures visited by the German patient who was suspected to have contracted dengue fever in Japan. TR: CDC miniature trap (CDC trap); HBC: human bait collection with insect net for 8 min. The numbers in parentheses show the total number of CDC traps and the total number of samples collected by HBC using an insect net for 8 min. \*In Ueda City, HBC was performed for 6 min for all of the mosquito collection sessions. \*\*Records of mosquito collection in the cities of Fuefuki and Hiroshima were not found, although the German patient visited these cities.

numbers of mosquitoes collected by HBC in a residential house in Ueda City was 4.8 in 2012 and 3.2 in 2013 (Figure 1). In Kyoto, more than 10 mosquitoes were collected by HBC in two Japanese temples. These temples are historical sightseeing points that are commonly visited by domestic and international tourists. The results suggest that a stationary human host at a shaded point under a tree canopy and near bushes or vegetation in the vicinity of the temple would be exposed to attacks by more than 10 female mosquitoes within 8 min. In Yokohama, the collection site was a

typical residential house, the garden of which was covered with plenty of vegetation. The average number of mosquitoes collected by HBC was almost 10, which is considerably high. It is hypothesized that rich vegetation in the gardens of residential houses and public spaces contributes to an environment that is suitable for harboring female mosquitoes, even when larval habitats are not constantly present in these environments.

*A. albopictus* was also collected by a CDC miniature trap enhanced with dry ice, which was set in public spaces in



**Figure 2.** Distribution of areas with annual mean temperatures ranging from below 10 °C to above 12 °C in 2013. All areas visited by the German patient were above 12 °C in annual mean temperature.

Nishinomiya City, on the ground in suburban areas in two cities in Hiroshima Prefecture, and on the ground in the National Institute of Infectious Diseases, Tokyo. While the numbers of *A. albopictus* mosquitoes collected by CDC miniature traps at most points were not high, a large number of mosquitoes were collected at one point in Tokyo (Figure 1). The results demonstrated that the population density of *A. albopictus* was considerably high and that there were marked differences in the population densities across areas, with discrepancies found between collection points even within a small public space or in different parts of a temple. This is likely related to the abundance of vegetation, as abundant vegetation harbors mosquitoes, and to the number of potential larval breeding habitats, including catch basins, located around the collection points.

### 3. Analysis of the recent distribution of *Aedes albopictus* with a focus on annual mean temperature by GIS in Japan

We previously reported that the annual mean temperature in Tohoku District of Honshu Island, Japan was clearly related to the distribution of *A. albopictus*.<sup>5</sup> However, the map used in the previous study was charted using old climate data, which were collected between 1971 and 2000. In this retrospective study, a map of 1-km mesh climate values was charted using data collected by the Japan Meteorological Agency in 2013. The annual mean temperature was above 12 °C in all five of the areas visited by the German patient (Figure 2). With the exception of higher altitude parts of the central mainland of Japan, the western parts of Tokyo were infested with *A. albopictus* and the population density was relatively high in comparison to northern Honshu.

### Acknowledgements

The authors acknowledge the Pest Control Division, Nishinomiya City Office, for their collection of mosquitoes in 2008 and Dr Meng Ling Moi for her valuable comments on an earlier version of the manuscript. The manuscript was proofread by Rise Japan LLC (Tokyo, Japan). This research was supported in part by the Environment Research and Technology Development Fund (S-8) of the Ministry of the Environment and Health Labor Sciences Research Grants (H21-Shinko-Ippan-005 and H24-Shinko-Ippan-007) from the Ministry of Health, Labor and Welfare, Japan.

*Conflict of interest:* None declared.

### References

1. Sukehiro N, Kida N, Umezawa M, Murakami T, Arai N, Jinnai T, et al. First report on invasion of yellow fever mosquito, *Aedes aegypti*, at Narita International Airport, Japan in August 2012. *Jpn J Infect Dis* 2013;**66**:189–94.
2. Hotta S. Dengue vector mosquitoes in Japan: the role of *Aedes albopictus* and *Aedes aegypti* in the 1942–1944 dengue epidemics of Japan main islands (in Japanese with English abstract). *Med Entomol Zool* 1998;**49**:267–74.
3. Effler PV, Pang L, Kitsutani P, Vorndam V, Makata M, Ayers T, et al. Dengue fever, Hawaii, 2001–2002. *Emerg Infect Dis* 2005;**11**:742–9.
4. Larrieu S, Dehacucq JS, Balleydier E, Jaffar MC, Michault A, Vilain P, et al. Re-emergence of dengue in Reunion, France, January to April 2012. *Euro Surveill* 2012;**17**. pii: 20173.
5. Kobayashi M, Nihei N, Kurihara T. Analysis of northern distribution of *Aedes albopictus* (Diptera: Culicidae) in Japan by geographical information system. *J Med Entomol* 2002;**39**:4–11.
6. European Center for Disease Prevention and Control (ECDC). Development of *Aedes albopictus* risk maps. ECDC Technical Report, 2009. Stockholm, Sweden: ECDC; 2009. Available at: [http://ecdc.europa.eu/en/publications/publications/0905\\_ter\\_development\\_of\\_aedes\\_albopictus\\_risk\\_maps.pdf](http://ecdc.europa.eu/en/publications/publications/0905_ter_development_of_aedes_albopictus_risk_maps.pdf) (accessed May 8, 2014).
7. Takasaki T. Imported dengue fever/dengue hemorrhagic fever cases in Japan. *Trop Med Health* 2011;**39**:13–5.
8. Schmidt-Chanasit J, Emmerich P, Tappe D, Gunther S, Schmidt S, Wolff D, et al. Autochthonous dengue virus infection in Japan imported into Germany, September 2013. *Euro Surveill* 2014;**19**. pii: 20681.