



Unparticle contribution to the hydrogen atom ground state energy



Michael F. Wondrak^{a,b}, Piero Nicolini^{a,b,*}, Marcus Bleicher^{a,b}

^a Frankfurt Institute for Advanced Studies (FIAS), Ruth-Moufang-Str. 1, 60438 Frankfurt am Main, Germany

^b Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität Frankfurt am Main, Max-von-Laue-Str. 1, 60438 Frankfurt am Main, Germany

ARTICLE INFO

Article history:

Received 29 April 2016

Accepted 6 June 2016

Available online 16 June 2016

Editor: A. Ringwald

Keywords:

Unparticle physics

Hydrogen ground state

Unparticle parameter constraints

ABSTRACT

In the present work we study the effect of unparticle modified static potentials on the energy levels of the hydrogen atom. By using Rayleigh–Schrödinger perturbation theory, we obtain the energy shift of the ground state and compare it with experimental data. Bounds on the unparticle energy scale Λ_U as a function of the scaling dimension d_U and the coupling constant λ are derived. We show that there exists a parameter region where bounds on Λ_U are stringent, signaling that unparticles could be tested in atomic physics experiments.

© 2016 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Unparticle physics is an extension of the Standard Model, consisting in the possibility of having a non-trivial scale invariant, yet undiscovered sector of particle physics.

At first sight, unparticles appear as a generalization of neutrinos because they share the following properties: scale invariance and only very weak interaction with other fields. Neutrinos enjoy the first property to a good approximation, although their oscillations disclose a small non-zero mass. The second property is a general requirement of any hypothetical particle sector because we want it to be, to a certain extent, hidden from current observations. On closer inspection, however, we find that unparticles differ drastically from neutrinos. Since we do not restrict the unparticle fields to be massless, we can no longer speak in terms of particle numbers as in the conventional manner. The unparticle field is controlled by a canonical scaling dimension d_U which is in general a non-integer number. Due to the unusual character, one refers to the matter described by such a theory as *unlike particles*, or unparticle stuff.

After Georgi's seminal paper [1] unparticle effects have been explored in many areas spanning collider physics [2–8], muonic

atoms [9], gauge and Higgs interactions [10,11], cosmology and astrophysics [12,13], AdS/CFT correspondence [14] as well as gravity short scale deviations [15,16] and black holes [17–21]. Unparticles also play a crucial role in the fractal properties of a quantum spacetime. A new fractality indicator, called un-spectral dimension, has recently been proposed to address the case of a random-walker problem in terms of an unparticle probe [22]. When the manifold topological dimension is 2, the un-spectral dimension turns out to be $2d_U$, *i.e.*, it depends only on the scaling dimension d_U . This fact explains the complete “fractalization” of the event horizon of un-gravity black holes [19,20], as well as of metallic plates for the Casimir effect in the presence of an un-photon field [23]. Finally, unparticles have been proposed to explain some anomalies in currents flowing in super-conductors [24] and transport phenomena in cuprates [25].

In this paper we want to address one of the basic parameters of unparticle physics, *i.e.*, the value of Λ_U , the typical energy scale of the theory. To achieve this goal, we consider the modifications of static potentials that emerge from virtual unparticle exchange. Specifically by considering the corrections to the Coulomb potential we calculate the deviations of the ground state energy of the hydrogen atom. We show that competitive bounds on Λ_U can be derived by a comparison with experimental data.

The paper is organized as follows. After a short review of the basic formalism of unparticle physics (Section 2), we present the calculation of the energy shift by a perturbative solution of the Schrödinger equation in the presence of an unparticle modified electrostatic potential (Section 3). Finally, in Section 4 we draw our conclusions.

* Corresponding author at: Frankfurt Institute for Advanced Studies (FIAS), Ruth-Moufang-Str. 1, 60438 Frankfurt am Main, Germany.

E-mail addresses: wondrak@fias.uni-frankfurt.de (M.F. Wondrak), nicolini@fias.uni-frankfurt.de (P. Nicolini), bleicher@fias.uni-frankfurt.de (M. Bleicher).




Fig. 1. Electron–proton interaction due to a vector field A_μ and an unparticle scalar field ϕ_U .

2. Unparticle physics and static potentials

We recap the basic motivations of unparticle physics along the lines of Georgi [1]. We start by considering that at some very high energy scale, the Standard Model is accompanied by an additional sector of Banks–Zaks fields (\mathcal{BZ}). The interaction between the two sectors takes place by exchange of mediating particles having a large mass scale M_U . If our energy scale of interest falls below M_U , we can apply effective field theory to integrate out the mediating field [26] and get the final interaction Lagrangian

$$\frac{1}{M_U^k} \mathcal{O}_{\text{SM}} \mathcal{O}_{\mathcal{BZ}} \quad (1)$$

where \mathcal{O}_{SM} denotes a Standard Model field operator of scaling dimension d_{SM} and $\mathcal{O}_{\mathcal{BZ}}$ is a Banks–Zaks field operator of scaling dimension $d_{\mathcal{BZ}}$. The factor M_U^{-k} guarantees the dimensional consistency of (1), being $k = d_{\text{SM}} + d_{\mathcal{BZ}} - D$ and D is the spacetime dimension.

If the energy is further decreased to $\Lambda_U < M_U$, the Banks–Zaks fields undergo a dimensional transmutation and exhibit a scale invariant behavior with a continuous mass distribution. For energies lower than Λ_U , the \mathcal{BZ} sector becomes unparticle operators \mathcal{O}_U . The matching conditions onto the Banks–Zaks operators are imposed at the energy scale Λ_U and determine the structure of the coupling between the Standard Model and the unparticle fields based on (1):

$$\frac{C_U \Lambda_U^{d_{\mathcal{BZ}} - d_U}}{M_U^k} \mathcal{O}_{\text{SM}} \mathcal{O}_U = \frac{\lambda}{\Lambda_U^{d_{\text{SM}} + d_U - D}} \mathcal{O}_{\text{SM}} \mathcal{O}_U \quad (2)$$

where d_U is the scaling dimension of the unparticle operator \mathcal{O}_U , C_U denotes a dimensionless constant and λ is a dimensionless coupling parameter defined by

$$\lambda = C_U \left(\frac{\Lambda_U}{M_U} \right)^k < 1. \quad (3)$$

The inequality holds if $C_U < 1$ and $d_{\mathcal{BZ}} > D - d_{\text{SM}}$. Any experimental bound on the interaction allows for constraints on the unparticle parameter space, i.e., Λ_U , λ and d_U . The scale hierarchy is given by $1 \text{ TeV} \leq \Lambda_U < M_U \leq M_{\text{Pl}}$, where M_{Pl} is the Planck mass [21]. In the rest of the present paper we assume $D = 4$ as well as the customary interval $1 < d_U < 2$ (cf. [2]). The case $d_U = 1$ does not give rise to fractalization or other continuous dimension effects of unparticle physics.

Unparticles have been largely employed in context of static potential emerging from virtual particle exchange [11,15,16,20,27]. Such results are instrumental to the working hypothesis of the current investigation. In view of the analysis of the hydrogen atom, we consider an additional contribution to the Coulomb potential for the presence of unparticle exchange in the interaction between electron and proton (cf. Fig. 1). In general, we assume electrons and protons to carry unparticle charges λ_e and λ_p , respectively, and a scalar unparticle field ϕ_U . At the same time, both particles also possess electric charges $\pm e$ allowing them to couple to the photon field A_μ . Both interactions are independent of each other since we assume the effective couplings between unparticle stuff




Fig. 2. Interaction energy per unit unparticle charge with respect to the radial distance r in the case $\Lambda_U = 1$ TeV. The unparticle scaling dimension d_U is raised from $d_U = 1.0$ to $d_U = 2.0$ in steps of 0.2 corresponding to growing dash lengths.

and photons to be negligible. Therefore the interaction Lagrangian can be written as

$$\mathcal{L}_{\text{int}} = J^\mu A_\mu + \frac{1}{(\Lambda_U)^{d_U - 1}} J_U \phi_U \quad (4)$$

where $J^\mu = j(\vec{x}) \delta_0^\mu$ and $J_U = j_U(\vec{x})$ and

$$j(\vec{x}) = -e \delta(\vec{x} - \vec{x}_e) + e \delta(\vec{x} - \vec{x}_p) \quad (5)$$

$$j_U(\vec{x}) = \lambda_e \delta(\vec{x} - \vec{x}_e) + \lambda_p \delta(\vec{x} - \vec{x}_p). \quad (6)$$

Alternatively one can consider the interaction Lagrangian¹ for a vector unparticle field $(A_U)_\mu$ and derive the static potential much in the same way as in the scalar case. The two results do not differ apart from a global sign (see e.g. [16]). We recall, however, that conformal invariance can be lost for vector fields if $d_U < 3$, although pure scale invariance can be preserved. For scalar unparticles the issue does not arise [26].

The expression for the unparticle interaction energy V_U between an electron and a proton in the static case reads [11,15,16,20,27]

$$V_U = -\xi_{d_U} \left(\frac{\lambda_e \lambda_p}{\Lambda_U^{2d_U - 2} r^{2d_U - 1}} \right) \quad (7)$$

where the coefficient is $\xi_{d_U} \equiv \frac{\sqrt{\pi}}{(2\pi)^{2d_U}} \frac{\Gamma(d_U - \frac{1}{2})}{\Gamma(d_U)}$ and $r \equiv |\vec{x}_e - \vec{x}_p|$ denotes the distance between the charges.

For the typical size of the hydrogen atom we can estimate the energy correction with the help of Fig. 2. For $r = 0.05$ nm the energy shift per unit unparticle charge varies between 3×10^2 eV and 6×10^{-19} eV in the range $1 < d_U < 2$ with $\Lambda_U = 1$ TeV.

3. Unparticle effects in the hydrogen atom

The hydrogen atom is a well investigated system. In the non-relativistic limit we can apply the Schrödinger formalism to describe electron dynamics in a static Coulomb potential of the proton, by using the particle reduced mass μ . The energy spectrum in Gaussian/natural units ($4\pi\epsilon_0 = \hbar = c = 1$) reads

$$E_{\text{th}, n}^S = -\frac{\mu \alpha^2}{2n^2} \quad (8)$$

¹ The Lagrangian in such a case reads

$$\mathcal{L}_{\text{int}} = J^\mu A_\mu + \frac{1}{(\Lambda_U)^{d_U - 1}} J_U^\mu (A_U)_\mu$$

where $J_U^\mu = j_U(\vec{x}) \delta_0^\mu$.

Table 1
Theoretical and experimental values of the hydrogen ground state energy.

Energy	Description	Value
E_{th}^S	Schrödinger, non-relativistic	−13.598 287 eV
$E_{\text{th}}^{S, \text{rel}}$	Schrödinger, incl. fine-structure	−13.598 468 eV
E_{th}^D	Dirac	−13.598 468 eV
$E_{\text{th}}^{\text{QED}}$	currently best theoretical value [29]	−13.598 434 49 (9) eV 3 288 086 857.127 6 (31) MHz · h
E_{exp}	currently best experimental value [30]	−13.598 434 48 (9) eV 3 288 086 856.8 (0.7) MHz · h

for all $n \in \mathbb{N}$. Here we used the fine-structure constant $\alpha \equiv e^2$. The biggest corrections to the Schrödinger description are of order α^4 and arise from relativistic effects (*i.e.* kinetic, spin–orbit and Darwin terms) that can be included via fine-structure modifications of the Hamiltonian.

There exist higher order corrections as well. The proton possesses a finite size which affects the form of the Coulomb potential for distances shorter than its radius. Since the proton has a spin which interacts with both the electron angular motion and the electron spin, hyperfine corrections arise. Finally, one can include QED corrections such as the Lamb shift. In Table 1, the ground state energy in different descriptions is displayed. Note that the uncertainty of the two most accurate values arises from the uncertainty in the Planck constant $h = 4.135\,667\,662\,(25) \times 10^{-15}$ eVs [28].

We start our analysis by considering the energy level shift due to unparticle effects within the non-relativistic description. The radial Schrödinger equation of the hydrogen atom in the presence of both electrostatic and unparticle potentials reads

$$H u_{nl}(r) = E_{nlm} u_{nl}(r) \quad (9)$$

with the Hamiltonian

$$H = -\frac{1}{2\mu} \frac{d^2}{dr^2} + \frac{l(l+1)}{2\mu r^2} - \frac{e^2}{r} - \xi_{d_U} \frac{\lambda_e \lambda_p}{\Lambda_U^{2d_U-2} r^{2d_U-1}} \quad (10)$$

where the complete wavefunction is given by

$$\psi_{nlm}(r, \vartheta, \varphi) = \frac{u_{nl}(r)}{r} Y_{lm}(\vartheta, \varphi). \quad (11)$$

The above equation is a second order, linear, ordinary differential equation with non-polynomial coefficients. To our best knowledge no exact solution is available in the literature but one can still rely on perturbation theory. This is justified since $\Lambda_U > 1$ TeV by hypothesis, *i.e.*, unparticle physics can lead only to subleading corrections to the Standard Model. As a result the energy shift of the ground state at the first order in perturbation theory reads

$$\begin{aligned} \Delta_{100}^{(1)} &= \left\langle 100^{(0)} \left| -\xi_{d_U} \frac{\lambda_e \lambda_p}{\Lambda_U^{2d_U-2} r^{2d_U-1}} \right| 100^{(0)} \right\rangle \\ &= -4 \xi_{d_U} \frac{\lambda_e \lambda_p}{\Lambda_U^{2d_U-2} a^3} \int_0^\infty dr r^{-2d_U+3} e^{-\frac{2r}{a}} \\ &= -\frac{1}{(2\pi)^{2d_U-2}} \frac{(d_U-1) \left(\frac{3}{2} - d_U\right)}{\sin(2\pi d_U) (\Gamma(d_U))^2} \\ &\quad \times \frac{\lambda_e \lambda_p}{\Lambda_U^{2d_U-2} a^{2d_U-1}}, \end{aligned} \quad (12)$$

where we used $\psi_{100}^{(0)} = \frac{2}{\sqrt{4\pi a^3}} e^{-\frac{r}{a}}$ with $a \equiv 1/(\alpha\mu) \simeq 5.29 \times 10^{-11}$ m.

For $1 < d_U < 2$, the sign of $\Delta_{100}^{(1)}$ is determined by the product of λ_e and λ_p . This means that like unparticle charges lead to a shift to lower energies while unlike charges cause a shift to higher energies.

The theoretical energy value E_{th} can be divided up into the Schrödinger contribution E_{th}^S and higher order terms $E_{\text{th}}^{\text{HO}}$. The same applies to the unparticle-modified quantities. As a result we can approximate the energy shift as

$$\begin{aligned} |E_{\text{th}, U} - E_{\text{th}}| &= \left| \left(E_{\text{th}, U}^S + E_{\text{th}, U}^{\text{HO}} \right) - \left(E_{\text{th}}^S + E_{\text{th}}^{\text{HO}} \right) \right| \\ &\simeq \left| E_{\text{th}, U}^S - E_{\text{th}}^S \right|. \end{aligned} \quad (13)$$

Here $E_{\text{th}, U}^{\text{HO}}$ stands for all the higher order contributions in the presence of unparticles, contributions we deliberately neglected in our first order description. This assumption introduces a theoretical error $\delta E_{\text{th}} \sim \left| E_{\text{th}, U}^{\text{HO}} - E_{\text{th}}^{\text{HO}} \right|$. Thus the bound on the energy shift can be written as

$$\left| E_{\text{th}, U}^S - E_{\text{th}}^S \right| = \left| \Delta_{100}^{(1)} \right| < \delta E_{\text{th}} + \delta E_{\text{exp}}, \quad (14)$$

where δE_{exp} is the experimental error. From Table 1, one can see that the Schrödinger description differs from the QED result by about 1.5×10^{-4} eV. The theoretical relative error can be estimated to be $\delta E_{\text{th}}/|E_{\text{th}}^S| \simeq 1.1 \times 10^{-5}$. It dominates over the experimental relative error that is of the order of 2.2×10^{-10} [30]. Accordingly we define $\delta_{\text{max}} \equiv (\delta E_{\text{th}} + \delta E_{\text{exp}})/|E_{\text{th}}^S| \simeq 1.1 \times 10^{-5}$ to get

$$\begin{aligned} \left| \frac{\Delta_{100}^{(1)}}{E_{\text{th}}^S} \right| &= \frac{2\alpha^{2d_U-3} \mu^{2d_U-2} (d_U-1) \left(\frac{3}{2} - d_U\right) |\lambda_e \lambda_p|}{(2\pi)^{2d_U-2} \sin(2\pi d_U) (\Gamma(d_U))^2 \Lambda_U^{2d_U-2}} \\ &< \delta_{\text{max}}. \end{aligned} \quad (15)$$

From (15) we obtain

$$\Lambda_U \geq \frac{\alpha \mu}{2\pi} \left(\frac{2}{\alpha} \frac{(d_U-1) \left(\frac{3}{2} - d_U\right) |\lambda_e \lambda_p|}{\sin(2\pi d_U) (\Gamma(d_U))^2 \delta_{\text{max}}} \right)^{\frac{1}{2d_U-2}}. \quad (16)$$

In Fig. 3 the lower bounds for Λ_U are illustrated for different values of the unparticle charge λ , where we assumed $|\lambda_e| = |\lambda_p| \equiv \lambda$. The area above each curve is the allowed parameter space for the chosen value of λ . If the parameter λ is larger than a threshold value, $\lambda > \lambda_{\text{th}} \equiv \sqrt{2\pi\alpha \delta_{\text{max}}} \simeq 7.1 \times 10^{-4}$, the lower bound on Λ_U is diverging for $d_U \rightarrow 1$ and exceeds 1 TeV in some parameter region. Here λ_{th} is obtained by requiring that the base of the exponentiation in (16) equals 1. The limit $\Lambda_U \rightarrow \infty$ corresponds to vanishing unparticle effects since the unparticle sector in the Lagrangian (4) disappears yielding the standard electrodynamics limit $\mathcal{L}_{\text{int}} \rightarrow J^\mu A_\mu$. This explains the divergent lower bound on Λ_U expected also in other experiments such as the electron $g-2$ anomaly [8] or the Casimir effect [23]. In contrast, for $\lambda \leq \lambda_{\text{th}}$, the




Fig. 3. Lower bounds for the energy scale Λ_U with respect to the unparticle scaling dimension d_U . The unparticle charge $\lambda \equiv |\lambda_e| = |\lambda_p|$ increases exponentially from $\lambda = 1.0 \times 10^{-6}$ to $\lambda = 1.0 \times 10^{-1}$ with decreasing dash lengths. The additional uppermost curve belongs to $\lambda = 9.0 \times 10^{-1}$.

scale Λ_U is weakly constrained in the limit $d_U \rightarrow 1$. This means that the unparticle contribution lies within δ_{\max} irrespectively of the value of Λ_U .

4. Discussion and conclusions

In the current work, we have derived the corrections to the hydrogen atom ground state energy due to the presence of an unparticle modified static potential. We obtained this result within a perturbative analysis of the non-relativistic theory of the hydrogen atom. Our result lets obtain compelling limits on the unparticle scale Λ_U . For $d_U \lesssim 1.3$, bounds on Λ_U exceed 1 TeV. Contrary to other proposed investigations (e.g. Newton's law correction [15], proton–proton collisions at the LHC [5,6]), the presented analysis can capture the key feature of unparticle physics, i.e., the dependence on the continuous scaling dimension d_U .

The presented results are filling a gap in the literature and are opening the route for further investigations. For instance one may be interested to find an exact solution of the Schrödinger equation (9) emerging from the inclusion of unparticle static potentials. Higher order corrections to the non-relativistic description can also be included in the analysis in order to approach the bounds currently offered by $g - 2$ analyses [8].

From the current non-relativistic analysis one can draw an important conclusion: Unparticle effects might be tested in atomic physics experiments.

Acknowledgements

This work has been supported by the Studienstiftung des deutschen Volkes and by the Helmholtz International Center for FAIR within the framework of the LOEWE program (Landesoffensive zur Entwicklung Wissenschaftlich-Ökonomischer Exzellenz) launched by the State of Hesse.

References

- [1] H. Georgi, Unparticle physics, *Phys. Rev. Lett.* 98 (2007) 221601.
- [2] H. Georgi, Another odd thing about unparticle physics, *Phys. Lett. B* 650 (2007) 275–278.
- [3] K. Cheung, W.-Y. Keung, T.-C. Yuan, Collider phenomenology of unparticle physics, *Phys. Rev. D* 76 (2007) 055003.
- [4] K. Cheung, W.-Y. Keung, T.-C. Yuan, Collider signals of unparticle physics, *Phys. Rev. Lett.* 99 (2007) 051803.
- [5] CMS Collaboration, Search for dark matter, extra dimensions, and unparticles in monojet events in proton–proton collisions at $\sqrt{s} = 8$ TeV, *Eur. Phys. J. C* 75 (2015) 235.
- [6] CMS Collaboration, Search for dark matter and unparticles produced in association with a Z boson in pp collisions at $\sqrt{s} = 8$ TeV, CMS Physics Analysis Summary CMS PAS EXO-12-054, CERN, 2015.
- [7] S. Kathrein, S. Knäpen, M.J. Strassler, Bounds from LEP on unparticle interactions with electroweak bosons, *Phys. Rev. D* 84 (2011) 015010.
- [8] Y. Liao, Bounds on unparticle couplings to electrons: from electron $g - 2$ to positronium decays, *Phys. Rev. D* 76 (2007) 056006.
- [9] A.M. Thalappilil, Bound states and fermiophobic unparticle oblique corrections to the photon, *Phys. Rev. D* 81 (2010) 035001.
- [10] P.J. Fox, A. Rajaraman, Y. Shirman, Bounds on unparticles from the Higgs sector, *Phys. Rev. D* 76 (2007) 075004.
- [11] P. Gaete, E. Spallucci, The Abelian Higgs model and a minimal length in an unparticle scenario, *Europhys. Lett.* 105 (2014) 21002.
- [12] H. Davoudiasl, Constraining unparticle physics with cosmology and astrophysics, *Phys. Rev. Lett.* 99 (2007) 141301.
- [13] S. Das, S. Mohanty, K. Rao, Test of unparticle long range forces from perihelion precession of Mercury, *Phys. Rev. D* 77 (2008) 076001.
- [14] G. Cacciapaglia, G. Marandella, J. Terning, The AdS/CFT/Unparticle correspondence, *J. High Energy Phys.* 02 (2009) 049.
- [15] H. Goldberg, P. Nath, Scalar modifications to gravity from unparticle effects may be testable, *Phys. Rev. Lett.* 100 (2008) 031803.
- [16] P. Gaete, E. Spallucci, Unparticle effective action, *Phys. Lett. B* 661 (2008) 319–324.
- [17] J.R. Mureika, Unparticle-enhanced black holes at the LHC, *Phys. Lett. B* 660 (2008) 561–566.
- [18] J.R. Mureika, Differentiating unparticles from extra dimensions via mini black hole thermodynamics, *Phys. Rev. D* 79 (2009) 056003.
- [19] P. Gaete, J.A. Helayel-Neto, E. Spallucci, Un-graviton corrections to the Schwarzschild black hole, *Phys. Lett. B* 693 (2010) 155–158.
- [20] J.R. Mureika, E. Spallucci, Vector unparticle enhanced black holes: exact solutions and thermodynamics, *Phys. Lett. B* 693 (2010) 129–133.
- [21] J.R. Mureika, Constraints on vector unparticle physics from cosmic censorship, *Int. J. Theor. Phys.* 51 (2012) 1259–1267.
- [22] P. Nicolini, E. Spallucci, Un-spectral dimension and quantum spacetime phases, *Phys. Lett. B* 695 (2011) 290–293.
- [23] A.M. Frassino, P. Nicolini, O. Panella, Un-Casimir effect, arXiv:1311.7173v2 [hep-ph], unpublished results.
- [24] J.P.F. LeBlanc, A.G. Grushin, Unparticle mediated superconductivity, *New J. Phys.* 17 (2015) 033039.
- [25] A. Karch, K. Limtragool, P.W. Phillips, Unparticles and anomalous dimensions in the cuprates, *J. High Energy Phys.* 03 (2016) 175.
- [26] B. Grinstein, K. Intriligator, I.Z. Rothstein, Comments on unparticles, *Phys. Lett. B* 662 (2008) 367–374.
- [27] N.G. Deshpande, S.D.H. Hsu, J. Jiang, Long range forces and limits on unparticle interactions, *Phys. Lett. B* 659 (2008) 888–890.
- [28] P.J. Mohr, D.B. Newell, B.N. Taylor, CODATA Recommended Values of the Fundamental Physical Constants, 2014, <http://dx.doi.org/10.5281/zenodo.22826>, Aug. 2015, accessed 11 January 2016.
- [29] U.D. Jentschura, S. Kotochigova, E.O. LeBigot, P.J. Mohr, B.N. Taylor, The Energy Levels of Hydrogen and Deuterium (version 2.1), National Institute of Standards and Technology, Gaithersburg, MD, 2005, <http://physics.nist.gov/HDEL>, 2005, accessed 11 January 2016.
- [30] A.E. Kramida, A critical compilation of experimental data on spectral lines and energy levels of hydrogen, deuterium, and tritium, *At. Data Nucl. Data Tables* 96 (2010) 586–644.