

Available online at www.sciencedirect.com**SciVerse ScienceDirect**

Advances in Mathematics 231 (2012) 1401–1416

**ADVANCES IN
Mathematics**

www.elsevier.com/locate/aim

A new series expansion for slice regular functions

Caterina Stoppato

Università degli Studi di Milano, Dipartimento di Matematica “F. Enriques”, Via Saldini 50, 20133 Milano, Italy

Received 5 September 2011; accepted 18 May 2012

Available online 10 August 2012

Communicated by Edoardo Vesentini

Abstract

A promising theory of quaternion-valued functions of one quaternionic variable, now called *slice regular* functions, has been introduced by Gentili and Struppa in 2006. The basic examples of slice regular functions are the power series of type $\sum_{n \in \mathbb{N}} q^n a_n$ on their balls of convergence $B(0, R) = \{q \in \mathbb{H} : |q| < R\}$. Conversely, if f is a slice regular function on a domain $\Omega \subseteq \mathbb{H}$ then it admits at each point $q_0 \in \Omega$ an expansion of type $f(q) = \sum_{n \in \mathbb{N}} (q - q_0)^{*n} a_n$ where $(q - q_0)^{*n}$ denotes the n th power of $q - q_0$ with respect to an appropriately defined multiplication $*$. However, the information provided by such an expansion is somewhat limited by a fact: if q_0 does not lie on the real axis then the set of convergence of the series in the previous equation needs not be a Euclidean neighborhood of q_0 . We are now able to construct a new type of expansion that is not affected by this phenomenon: an expansion into series of polynomials valid in open subsets of the domain. Along with this construction, we present applications to the computation of the multiplicities of zeros and of partial derivatives.

© 2012 Elsevier Inc. All rights reserved.

Keywords: Functions of a hypercomplex variable; Slice regular function; Quaternionic series expansion; Quaternionic analytic function

1. Introduction

Let \mathbb{H} denote the real algebra of quaternions, that is the vector space \mathbb{R}^4 endowed with the multiplication constructed as follows: if $1, i, j, k$ denotes the standard basis, define

$$i^2 = j^2 = k^2 = -1,$$

$$ij = -ji = k, \quad jk = -kj = i, \quad ki = -ik = j,$$

E-mail address: caterina.stoppato@unimi.it.

0001-8708/\$ - see front matter © 2012 Elsevier Inc. All rights reserved.

doi:10.1016/j.aim.2012.05.023

let 1 be the neutral element and extend the operation by distributivity and linearity to all quaternions $q = x_0 + x_1i + x_2j + x_3k$. The *conjugate* of such a q is defined as $\bar{q} = x_0 - x_1i - x_2j - x_3k$, its *real* and *imaginary parts* as $\text{Re}(q) = x_0$ and $\text{Im}(q) = x_1i + x_2j + x_3k$, and its *modulus* as $|q| = \sqrt{q\bar{q}} = \sqrt{\text{Re}(q)^2 + |\text{Im}(q)|^2}$. The multiplicative inverse of each $q \neq 0$ is computed as

$$q^{-1} = \frac{\bar{q}}{|q|^2}.$$

Much literature has been devoted to the possibility of defining for quaternionic functions a notion of regularity playing the same role as holomorphy for complex functions. This is by no means an elementary question, as neatly explained in [19]. First of all, let us consider the following fact.

Theorem 1.1. *A function $f : \mathbb{H} \rightarrow \mathbb{H}$ is left q -differentiable at every $q \in \mathbb{H}$, i.e.*

$$\lim_{h \rightarrow 0} h^{-1}[f(q + h) - f(q)]$$

exists at every $q \in \mathbb{H}$, if and only if there exist $a, b \in \mathbb{H}$ such that $f(q) = qa + b$ for all $q \in \mathbb{H}$.

A proof can be found in [19]. The class of functions encompassed does not grow significantly if we vary the domain of definition or consider the analogous notion of right q -differentiability. The next corollary can be easily derived.

Corollary 1.2. *The class of functions $f : \mathbb{H} \rightarrow \mathbb{H}$ admitting at every $q_0 \in \mathbb{H}$ a power series expansion*

$$f(q) = \sum_{n \in \mathbb{N}} (q - q_0)^n a_n$$

with $\{a_n\}_{n \in \mathbb{N}}$ coincides with the class of affine functions $f(q) = qa + b$ with $a, b \in \mathbb{H}$.

Once again, the situation changes but little if we consider an expansion of type $f(q) = \sum_{n \in \mathbb{N}} a_n (q - q_0)^n$ instead. It should be noticed, however, that $(q - q_0)^n a_n$ and $a_n (q - q_0)^n$ are not the only possible terms of degree n in $q - q_0$: in fact, in an (associative) non-commutative framework, the generic form of such a term is

$$\alpha_0(q - q_0)\alpha_1(q - q_0) \dots \alpha_{n-1}(q - q_0)\alpha_n. \tag{1}$$

We may try to define a notion of analyticity based on series of terms of this type, but we are immediately discouraged by the following fact.

Proposition 1.3. *The class of functions $f : \mathbb{H} \rightarrow \mathbb{H}$ admitting at every $q_0 \in \mathbb{H}$ an expansion into series of monomials of type (1) coincides with the class of functions $f : \mathbb{H} \rightarrow \mathbb{H}$ that are analytic in the four real variables x_0, \dots, x_3 .*

The proof of the preceding proposition relies upon the fact that if $q = x_0 + ix_1 + jx_2 + kx_3$, then

$$\begin{aligned} x_0 &= \frac{1}{4}(q - iqi - jqj - kqk), & x_1 &= \frac{1}{4i}(q - iqi + jqj + kqk), \\ x_2 &= \frac{1}{4j}(q + iqi - jqj + kqk), & x_3 &= \frac{1}{4k}(q + iqi + jqj - kqk). \end{aligned}$$

The facts just mentioned encouraged to base the study of quaternionic functions on analogs of the Cauchy–Riemann equations rather than on some notion of analyticity. The best known of such analogs is due to Fueter [5,6] and it gave rise to a renowned function theory on Clifford algebras, for which we refer the reader to [1,4,15] and references therein. Let us also mention the work [16] on the specific topic of analyticity in this context.

A different notion of quaternionic analyticity arose from the theory introduced by Gentili and Struppa in [10,11]. They gave the following definition, where $\mathbb{S} = \{q \in \mathbb{H} : q^2 = -1\}$ denotes the 2-sphere of *imaginary units*.

Definition 1.4. Let Ω be a domain in \mathbb{H} and let $f : \Omega \rightarrow \mathbb{H}$ be a function. For all $I \in \mathbb{S}$, let us denote $L_I = \mathbb{R} + I\mathbb{R}$, $\Omega_I = \Omega \cap L_I$ and $f_I = f|_{\Omega_I}$. The function f is called (*slice*) *regular* if, for all $I \in \mathbb{S}$, the restriction f_I is holomorphic, i.e. the function $\bar{\partial}_I f : \Omega_I \rightarrow \mathbb{H}$ defined by

$$\bar{\partial}_I f(x + Iy) = \frac{1}{2} \left(\frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f_I(x + Iy)$$

vanishes identically.

The definition immediately implies that any power series $f(q) = \sum_{n \in \mathbb{N}} q^n a_n$ defines a regular function on its ball of convergence

$$B(0, R) = \{q \in \mathbb{H} : |q| < R\}$$

(a perfect analog of Abel’s theorem holds). Moreover, the set of such series forms a real algebra when endowed with the usual addition $+$ and the multiplication $*$ defined in the following manner:

$$\left(\sum_{n \in \mathbb{N}} q^n a_n \right) * \left(\sum_{n \in \mathbb{N}} q^n b_n \right) = \sum_{n \in \mathbb{N}} q^n \sum_{k=0}^n a_k b_{n-k}. \tag{2}$$

In [9], we considered series of the form

$$f(q) = \sum_{n \in \mathbb{N}} (q - q_0)^{*n} a_n \tag{3}$$

where $(q - q_0)^{*n} = (q - q_0) * \dots * (q - q_0)$ denotes the $*$ -product of n copies of $q \mapsto q - q_0$. We were able to prove that the sets of convergence of such series are balls with respect to a distance $\sigma : \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{R}$ defined in the following fashion.

Definition 1.5. For all $p, q \in \mathbb{H}$, we set

$$\sigma(q, p) = \begin{cases} |q - p| & \text{if } p, q \text{ lie on the same complex plane } L_I \\ \omega(q, p) & \text{otherwise} \end{cases} \tag{4}$$

where

$$\omega(q, p) = \sqrt{[\operatorname{Re}(q) - \operatorname{Re}(p)]^2 + [|\operatorname{Im}(q)| + |\operatorname{Im}(p)|]^2}. \tag{5}$$

More precisely, we proved the next theorem.

Theorem 1.6. *If Ω is a domain in \mathbb{H} , a function $f : \Omega \rightarrow \mathbb{H}$ is regular if and only if it is σ -analytic, i.e. it admits at every $q_0 \in \Omega$ an expansion of type (3) that is valid in a σ -ball $\Sigma(q_0, R) = \{q \in \mathbb{H} : \sigma(q, q_0) < R\}$.*

The analogy with the complex case is remarkable, but it should be taken into account that the topology induced by σ is finer than the Euclidean: if $q_0 = x_0 + Iy_0$ does not lie on the real axis then for $R < 2y_0$ the σ -ball $\Sigma(q_0, R)$ reduces to a (2-dimensional) disk $\{z \in L_I : |z - q_0| < R\}$ in the complex plane L_I through q_0 (see [9] for a detailed account on σ -balls). Hence the series expansion (3), in general, may not predict the behavior of f in a Euclidean neighborhood of q_0 , but only along the complex plane L_I containing q_0 . This curious phenomenon is partly explained by the fact that for a generic domain Ω in \mathbb{H} , a regular function $f : \Omega \rightarrow \mathbb{H}$ needs not be continuous, as shown by the next example.

Example 1.7. For a fixed $I \in \mathbb{S}$, we can define a regular function $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ by setting

$$f(q) = \begin{cases} 0 & \text{if } q \in \mathbb{H} \setminus L_I \\ 1 & \text{if } q \in L_I \setminus \mathbb{R}. \end{cases}$$

Clearly, f is not continuous.

However, [3] explained that real differentiability (and other interesting properties) are granted if Ω is carefully chosen. First, let us consider the following class of domains.

Definition 1.8. Let Ω be a domain in \mathbb{H} , intersecting the real axis. If, for all $I \in \mathbb{S}$, $\Omega_I = \Omega \cap L_I$ is a domain in $L_I \simeq \mathbb{C}$ then Ω is called a *slice domain*.

Theorem 1.9 (Identity Principle). Let Ω be a slice domain and let $f, g : \Omega \rightarrow \mathbb{H}$ be slice regular. Suppose that f and g coincide on a subset C of Ω_I , for some $I \in \mathbb{S}$. If C has an accumulation point in Ω_I , then $f \equiv g$ in Ω .

Second, let us consider slice domains having the additional property of axial symmetry with respect to the real axis, i.e. those slice domains Ω such that

$$\Omega = \bigcup_{x+Iy \in \Omega} x + y\mathbb{S}.$$

The word *symmetric* will refer to this type of symmetry throughout the paper. A very peculiar property holds for regular functions f on symmetric slice domains: they are affine when restricted to a single 2-sphere $x + y\mathbb{S}$. Indeed, the Representation Formula proven in [3] can be restated as follows.

Theorem 1.10. Let f be a regular function on a symmetric slice domain Ω and let $x_0 + y_0\mathbb{S} \subset \Omega$. For all $q, q_0, q_1, q_2 \in x_0 + y_0\mathbb{S}$ with $q_1 \neq q_2$

$$f(q) = (q_2 - q_1)^{-1} [\bar{q}_1 f(q_1) - \bar{q}_2 f(q_2)] + q(q_2 - q_1)^{-1} [f(q_2) - f(q_1)] \tag{6}$$

and

$$f(q) = f(q_0) + (q - q_0)(q_1 - q_2)^{-1} [f(q_1) - f(q_2)] \tag{7}$$

where $(q_2 - q_1)^{-1} [f(q_2) - f(q_1)]$ and $(q_2 - q_1)^{-1} [\bar{q}_1 f(q_1) - \bar{q}_2 f(q_2)]$ do not depend on the choice of q_1, q_2 , but only on x_0, y_0 .

As an immediate consequence of Theorem 1.10, if f is a regular function on a symmetric slice domain then its values can all be recovered from those of one of its restrictions f_I . This allowed a further study of these functions in [3], including the construction of a structure of real algebra for regular functions on a symmetric slice domain Ω (with the usual addition $+$ and a

multiplication $*$ that extends the one defined in Eq. (2)) and the observation that a regular f on a symmetric slice domain is $C^\infty(\Omega)$. This last result has been improved to $f \in C^\omega(\Omega)$ in [13]. This makes it reasonable to expect, in the case of symmetric slice domains, a stronger form of analyticity than the one presented in Theorem 1.6. Finding this stronger notion of analyticity is the aim of the present paper, which introduces a new type of series expansions and proves their validity in Euclidean open sets.

We start in Section 2 with a regular function f on a symmetric slice domain Ω , and we construct a formal expansion of the form

$$f(q) = \sum_{n \in \mathbb{N}} [(q - x_0)^2 + y_0^2]^n [A_{2n} + (q - q_0)A_{2n+1}] \tag{8}$$

at each point $q_0 = x_0 + Iy_0 \in \Omega$. We then prove what follows.

Proposition 1.11. *Let $\{a_n\}_{n \in \mathbb{N}} \subset \mathbb{H}$ and suppose*

$$\limsup_{n \rightarrow +\infty} |a_n|^{1/n} = 1/R \tag{9}$$

for some $R > 0$. Let $q_0 = x_0 + Iy_0 \in \mathbb{H}$ with $x_0 \in \mathbb{R}, y_0 > 0, I \in \mathbb{S}$ and set $P_{2n}(q) = [(q - x_0)^2 + y_0^2]^n$ and $P_{2n+1}(q) = [(q - x_0)^2 + y_0^2]^n (q - q_0)$ for all $n \in \mathbb{N}$. Then the function series $\sum_{n \in \mathbb{N}} P_n(q)a_n$ converges absolutely and uniformly on compact sets in

$$U(x_0 + y_0\mathbb{S}, R) = \{q \in \mathbb{H} : |(q - x_0)^2 + y_0^2| < R^2\}, \tag{10}$$

where it defines a regular function. Furthermore, the series diverges at every $q \in \mathbb{H} \setminus \overline{U(x_0 + y_0\mathbb{S}, R)}$.

In Section 3, we obtain estimates for the coefficients of our new formal expansion (8). More precisely, we prove that for every $U = U(x_0 + y_0\mathbb{S}, R)$ such that $\overline{U} \subset \Omega$, there exists a constant $C > 0$ such that

$$|A_n| \leq C \cdot \frac{\max_U |f|}{R^n}.$$

This result is achieved by means of new integral representations.

In Section 4 we define f to be *symmetrically analytic* if it admits at any $q_0 \in \Omega$ an expansion of type (8) valid in a neighborhood of q_0 . We are able to prove that regularity is equivalent to symmetric analyticity.

In Section 5, we apply the new series expansion (8) to the computation of the multiplicities of the zeros of regular functions. Finally, in Section 6, we apply it to the computation of directional derivatives.

Theorem 1.12. *Let f be a regular function on a symmetric slice domain Ω , and let $q_0 = x_0 + Iy_0 \in \Omega$. For all $v \in \mathbb{H}, |v| = 1$ the derivative of f along v can be computed at q_0 as*

$$\lim_{t \rightarrow 0} \frac{f(q_0 + tv) - f(q_0)}{t} = vA_1 + (q_0v - v\bar{q}_0)A_2. \tag{11}$$

We conclude by studying the bearings of this result in (real and) complex coordinates: it turns out that if we fix q_0 and choose appropriate complex coordinates on \mathbb{H} , then f is complex differentiable at q_0 and its Jacobian at the same point can be easily computed in terms of A_1 and A_2 .

2. A new series expansion

Let us recall a result from [18] (extending [7]) and derive from it the subsequent theorem.

Proposition 2.1. *Let f be a regular function on a symmetric slice domain Ω . A point $q_0 \in \Omega$ is a zero of f if and only if there exists a regular function $g : \Omega \rightarrow \mathbb{H}$ such that $f(q) = (q - q_0) * g(q)$.*

Theorem 2.2. *Let f be a regular function on a symmetric slice domain Ω . For each $q_0 \in \Omega$, let us denote as $R_{q_0}f : \Omega \rightarrow \mathbb{H}$ the function such that*

$$f(q) = f(q_0) + (q - q_0) * R_{q_0}f(q).$$

If $q_0 = x_0 + Iy_0$ for $x_0, y_0 \in \mathbb{R}$ and for some $I \in \mathbb{S}$, then

$$f(q) = f(q_0) + (q - q_0)R_{q_0}f(\bar{q}_0) + [(q - x_0)^2 + y_0^2]R_{\bar{q}_0}R_{q_0}f(q)$$

for all $q \in \Omega$.

Proof. The existence of a regular $R_{q_0}f : \Omega \rightarrow \mathbb{H}$ such that

$$f(q) = f(q_0) + (q - q_0) * R_{q_0}f(q),$$

is granted by the previous proposition, since $f - f(q_0)$ is a regular function on Ω vanishing at q_0 . Applying the same procedure to $R_{q_0}f$ at the point \bar{q}_0 yields

$$\begin{aligned} f(q) &= f(q_0) + (q - q_0) * [R_{q_0}f(\bar{q}_0) + (q - \bar{q}_0) * R_{\bar{q}_0}R_{q_0}f(q)] \\ &= f(q_0) + (q - q_0)R_{q_0}f(\bar{q}_0) + [(q - x_0)^2 + y_0^2]R_{\bar{q}_0}R_{q_0}f(q) \end{aligned}$$

where we have taken into account that

$$\begin{aligned} (q - q_0) * (q - \bar{q}_0) &= q^2 - q(q_0 + \bar{q}_0) + q_0\bar{q}_0 \\ &= q^2 - q2x_0 + x_0^2 + y_0^2 = (q - x_0)^2 + y_0^2. \quad \square \end{aligned}$$

If we repeatedly apply the previous theorem, we get the formal expansion

$$\begin{aligned} f(q) &= f(q_0) + (q - q_0) R_{q_0}f(\bar{q}_0) + [(q - x_0)^2 + y_0^2] \\ &\quad \times [R_{\bar{q}_0}R_{q_0}f(q_0) + (q - q_0)R_{q_0}R_{\bar{q}_0}R_{q_0}f(\bar{q}_0)] + \dots \\ &\quad + [(q - x_0)^2 + y_0^2]^n [(R_{\bar{q}_0}R_{q_0})^n f(q_0) + (q - q_0)R_{q_0}(R_{\bar{q}_0}R_{q_0})^n f(\bar{q}_0)] + \dots \end{aligned}$$

where $(R_{\bar{q}_0}R_{q_0})^n$ denotes the n th iterate of $R_{\bar{q}_0}R_{q_0}$. If $A_{2n} = (R_{\bar{q}_0}R_{q_0})^n f(q_0)$ and $A_{2n+1} = R_{q_0}(R_{\bar{q}_0}R_{q_0})^n f(\bar{q}_0)$ for all $n \in \mathbb{N}$, our new formal expansion reads as

$$f(q) = \sum_{n \in \mathbb{N}} P_n(q)A_n \tag{12}$$

where $P_{2n}(q) = [(q - x_0)^2 + y_0^2]^n$ and $P_{2n+1}(q) = [(q - x_0)^2 + y_0^2]^n (q - q_0)$ for all $n \in \mathbb{N}$. Let us study the sets of convergence of function series of this type.

Lemma 2.3. *Let $x_0 \in \mathbb{R}, y_0 > 0, q \in \mathbb{H}$ and let $r \geq 0$ be such that $|(q - x_0)^2 + y_0^2| = r^2$. If $q_0 = x_0 + Iy_0$ for some $I \in \mathbb{S}$ then*

$$\sqrt{r^2 + y_0^2} - y_0 \leq |q - q_0| \leq \sqrt{r^2 + y_0^2} + y_0.$$

Proof. If $r = 0$, i.e. $q \in x_0 + y\mathbb{S}$, then clearly $0 \leq |q - q_0| \leq 2y_0$. Else $q \notin x_0 + y\mathbb{S}$ and

$$|(q - x_0)^2 + y_0^2| = |q - q_0| |(q - q_0)^{-1} q (q - q_0) - \bar{q}_0| = |q - q_0| |q - \tilde{q}_0|$$

where $\tilde{q}_0 = (q - q_0)\bar{q}_0(q - q_0)^{-1} \in x_0 + y_0\mathbb{S}$. If $|q - q_0| > \sqrt{r^2 + y_0^2} + y_0$ then

$$|q - \tilde{q}_0| \geq |q - q_0| - |q_0 - \tilde{q}_0| \geq |q - q_0| - 2y_0 > \sqrt{r^2 + y_0^2} - y_0$$

so that

$$|(q - x_0)^2 + y_0^2| = |q - q_0| |q - \tilde{q}_0| > r^2 + y_0^2 - y_0^2 = r^2$$

a contradiction with the hypothesis. A similar reasoning excludes that $|q - q_0| < \sqrt{r^2 + y_0^2} - y_0$. \square

Proposition 2.4. Let $\{a_n\}_{n \in \mathbb{N}} \subset \mathbb{H}$ and suppose

$$\limsup_{n \rightarrow +\infty} |a_n|^{1/n} = 1/R \tag{13}$$

for some $R > 0$. Let $q_0 = x_0 + Iy_0 \in \mathbb{H}$ with $x_0 \in \mathbb{R}$, $y_0 > 0$, $I \in \mathbb{S}$ and set $P_{2n}(q) = [(q - x_0)^2 + y_0^2]^n$ and $P_{2n+1}(q) = [(q - x_0)^2 + y_0^2]^n (q - q_0)$ for all $n \in \mathbb{N}$. Then the function series

$$\sum_{n \in \mathbb{N}} P_n(q) a_n \tag{14}$$

converges absolutely and uniformly on compact sets in

$$U(x_0 + y_0\mathbb{S}, R) = \{q \in \mathbb{H} : |(q - x_0)^2 + y_0^2| < R^2\}, \tag{15}$$

where it defines a regular function. Furthermore, the series (14) diverges at every $q \in \mathbb{H} \setminus U(x_0 + y_0\mathbb{S}, R)$.

Proof. If K is a compact subset of $U(x_0 + y_0\mathbb{S}, R)$ then there exists $r < R$ such that $|(q - x_0)^2 + y_0^2| \leq r^2$ for all $q \in K$. Thus, for all $q \in K$

$$|P_{2n}(q) a_{2n}| = |(q - x_0)^2 + y_0^2|^n |a_{2n}| \leq r^{2n} |a_{2n}|$$

while (thanks to the previous lemma)

$$\begin{aligned} |P_{2n+1}(q) a_{2n+1}| &= |(q - x_0)^2 + y_0^2|^n |q - q_0| |a_{2n+1}| \\ &\leq r^{2n} \left(\sqrt{r^2 + y_0^2} + y_0 \right) |a_{2n+1}|. \end{aligned}$$

Hence (14) is dominated on K by a number series $\sum_{n \in \mathbb{N}} c_n$ with

$$\limsup_{n \rightarrow +\infty} |c_n|^{1/n} = r/R < 1.$$

This guarantees absolute and uniform convergence of (14) in K . This, in turn, proves the regularity of the sum (since all the addends in (14) are regular polynomials).

Finally, if $q \in \mathbb{H} \setminus U(x_0 + y_0\mathbb{S}, R)$ then $|(q - x_0)^2 + y_0^2| = r^2$ for some $r > R$. Reasoning as before, we get $|P_{2n}(q) a_{2n}| \geq r^{2n} |a_{2n}|$ and

$$|P_{2n+1}(q) a_{2n+1}| \geq r^{2n} \left(\sqrt{r^2 + y_0^2} - y_0 \right) |a_{2n+1}|.$$




Fig. 1. Examples of $U(\mathbb{S}, R)$, intersected with a plane L_I , for different values of R .

Thus, $\sum_{n \in \mathbb{N}} P_n(q) a_n$ dominates a number series $\sum_{n \in \mathbb{N}} C_n$ with

$$\limsup_{n \rightarrow +\infty} |C_n|^{1/n} = r/R > 1$$

and it must diverge. \square

Let us now describe the sets $U(x_0 + y_0\mathbb{S}, R)$. The intersection between the hypersurface $|(q - x_0)^2 + y_0^2| = R^2$ and any complex plane L_I is a polynomial lemniscate, connected for $R \geq y_0$ and having two connected components for $R < y_0$. If $y_0 \neq 0$ then at the critical value $R = y_0$ the lemniscate is of figure-eight type. Some examples are portrayed in Fig. 1.

Remark 2.5. For $R > y_0$ the set $U(x_0 + y_0\mathbb{S}, R)$ is a symmetric slice domain; for $0 < R \leq y_0$ the set $U(x_0 + y_0\mathbb{S}, R)$ is a symmetric domain whose intersection with any complex plane L_I has two connected components.

So far, we have introduced the formal expansion (12), and we have studied what its set of convergence would be if estimates for the coefficients A_n were provided. In the next section, we will search exactly for such estimates by means of Cauchy-type integral representations.

3. Integral formulas and Cauchy estimates

Let us recall the “slice-wise” Cauchy integral formula proven in [11]. We first introduce some notations. Suppose $\gamma_I : [0, 1] \rightarrow L_I$ to be a rectifiable curve whose support lies in a complex plane L_I for some $I \in \mathbb{S}$; let Γ_I be a neighborhood of γ_I in L_I and let $f, g : \Gamma_I \rightarrow \mathbb{H}$ be continuous functions. If $J \in \mathbb{S}$ is such that $J \perp I$ then $L_I + L_I J = \mathbb{H} = L_I + J L_I$ and there

exist continuous functions $F, G, H, K : \Gamma_I \rightarrow L_I$ such that $f = F + GJ$ and $g = H + JK$ in Γ_I . Then

$$\int_{\gamma_I} g(s)dsf(s) := \int_{\gamma_I} H(s)dsF(s) + \int_{\gamma_I} H(s)dsG(s)J + J \int_{\gamma_I} K(s)dsF(s) + J \int_{\gamma_I} K(s)dsG(s)J.$$

The aforementioned Cauchy-type formula reads as follows.

Lemma 3.1. *Let f be a regular function on a symmetric slice domain Ω , let $I \in \mathbb{S}$ and let U_I be a bounded Jordan domain in L_I , with $\overline{U_I} \subset \Omega_I$. If ∂U_I is rectifiable then*

$$f(z) = \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{s - z} f(s) \tag{16}$$

for all $z \in U_I$.

The previous lemma was the basis for the proof of a more complete Cauchy formula in [2]. It will now provide a new integral representation and lead us to the desired estimates for the coefficients A_n in the formal expansion (12).

Theorem 3.2. *Let f be a regular function on a symmetric slice domain Ω , let $I \in \mathbb{S}$ and let U_I be a symmetric bounded Jordan domain in L_I , with $\overline{U_I} \subset \Omega_I$. If ∂U_I is rectifiable then for each $z_0 = x_0 + Iy_0 \in U_I$ and for all $z \in U_I$*

$$f(z) = f(z_0) + (z - z_0) \cdot \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z)(s - z_0)} f(s). \tag{17}$$

Furthermore,

$$(R_{\bar{z}_0} R_{z_0})^n f(z) = \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z)[(s - x_0)^2 + y_0^2]^n} f(s) \tag{18}$$

$$R_{z_0} (R_{\bar{z}_0} R_{z_0})^n f(z) = \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z)(s - z_0)[(s - x_0)^2 + y_0^2]^n} f(s) \tag{19}$$

for all $n \in \mathbb{N}$ and all $z \in U_I$.

Proof. The previous lemma is equivalent to (18) in the special case $n = 0$. That immediately implies

$$\begin{aligned} R_{z_0} f(z) &= (z - z_0)^{-1} [f(z) - f(z_0)] \\ &= \frac{1}{2\pi I} \int_{\partial U_I} \frac{1}{z - z_0} \left[\frac{1}{s - z} - \frac{1}{s - z_0} \right] ds f(s) \\ &= \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z)(s - z_0)} f(s), \end{aligned}$$

that is (19) in the special case $n = 0$. This last equality, in turn, implies (17). Moreover,

$$R_{z_0} f(\bar{z}_0) = \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - x_0)^2 + y_0^2} f(s)$$

and

$$\begin{aligned}
 R_{\bar{z}_0} R_{z_0} f(z) &= (z - \bar{z}_0)^{-1} [R_{z_0} f(z) - R_{z_0} f(\bar{z}_0)] \\
 &= \frac{1}{2\pi I} \int_{\partial U_I} \frac{1}{z - \bar{z}_0} \left[\frac{1}{(s - z)(s - z_0)} - \frac{1}{(s - x_0)^2 + y_0^2} \right] ds f(s) \\
 &= \frac{1}{2\pi I} \int_{\partial U_I} \frac{1}{z - \bar{z}_0} \frac{s - \bar{z}_0 - (s - z)}{(s - z)[(s - x_0)^2 + y_0^2]} ds f(s) \\
 &= \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z)[(s - x_0)^2 + y_0^2]} f(s),
 \end{aligned}$$

that is (18) with $n = 1$. An analogous reasoning proves that if (18) holds for $n = k$, then (19) holds for $n = k$, which in turn implies (18) for $n = k + 1$ completing the proof. \square

We are now in a position to get the desired estimates for the coefficients A_n in the formal expansion (12).

Corollary 3.3. *Let f be a regular function on a symmetric slice domain Ω , let $x_0 + y_0\mathbb{S} \subset \Omega$ and let $q_0 \in x_0 + y_0\mathbb{S}$. Set $A_{2n} = (R_{\bar{q}_0} R_{q_0})^n f(q_0)$ and $A_{2n+1} = R_{q_0} (R_{\bar{q}_0} R_{q_0})^n f(\bar{q}_0)$. For every $U = U(x_0 + y_0\mathbb{S}, R)$ such that $\bar{U} \subset \Omega$, there exists a constant $C > 0$ such that*

$$|A_n| \leq C \cdot \frac{\max_{\partial U} |f|}{R^n} \tag{20}$$

for all $n \in \mathbb{N}$.

Proof. If we choose $U = U(x_0 + y_0\mathbb{S}, R)$ then the previous theorem implies

$$\begin{aligned}
 |(R_{\bar{q}_0} R_{q_0})^n f(q_0)| &= \left| \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{(s - z_0)[(s - x_0)^2 + y_0^2]^n} f(s) \right| \\
 &\leq \frac{1}{2\pi} \int_{\partial U_I} \frac{|f(s)|}{\left(\sqrt{R^2 + y_0^2} - y_0\right) R^{2n}} d|s| \leq C \cdot \frac{\max_{\partial U} |f|}{R^{2n}}
 \end{aligned}$$

if we set $C = \frac{\text{length}(\partial U_I)}{2\pi(\sqrt{R^2 + y_0^2} - y_0)}$. On the other hand

$$\begin{aligned}
 |R_{q_0} (R_{\bar{q}_0} R_{q_0})^n f(\bar{q}_0)| &= \left| \frac{1}{2\pi I} \int_{\partial U_I} \frac{ds}{[(s - x_0)^2 + y_0^2]^{n+1}} f(s) \right| \\
 &\leq \frac{1}{2\pi} \int_{\partial U_I} \frac{|f(s)|}{R^{2n+2}} d|s| \leq K \cdot \frac{\max_{\partial U} |f|}{R^{2n+1}}
 \end{aligned}$$

where $K = \frac{\text{length}(\partial U_I)}{2\pi R} \leq C$. \square

4. A new notion of analyticity

We are now ready to prove the desired result concerning the expansion of slice regular functions.

Theorem 4.1. Let f be a regular function on a symmetric slice domain Ω , and let $x_0, y_0 \in \mathbb{R}$ and $R > 0$ be such that $U(x_0 + y_0\mathbb{S}, R) \subseteq \Omega$. For all $q_0 \in x_0 + y_0\mathbb{S}$, setting $A_{2n} = (R_{\bar{q}_0} R_{q_0})^n f(q_0)$ and $A_{2n+1} = R_{q_0} (R_{\bar{q}_0} R_{q_0})^n f(\bar{q}_0)$, we have that

$$f(q) = \sum_{n \in \mathbb{N}} [(q - x_0)^2 + y_0^2]^n [A_{2n} + (q - q_0)A_{2n+1}] \quad (21)$$

for all $q \in U(x_0 + y_0\mathbb{S}, R)$.

Proof. Thanks to Proposition 2.4 and Corollary 3.3, the function series in Eq. (21) converges in $U = U(x_0 + y_0\mathbb{S}, R)$, where it defines a regular function. Let us consider the difference

$$g(q) = f(q) - \sum_{n \in \mathbb{N}} [(q - x_0)^2 + y_0^2]^n [A_{2n} + (q - q_0)A_{2n+1}].$$

By construction, $g(q) = [(q - x_0)^2 + y_0^2]^n (R_{\bar{q}_0} R_{q_0})^n f(q)$ for all $n \in \mathbb{N}$. For any choice of $I \in \mathbb{S}$ and for all $z \in U_I$, we derive that

$$g_I(z) = [(z - x_0)^2 + y_0^2]^n (R_{\bar{q}_0} R_{q_0})^n f_I(z) = [z - (x_0 + Iy_0)]^n h_I^{[n]}(z)$$

where $h_I^{[n]}(z) = [z - (x_0 - Iy_0)]^n (R_{\bar{q}_0} R_{q_0})^n f_I(z)$ is holomorphic in U_I . The identity principle for holomorphic functions of one complex variable implies that $g_I \equiv 0$. Since I can be arbitrarily chosen in \mathbb{S} , the function g must be identically zero in U . This is equivalent to Eq. (21). \square

Definition 4.2. Let f be a regular function on a symmetric slice domain Ω . We say that f is *symmetrically analytic* if it admits at any $q_0 \in \Omega$ an expansion of type (12) valid in a neighborhood of q_0 .

The previous theorem, along with Proposition 2.4, proves what follows.

Corollary 4.3. Let Ω be a symmetric slice domain. A function $f : \Omega \rightarrow \mathbb{H}$ is regular if, and only if, it is symmetrically analytic.

We conclude this section reformulating expansion (21) as follows, thanks to Theorem 1.10.

Corollary 4.4. Let f be a regular function on a symmetric slice domain Ω , and let $x_0, y_0 \in \mathbb{R}$ and $R > 0$ be such that $U(x_0 + y_0\mathbb{S}, R) \subseteq \Omega$. Then, for all $q \in U(x_0 + y_0\mathbb{S}, R)$

$$f(q) = \sum_{n \in \mathbb{N}} [(q - x_0)^2 + y_0^2]^n [C_{2n} + qC_{2n+1}] \quad (22)$$

where the coefficients C_n depend only on f, x_0, y_0 and can be computed as

$$\begin{aligned} C_{2n} &= (q_2 - q_1)^{-1} [\bar{q}_1 (R_{\bar{q}_0} R_{q_0})^n f(q_1) - \bar{q}_2 (R_{\bar{q}_0} R_{q_0})^n f(q_2)], \\ C_{2n+1} &= (q_2 - q_1)^{-1} [(R_{\bar{q}_0} R_{q_0})^n f(q_2) - (R_{\bar{q}_0} R_{q_0})^n f(q_1)] \end{aligned} \quad (23)$$

for all $q_0, q_1, q_2 \in x_0 + y_0\mathbb{S}$ with $q_1 \neq q_2$.

We notice that the odd-indexed coefficients in expansions (21) and (22) coincide.

5. Computing multiplicities of zeros

As a first application of our new type of series expansions, let us look at its bearings in the computation of the multiplicities of the zeros. After the study of the roots of polynomials of [17], the following result was proven for power series in [7], and in its present form in [8] (for generalizations to other real alternative algebras, see [13,14]).

Theorem 5.1 (*Structure of the Zero Set*). *Let f be a regular function on a symmetric slice domain Ω . If f does not vanish identically, then the zero set of f consists of isolated points or isolated 2-spheres of the form $x + y\mathbb{S}$.*

The following notion of multiplicity was introduced in [7].

Definition 5.2. Let f be a regular function on a symmetric slice domain Ω and let $q_0 \in \Omega$. We define the (classical) multiplicity of q_0 as a zero of f and denote by $m_f(q_0)$ the largest $n \in \mathbb{N}$ such that $f(q) = (q - q_0)^{*n} * g(q)$ for some regular $g : \Omega \rightarrow \mathbb{H}$.

In other words, $m_f(q_0)$ is the index n of the first non vanishing coefficient a_n in the expansion (3). Despite its analogy with complex multiplicity, this notion of multiplicity is not completely satisfactory, since the sum of the classical multiplicities of the zeros of a polynomial is unrelated to its degree.

Example 5.3. Let us choose $I \in \mathbb{S}$ and let

$$P(q) = (q - I) * (q + I) = q^2 + 1.$$

P has multiplicity $m_P(I) = 1$ at all $I \in \mathbb{S}$.

Example 5.4. Let us choose $I, J \in \mathbb{S}$ with $I \neq -J$ and let

$$P(q) = (q - I) * (q - J) = q^2 - q(I + J) + IJ.$$

The zero set of P is $\{I\}$. If $I \neq J$ then $m_P(I) = 1$, while f has degree 2.

For this reason [12] introduced alternative notions of multiplicity for the roots of regular polynomials. We present them as generalized in [18] to all regular functions on symmetric slice domains.

Theorem 5.5. *Let f be a regular function on a symmetric slice domain Ω ; suppose $f \not\equiv 0$ and let $x + y\mathbb{S} \subset \Omega$. There exist $m \in \mathbb{N}$ and a regular function $\tilde{f} : \Omega \rightarrow \mathbb{H}$ not identically zero in $x + y\mathbb{S}$ such that*

$$f(q) = [(q - x)^2 + y^2]^m \tilde{f}(q). \tag{24}$$

If \tilde{f} has a zero $p_1 \in x + y\mathbb{S}$ then such a zero is unique and there exist $n \in \mathbb{N}$, $p_2, \dots, p_n \in x + y\mathbb{S}$ (with $p_i \neq \bar{p}_{i+1}$ for all $i \in \{1, \dots, n - 1\}$) such that

$$\tilde{f}(q) = (q - p_1) * (q - p_2) * \dots * (q - p_n) * g(q) \tag{25}$$

for some regular function $g : \Omega \rightarrow \mathbb{H}$ which does not have zeros in $x + y\mathbb{S}$.

Definition 5.6. In the situation of Theorem 5.5, we say that f has spherical multiplicity $2m$ at $x + y\mathbb{S}$ and that f has isolated multiplicity n at p_1 .

As observed in [12], the degree of a polynomial equals the sum of the spherical multiplicities and of the isolated multiplicities of its zeros. For instance: in the previous example, $q^2 + 1$ has

spherical multiplicity 2 at \mathbb{S} , $(q - I) * (q - J)$ (with $I \neq -J$) has spherical multiplicity 0 at \mathbb{S} and isolated multiplicity 2 at I .

Our new series expansion allows an immediate computation of the spherical multiplicity and it gives some information on the isolated multiplicity.

Remark 5.7. Let f be a regular function on a symmetric slice domain Ω and let $q_0 = x_0 + Iy_0 \in \Omega$. If A_{2n} or A_{2n+1} is the first non vanishing coefficient in the expansion (21) then $2n$ is the spherical multiplicity of f at $x_0 + y_0\mathbb{S}$. Moreover, q_0 has positive isolated multiplicity if and only if $A_{2n} = 0$.

Remark 5.8. Let f be a regular function on a symmetric slice domain Ω and let $x_0 + y_0\mathbb{S} \subset \Omega$. If the first non vanishing coefficient in the expansion (22) is C_{2n} or C_{2n+1} then $2n$ is the spherical multiplicity of f at $x_0 + y_0\mathbb{S}$. Moreover, there exists a point with positive isolated multiplicity in $x_0 + y_0\mathbb{S}$ if, and only if, $C_{2n+1}^{-1}C_{2n} \in x_0 + y_0\mathbb{S}$.

6. Differentiating regular functions

In Section 4 we proved that a regular function f on a symmetric slice domain Ω admits an expansion of the form

$$f(q) = \sum_{n \in \mathbb{N}} [(q - x_0)^2 + y_0^2]^n [A_{2n} + (q - q_0)A_{2n+1}]$$

at each $q_0 \in \Omega$. This new expansion allows the computation of the real partial derivatives of f .

Theorem 6.1. Let f be a regular function on a symmetric slice domain Ω , and let $q_0 = x_0 + Iy_0 \in \Omega$. For all $v \in \mathbb{H}$, $|v| = 1$ the derivative of f along v can be computed at q_0 as

$$\lim_{t \rightarrow 0} \frac{f(q_0 + tv) - f(q_0)}{t} = vA_1 + (q_0v - v\bar{q}_0)A_2 \tag{26}$$

where $A_1 = R_{q_0}f(\bar{q}_0)$, $A_2 = R_{\bar{q}_0}R_{q_0}f(q_0)$. In particular if $e_0, e_1, e_2, e_3 \in \mathbb{H}$ form a basis for \mathbb{H} and if x_0, x_1, x_2, x_3 denote the corresponding coordinates, then

$$\frac{\partial f}{\partial x_i}(q_0) = e_i R_{q_0}f(\bar{q}_0) + (q_0e_i - e_i\bar{q}_0)R_{\bar{q}_0}R_{q_0}f(q_0). \tag{27}$$

Proof. We observe that

$$(q - x_0)^2 + y_0^2 = (q - q_0) * (q - \bar{q}_0) = q(q - q_0) - (q - q_0)\bar{q}_0$$

and for $q = q_0 + tv$

$$(q - x_0)^2 + y_0^2 = (q_0 + tv)tv - tv\bar{q}_0 = t(tv^2 + q_0v - v\bar{q}_0)$$

whence

$$f(q_0 + tv) = \sum_{n \in \mathbb{N}} t^n (tv^2 + q_0v - v\bar{q}_0)^n [A_{2n} + tvA_{2n+1}].$$

Hence,

$$f(q_0 + tv) - f(q_0) = tA_1 + t(tv^2 + q_0v - v\bar{q}_0)[A_2 + tvA_3] + o(t)$$

and the thesis immediately follows. \square

We notice that similar reasonings allow the computation of higher order derivatives. Furthermore, we make the following observation.

Remark 6.2. Let f be a regular function on a symmetric slice domain Ω and let $q_0 \in \Omega$. If v lies in the same L_I as q_0 , then v commutes with q_0 and

$$\begin{aligned} vR_{q_0}f(\bar{q}_0) + (q_0v - v\bar{q}_0)R_{\bar{q}_0}R_{q_0}f(q_0) &= v[R_{q_0}f(\bar{q}_0) + (q_0 - \bar{q}_0)R_{\bar{q}_0}R_{q_0}f(q_0)] \\ &= v[R_{q_0}f(\bar{q}_0) + R_{q_0}f(q_0) - R_{q_0}f(\bar{q}_0)] \\ &= vR_{q_0}f(q_0). \end{aligned}$$

On the other hand, if $q_0 = x_0 + Iy_0$ with $y_0 \neq 0$ and v is tangent to the 2-sphere $x_0 + y_0\mathbb{S}$ at q_0 , then $q_0v = v\bar{q}_0$ and

$$vR_{q_0}f(\bar{q}_0) + (q_0v - v\bar{q}_0)R_{\bar{q}_0}R_{q_0}f(q_0) = vR_{q_0}f(\bar{q}_0).$$

As a consequence, the notions of derivatives of regular functions already introduced in the literature can be recovered as special cases of formula (26). Indeed, $R_{q_0}f(q_0)$ coincides with $\partial_c f(q_0)$, where $\partial_c f$ denotes the Cullen (or slice) derivative of f , defined in [11] to equal

$$\partial_I f(x + Iy) = \frac{1}{2} \left(\frac{\partial}{\partial x} - I \frac{\partial}{\partial y} \right) f_I(x + Iy)$$

at each point $x + Iy \in \Omega_I$. On the other hand, $R_{q_0}f(\bar{q}_0)$ equals $\frac{\partial_s f(q_0)}{y_0}$, where $\partial_s f$ denotes the spherical derivative of f , defined in [13] setting

$$\partial_s f(q_0) = \frac{1}{2} \text{Im}(q_0)^{-1} (f(q_0) - f(\bar{q}_0)).$$

We conclude by looking at the implications of our result in complex coordinates. In the hypotheses of the previous theorem, let us choose $I \in \mathbb{S}$ so that $q_0 \in L_I$, choose $J \in \mathbb{S}$ with $I \perp J$ and set $e_0 = 1, e_1 = I, e_2 = J, e_3 = IJ$. Then $\mathbb{H} = (\mathbb{R} + I\mathbb{R}) + (\mathbb{R} + I\mathbb{R})J$ can be identified with \mathbb{C}^2 setting $z_1 = x_0 + Ix_1, z_2 = x_2 + Ix_3, \bar{z}_1 = x_0 - Ix_1, \bar{z}_2 = x_2 - Ix_3$; we may as well split $f = f_1 + f_2J$ for some $f_1, f_2 : \Omega \rightarrow L_I$.

Theorem 6.3. Let Ω be a symmetric slice domain, let $f : \Omega \rightarrow \mathbb{H}$ be a regular function and let $q_0 \in \Omega$. Choose $I, J \in \mathbb{S}$ so that $q_0 \in L_I$ and $I \perp J$, let $z_1, z_2, \bar{z}_1, \bar{z}_2$ be the induced coordinates and let $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2$ be the corresponding derivations. Then

$$\begin{pmatrix} \bar{\partial}_1 f_1 & \bar{\partial}_2 f_1 \\ \bar{\partial}_1 f_2 & \bar{\partial}_2 f_2 \end{pmatrix} \Big|_{q_0} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}. \tag{28}$$

Furthermore, if $R_{q_0}f$ splits as $R_{q_0}f = R_1 + R_2J$ with R_1, R_2 ranging in L_I then

$$\begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 \\ \partial_1 f_2 & \partial_2 f_2 \end{pmatrix} \Big|_{q_0} = \begin{pmatrix} R_1(q_0) & -\overline{R_2(\bar{q}_0)} \\ R_2(q_0) & \overline{R_1(\bar{q}_0)} \end{pmatrix}. \tag{29}$$

Proof. Setting $e_0 = 1, e_1 = I, e_2 = J, e_3 = IJ$ and denoting as x_0, x_1, x_2, x_3 the corresponding real coordinates, the derivations $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2$ are defined by

$$\partial_1 = \frac{1}{2} \left(\frac{\partial}{\partial x_0} - I \frac{\partial}{\partial x_1} \right), \quad \bar{\partial}_1 = \frac{1}{2} \left(\frac{\partial}{\partial x_0} + I \frac{\partial}{\partial x_1} \right)$$

$$\partial_2 = \frac{1}{2} \left(\frac{\partial}{\partial x_2} - I \frac{\partial}{\partial x_3} \right), \quad \bar{\partial}_2 = \frac{1}{2} \left(\frac{\partial}{\partial x_2} + I \frac{\partial}{\partial x_3} \right).$$

By means of the previous proposition, we compute

$$\begin{aligned} \frac{\partial f}{\partial x_0}(q_0) &= R_{q_0} f(q_0) = R_1(q_0) + R_2(q_0)J, \\ \frac{\partial f}{\partial x_1}(q_0) &= IR_{q_0} f(q_0) = IR_1(q_0) + IR_2(q_0)J, \\ \frac{\partial f}{\partial x_2}(q_0) &= JR_{q_0} f(\bar{q}_0) + (q_0J - J\bar{q}_0)R_{\bar{q}_0}R_{q_0} f(q_0) = JR_{q_0} f(\bar{q}_0) \\ &= J(R_1(\bar{q}_0) + R_2(\bar{q}_0)J) = -\overline{R_2(\bar{q}_0)} + \overline{R_1(\bar{q}_0)}J, \\ \frac{\partial f}{\partial x_3}(q_0) &= KR_{q_0} f(\bar{q}_0) + (q_0K - K\bar{q}_0)R_{\bar{q}_0}R_{q_0} f(q_0) = KR_{q_0} f(\bar{q}_0) \\ &= IJ(R_1(\bar{q}_0) + R_2(\bar{q}_0)J) = -I\overline{R_2(\bar{q}_0)} + I\overline{R_1(\bar{q}_0)}J, \end{aligned}$$

where we have taken into account that $Jz = \bar{z}J$ for all $z \in L_I$. The thesis follows by direct computation. \square

Eq. (28) tells us that fixing q_0 and choosing appropriate complex coordinates on \mathbb{H} , f is complex differentiable at q_0 ; Eq. (29) then allows to compute the complex Jacobian of f at q_0 . These tools open the possibility of using complex variables in the study of slice regular functions. We also point out that in the special case where $q_0 \in \mathbb{R}$ the complex Jacobian has the form

$$\begin{pmatrix} a & -\bar{b} \\ b & \bar{a} \end{pmatrix},$$

which implies $\lim_{h \rightarrow 0} h^{-1}[f(q_0 + h) - f(q_0)] = a + bJ$ and proves what follows.

Corollary 6.4. *Let Ω be a symmetric slice domain. If $f : \Omega \rightarrow \mathbb{H}$ is a regular function then it is left q -differentiable at each $q_0 \in \Omega \cap \mathbb{R}$, where $\lim_{h \rightarrow 0} h^{-1}[f(q_0 + h) - f(q_0)] = R_{q_0} f(q_0) = \partial_c f(q_0)$.*

Acknowledgments

The author is partially supported by FSE, Regione Lombardia, GNSAGA INdAM and FIRB “Geometria Differenziale Complessa e Dinamica Olomorfa”.

References

- [1] F. Brackx, R. Delanghe, F. Sommen, Clifford Analysis, in: Research Notes in Mathematics, vol. 76, Pitman (Advanced Publishing Program), Boston, MA, 1982.
- [2] F. Colombo, G. Gentili, I. Sabadini, A Cauchy kernel for slice regular functions, Ann. Global Anal. Geom. 37 (4) (2010) 361–378.
- [3] F. Colombo, G. Gentili, I. Sabadini, D. Struppa, Extension results for slice regular functions of a quaternionic variable, Adv. Math. 222 (5) (2009) 1793–1808.
- [4] F. Colombo, I. Sabadini, F. Sommen, D.C. Struppa, Analysis of Dirac Systems and Computational Algebra, in: Progress in Mathematical Physics, vol. 39, Birkhäuser Boston Inc., Boston, MA, 2004.
- [5] R. Fueter, Die Funktionentheorie der Differentialgleichungen $\Delta u = 0$ und $\Delta \Delta u = 0$ mit vier reellen Variablen, Comment. Math. Helv. 7 (1) (1934) 307–330.
- [6] R. Fueter, Über die analytische Darstellung der regulären Funktionen einer Quaternionenvariablen, Comment. Math. Helv. 8 (1) (1935) 371–378.

- [7] G. Gentili, C. Stoppato, Zeros of regular functions and polynomials of a quaternionic variable, *Michigan Math. J.* 56 (3) (2008) 655–667.
- [8] G. Gentili, C. Stoppato, The zero sets of slice regular functions and the open mapping theorem, in: I. Sabadini, F. Sommen (Eds.), *Hypercomplex Analysis and Applications*, in: *Trends in Mathematics*, Birkhäuser Verlag, Basel, 2011, pp. 95–107.
- [9] G. Gentili, C. Stoppato, Power series and analyticity over the quaternions, *Math. Ann.* 352 (1) (2012) 113–131.
- [10] G. Gentili, D.C. Struppa, A new approach to Cullen-regular functions of a quaternionic variable, *C. R. Math. Acad. Sci. Paris* 342 (10) (2006) 741–744.
- [11] G. Gentili, D.C. Struppa, A new theory of regular functions of a quaternionic variable, *Adv. Math.* 216 (1) (2007) 279–301.
- [12] G. Gentili, D.C. Struppa, On the multiplicity of zeroes of polynomials with quaternionic coefficients, *Milan J. Math.* 76 (2008) 15–25.
- [13] R. Ghiloni, A. Perotti, Slice regular functions on real alternative algebras, *Adv. Math.* 226 (2) (2011) 1662–1691.
- [14] R. Ghiloni, A. Perotti, Zeros of regular functions of quaternionic and octonionic variable: a division lemma and the camshaft effect, *Ann. Mat. Pura Appl.* (4) 190 (2011) 539–551.
- [15] V.V. Kravchenko, M.V. Shapiro, *Integral Representations for Spatial Models of Mathematical Physics*, in: *Pitman Research Notes in Mathematics Series*, vol. 351, Longman, Harlow, 1996.
- [16] G. Laville, E. Lehman, Analytic Cliffordian functions, *Ann. Acad. Sci. Fenn. Math.* 29 (2) (2004) 251–268.
- [17] A. Pogorui, M. Shapiro, On the structure of the set of zeros of quaternionic polynomials, *Complex Var. Theory Appl.* 49 (6) (2004) 379–389.
- [18] C. Stoppato, Singularities of slice regular functions, *Math. Nachr.* 285 (10) (2012) 1274–1293.
- [19] A. Sudbery, Quaternionic analysis, *Math. Proc. Cambridge Philos. Soc.* 85 (2) (1979) 199–224.