

JOURNAL OF APPROXIMATION THEORY 52, 58–81 (1988)

Nonsmooth Analysis and Approximation

DIMITRIOS KANDILAKIS AND NIKOLAOS S. PAPAGEORGIOU*

*Department of Mathematics, University of Illinois,
Urbana, Illinois 61801, U.S.A.*

Communicated by P. L. Butzer

Received July 23, 1984; revised November 27, 1985

1. INTRODUCTION

Initially, optimization and the theory of best approximation developed independently. However, in the 1960s, with the appearance of convex analysis, it was realized that best approximation problems can be regarded as special problems of optimization. This led to a systematic effort to obtain approximation results as special cases of more general theorems of optimization theory. This parallel treatment is presented in the monographs of Holmes [14] and Laurent [17], which illustrate that there is a strong interaction between approximation theory and what is known by now as “nonsmooth analysis.”

This paper develops along these lines and concentrates on problems of stability (sensitivity) and stochastic approximation.

In the study of stability our main tool is the so called Kuratowski–Mosco convergence of sets and the corresponding τ -convergence of proper functions. So we perturb the data determining the f -best approximations and the f -farthest points and we examine how the sets of these points vary. Such sensitivity analysis is, among other things, very important in designing efficient numerical algorithms. Additional results in this direction were recently obtained by the authors in [20].

In stochastic approximation, which is studied in Section 4, we allow both the set and the function to depend measurably on a parameter ω and we examine the dependence on ω of the various notions of approximation theory. We also study the approximation problem in which the function is the integral functional determined by $f(\cdot, \cdot)$. In all these our main tools are the theory of normal integrands of Rockafellar [23, 24] and the theory of measurable multifunctions.

Finally in Section 5 we have gathered some general results which illustrate the strong interaction between approximation theory and non-linear analysis.

* Research supported by N.S.F. Grant D.M.S.-8403135.

2. PRELIMINARIES

Let (Ω, Σ) be a measurable space and X a Polish space. Let $F: \Omega \rightarrow 2^X \setminus \{\emptyset\}$ be a multifunction (set-valued function) with closed values. Then the following statements are equivalent:

- (i) $F^-(U) = \{\omega \in \Omega: F(\omega) \cap U \neq \emptyset\} \in \Sigma$ for all $U \subseteq X$ open,
- (ii) $\omega \rightarrow d_{F(\omega)}(x) = \inf_{z \in F(\omega)} \|x - z\|$ is measurable for all $x \in X$,
- (iii) there exist measurable functions $f_n: \Omega \rightarrow X$ s.t.

$$F(\omega) = \text{cl} \{f_n(\omega)\}_{n \geq 1} \quad \text{for all } \omega \in \Omega \quad (\text{Castaing's representation}).$$

A multifunction satisfying any of the above statements is said to be measurable. If there exists a complete, σ -finite measure $\mu(\cdot)$ on Σ , (i) \rightarrow (iii) are all equivalent to

(iv) $\text{Gr } F = \{(\omega, x) \in \Omega \times X: x \in F(\omega)\} \in \Sigma \times B(X)$, where $B(X)$ is the Borel σ -field of X .

From now on assume that (Ω, Σ, μ) is a complete probability space and X a separable Banach space. By X^* we will denote its topological dual. We will use the notations

$$P_{f(\cdot)}(X) = \{A \subseteq X: \text{nonempty, closed, (convex)}\},$$

$$P_{(w)k(\cdot)}(X) = \{A \subseteq X: \text{nonempty, (}w\text{-)compact, (convex)}\},$$

where w denotes the weak topology on X . If $A \subseteq X$ we will denote by $\sigma_A(\cdot)$ the support function of A , i.e., for all $x^* \in X^*$ $\sigma_A(x^*) = \sup_{x \in A} \langle x^*, x \rangle$.

Consider the set $S_F^1 = \{f(\cdot) \in L_X^1(\Omega): f(\omega) \in F(\omega) \mu\text{-a.e.}\}$, i.e., S_F^1 contains all selectors of $F(\cdot)$ belonging to the Lebesgue–Bochner space $L_X^1(\Omega)$. Clearly S_F^1 is a closed (maybe empty) subset of $L_X^1(\Omega)$. It is nonempty if and only if $\inf_{x \in F(\omega)} \|x\| \in L_+^1(\Omega)$. We will say that $F(\cdot)$ is integrably bounded if and only if $\sup_{x \in F(\omega)} \|x\| = |F(\omega)| \in L_+^1(\Omega)$. Using S_F^1 we can define an integral for $F(\cdot)$,

$$\int_{\Omega} F(\omega) d\mu(\omega) = \left\{ \int_{\Omega} f(\omega) d\mu(\omega): f(\cdot) \in S_F^1 \right\},$$

where $\int_{\Omega} f(\omega) d\mu(\omega)$ is the usual Bochner integral. This set-valued integral is known as Aumann's integral. For more details on measurable multifunctions and their integral we refer to Castaing–Valadier [4], Himmelberg [10], and Rockafellar [23].

Now let us pass to normal integrands. These were introduced and studied by Rockafellar [23, 24], as the appropriate generalization to accommodate the needs of optimization and optimal control, of the

Caratheodory integrands from the calculus of variations. So assume that $f: \Omega \times X \rightarrow \overline{\mathbb{R}}$ is a proper integrand (i.e., $f(\cdot, \cdot)$ takes values in $(-\infty, +\infty]$ and $f \neq +\infty$). We say that $f(\cdot, \cdot)$ is a normal integrand if and only if $\omega \rightarrow \text{epif}(\omega, \cdot) = \{(x, \lambda) \in X \times \mathbb{R} : f(\omega, x) \leq \lambda\}$ is closed valued and measurable. A straightforward application of Von Neumann's projection theorem tells us that the above definition is equivalent to saying that $f(\cdot, \cdot)$ is $\Sigma \times B(X)$ -measurable and $f(\omega, \cdot)$ is l.s.c. for all $\omega \in \Omega$. Recall that normality is preserved by the Fenchel transform. For more details the reader can look at the excellent survey paper of Rockafellar [23].

As we already mentioned in the Introduction, in the next section we will be using the Kuratowski–Mosco convergence of sets and the corresponding τ -convergence of proper functions. Very briefly we will recall those notions. Let $\{A_n, A\}_{n \geq 1} \subseteq 2^X$ and set $w\text{-}\overline{\lim}_{n \rightarrow \infty} A_n = \{x \in X : x = w\text{-}\lim_{k \rightarrow \infty} x_k, x_k \in A_{n_k}, k \geq 1\}$ and $s\text{-}\underline{\lim}_{n \rightarrow \infty} A_n = \{x \in X : x = s\text{-}\lim_{n \rightarrow \infty} x_n, x_n \in A_n, n \geq 1\}$. We say that the A_n 's converge to A in the Kuratowski–Mosco sense (denoted by $A_n \rightarrow^{\text{K-M}} A$) if and only if $w\text{-}\overline{\lim} A_n = A = s\text{-}\underline{\lim} A_n$. If $\{f_n, f\}_{n \geq 1} \subseteq \overline{\mathbb{R}}^X$ are proper functions then $\{f_n\}_{n \geq 1}$ τ -converges to f (denoted by $f_n \rightarrow^\tau f$) if and only if $\text{epif}_n \rightarrow^{\text{K-M}} \text{epif}$. For more details we refer to Mosco [19] and Salinetti–Wets [25].

A last piece of terminology. If $f \in \overline{\mathbb{R}}^X$ is proper, by $\text{dom } f$ we denote the effective domain of $f(\cdot)$, i.e., $\text{dom } f = \{x \in X : f(x) < +\infty\}$. Moreover, all $\overline{\mathbb{R}}$ -valued functions will be assumed to be proper.

3. STABILITY RESULTS

If $f \in \overline{\mathbb{R}}^X$ is a proper function and $A \subseteq X$ nonempty, then we set $f_A(x) = \inf_{y \in A} f(x - y)$ and $P_{f,A}(x) = \{h \in A : f_A(x) = f(x - h)\}$. The elements of $P_{f,A}(x)$ are said to be elements of f -best approximation (f -b.a.) to x from the set A . Throughout this paper we will assume that $f_A(\cdot)$ is proper.

A sensitivity analysis was first conducted by Brosowski–Deutch–Nürnberger [3], who considered a family $\{A_t\}_{t \in T}$ of subsets of a normed space X parametrized by a topological space T and studied the continuity of $t \rightarrow P_{A_t}(x)$ (here $f(\cdot) = \|\cdot\|$). Recently Tsukada [28] addressed the same problem but with a nonparametrized method. Namely he allowed the sets $\{A_n\}_{n \geq 1}$ to converge to A in the K–M sense and then examined what happens to the sequence $\{P_{A_n}(x)\}_{n \geq 1}$. His study was limited to strictly convex, reflexive Banach spaces.

Our first result examines the behavior of $P_{f,A}(x)$ under variations of the set A . Assume that X is a Banach space.

THEOREM 3.1. *If $f: X \rightarrow \mathbb{R}$ is continuous and w -sequentially l.s.c. and*

$$\{A_n, A\}_{n \geq 1} \subseteq P_{fc}(X) \quad \text{s.t. } A_n \xrightarrow{\text{K-M}} A,$$

then for all $x \in X$, $w\text{-}\overline{\lim} P_{f, A_n}(x) \subseteq P_{f, A}(x)$.

Proof. Let $h \in w\text{-}\overline{\lim} P_{f, A_n}(x)$. Then by definition we can find $h_k \in P_{f, A_{n_k}}(x)$ s.t. $h_k \rightarrow^w h$. Let $y \in A$ and let $y_k \in A_{n_k}$ s.t. $y_k \rightarrow^s y$. This is possible since $A_n \xrightarrow{\text{K-M}} A$. Then using the properties of $f(\cdot)$ we have

$$\begin{aligned} f(x - h_k) &\leq f(x - y_k) \\ \Rightarrow \underline{\lim} f(x - h_k) &\leq \lim f(x - y_k) \\ \Rightarrow f(x - h) &\leq f(x - y). \end{aligned}$$

Note that $h \in A$ and since $y \in A$ was arbitrary we conclude that $h \in P_{f, A}(x)$. Q.E.D.

Remark. Note that with our assumptions on $f(\cdot)$ we cover the case where $f(\cdot) = \|\cdot\|$. We could have also assumed that $f(\cdot)$ is w -sequentially continuous.

Let $f(\cdot) = \|\cdot\|$. Recall that if X is reflexive and strictly convex then every $A \in P_{fc}(X)$ is a Chebyshev set, i.e., $P_A(x)$ is a singleton for all $x \in X$. Also X is said to have property (H) if and only if for every $x_n \rightarrow^w x$ with $\|x_n\| \rightarrow \|x\|$ we have $x_n \rightarrow^s x$. Locally uniformly convex spaces (in particular Hilbert spaces) have property (H). Using Theorem 3.1 we can have the following corollary which is Theorem 3.2(i) of Tsukada [28].

COROLLARY (28). *If X is reflexive and strictly convex and $\{A_n, A\}_{n \geq 1} \subseteq P_{fc}(X)$ with*

$$A_n \xrightarrow{\text{K-M}} A \quad \text{as } n \rightarrow \infty,$$

then for all $x \in X$, $P_{A_n}(x) \rightarrow^w P_A(x)$ as $n \rightarrow \infty$.

If in addition X has property (H) then $P_{A_n}(x) \rightarrow^s P_A(x)$.

Proof. Since $P_{A_n}(x)$, $P_A(x)$ $n \geq 1$ are singletons, the first claim of the corollary follows directly from Theorem 3.1. Then $\|x - P_{A_n}(x)\| \leq \underline{\lim} \|x - P_{A_n}(x)\|$. Let $y_n \in A_n$ s.t. $y_n \rightarrow^s P_A(x)$. Then $\|x - y_n\| \rightarrow \|x - P_A(x)\|$. Since $\|x - P_{A_n}(x)\| \leq \|x - y_n\|$ we get that $\overline{\lim} \|x - P_{A_n}(x)\| \leq \|x - P_A(x)\|$. Thus $\|x - P_{A_n}(x)\| \rightarrow \|x - P_A(x)\|$ and because X has property (H) we get that $P_{A_n}(x) \rightarrow^s P_A(x)$. Q.E.D.

Remark. In the above proof we also got that $d_{A_n}(\cdot) \rightarrow d_A(\cdot)$.

Next we will allow $f(\cdot)$ to vary too. Then we have the following variant of Theorem 3.1. Assume X if finite dimensional. By $\delta_A(\cdot)$ we will denote the indicator function of $A \subseteq X$.

THEOREM 3.2. *If $\{f_n, f\}_{n \geq 1} \subseteq \mathbb{R}^X$ are continuous, convex, $f_n \rightarrow f$ as $n \rightarrow \infty$ and $\{A_n, A\}_{n \geq 1} \subseteq P_{fc}(X)$ s.t. $A_n \rightarrow^{K-M} A$ then for all $x \in X$, $\overline{\lim} P_{f_n, A_n}(x) \subseteq P_{f, A}(x)$.*

Proof. For any $x \in X$, let $f_x: X \rightarrow \mathbb{R}$ be defined by $f_x(y) = f(x - y)$. Then from Corollary 2E of Salinetti–Wets [25] we have $f_{n,x}(\cdot) \rightarrow^t f_x(\cdot)$ as $n \rightarrow \infty$. Also since $A_n \rightarrow^{K-M} A$, we have that $\delta_{A_n}(\cdot) \rightarrow^t \delta_A(\cdot)$. Note that $\text{dom } f_x - \text{dom } \delta_A = \mathbb{R}^n - A = \mathbb{R}^n$. So by Theorem 5 of McLinden–Bergstrom [18] we get that $(f_{n,x} + \delta_{A_n})(\cdot) \rightarrow^t (f_x + \delta_A)(\cdot) \Rightarrow (f_{n,x} + \delta_{A_n})^*(\cdot) \rightarrow^{t^*} (f_x + \delta_A)^*(\cdot) \Rightarrow \text{Gr}(f_{n,x} + \delta_{A_n})^* \rightarrow^{K-M} \text{Gr } \partial(f_x + \delta_A)^*(\cdot)$. But recall that $P_{f_n, A_n}(x) = \partial[f_{n,x} + \delta_{A_n}]^*(0)$ and $P_{f, A}(x) = \partial[f_x + \delta_A]^*(0)$. Hence it follows easily that $\overline{\lim} P_{f_n, A_n}(x) \subseteq P_{f, A}(x)$. Q.E.D.

We will close this section, with a result analogous to Theorem 3.1 but for f -farthest points. If $f \in \mathbb{R}^X$ is a proper function, $A \subseteq X$ is nonempty then we define $\hat{f}_A(x) = \sup_{y \in A} f(x - y)$ and $Q_{f, A}(x) = \{h \in A: \hat{f}_A(x) = f(x - h)\}$. To avoid trivialities we will always assume that $\hat{f}_A(\cdot)$ is proper. Let X be a Banach space. By $h(\cdot, \cdot)$ we will denote the Hausdorff distance on 2^X .

THEOREM 3.3. *If $f: X \rightarrow \mathbb{R}$ is continuous, $\{A_n\}_{n \geq 1} \subseteq P_k(X)$ and $A_n \rightarrow^h A$ as $n \rightarrow \infty$ then for all $x \in X$, $\hat{f}_{A_n}(x) \rightarrow \hat{f}_A(x)$ and $s\text{-}\overline{\lim} Q_{f, A_n}(x) \subseteq Q_{f, A}(x)$.*

Proof. First note that $A \in P_k(X)$. For any $y \in A$, Let $y_n \in A_n$ s.r. $y_n \rightarrow^s y$. Then we have $f(x - y_n) \leq \hat{f}_{A_n}(x) \Rightarrow f(x - y) \leq \liminf \hat{f}_{A_n}(x) \Rightarrow \hat{f}_A(x) \leq \liminf \hat{f}_{A_n}(x)$. Let $h_n \in A_n$ $n \geq 1$ s.t. $\hat{f}_{A_n}(x) = f(x - h_n)$. Observe that $d_A(h_n) \leq d_{A_n}(h_n) + h(A_n, A) = h(A_n, A) \rightarrow 0$ as $n \rightarrow \infty$. So $d_A(h_n) \rightarrow 0$. Let $a_n \in A$ s.t. $d_A(h_n) = \|h_n - a_n\|$. By passing to a subsequence if necessary, we may assume that $a_n \rightarrow^s h \in A$. Then $\|h_n - h\| \leq \|h_n - a_n\| + \|a_n - h\| \rightarrow 0$. So $h_n \rightarrow^s h \in A$. This implies that

$$\overline{\lim} \hat{f}_{A_n}(x) = \lim f(x - h_n) = f(x - h) \leq \hat{f}_A(x).$$

Hence we finally have that $\hat{f}_{A_n}(\cdot) \rightarrow \hat{f}_A(\cdot)$. Next let $h_k \in Q_{f, A_{n_k}}(x)$ $k \geq 1$ and $h_k \rightarrow^s h$. Then $\hat{f}_{A_{n_k}}(x) = f(x - h_k) \rightarrow f(x - h) = \hat{f}_A(x)$ as $k \rightarrow \infty \Rightarrow h \in Q_{f, A}(x) \Rightarrow s\text{-}\overline{\lim} Q_{f, A_n}(x) \subseteq Q_{f, A}(x)$. Q.E.D.

4. STOCHASTIC APPROXIMATION

Throughout this section assume that (Ω, Σ, μ) is a complete probability space and X a separable Banach space.

In the first result on the f -approximation of random sets we examine how the pointwise approximations are related to the aggregate (integral) approximation. So we will obtain a relation between the functions

$$f_{F(\omega)}(x) = \inf\{f(x - y) : y \in F(\omega)\}$$

and

$$f_{\int F} (x) = \inf \left\{ f(x - y) : y \in \int_{\Omega} F(\omega) \right\}.$$

THEOREM 4.1. $F: \Omega \rightarrow P_f(X)$ is measurable with $S_F^1 \neq \emptyset$, $f: X \rightarrow \mathbb{R}$ is l.s.c., convex and for all $x(\cdot) \in \mathcal{L}^1(X) = \{x(\cdot) \in L^1_X(\Omega) : x = \int_{\Omega} x(\omega) d\mu(\omega)\}$ and all $y(\cdot) \in S_F^1$, $f(x(\cdot) - y(\cdot))$ is quasintegrable and integrable for one such pair $(x'(\cdot), y'(\cdot))$ then $f_{\int F}(x) = [\int_{\Omega} f_{F(\omega)}](x)$, where \int_{Ω} denotes the operation of continuous infimal convolution.

Proof. From Lemma 2.1 of Hiai–Umegaki [9] we know that $\omega \rightarrow f_{F(\omega)}(x)$ is measurable. So the continuous infimal convolution in the conclusion of the theorem is well defined. Also from the definition of the Aumann integral we have that

$$f_{\int F}(x) = \inf_{y \in \int F} f(x - y) = \inf_{y(\cdot) \in S_F^1} f\left(x - \int_{\Omega} y(\omega) d\mu(\omega)\right).$$

Let $x(\cdot) \in \mathcal{L}^1(X)$. Then we have

$$f_{\int F}(x) = \inf_{y(\cdot) \in S_F^1} f\left(\int_{\Omega} (x(\omega) - y(\omega)) d\mu(\omega)\right).$$

Applying Jensen’s inequality (see Kozek–Suchanecki [15, Corollary 7.1]) we get that

$$f_{\int F}(x) \leq \inf_{y(\cdot) \in S_F^1} \int_{\Omega} f(x(\omega) - y(\omega)) d\mu(\omega).$$

Since $x(\cdot) \in \mathcal{L}^1(x)$ was arbitrary we have that

$$f_{\int F}(x) \leq \inf_{\substack{y(\cdot) \in S_F^1 \\ x(\cdot) \in \mathcal{L}^1(x)}} \int_{\Omega} f(x(\omega) - y(\omega)) d\mu(\omega).$$

Applying Theorem 2.2 of [9] we have that

$$\begin{aligned} f_{\int F}(x) &\leq \inf_{x(\cdot) \in \mathcal{L}^1(x)} \int_{\Omega} \inf_{y \in F(\omega)} f(x(\omega) - y) d\mu(\omega) \\ &= \inf_{x(\cdot) \in \mathcal{L}^1(x)} \int_{\Omega} f_{F(\omega)}(x(\omega)) d\mu = \left[\oint_{\Omega} f_{F(\omega)} \right] (x). \end{aligned}$$

Hence we have shown that

$$f_{\int F}(x) \leq \left[\oint_{\Omega} f_{F(\omega)} \right] (x). \quad (1)$$

Next let $h \in \int_{\Omega} F(\omega) d\mu(\omega)$ s.t. $f(x-h) \leq f_{\int F}(x) + \varepsilon$, $\varepsilon > 0$. Note that $h = \int_{\Omega} h(\omega) d\mu(\omega)$ with $h(\cdot) \in S_F^1$. Let $x(\cdot) = x - h + h(\cdot)$. Clearly $x(\cdot) \in \mathcal{L}^1(x)$ and $f(x(\omega) - h(\omega)) = f(x-h) \leq f_{\int F}(x) + \varepsilon$. Also $f_{F(\omega)}(x(\omega)) \leq f(x(\omega) - h(\omega))$. So we have $f_{F(\omega)}(x(\omega)) \leq f_{\int F}(x) + \varepsilon$. Integrating both sides over Ω we get that

$$\begin{aligned} \int_{\Omega} f_{F(\omega)}(x(\omega)) d\mu(\omega) &\leq f_{\int F}(x) + \varepsilon \\ \Rightarrow \inf_{x(\cdot) \in \mathcal{L}^1(x)} \int_{\Omega} f_{F(\omega)}(x(\omega)) d\mu(\omega) &\leq f_{\int F}(x) + \varepsilon. \\ \Rightarrow \left[\oint_{\Omega} f_{F(\omega)} \right] (x) &\leq f_{\int F}(x) + \varepsilon. \end{aligned}$$

Let $\varepsilon \downarrow 0$. We have that

$$\left[\oint_{\Omega} f_{F(\omega)} \right] (x) \leq f_{\int F}(x). \quad (2)$$

From (1) and (2) we conclude that $f_{\int F}(x) = \left[\oint_{\Omega} f_{F(\omega)} \right] (x)$. Q.E.D.

Remark. If $f(\cdot)$ is such that $f(x(\cdot))$ is quasintegrable for all $x(\cdot) \in L_X^1(\Omega)$, then the theorem is true for all $x \in X$. This is the case if $f(\cdot)$ is bounded from below.

When $f(\cdot) = \|\cdot\|$, then the quasintegrability hypothesis is automatically satisfied and if $F: \Omega \rightarrow P_f(X)$ is as above then:

COROLLARY. For all $x \in X$, $d_{\int F}(x) = \left[\oint_{\Omega} d_{F(\omega)} \right] (x)$.

For the pointwise approximation problem we can say more. By S_F we will denote the set of measurable selectors of $F(\cdot)$. Also a set $A \subseteq X$ non-empty is said to be f -proximal if and only if for all $x \in X$, $P_{f,A}(x) \neq \emptyset$.

THEOREM 4.2. *If $f: X \rightarrow \overline{\mathbb{R}}$ is l.s.c. and $F: \Omega \rightarrow P_f(X)$ is measurable with $F(\omega)$ being f -proximal for all $\omega \in \Omega$ then for all $x \in X$, there exists $h(\cdot) \in S_F$ s.t. for all $\omega \in \Omega$, $f_{F(\omega)}(x) = f(x - h(\omega))$.*

Proof. For $x \in X$ consider the multifunction $\omega \rightarrow P_{f, F(\omega)}(x)$. By definition we have that

$$P_{f, F(\omega)}(x) = \{h \in X: f(x - h) - f_{F(\omega)}(x) = g_x(\omega, h) = 0\} \cap F(\omega).$$

Recall that $\omega \rightarrow f_{F(\omega)}(x)$ is measurable, while $f(x - \cdot)$ is l.s.c. Hence $g_x(\omega, h)$ being the sum of two normal integrands is normal. Thus

$$\text{Gr } P_{f, F(\cdot)}(x) = \{(\omega, h): g_x(\omega, h) = 0\} \cap \text{Gr } F \in \Sigma \times B(X).$$

Apply Aumann's selection theorem to find $h: \Omega \rightarrow X$ measurable s.t. $h(\omega) \in P_{f, F(\omega)}(x)$ for all $\omega \in \Omega$. Therefore $f(x - h(\omega)) = f_{F(\omega)}(x)$. Q.E.D.

Remark. The above result is still true if instead of $x \in X$ we have a measurable function $x: \Omega \rightarrow X$.

Next consider the following integral functional

$$I_f(x(\cdot)) = \int_{\Omega} f(\omega, x(\omega)) \, d\mu(\omega),$$

where $f: \Omega \times X \rightarrow \overline{\mathbb{R}}$ is a measurable integrand and $x: \Omega \rightarrow X$ is measurable. Additional hypotheses will be introduced later. Let $M \subseteq L^1_X(\Omega)$ and define

$$I_f^M(x(\cdot)) = \inf_{y(\cdot) \in M} I_f(x(\cdot) - y(\cdot)).$$

Having established this notation consider a multifunction $F: \Omega \rightarrow 2^X \setminus \{\emptyset\}$ s.t. $S_F^1 \neq \emptyset$. We will examine the following two problems.

- (1) For $\omega \in \Omega$, find $h \in F(\omega)$ s.t. $f_{F(\omega)}(\omega, x(\omega)) = f(\omega, x(\omega) - h)$.
- (2) Find $h(\cdot) \in S_F^1$ s.t. $I_f^{S_F^1}(x(\cdot)) = I_f(x(\cdot) - h(\cdot))$.

By $P_{f(\omega, \cdot), F(\omega)}(x(\omega))$, $\omega \in \Omega$ we will denote the solution set of problem (1), while by $P_{I_f, S_F^1}(x(\cdot))$ we will denote the solution set of (2).

Under normality and measurability hypotheses on $f(\cdot, \cdot)$ and $F(\cdot)$, respectively, we can show as in Theorem 4.2 that $\omega \rightarrow P_{f(\omega, \cdot), F(\omega)}(x(\omega))$ is nonempty, closed valued, and measurable.

Our next theorem compares those two solution sets.

THEOREM 4.3. *If $f: \Omega \times X \rightarrow \overline{\mathbb{R}}$ is a normal, convex integrand s.t. for all $x(\cdot) \in L^1_X(\Omega)$, $f(\cdot, x(\cdot))$ is integrable and $F: \Omega \rightarrow P_{wkc}(X)$ is integrably bounded, then for any $x(\cdot) \in L^1_X(\Omega)$ we have $P_{I_f, S_F^1}(x(\cdot)) = S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1$ and is a w -compact subset of $L^1_X(\Omega)$.*

Proof. First note that since for all $\omega \in \Omega, f(\omega, \cdot) \in \Gamma_0(x) = \{\text{proper, l.s.c., convex, } \mathbb{R}\text{-valued functions defined on } X\}$ and $F(\omega) \in P_{wkc}(X)$, then by Weierstrass theorem we have that $P_{f(\omega, \cdot), F(\omega)}(x(\omega)) \neq \emptyset$. Furthermore as we already pointed out $\omega \rightarrow P_{f(\omega, \cdot), F(\omega)}(x(\omega))$ is a measurable multifunction, with w -compact, convex values. Also it is integrably bounded since $F(\cdot)$ is. So from Proposition 3.1 of [21] we deduce that $S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1 \neq \emptyset$, is w -compact and convex in $L_X^1(\Omega)$.

It is clear that $S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1 \subseteq P_{I_f, S_f^1}(x(\cdot))$. Let $h(\cdot) \in P_{I_f, S_f^1}(x(\cdot))$. Then $I_f^{S_f^1}(x(\cdot)) = I_f(x(\cdot) - h(\cdot))$. So we have

$$\int_{\Omega} [f(\omega, x(\omega) - h(\omega)) - f_{F(\omega)}(\omega, x(\omega))] d\mu(\omega) = 0.$$

Since $f(\omega, x(\omega)) - h(\omega) \geq f_{F(\omega)}(\omega, x(\omega))$ μ -a.e. we get that

$$\begin{aligned} f(\omega, x(\omega)) - h(\omega) &= f_{F(\omega)}(\omega, x(\omega)) \mu\text{-a.e.} \\ \Rightarrow h(\cdot) &\in S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1. \end{aligned}$$

Thus the claim of the theorem follows.

Q.E.D.

Remark. When $f(\cdot, \cdot) = \|\cdot\|$, then the above theorem tells us that for any $x(\cdot) \in L_X^1(\Omega)$, the best approximation from S_f^1 is also a pointwise best approximation to $x(\omega)$ from $F(\omega)$ and vice versa. Furthermore the set of such best approximations is w -compact in $L_X^1(\Omega)$.

Working as above we can have an analogous result for f -farthest points.

THEOREM 4.4. *If $f: \Omega \times X \rightarrow \bar{\mathbb{R}}$ is $\Sigma \times B(X)$ -measurable and w -u.s.c. in x and for all $x(\cdot) \in L_X^1(\Omega), f(\cdot, x(\cdot))$ is integrable, while $F: \Omega \rightarrow P_{wkc}(X)$ is integrably bounded, then $Q_{I_f, S_f^1}(x(\cdot)) = S_{Q_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1$ for all $x(\cdot) \in L_X^1(\Omega)$.*

Theorem 4.3 is useful in obtaining interesting information about the structure of certain f -proximal sets.

If $f \in \mathbb{R}^X$ is a proper function and A a nonempty subset of X , then we say that A is an f -sun if for each $x \in X \setminus A$ there exists $h \in P_{f, A}(x)$ s.t. $h \in P_{f, A}(h + \lambda(x - h))$ for all $\lambda > 0$. We will say that A is a strict f -sun if this is true for all $h \in P_{f, A}(x)$. Note that if $f(\cdot) = \|\cdot\|$, then the above definition reduces to the classical definition of solarly of a set (see Vlasov [29]).

As before X is a separable Banach space.

THEOREM 4.5. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a Caratheodory, sublinear integrand s.t. for all $x(\cdot) \in L_X^1(\Omega), f(\cdot, x(\cdot))$ is integrable and $F: \Omega \rightarrow P_{wkc}(X)$ is integrably bounded, then $F(\omega)$ is an $f(\omega, \cdot)$ -sun μ -a.e. if and only if S_f^1 is an I_f -sun.*

Proof. First assume that S_F^1 is a I_f -sun. This means that for all $x(\cdot) \in L_X^1(\Omega)$ there exists $h(\cdot) \in P_{I_f, S_F^1}(x(\cdot))$ s.t. $h(\cdot) \in P_{I_f, S_F^1}(h(\cdot) + \lambda(x(\cdot) - h(\cdot)))$ for all $\lambda > 0$. Let $x(\omega) \equiv x$ and use Theorem 4.3, to get that

$$h(\omega) \in P_{f(\omega, \cdot), F(\omega)}(x) \mu\text{-a.e.} \Rightarrow h(\omega) \in P_{f(\omega, \cdot), F(\omega)}(h(\omega) + \lambda(x - h(\omega))) \mu\text{-a.e.}$$

for all $\lambda > 0$, which means that $P_{f(\omega, \cdot), F(\omega)}(x)$ is an $f(\omega, \cdot)$ sun μ -a.e.

Now assume that $F(\omega)$ is an $f(\omega, \cdot)$ -sun μ -a.e. By definition this means that for μ -almost all $\omega \in \Omega$ we have that, for all $x \in X$ there exists $h \in P_{f(\omega, \cdot), F(\omega)}(x)$ s.t. $h \in P_{f(\omega, \cdot), F(\omega)}(h + \lambda(x - h))$ for all $\lambda > 0$. Let $x(\cdot) \in L_X^1(\Omega)$ and consider the multifunction $\Gamma(\cdot)$ defined by

$$\Gamma(\omega) = \{h \in P_{f(\omega, \cdot), F(\omega)}(x(\omega)): h \in P_{f(\omega, \cdot), F(\omega)}(h + \lambda(x(\omega) - h)), \lambda \geq 0\}.$$

From Govindarajulu-Pai [7] we know that $x \rightarrow P_{f(\omega, \cdot), F(\omega)}(x)$ is u.s.c. So we can write that

$$\Gamma(\omega) = \bigcap_{\substack{\lambda \geq 0 \\ \lambda = \text{rational}}} P_{f(\omega, \cdot), F(\omega)}(h + \lambda(x(\omega) - h)).$$

$\Rightarrow \Gamma(\cdot)$ has a measurable graph (see [4, Theorem III-40]).

Applying Aumann's selection theorem to find $h: \Omega \rightarrow X$ measurable s.t. $h(\omega) \in \Gamma(\omega)$ for all $\omega \in \Omega$. So we have that

$$\begin{aligned} h(\omega) &\in P_{f(\omega, \cdot), F(\omega)}(h(\omega) + \lambda(x(\omega) - h(\omega))), \quad \omega \in \Omega \\ &\Rightarrow h(\cdot) \in S_{f(\cdot, \cdot), F(\cdot)}^1(h(\cdot) + \lambda(x(\cdot) - h(\cdot))) \\ &\Rightarrow h(\cdot) \in P_{I_f, S_F^1}(h(\cdot) + \lambda(x(\cdot) - h(\cdot))), \quad \lambda \geq 0. \end{aligned}$$

which proves that S_F^1 is an I_f -sun.

Q.E.D.

Now we will pass to the integral functional $F: X \rightarrow \bar{\mathbb{R}}$ defined by $F(x) = \int_{\Omega} f(\omega, x) d\mu(\omega)$, where $f(\cdot, \cdot)$ is a convex integrand. The next result provides a necessary and sufficient condition for the existence of f -b.a.

Assume that X is a reflexive, separable Banach space.

PROPOSITION 4.1. *If $f: \Omega \times X \rightarrow \bar{\mathbb{R}}$ is a normal, convex integrand s.t. for all $x: \Omega \rightarrow X$ measurable and bounded, $f(\cdot, x(\cdot))$ is integrable and A is a non-empty, closed, convex subset of X , then for all $x \in X$ we have, $h \in P_{F, A}(x)$ if and only if there exists*

$$x^*(\cdot) \in S_{\partial f(\cdot, x-h)}^1 \quad \text{s.t.} \quad x^* = \int_{\Omega} x^*(\omega) d\mu(\omega) \quad \text{and} \quad \sigma_A(x^*) = (x^*, h).$$

Proof. Note that $\inf_{y \in A} F(x - y) = \inf_{y \in X} [F_x + \delta_A](y)$, where $F_x(y) = F(x - y)$. From convex analysis (see [24]) we know that $h \in P_{f,A}(x)$ if and only if $0 \in \partial(F_x + \delta_A)(h)$. But from Theorem 23(a) of Rockafellar [24] we know that $F(\cdot)$ is continuous, convex. So applying the Moreau–Rockafellar theorem (see Laurent [17]) we get that

$$0 \in \partial(F_x + \delta_A)(h) = \partial F_x(h) + \partial \delta_A(h).$$

Observe that $\partial F_x(h) = -\partial F(x - h)$. Also from Theorem 23 (b) of Rockafellar [24] we know that

$$\partial F(x - h) = \int_{\Omega} \partial f(\omega, x - h) d\mu(\omega).$$

So there exists $x^*(\cdot) \in S_{\partial f(\cdot, x-h)}^1$ s.t. $x^* = \int_{\Omega} x^*(\omega) d\mu(\omega)$ and $x^* \in \partial \delta_A(h)$. Hence $(x^*, y - h) \leq 0$ for all $y \in A$. Thus $\sigma_A(x^*) = (x^*, h)$. Q.E.D.

Useful for the purposes of numerical analysis is the concept of ε - f -best approximation. We will say that $h \in A$ is an ε - f -b.a. to x from A if we have $f(x - h) \leq f_A(x) + \varepsilon$. We will denote the set of ε - f -best approximations to A by $P_{f,A}^\varepsilon$. For those points we have a result analogous to Proposition 4.1.

Assume that X is a finite dimensional Banach space.

PROPOSITION 4.2. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a Caratheodory, convex integrand s.t. for all $x: \Omega \rightarrow X$ measurable and bounded $f(\cdot, x(\cdot))$ is integrable and $A \subseteq X$ is nonempty, closed, and convex, then $h \in P_{f,A}^\varepsilon(x)$ if and only if there exists $\varepsilon: \Omega \rightarrow \mathbb{R}_+$ measurable, $\varepsilon' \geq 0$, $x^*: \Omega \rightarrow X^*$ measurable s.t. $x^*(\omega) \in \partial f(\omega, x - h)$ μ -a.e.,*

$$x^* = \int_{\Omega} x^*(\omega) d\mu(\omega), \quad \sigma_A(x^*) - \varepsilon' \leq (x^*, h)$$

and

$$\int_{\Omega} \varepsilon(\omega) d\mu(\omega) + \varepsilon' = \varepsilon.$$

Proof. Note that

$$P_{f,A}^\varepsilon(x) = \{h \in A: F(x - h) \leq F_A(x) + \varepsilon\}.$$

So $h \in P_{f,A}^\varepsilon(x)$ is equivalent to saying that for all $y \in A$ we have $F(x - h) \leq F(x - y) + \varepsilon \Leftrightarrow -\varepsilon \leq F_x(y) - F_x(h) \Leftrightarrow -\varepsilon \leq (F_x + \delta_A)(y) - (F_x + \delta_A)(h) \Leftrightarrow 0 \in \partial_\varepsilon(F_x + \delta_A)(h)$. From Theorem 23(a) of Rockafellar

[24] we know that $F_x(\cdot)$ is continuous, convex on X . Also from Hiriart-Urruty [12] [13] we have

$$\partial_\varepsilon(F_x + \delta_A)(h) = \bigcup_{\substack{\varepsilon_1 \geq 0, \varepsilon_2 \geq 0 \\ \varepsilon_1 + \varepsilon_2 = \varepsilon}} \{ \partial_{\varepsilon_1} F_x(h) + \partial_{\varepsilon_2} \delta_A(h) \}.$$

Moreover, again from [13] we have that

$$\partial_{\varepsilon_1} F_x(h) = \bigcup_{\substack{\varepsilon(\cdot) \in \mathcal{L}^1(\varepsilon) \\ \varepsilon(\omega) \geq 0}} \int_{\Omega} \partial_{\varepsilon(\omega)} f_x(\omega, h) \, d\mu(\omega).$$

So $0 \in \partial_\varepsilon(F_x + \delta_A)(h)$ is equivalent to saying that there exists $z^* \in \int_{\Omega} \partial_{\varepsilon(\omega)} f_x(\omega, h) \, d\mu(\omega)$ for which $-z^* \in \partial_{\varepsilon'} \delta_A(h)$ and $\int_{\Omega} \varepsilon(\omega) \, d\mu(\omega) + \varepsilon' = \varepsilon$, where $\varepsilon(\omega) \geq 0$. Hence $z^* = \int_{\Omega} z^*(\omega) \, d\mu(\omega)$ with $z^*(\omega) \in \partial_{\varepsilon(\omega)} f_x(\omega, h)$ μ -a.e. But note that $\partial_{\varepsilon(\omega)} f_x(\omega, h) = -\partial_{\varepsilon(\omega)} f(\omega, x - h)$. So finally we can find $x^*: \Omega \rightarrow X^*$ measurable s.t. $x^*(\omega) \in \partial_{\varepsilon(\omega)} f(\omega, x - h)$ μ -a.e., $x^* = \int_{\Omega} x^*(\omega) \, d\mu(\omega)$ and $x^* \in \partial_{\varepsilon'} \delta_A(h)$ which is equivalent to saying that $\sigma_A(x^*) - \varepsilon' \leq \langle x^*, h \rangle$, with $\int_{\Omega} \varepsilon(\omega) \, d\mu(\omega) + \varepsilon' = \varepsilon$ and $\varepsilon(\omega) \geq 0$ for all $\omega \in \Omega$. Q.E.D.

Next we will have a result analogous to Proposition 4.1 for the integral functional $I_f(\cdot)$.

Assume X is a reflexive, separable Banach space.

PROPOSITION 4.3. *If $f: \Omega \times X \rightarrow \bar{\mathbb{R}}$ is a normal, convex, integrand s.t. for all $x(\cdot) \in L^1_X(\Omega)$, $I_f(x(\cdot))$ is finite and $F: \Omega \rightarrow P_{fc}(X)$ is integrably bounded, then for any $x(\cdot) \in L^1_X(\Omega)$ we have $h(\cdot) \in P_{I_f, S^1_F}(x(\cdot))$ if and only if*

$$\min_{x^*(\cdot) \in S^{\infty}_{\text{ext}} \partial_{I_f}(x(\cdot) - h(\cdot))} \int_{\Omega} (x^*(\omega), y(\omega) - h(\omega)) \, d\mu(\omega) \leq 0 \quad \text{for all } y(\cdot) \in S^1_F.$$

Proof. As before $h(\cdot) \in P_{I_f, S^1_F}(x(\cdot))$ if and only if $0 \in \partial(I_{f_{x(\cdot)}} + \partial_{S^1_F})(h(\cdot))$.

Recall that $I_{f_{x(\cdot)}}(\cdot)$ is continuous, convex on $L^1_X(\Omega)$. So we can apply the Moreau-Rockafellar theorem and get that $0 \in \partial I_{f_{x(\cdot)}}(h(\cdot)) + \partial \delta_{S^1_F}(h(\cdot))$. This means that there exists $-\hat{x}^*(\cdot) \in \partial I_{f_{x(\cdot)}}(h(\cdot))$ s.t. $\hat{x}^*(\cdot) \in \partial \delta_{S^1_F}(h(\cdot))$. Since $\partial I_{f_{x(\cdot)}}(h(\cdot)) = -\partial(I_f(x(\cdot) - h(\cdot)))$ we have $\hat{x}^*(\cdot) \in \partial I_f(x(\cdot) - h(\cdot))$ and $\langle \hat{x}^*(\cdot), y(\cdot) - h(\cdot) \rangle \leq 0$ for all $y(\cdot) \in S^1_F$. Recalling that $\partial I_f(x(\cdot) - h(\cdot))$ is w^* -compact we can write that

$$\begin{aligned} & \min_{x^*(\cdot) \in \partial I_f(x(\cdot) - h(\cdot))} (x^*(\cdot), y(\cdot) - h(\cdot)) \\ &= \min_{x^*(\cdot) \in \text{ext } \partial I_f(x(\cdot) - h(\cdot))} (x^*(\cdot), y(\cdot) - h(\cdot)) \leq 0. \end{aligned}$$

From Theorem 2.1(c) of Rockafellar [24] we know that

$$\begin{aligned}\partial I_f(x(\cdot) - h(\cdot)) &= S_{\partial f(\cdot, x(\cdot) - h(\cdot))}^\infty \\ &\Rightarrow \text{ext } \partial I_f(x(\cdot) - h(\cdot)) = \text{ext } S_{\partial f(\cdot, x(\cdot) - h(\cdot))}^\infty\end{aligned}$$

Then from Benamara [2] we get that

$$\text{ext } \partial I_f(x(\cdot) - h(\cdot)) = S_{\text{ext } \partial f(\cdot, x(\cdot) - h(\cdot))}^\infty.$$

Therefore we have that

$$\min_{x^*(\cdot) \in S_{\text{ext } \partial f(\cdot, x(\cdot) - h(\cdot))}^\infty} \int_{\Omega} (x^*(\omega), y(\omega) - h(\omega)) \, d\mu(\omega) \leq 0$$

for all $y(\cdot) \in S_F^1$.

Q.E.D.

Remark. In the above result instead of $p = 1$, $q = +\infty$ we could have used a pair $p, q \geq 1$ of conjugate exponents.

We will close our investigation of the multifunction $P_{I_f, S_F^1}(\cdot)$ with a theorem concerning its continuity properties.

Assume X is a reflexive, separable Banach space.

THEOREM 4.6. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a Caratheodory, convex integrand s.t. for all $x: \Omega \rightarrow X$ bounded and measurable, $f(\cdot, x(\cdot))$ is integrable and $F: \Omega \rightarrow P_{\text{wkc}}(X)$ is integrably bounded, then $P_{I_f, S_F^1}(\cdot)$ is u.s.c. from L_X^1 into (L_X^1, w) .*

Proof. We know that for all $x(\cdot) \in L_X^1(\Omega)$, $P_{I_f, S_F^1}(x(\cdot)) = S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1$. So for any $x^* \in X^*$ we have $\sigma_{P_{I_f, S_F^1}(x^*)} = \sigma_{S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1}$. Then by definition

$$\sigma_{S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1}(x^*) = \sup_{h(\cdot) \in S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1} (x^*, h(\cdot)),$$

where (\cdot, \cdot) denotes the duality brackets between L_X^1 and L_X^∞ . Thus we have that

$$\begin{aligned}\sigma_{S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1}(x^*) &= \sup_{h(\cdot) \in S_{P_{f(\cdot, \cdot), F(\cdot)}(x(\cdot))}^1} \int_{\Omega} (x^*, h(\omega)) \, d\mu(\omega) \\ &= \int_{\Omega} \sup_{h \in P_{f(\omega, \cdot), F(\omega)}(x(\omega))} d\mu(\omega) = \int_{\Omega} \sigma_{P_{f(\omega, \cdot), F(\omega)}(x(\omega))}(x^*) \, d\mu(\omega). \quad (*)\end{aligned}$$

Next we will show that $x(\cdot) \rightarrow \sigma_{P_{I_f, S_F^1}(x(\cdot))}(x^*)$ is u.s.c. on L_X^1 . So let $x_n(\cdot) \rightarrow^s L_X^1 x(\cdot)$. Then by passing to a subsequence if necessary we may

assume that $x_n(\omega) \rightarrow^s x(\omega)$ μ -a.e. Using Fatou's lemma and (*) we get that

$$\overline{\lim} \sigma_{P_{I_f, S_F^1(x_n(\cdot))}}(x^*) \leq \int_{\Omega} \overline{\lim} \sigma_{P_{f(\omega, \cdot), F(\omega)}(x_n(\omega))}(x^*) d\mu(\omega).$$

From Govinarajulu-Pai [7] (Proposition 2.4), we know that $x \rightarrow P_{f(\omega, \cdot), F(\omega)}(x)$ is w -u.s.c. So Proposition 2 (p. 122) of Aubin-Ekeland [1] tells us that $x \rightarrow \sigma_{P_{f(\omega, \cdot), F(\omega)}(x)}(x^*)$ is w -u.s.c. So we have

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \sigma_{P_{f(\omega, \cdot), F(\omega)}(x_n(\cdot))}(x^*) &\leq \int_{\Omega} \sigma_{P_{f(\omega, \cdot), F(\omega)}(x(\omega))}(x^*) \\ \Rightarrow \overline{\lim}_{n \rightarrow \infty} \sigma_{P_{I_f, S_F^1(x_n(\cdot))}}(x^*) &\leq \int_{\Omega} \sigma_{P_{f(\omega, \cdot), F(\omega)}(x(\omega))}(x^*) d\mu(\omega) \\ &= \sigma_{S_{P_{f(\cdot, \cdot), F(\cdot)}}^1(x(\cdot))}(x^*) = \sigma_{P_{I_f, S_F^1(x(\cdot))}}(x^*) \\ \Rightarrow x(\cdot) \rightarrow \sigma_{P_{I_f, S_F^1(x(\cdot))}}(x^*) &\text{ is u.s.c.} \end{aligned}$$

Since $P_{I_f, S_F^1(x(\cdot))}$ has w -compact, convex values in L_X^1 , Theorem 10 (p. 128) of [1] tells us that $x(\cdot) \rightarrow P_{I_f, S_F^1(x(\cdot))}$ is u.s.c. from L_X^1 into (L_X^1, w) . Q.E.D.

Remark. If for all $\omega \in \Omega$, $f(\omega, x) = \|x\|$ and $F(\omega)$ is Chebyshev and if $x_n(\cdot) \rightarrow^{s-L_X^1} x(\cdot)$ and $h_n(\cdot)$ are the best approximations to $x_n(\cdot)$ from S_F^1 , then $h_n(\cdot) \rightarrow^{w-L_X^1} h(\cdot) =$ best approximation to $x(\cdot)$ from S_F^1 .

We will close our study on stochastic f -best approximations, by examining what happens when we consider the conditional expectation of the integrand $f(\cdot, \cdot)$ with respect to a sub- σ -field Σ_0 of Σ .

Assume X is finite dimensional.

THEOREM 4.7. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a Caratheodory, inf-compact integrand s.t. for all $x \in X$, $\int_{\Omega} f(\omega, x) d\mu(\omega) \leq +\infty$ and $f(\omega, x) \geq a(\omega)$ μ -a.e., where $a(\cdot)$ is integrable and if $A \subseteq X$ is nonempty, closed, bounded, and convex, then for all $x \in X$, $(E^{\Sigma_0} f)_A(\omega, x) = E^{\Sigma_0} f_A(\omega, x)$ μ -a.e.*

Proof. First recall that for all $\omega \in \Omega$, $f_A(\omega, \cdot) = [f(\omega, \cdot) \square \delta_A](\cdot)$ and since by hypothesis $f(\omega, \cdot)$ is inf-compact, using Proposition 6.5.5 of Laurent [17] we deduce that $f_A(\omega, \cdot)$ is proper, l.s.c., and convex. Also Proposition 2R of Rockafellar [23] tells us that $f_A(\cdot, \cdot)$ is $\Sigma \times B(X)$ -

measurable. Furthermore it is μ -a.e. bounded from below by $a(\omega)$ which is integrable. So $f_A(\cdot, \cdot)$ is a quasintegrable integrand and so we can consider its conditional expectation with respect to Σ_0 . Let $B \in \Sigma_0$, $x: \Omega \rightarrow X$ bounded, Σ_0 -measurable and $y \in A$. We have

$$\begin{aligned} \int_B E^{\Sigma_0} f_A(\omega, x(\omega)) d\mu(\omega) &= \int_B f_A(\omega, x(\omega)) d\mu(\omega) \leq \int_B f(\omega, x(\omega) - y) d\mu(\omega). \\ &= \int_B E^{\Sigma_0} f(\omega, x(\omega) - y) d\mu(\omega). \end{aligned}$$

Invoking Lemma 6 of Thibault [27] we get that

$$E^{\Sigma_0} f_A(\omega, x) \leq E^{\Sigma_0} f(\omega, x - y)$$

for all $\omega \in \Omega \setminus N_y$, $\mu(N_y) = 0$. Let $\{y_n\}_{n \geq 1}$ be dense in X . Then clearly $E^{\Sigma_0} f_A(\omega, x) \leq E^{\Sigma_0} f(\omega, x - y_n)$ μ -a.e. and so exploiting the continuity of $E^{\Sigma_0} f(\omega, \cdot)$ (see [27]) we have that $E^{\Sigma_0} f_A(\omega, x) \leq E^{\Sigma_0} f(\omega, x - y)$ μ -a.e. $\Rightarrow E^{\Sigma_0} f_A(\omega, x) \leq (E^{\Sigma_0} f)_A(\omega, x)$ μ -a.e.

On the other hand for all $y: \Omega \rightarrow A$ Σ_0 -measurable and all $x: \Omega \rightarrow X$, bounded, Σ_0 -measurable and for $B \in \Sigma_0$, we have that

$$\begin{aligned} \int_B (E^{\Sigma_0} f)_A(\omega, x(\omega)) d\mu(\omega) &\leq \int_B E^{\Sigma_0} f(\omega, x(\omega) - y(\omega)) d\mu(\omega) \\ &= \int_B f(\omega, x(\omega) - y(\omega)) d\mu(\omega) \\ &\Rightarrow \int_B (E^{\Sigma_0} f)_A(\omega, x(\omega)) d\mu(\omega) \\ &\leq \inf_{y(\cdot) \in S_A^1(\Sigma_0)} \int_B f(\omega, x(\omega) - y(\omega)) d\mu(\omega) \\ &= \int_B \inf_{y \in A} f(\omega, x(\omega) - y) d\mu(\omega) \\ &= \int_B f_A(\omega, x(\omega)) d\mu(\omega) \\ &= \int_B E^{\Sigma_0} f_A(\omega, x(\omega)) d\mu(\omega). \end{aligned}$$

As before we get that $(E^{\Sigma_0} f)_A(\omega, x) \leq E^{\Sigma_0} f_A(\omega, x)$ μ -a.e. (exceptional set is independent of x)

$$\Rightarrow (E^{\Sigma_0} f)_A(\omega, x) = E^{\Sigma_0} f_A(\omega, x) \mu\text{-a.e.} \quad \text{Q.E.D.}$$

Now we change direction and pass to the study of stochastic f -farthest points. The first result shows that $S_{I_f, S_F^1}(x(\cdot))$ has a solarity type property. Assume X is any separable Banach space.

PROPOSITION 4.4. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a Caratheodory, sublinear integrand, $F: \Omega \rightarrow P_f(X)$ is a measurable multifunction with $S_F^1 \neq \emptyset$ and there is a $y_0(\cdot) \in S_F^1$ s.t. $I_f(y_0(\cdot)) < \infty$, then $h(\cdot) \in Q_{I_f, S_F^1}(x(\cdot))$ implies that $h(\cdot) \in Q_{I_f, S_F^1}(h(\cdot) + \lambda(x(\cdot) - h(\cdot)))$ for all $\lambda > 0$.*

Proof. By definition $h(\cdot) \in Q_{I_f, S_F^1}(x(\cdot))$ means that

$$\tilde{I}_f^{S_F^1}(x(\cdot)) = I_f(x(\cdot) - h(\cdot)) = \int_{\Omega} f(\omega, x(\omega) - h(\omega)) \, d\mu(\omega).$$

Also

$$\begin{aligned} \tilde{I}_f^{S_F^1}(x(\cdot)) &= \sup_{y(\cdot) \in S_F^1} I_f(x(\cdot) - y(\cdot)) \\ &= \sup_{y(\cdot) \in S_F^1} \int_{\Omega} f(\omega, x(\omega) - y(\omega)) \, d\mu(\omega) \\ &= \int_{\Omega} \sup_{y \in F(\omega)} f(\omega, x(\omega) - y) \, d\mu(\omega). \end{aligned}$$

So we have that

$$\int_{\Omega} \sup_{y \in F(\omega)} f(\omega, x(\omega) - y) \, d\mu(\omega) = \int_{\Omega} f(\omega, x(\omega) - h(\omega)) \, d\mu(\omega).$$

Since $h(\omega) \in F(\omega)$ μ -a.e. we deduce that

$$\begin{aligned} f(\omega, x(\omega) - h(\omega)) &= \sup_{y \in F(\omega)} f(\omega, x(\omega) - y) \, \mu\text{-a.e.} \\ &\Rightarrow h(\omega) \in Q_{f(\omega, \cdot), F(\omega)}(x(\omega)) \, \mu\text{-a.e.} \end{aligned}$$

From Proposition 2.8 of Govindarajulu–Pai [8] we know that for all $\lambda > 0$, $h(\omega) \in Q_{f(\omega, \cdot), F(\omega)}(h(\omega) + \lambda(x(\omega) - h(\omega)))$ μ -a.e. So for all $y(\cdot) \in S_F^1$ we have

$$\begin{aligned} f(\omega, h(\omega) + \lambda(x(\omega) - h(\omega)) - y(\omega)) &\leq f(\omega, \lambda(x(\omega) - h(\omega))) \, \mu\text{-a.e.} \\ &\Rightarrow I_f(h(\cdot) + \lambda(x(\cdot) - h(\cdot)) - y(\cdot)) \leq I_f(\lambda(x(\cdot) - h(\cdot))) \\ &\Rightarrow h(\cdot) \in Q_{I_f, S_F^1}(h(\cdot) + \lambda(x(\cdot) - h(\cdot))) \text{ for all } \lambda > 0. \quad \text{Q.E.D.} \end{aligned}$$

The next theorem provides a pointwise necessary condition for $h(\cdot)$ to be in $Q_{I_f, S_F^1}(x(\cdot))$.

Assume as before that X is a separable Banach space. If $A \subseteq X$, by $N_A(x)$ we will denote the normal cone to A at the point x (see Clarke [5]).

THEOREM 4.8. *If $f: \Omega \times X \rightarrow \mathbb{R}$ is a $k(\cdot)$ -Lipschitz, L^1 -measurable integrand with $k(\cdot) \in L^\infty(\Omega)$ and $F: \Omega \rightarrow P_{wk_c}(X)$ is integrably bounded, then $h(\cdot) \in Q(x(\cdot))$ implies that $(-\partial f(\omega, x(\omega) - h(\omega))) \cap N_{F(\omega)}(h(\omega)) = \emptyset$ μ -a.e.*

Proof. We will start by showing that if $h(\cdot) \in Q_{I_f, S_F^1}(x(\cdot)_1)$, then for some $l > 0$, $h(\cdot)$ also solves locally the following maximization problem

$$\sup_{z(\cdot) \in L_X^1} [I_{f_{x(\cdot)}} - ld_{S_F^1}](z(\cdot)). \quad (*)$$

Suppose not. Then for every $n \geq 1$ there exists $z_n(\cdot) \rightarrow^{s-L^1} h(\cdot)$ s.t.

$$I_{f_{x(\cdot)}}(z_n(\cdot)) - nd_{S_F^1}(z_n(\cdot)) > I_f(x(\cdot) - h(\cdot)) = \hat{I}_f^{S_F^1}(x(\cdot)).$$

Then $I_{f_{x(\cdot)}}(z_n(\cdot)) - I_{f_{x(\cdot)}}(h(\cdot)) > nd_{S_F^1}(z_n(\cdot))$ and so $\beta_n = d_{S_F^1}(z_n(\cdot)) \rightarrow 0$ as $n \rightarrow \infty$. Recall that S_F^1 is w -compact in L_X^1 . So we can find $h_n(\cdot) \in S_F^1$ s.t. $d_{S_F^1}(z_n(\cdot)) = \beta_n = \|z_n - h_n\|_1$ for all $n \geq 1$. So we have

$$I_f(x(\cdot) - h_n(\cdot)) \leq I_f(x(\cdot) - h(\cdot)) = \hat{I}_f^{S_F^1}(x(\cdot)) < I_f(x(\cdot) - z_n(\cdot)) - n\beta_n. \quad (1)$$

On the other hand, from the Lipschitzness hypothesis we have that

$$\begin{aligned} I_f(x(\cdot) - z_n(\cdot)) &\leq I_f(x(\cdot) - h_n(\cdot)) + \int_{\Omega} k(\omega) \|z_n(\omega) - h_n(\omega)\| d\mu(\omega) \\ &\leq I_f(x(\cdot) - h_n(\cdot)) + \|k\|_{\infty} \beta_n. \end{aligned}$$

Let $n \geq 1$ be such that $\|k\|_{\infty} \leq n$. Then we have

$$I_f(x(\cdot) - z_n(\cdot)) - \eta\beta_n \leq I_f(x(\cdot) - h_n(\cdot)). \quad (2)$$

From (1) and (2) above we produce the desired contradiction. Hence, knowing that $h(\cdot)$ solves (*) locally, we can write that

$$0 \in \partial[I_{f_{x(\cdot)}} - ld_{S_F^1}](h(\cdot)),$$

where the subdifferential here is the generalized subdifferential in the sense

of Clarke [5]. Recall that Clarke's subdifferential is subadditive. So we have that

$$\begin{aligned} 0 &\in \partial I_{f(x(\cdot))}(h(\cdot)) + \partial[-l d_{S_F^1}](h(\cdot)) \\ &\Rightarrow 0 \in \partial I_{f(x(\cdot))}(h(\cdot)) - l \cdot \partial d_{S_F^1}(h(\cdot)) = -\partial I_f(x(\cdot) - h(\cdot)) - l \cdot \partial d_{S_F^1}(h(\cdot)) \\ &\Rightarrow -\partial I_f(x(\cdot) - h(\cdot)) \cap (l \partial d_{S_F^1}(h(\cdot))) \neq \emptyset. \end{aligned}$$

From Clarke [6] we know that $\partial I_f(x(\cdot) - h(\cdot)) \subseteq S_{\partial f(\cdot, x(\cdot) - h(\cdot))}^1$. Also note that $d_{S_F^1}(\cdot)$ and $d_{F(\omega)}(\cdot)$ ($\omega \in \Omega$), are Lipschitz, convex functions. Moreover, it is easy to see that for any $v(\cdot) \in L^1_X(\Omega)$

$$d_{S_F^1}(v) \leq \int_{\Omega} d_{F(\omega)}(v(\omega)) \, d\mu(\omega) = I_{d_F}(v(\cdot)).$$

Thus for $h(\cdot) \in S_F^1$, we have that

$$\partial d_{S_F^1}(h) \subseteq \partial I_{d_F}(h) = S_{\partial_{F(\cdot)} d(h(\cdot))}^1.$$

Combining all the above observations we deduce that there exists $x^*: \Omega \rightarrow X^*$ s.t.

$$-x^*(\omega) \in \partial f(\omega, x(\omega) - h(\omega)) \, \mu\text{-a.e.} \quad \text{and} \quad x^*(\omega) \in l \partial d_{F(\omega)}(h(\omega)) \, \mu\text{-a.e.}$$

But recall that $l \partial d_{F(\omega)}(h(\omega)) \subseteq N_{F(\omega)}(h(\omega))$ for all $\omega \in \Omega$. So $x^*(\omega) \in N_{F(\omega)}(h(\omega)) \, \mu\text{-a.e.}$ Therefore finally we have

$$(-\partial f(\omega, x(\omega) - h(\omega))) \cap N_{F(\omega)}(h(\omega)) \subseteq \emptyset \, \mu\text{-a.e.} \quad \text{Q.E.D.}$$

Now we turn for a while our attention to the pointwise maximization problem and we will examine the multifunction $\omega \rightarrow Q_{f(\omega, \cdot), F(\omega)}(x(\omega))$.

As always X is a separable Banach space.

PROPOSITION 4.5. *If $f: \Omega \times X \rightarrow \overline{\mathbb{R}}$ is a measurable integrand and $F: \Omega \rightarrow P_f(X)$ is a measurable multifunction then for all $x: \Omega \rightarrow X$ measurable, $\omega \rightarrow Q_{f(\omega, \cdot), F(\omega)}(x(\omega))$ is graph measurable.*

Proof. By definition we have that

$$\begin{aligned} Q_{f(\omega, \cdot), F(\omega)}(x(\omega)) &= \{h \in F(\omega): f(\omega, x(\omega) - h) = \hat{f}_{F(\omega)}(x(\omega))\} \\ &= \{h \in X: f(\omega, x(\omega) - h) \geq \hat{f}_{F(\omega)}(x(\omega))\} \cap F(\omega). \end{aligned}$$

We claim that $\omega \rightarrow \hat{f}_{F(\omega)}(x(\omega))$ is measurable. To see that let $\lambda > 0$. Then we have that $\hat{f}_{F(\omega)}(x(\omega)) > \lambda$ if and only if there exists $y \in F(\omega)$ s.t.

$f(\omega, x(\omega) - y) > \lambda$. So we can write that $\{\omega \in \Omega: f(x(\omega)) > \lambda\} = \text{proj}_{\Omega}[\{(\omega, y) \in \Omega \times X: f(\omega, x(\omega) - y) > \lambda\} \cap \text{Gr } F]$. Recall that $\text{Gr } F \in \Sigma \times \mathcal{B}(X)$. Also since $f(\cdot, \cdot)$ is $\Sigma \times \mathcal{B}(X)$ -measurable, we have $\{(\omega, y) \in \Omega \times X: f(\omega, x(\omega)) - y > \lambda\} \in \Sigma \times \mathcal{B}(X)$. So their intersection is in $\Sigma \times \mathcal{B}(X)$. Then the projection theorem tells us that $\{\omega \in \Omega: \hat{f}_{F(\omega)}(x(\omega)) > \lambda\} \in \Sigma$. Hence $\omega \rightarrow \hat{f}_{F(\omega)}(x(\omega))$ is measurable. From this we deduce that $(\omega, h) \rightarrow \phi(\omega, h) = f(\omega, x(\omega) - h) - \hat{f}_{F(\omega)}(x(\omega))$ is $\Sigma \times \mathcal{B}(X)$ -measurable. Now observe that

$$\text{Gr } Q_{f(\cdot, \cdot), F(\cdot)}(x(\cdot)) = \{(\omega, h) \in \Omega \times X: \phi(\omega, h) \geq 0\} \cap \text{Gr } F \in \Sigma \times \mathcal{B}(X).$$

Q.E.D.

An interesting consequence of this proposition is the following result. Assume that the same set of hypotheses is still in effect.

COROLLARY. *If for all $\omega \in \Omega$, $Q_{f(\omega, \cdot), F(\omega)}(x(\omega)) \neq \emptyset$, then there exists $h: \Omega \rightarrow X$ measurable selector of $F(\cdot)$ s.t.*

$$\hat{f}_{F(\omega)}(x(\omega)) = f(x(\omega) - h(\omega)).$$

Proof. From the previous proposition we know that $\text{Gr } Q_{f(\cdot, \cdot), F(\cdot)}(x(\cdot)) \in \Sigma \times \mathcal{B}(X)$. Apply Aumann's selection theorem to get $h: \Omega \rightarrow X$ measurable s.t. $h(\omega) \in F(\omega)$ and $\hat{f}_{F(\omega)}(x(\omega)) = f(\omega, x(\omega) - h(\omega))$ for all $\omega \in \Omega$. Q.E.D.

Under mild regularity assumptions on $f(\cdot, \cdot)$, we can have the following interesting characterization of the pointwise stochastic f -farthest points.

PROPOSITION 4.6. *If $f: \Omega \times X \rightarrow \bar{\mathbb{R}}$ is a measurable integrand s.t. for all $\omega \in \Omega$, $f(\omega, \cdot)$ is proper, convex, u.s.c., and $F: \Omega \rightarrow P_{wkc}(X)$ is measurable, then given any $x \in X$, we can find $h: \Omega \rightarrow X$ measurable s.t. for all $\omega \in \Omega$, $h(\omega) \in \text{ext } F(\omega)$ and $h(\omega) \in Q_{f(\omega, \cdot), F(\omega)}(x(\omega))$.*

Proof. We saw in the proof of Proposition 4.5 that $\omega \rightarrow \hat{f}_{F(\omega)}(x)$ is measurable. Consider the multifunction $G(\cdot)$ defined by

$$G(\omega) = \{h \in \text{ext } F(\omega): f(\omega, x - h) = \hat{f}_{F(\omega)}(x)\}.$$

From Bauer's maximum principle, we get that for all $\omega \in \Omega$, $G(\omega) \neq \emptyset$. Also $\text{Gr } G = \{(\omega, h) \in \Omega \times X: f(\omega, x - h) - \hat{f}_{F(\omega)}(x) = 0\} \cap \text{Gr}(\text{ext } F)$. Recall that the first set in the intersection is in $\Sigma \times \mathcal{B}(X)$, while from Benamara [2] we know that $\text{Gr}(\text{ext } F) \in \Sigma \times \mathcal{B}(X)$. So we can apply Aumann's selection theorem to find $h: \Omega \rightarrow X$ measurable s.t. for all $\omega \in \Omega$, $h(\omega) \in G(\omega)$. This is the desired $h(\cdot)$. Q.E.D.

Remark. It is easy to check that this result holds true if instead of a fixed $x \in X$, we have $x: \Omega \rightarrow X$ measurable.

As with stochastic f -approximations, we will conclude our study of stochastic f -farthest points, by looking at the conditional expectation of $\hat{f}_A(\cdot)$ with respect to a sub- σ -field Σ_0 of Σ . The result is analogous to Theorem 4.7 but our assumptions on the space X and on the integrand $f(\cdot, \cdot)$ are less restrictive. The space X is as always a separable Banach space.

THEOREM 4.9. *If $f: \Omega \times X \rightarrow \bar{\mathbb{R}}$ is a normal integrand s.t. for all $x(\cdot) \in L^1_X(\Omega)$, $f(\cdot, x(\cdot))$ is integrable and there exists $a(\cdot)$ integrable s.t. $f(\omega, x) \geq a(\omega)$ μ -a.e. for all $x \in X$ and if $F: \Omega \rightarrow P_f(X)$ is Σ_0 -measurable and integrably bounded then for all $x(\cdot) \in L^1_X(\Sigma_0)$ we have*

$$[E^{\Sigma_0} f]_{F(\omega)}(\omega, x(\omega)) = E^{\Sigma_0} \hat{f}_{F(\omega)}(\omega, x(\omega)) \mu\text{-a.e.}$$

Proof. From Thibault [27] we know that there exists an increasing sequence of Caratheodory integrands $\{f_n(\cdot, \cdot)\}_{n \geq 1}$ s.t. for all $x \in X$ we have that

$$f(\omega, x) = \sup_{n \geq 1} f_n(\omega, x) \mu\text{-a.e.}$$

So we can write that

$$\begin{aligned} \hat{f}_{F(\omega)}(\omega, x(\omega)) &= \sup_{y \in F(\omega)} f(\omega, x(\omega) - y) \\ &= \sup_{y \in F(\omega)} \sup_{n \geq 1} f_n(\omega, x(\omega) - y) \\ &= \sup_{n \geq 1} \sup_{y \in F(\omega)} f_n(\omega, x(\omega) - y) = \sup_{n \geq 1} (\hat{f}_n)_{F(\omega)}(\omega, x(\omega)) \mu\text{-a.e.} \end{aligned}$$

Since X is separable and $f_n(\cdot, \cdot)$ are Caratheodory integrands, we have that $(\hat{f}_n)_{F(\cdot)}(\cdot, \cdot)$ are $\Sigma \times B(X)$ -measurable. So $\hat{f}_{F(\cdot)}(\cdot, \cdot) = \sup_{n \geq 1} (\hat{f}_n)_{F(\cdot)}(\cdot, \cdot)$ is a normal integrand. Furthermore for all $x \in X$, $\hat{f}_{F(\omega)}(\omega, x) \geq a(\omega)$ μ -a.e. Hence $\hat{f}_{F(\cdot)}(\cdot, \cdot)$ is a quasintegrable, normal integrand and so $E^{\Sigma_0} \hat{f}(\cdot, \cdot)$ exists. Using Proposition 12 of Thibault [27] we have that for any $A \in \Sigma_0$ and any $y(\cdot) \in S^1_F(\Sigma_0)$,

$$\begin{aligned} \int_A E^{\Sigma_0} f(\omega, x(\omega) - y(\omega)) d\mu(\omega) &= \int_A f(\omega, x(\omega) - y(\omega)) d\mu(\omega) \\ &\Rightarrow \sup_{y(\cdot) \in S^1_F(\Sigma_0)} \int_A E^{\Sigma_0} f(\omega, x(\omega) - y(\omega)) \end{aligned}$$

$$\begin{aligned}
&= \sup_{y(\cdot) \in S_F^1(\Sigma_0)} \int_A f(\omega, x(\omega) - y(\omega)) \, d\mu(\omega) \\
&= \int_A \sup_{y \in F(\omega)} f(\omega, x(\omega) - y) \, d\mu(\omega) \\
&= \int_A \hat{f}_{F(\omega)}(\omega, x(\omega)) \, d\mu(\omega) \\
&= \int_A E^{\Sigma_0} \hat{f}_{F(\omega)}(\omega, x(\omega)) \, d\mu(\omega). \tag{1}
\end{aligned}$$

On the other hand note that

$$\begin{aligned}
&\sup_{y(\cdot) \in S_F^1(\Sigma_0)} \int_A E^{\Sigma_0} f(\omega, x(\omega) - y(\omega)) \, d\mu(\omega) \\
&= \int_A \sup_{y \in F(\omega)} E^{\Sigma_0} f(\omega, x(\omega) - y) \, d\mu(\omega) \\
&= \int_A (E^{\Sigma_0} f)(\omega, x(\omega)) \, d\mu(\omega). \tag{2}
\end{aligned}$$

From (1) and (2) above we get that for all $A \in \Sigma_0$ and all $x(\cdot) \in L_X^1(\Sigma_0)$,

$$\int_A E^{\Sigma_0} \hat{f}_{F(\omega)}(\omega, x(\omega)) \, d\mu(\omega) = \int_A (E^{\Sigma_0} f)(\omega, x(\omega)) \, d\mu(\omega).$$

Invoking Proposition 7 of [27] we conclude that

$$E^{\Sigma_0} \hat{f}_{F(\omega)}(\omega, x(\omega)) = (E^{\Sigma_0} f)(\omega, x(\omega)) \, \mu\text{-a.e.} \quad \text{Q.E.D.}$$

5. GENERAL RESULTS

In this section we have gathered some useful general results about f -best approximations.

The first result illustrates how fixed point theory can be instrumental in obtaining interesting results about f -best approximations. Our theorem generalizes earlier results obtained by Ky Fan [16] and Reich [22].

Assume that X is a locally convex space. We recall that a set $A \subseteq X$ is said to be f -approximatively compact if and only if for all $x \in X$, every minimizing net $\{h_\alpha\}$ (i.e., $f(x - h_\alpha) \rightarrow f_A(x)$) has a convergent subnet in A .

THEOREM 5.1. *If $f: X \rightarrow \mathbb{R}$ is continuous, sublinear, $A \subseteq X$ is a nonempty f -approximatively compact, convex set, and $\phi: A \rightarrow X$ is continuous with $\phi(A)$ compact, then there exists $h \in A$ s.t. $f_A(\phi(h)) = f(\phi(h) - h)$.*

Proof. Consider the multifunction $\Gamma: A \rightarrow P_{fc}(A)$ defined by $\Gamma(y) = (P_{f,A} \circ \phi)(y) = P_{f,A}(\phi(y))$. From [7, Proposition 2.4] we know that $P_{f,A}(\cdot)$ is u.s.c. while $\phi(\cdot)$ is by hypothesis continuous. So we deduce that $(P_{f,A} \circ \phi)(\cdot)$ is u.s.c. Moreover, we claim that $P_{f,A}(\cdot)$ has nonempty, compact, convex values. Nonemptiness follows from Proposition 2.1 of [7], while convexity is a straightforward consequence of the sublinearity of $f(\cdot)$. For compactness let $\{z_a\}$ be a net in $P_{f,A}(x)$. Then by definition we have that $f_A(x) = f(x - z_a)$. So $\{z_a\}$ is trivially a minimizing net in A . Because by hypothesis A is f -approximately compact, we can find a subnet $\{z_b\}$ s.t. $z_b \rightarrow z \in A$. Also because of the continuity of $f(\cdot)$ we get that $f_A(x) = f(x - z)$. So $z \in P_{f,A}(x)$ and this proves that $P_{f,A}(\cdot)$ is compact valued. Then since $\phi(A)$ is compact, we have that $P_{f,A}(\phi(A))$ is compact. Applying Himmelberg's fixed point theorem [11], we get that there exists $h \in A$ s.t. $h \in \Gamma(h) = P_{f,A}(\phi(h)) \Rightarrow f_A(\phi(h)) = f(\phi(h) - h)$. Q.E.D.

Remark. When $f(\cdot) = p(\cdot)$, a continuous seminorm on X , then our theorem recovers the result of Reich [22].

We will conclude with two propositions on the properties of the multifunction $P_{f,A}(\cdot)$. In both X is assumed to be a locally convex space.

PROPOSITION 5.1. *If $f: X \rightarrow \mathbb{R}$ is continuous sublinear and $A \subseteq X$ is nonempty, f -approximately compact, closed, and convex, then for any $K \subseteq X$ nonempty, connected, $P_{f,A}(K) = \bigcup_{x \in K} P_{f,A}(x)$ is connected too.*

Proof. For all $x \in X$, $P_{f,A}(x)$ is convex and so connected. Also recall that $P_{f,A}(\cdot)$ is u.s.c. Hence it maps connected sets to connected set. Thus $P_{f,A}(K)$ is connected. Q.E.D.

PROPOSITION 5.2. *If $f: X \rightarrow \mathbb{R}$ is continuous, sublinear and $A \subseteq X$ is nonempty closed, f -approximately compact on $K \subseteq X$ nonempty and compact, then $P_{f,A}(K)$ is nonempty, compact and $\{x, h\} \in K \times A: h \in P_{f,A}(x)\}$ is compact in $X \times X$.*

Proof. Since A is f -approximately compact on K , for all $x \in K$ $P_{f,A}(x) \in P_K(X)$. Also $P_{f,A}(\cdot)$ is u.s.c. on K . Then the claims of the proposition follow from the results of Smithson [26]. Q.E.D.

ACKNOWLEDGMENT

The authors are very grateful to the referee for his(her) invaluable comments and suggestions.

REFERENCES

1. J. P. AUBIN AND I. EKELAND, "Applied Nonlinear Analysis," Wiley, New York, 1984.
2. M. BENAMARA, Sections mesurables extrémales d'une multiapplication, *C. R. Acad. Sci. Paris* **278** (1974), 1249–1252.
3. B. BROSIOWSKI, F. DEUTSCH, AND G. NÜRNBERGER, Parametric approximation, *J. Approx. Theory* **29** (1980), 261–277.
4. C. CASTAING AND M. VALADIER, "Convex Analysis and Measurable Multifunctions," Lecture Notes in Math., Vol. 580, Springer, Berlin, 1977.
5. F. H. CLARKE, "Optimization and Nonsmooth Analysis," Wiley, New York, 1983.
6. F. H. CLARKE, Generalized gradients of Lipschitz functionals, *Advan. Math.* **40** (1981), 52–67.
7. P. GOVINDARAJULU AND D. PAI, On properties of sets related to f -projections, *J. Math. Anal. Appl.* **73** (1980), 457–465.
8. P. GOVINDARAJULU AND D. PAI, On f -farthest points of sets, *Indian J. Pure Appl. Math.* **14** (1983), 873–882.
9. F. HIAI AND H. UMEGAKI, Integrals, conditional expectations and martingales of multivariable functions, *J. Multivariate Anal.* **7** (1977), 149–182.
10. C. HIMMELBERG, Measurable relations, *Fund. Math.* **87** (1975), 53–72.
11. C. HIMMELBERG, Fixed points of compact multifunctions, *J. Math. Anal. Appl.* **38** (1972), 205–207.
12. J. B. HIRIART-URRUTY, Lipschitz r -continuity of the approximate subdifferential of a convex function, *Math. Scand.* **47** (1980), 123–341.
13. J. B. HIRIART-URRUTY, ε -subdifferential, in "Convex Analysis and Its Applications" (J. P. Aubin and R. Vinter, Eds.), Pitman, London, 1982.
14. R. HOLMES, "A Course in Optimization and Approximation," Lecture Notes in Math, Vol. 257, Springer, Berlin, 1972.
15. A. KOZEK AND Z. SUCHANECKI, Multifunctions of faces for conditional expectations of selectors and Jensen's inequality, *J. Multivariate Anal.* **10** (1980), 579–598.
16. KY FAN, Extensions of two fixed point theorems of F. E. Browder, *Math. Z.* **112** (1969), 234–240.
17. P. J. LAURENT, "Approximation et Optimisation," Hermann, Paris, 1972.
18. L. McLINDEN AND R. BERGSTROM, Preservation of convergence of convex sets and functions in finite dimensions, *Trans. Amer. Math. Soc.* **268** (1981), 127–142.
19. U. MOSCO, On the continuity of the Young–Fenchel transform, *J. Math. Anal. Appl.* **35** (1971), 518–535.
20. N. S. PAPAGEORGIOU AND D. KANDILAKIS, Convergence in approximation and nonsmooth analysis, *J. Approx. Theory* **49** (1987), 41–54.
21. N. S. PAPAGEORGIOU, On the theory of Banach space valued multifunctions. Part I: Integration and conditional expectation, *J. Multivariate Anal.* **17** (1985), 185–206.
22. S. REICH, Approximate selections, best approximations, fixed points and invariant sets, *J. Math. Anal. Appl.* **62** (1978), 104–131.
23. R. T. ROCKAFELLAR, Integral functionals, normal integrands and measurable selections, in "Nonlinear Operators and the Calculus of Variations" (J. P. Gossez et al., Eds.), Lecture Notes in Math., Vol. 543, Springer, Berlin, 1976.
24. R. T. ROCKAFELLAR, "Conjugate Duality and Optimization," C.B.M.S. No. 16, S.I.A.M., Philadelphia, 1974.
25. G. SALINETTI AND R. WETS, On the relations between two types of convergence of convex functions, *J. Math. Anal. Appl.* **60** (1977), 211–262.
26. R. SMITHSON, Multifunctions, *Nieuw Arch. Wisk.* **20** (1972), 31–53.
27. L. THIBAUT, Esperances conditionnelles d'intégrandes semi-continues, *Ann. Inst. Henri Poincaré* **17** (1981), 337–350.

28. M. TSUKADA, Convergence of best approximations in a smooth Banach space, *J. Approx. Theory* **40** (1984), 301–309.
29. L. P. VLASOV, Approximative properties of sets in normed linear spaces, *Russian Math. Surveys* **28** (1973), 1–95.