

## Coronary Artery Disease

# Predictors of Cardiac Rehabilitation Referral in Coronary Artery Disease Patients

## Findings From the American Heart Association's Get With The Guidelines Program

Todd M. Brown, MD, MSPH,\* Adrian F. Hernandez, MD, MHS,† Vera Bittner, MD, MSPH,\* Christopher P. Cannon, MD,‡ Gray Ellrodt, MD,§ Li Liang, PhD,† Eric D. Peterson, MD, MPH† Ileana L. Piña, MD,|| Monika M. Safford, MD,\* Gregg C. Fonarow, MD,¶ on behalf of the American Heart Association Get With The Guidelines Investigators

*Birmingham, Alabama; Durham, North Carolina; Boston and Pittsfield, Massachusetts; Cleveland, Ohio; and Los Angeles, California*

**Objectives**

Our purpose was to determine factors independently associated with cardiac rehabilitation referral, which are currently not well described at a national level.

**Background**

Substantial numbers of eligible patients are not referred to cardiac rehabilitation at hospital discharge despite proven reductions in mortality and national guideline recommendations.

**Methods**

We used data from the American Heart Association's Get With The Guidelines program, analyzing 72,817 patients discharged alive after a myocardial infarction, percutaneous coronary intervention, or coronary artery bypass graft surgery between January 2000 and September 2007 from 156 hospitals. We identified factors associated with cardiac rehabilitation referral at discharge and performed multivariable logistic regression, adjusted for clustering, to identify which factors were independently associated with cardiac rehabilitation referral.

**Results**

Mean age was  $64.1 \pm 13.0$  years, 68% were men, 79% were white, and 30% had diabetes, 66% hypertension, and 52% dyslipidemia; mean body mass index was  $29.1 \pm 6.3$  kg/m<sup>2</sup>, and mean ejection fraction  $49.0 \pm 13.6\%$ . All patients were admitted for coronary artery disease (CAD), with 71% admitted for myocardial infarction. Overall, only 40,974 (56%) were referred to cardiac rehabilitation at discharge, ranging from 53% for myocardial infarction to 58% for percutaneous coronary intervention and to 74% for coronary artery bypass graft patients. Older age, non-ST-segment elevation myocardial infarction, and the presence of most comorbidities were associated with decreased odds of cardiac rehabilitation referral.

**Conclusions**

Despite strong evidence for benefit, only 56% of eligible CAD patients discharged from these hospitals were referred to cardiac rehabilitation. Increased physician awareness about the benefits of cardiac rehabilitation and initiatives to overcome barriers to referral are critical to improve the quality of care of patients with CAD. (J Am Coll Cardiol 2009;54:515–21) © 2009 by the American College of Cardiology Foundation

Cardiac rehabilitation reduces morbidity and mortality in coronary artery disease (CAD) patients after myocardial infarction, improves risk factor management, and is a class I indication in numerous national guidelines after myocardial

infarction or revascularization (1–8). However, despite the proven benefits of cardiac rehabilitation and these national guideline recommendations, cardiac rehabilitation continues to be significantly underutilized, based predominately on

From the \*University of Alabama at Birmingham, Birmingham, Alabama; †Duke Clinical Research Institute, Durham, North Carolina; ‡Brigham and Women's Hospital, Boston, Massachusetts; §Berkshire Medical Center, Pittsfield, Massachusetts; ||Case Western Reserve University, Cleveland, Ohio; and the ¶UCLA Medical Center, Los Angeles, California. Get With The Guidelines-CAD is sponsored by the American Heart Association with funding, in part, from an unrestricted education grant from the Merck/Schering-Plough Partnership. At the time of this work, Dr. Brown was supported, in part, by grant 5 T32 HS013852 from the Agency for Healthcare Research and Quality, Rockville, Maryland. Dr. Hernandez has received research funding from Johnson & Johnson, GlaxoSmithKline, Merck, and Medtronic,

and honoraria from Novartis and AstraZeneca. Dr. Cannon has received research grants/support from Accumetrics, AstraZeneca, Bristol-Myers Squibb/Sanofi Partnership, GlaxoSmithKline, Merck, and Merck/Schering-Plough Partnership, and has been a clinical advisor for and has equity in Automedics Medical Systems. Dr. Piña is on the Speakers' Bureau of Solvay, AstraZeneca, Innovia, Merck, Sanofi-Aventis, and Novartis; is a consultant for the Food and Drug Administration; and has received a grant/research from the National Institutes of Health. Dr. Fonarow has received honorarium as a consultant for Merck, Schering-Plough, and GlaxoSmithKline.

Manuscript received December 18, 2008; revised manuscript received February 2, 2009, accepted February 6, 2009.

## Abbreviations and Acronyms

<b>ACC</b>	= American College of Cardiology
<b>AHA</b>	= American Heart Association
<b>AOR</b>	= adjusted odds ratio
<b>CABG</b>	= coronary artery bypass graft
<b>CAD</b>	= coronary artery disease
<b>CI</b>	= confidence interval
<b>GWTG</b>	= Get With The Guidelines
<b>OR</b>	= odds ratio
<b>PCI</b>	= percutaneous coronary intervention

single-center evaluations (1,9-17). A recent analysis of national Medicare claims data also indicated that cardiac rehabilitation use is suboptimal, as only 19% of Medicare patients enrolled in cardiac rehabilitation after a myocardial infarction or coronary artery bypass graft (CABG) surgery (9). These claims-based analyses, however, lacked the detailed clinical data necessary to accurately determine eligibility for cardiac rehabilitation or to determine what percentage had actually been referred for cardiac rehabilitation.

In response to these data concerning the underutilization of cardiac rehabilitation, the American Association of Cardiovascular and Pulmonary Rehabilitation, the American College of Cardiology (ACC), and the American Heart Association (AHA) recently published performance measures for the referral to and delivery of cardiac rehabilitation services (18). These performance measures state that all hospitalized patients with a qualifying cardiovascular disease event should be referred to outpatient cardiac rehabilitation before hospital discharge. Qualifying cardiovascular disease events include myocardial infarction, percutaneous coronary intervention (PCI), CABG surgery, stable angina, heart transplant, and heart valve surgery. As the authors of these performance measures emphasize, numerous barriers preventing attendance in cardiac rehabilitation exist, but the most important and easily overcome barrier is the failure of health care providers to refer eligible patients to cardiac rehabilitation (18). Understanding referral patterns and barriers to cardiac rehabilitation referral is necessary to improve the utilization of cardiac rehabilitation in eligible CAD patients.

Our current study extends these prior works investigating cardiac rehabilitation underutilization by analyzing cardiac rehabilitation referral among eligible patients with myocardial infarction, PCI, or CABG surgery discharged from hospitals participating in the AHA's Get With The Guidelines (GWTG) program. Furthermore, we identified factors that were independently associated with cardiac rehabilitation referral.

Our current study extends these prior works investigating cardiac rehabilitation underutilization by analyzing cardiac rehabilitation referral among eligible patients with myocardial infarction, PCI, or CABG surgery discharged from hospitals participating in the AHA's Get With The Guidelines (GWTG) program. Furthermore, we identified factors that were independently associated with cardiac rehabilitation referral.

## Methods

**Data source.** The AHA GWTG program is a voluntary, observational data collection and quality-improvement initiative that began in 2000 and has been described in detail previously (19). Participating hospitals use the point-of-service, interactive, Internet-based Patient Management Tool (Outcome Sciences, Inc., Cambridge, Massachusetts)

to submit clinical information regarding in-hospital care and outcomes of patients hospitalized for CAD, stroke, or heart failure. Participating hospitals submit consecutive, eligible patients to the database, and compliance with local regulatory and privacy guidelines is required.

Trained personnel abstract the data, and patients are assigned to race/ethnicity categories using options defined by the electronic case report form. Other variables collected in the CAD module include sociodemographic and clinical characteristics, medical history, admitting diagnosis, inpatient medical therapies and procedures, ejection fraction, hospital characteristics, compliance with CAD-related performance measures including referral to cardiac rehabilitation, contraindications for evidence-based therapies, and in-hospital outcomes. Referral to cardiac rehabilitation is recorded as: 1) yes; 2) no; 3) not documented; or 4) not applicable.

Outcome Sciences, Inc. serves as the data collection and coordination center for GWTG. The Duke Clinical Research Institute serves as the data analysis center and has an agreement and institutional review board approval to analyze the aggregate de-identified data for research purposes. The Internet-based data entry system includes edit checks to insure the completeness of the reported data. Data quality is monitored and reports are generated to ensure the completeness and accuracy of the submitted data. Only sites and variables with a high degree of completeness are included in analyses.

**Study population.** We identified 310,563 patients with CAD-related diagnoses discharged from 555 GWTG participating hospitals between January 2000 and September 2007 in the GWTG CAD module. We limited our analysis to individuals with class I indications for cardiac rehabilitation by only including patients discharged alive who had an acute myocardial infarction, PCI, or CABG surgery during their hospitalization. We excluded patients whose primary admission diagnoses were non-CAD related (n = 26,932), those without an acute myocardial infarction, PCI, or CABG surgery (n = 59,121), and those discharged to skilled nursing facilities, acute care facilities, hospice, or against medical advice; those who expired; or those whose discharge status was not documented or missing (n = 48,302). Additionally, to ensure the accuracy of our analysis, patients discharged from hospitals failing to report data on past medical history or in-hospital procedures >10% of the time (n = 52,483 patients from 201 hospitals) or cardiac rehabilitation referral >25% of the time (n = 50,908 patients from 170 hospitals) were also excluded. This resulted in a final sample size of 72,817 patients from 156 hospitals. Compared with those included in the analysis, hospitals excluded from the analysis had fewer beds, were less likely to be a teaching hospital, less likely to have interventional and heart transplant capabilities, and more likely to be from the northeast.

**Statistical analysis.** The main outcome measure was the proportion of patients referred to cardiac rehabilitation at discharge. Individuals with missing values for cardiac reha-

bilitation referral, cardiac rehabilitation referral not documented, or those with cardiac rehabilitation recorded as “not applicable” were considered not referred for this analysis. Because “not applicable” might be used to document those felt to have a contraindication for cardiac rehabilitation, we also calculated the proportion of individuals referred to cardiac rehabilitation after excluding those individuals where “not applicable” was recorded ( $n = 7,454$ ).

Using chi-square tests for categorical variables and Wilcoxon rank sum tests for continuous variables, we compared the baseline characteristics of patients referred to cardiac rehabilitation with those not referred. Additionally, we were interested in whether individuals referred to cardiac rehabilitation were more likely to receive other evidence-based therapies during their hospitalization. To investigate this, we compared compliance with established CAD performance measures (20,21) between those referred and not referred to cardiac rehabilitation using the chi-square test. Percentages and medians with interquartile ranges were reported for categorical and continuous variables, respectively.


A multivariable logistic regression analysis using the generalized estimating equations method, adjusting for within-hospital clustering, was performed to identify factors associated with cardiac rehabilitation referral at discharge. The initial model included age, sex, race, body mass index, systolic blood pressure, insurance status, admitting diagnosis, procedures (PCI or CABG), hospital characteristics (bed size, region, teaching status, and transplant capability), and patient comorbidities including anemia, stroke, heart failure, prior myocardial infarction, diabetes, hyperlipidemia, hypertension, chronic obstructive pulmonary disease, peripheral arterial disease, chronic kidney disease (serum creatinine  $>2.0$  mg/dl), dialysis, atrial fibrillation or flutter, depression, pacemaker or implantable cardioverter-defibrillator, and alcohol or tobacco use. Factors with a  $p$  value  $<0.05$  were kept in the reduced model,

and the odds ratios (ORs) for the significant factors in the reduced model were reported. A secondary analysis adding a variable for the presence of a left ventricular ejection fraction  $<40\%$  into the reduced model was performed to evaluate the influence of left ventricular systolic dysfunction on the probability of cardiac rehabilitation referral. A  $p$  value  $<0.05$  was considered statistically significant for all tests. All analyses were performed using SAS software version 9.1 (SAS Institute, Cary, North Carolina).

## Results

Of the 72,817 individuals discharged from these 156 GWTG participating hospitals after a myocardial infarction, PCI, or CABG surgery from January 2000 to September 2007, only 40,974 (56%) were referred to cardiac rehabilitation at hospital discharge. This ranged from 53% for patients admitted with a myocardial infarction, to 58% for those undergoing PCI, to 74% for those undergoing CABG surgery. After excluding individuals where cardiac rehabilitation was deemed to be “not applicable,” still only 63% (40,974 of 65,363) were referred. The distribution of referral rates by hospital is displayed in Figure 1. The median referral rate by hospital was 43% and ranged from 0% to 100%. As shown in Figure 1, 35% of hospitals referred fewer than 20% of eligible patients, and only one-third of hospitals referred over 60% of eligible patients.

Table 1 lists the demographic and clinical characteristics in the overall population as well as in those referred and not referred to cardiac rehabilitation. When compared with those not referred, individuals referred to cardiac rehabilitation were younger, more likely to be male and white race, less likely to have an unspecified or non-ST-segment elevation myocardial infarction, less likely to have Medicare, less likely to have most medical comorbidities, and more likely to have undergone PCI or CABG. Table 2 lists the proportion of eligible individuals



**Figure 1** Cardiac Rehabilitation Referral Patterns Among 156 GWTG Participating Hospitals, 2000 to 2007

GWTG = Get With The Guidelines.

Table 1	Baseline Demographic and Clinical Characteristics in the Overall Population and in Those Referred and Not Referred to CR			
	Overall Population (n = 72,817)	Not Referred to CR (n = 31,843)	Referred to CR (n = 40,974)	p Value
Age (yrs)	64 (55–74)	65 (55–75)	63 (54–73)	<0.0001
Sex (%)				
Female	32.1	33.7	30.8	<0.0001
Race (%)				
White	79	76	82	
African American	7	7	7	<0.0001
Hispanic	7	10	5	
Body mass index (kg/m <sup>2</sup> )	28.2 (25.0–32.2)	27.9 (24.7–32.0)	28.4 (25.2–32.4)	<0.0001
Ejection fraction (%)	50 (40–60)	50 (40–60)	50 (40–60)	0.0039
Admitting diagnosis (%)				
Unspecified MI	45.3	51.1	40.7	
CAD	20.7	15.8	24.6	
NSTEMI	16.3	17.4	15.4	<0.0001
STEMI	9.1	7.4	10.4	
Unstable angina	8.7	8.3	8.9	
Insurance status (%)				
Other	56.3	50.9	60.5	
Medicare	32.3	37.2	28.5	<0.0001
None	8.2	8.4	8.1	
Medicaid	3.2	3.4	3.0	
Comorbidities (%)				
Hypertension	66.3	66.8	65.9	0.0192
Dyslipidemia	51.6	47.8	54.6	<0.0001
Smoking	31.4	29.0	33.3	<0.0001
Diabetes	30.1	31.0	29.4	<0.0001
Prior MI	19.2	19.7	18.8	0.0029
COPD	11.5	12.4	10.9	<0.0001
PAD	7.5	8.1	7.1	<0.0001
Stroke or TIA	5.8	6.9	4.9	<0.0001
Chronic dialysis	1.5	1.9	1.2	<0.0001
ICD	0.2	0.3	0.2	0.0086
In-hospital procedures (%)				
PCI	66.8	64.5	68.6	<0.0001
CABG	15.3	9.0	20.2	<0.0001

All values listed as median (interquartile range) or %. Wilcoxon 2-sample test performed for continuous variables; chi-square test performed for categorical variables.

CABG = coronary artery bypass graft surgery; CAD = coronary artery disease; COPD = chronic obstructive pulmonary disease; CR = cardiac rehabilitation; ICD = implantable cardioverter-defibrillator; MI = myocardial infarction; NSTEMI = non-ST-segment elevation myocardial infarction; PAD = peripheral arterial disease; PCI = percutaneous coronary intervention; STEMI = ST-segment elevation myocardial infarction; TIA = transient ischemic attack.

who received therapies in compliance with established performance measures in the overall population and among those referred or not referred to cardiac rehabilitation. Individuals referred to cardiac rehabilitation were also more likely to receive therapies in accordance with each of these performance measures.

Factors independently associated with cardiac rehabilitation referral in the final multivariable model are listed in Table 3. Undergoing CABG surgery (adjusted odds ratio [AOR]: 3.00) and PCI (AOR: 1.67) were the factors most strongly associated with cardiac rehabilitation referral. Other factors independently associated with referral included younger age, ST-segment elevation myocardial infarction, and a history of dyslipidemia or smoking. Most comorbidities were associated with decreased odds for referral. Adding left ventricular systolic

dysfunction (left ventricular ejection fraction <40%) to the model did not significantly alter the factors associated with cardiac rehabilitation referral. Patients with left ventricular systolic dysfunction were slightly less likely to be referred to cardiac rehabilitation (AOR: 0.96, 95% confidence interval [CI]: 0.92 to 1.01), although this association was not statistically significant.

### Discussion

Participation in cardiac rehabilitation is associated with reductions in mortality and recurrent myocardial infarction. Taylor et al. (22), in a meta-analysis of randomized clinical trials of exercise-based cardiac rehabilitation, demonstrated that participation in cardiac rehabilitation was associated



**Table 2** Proportion of Eligible Individuals in the Overall Population and in Those Referred and Not Referred to CR Who Received Therapies in Compliance With Established Performance Measures

	Overall Population (n = 72,817)	Not Referred to CR (n = 31,843)	Referred to CR (n = 40,974)	p Value
During hospitalization				
AMI and angina patients without contraindication receiving aspirin <24 h	95.6	94.3	96.7	<0.0001
Current smokers that receive smoking cessation advice	92.1	88.0	94.9	<0.0001
Patients with an LDL-C >100 mg/dl receiving lipid-lowering drugs	89.6	85.3	92.5	<0.0001
At discharge				
Patients with documented LVSD discharged on an ACE-I or ARB	83.5	80.9	85.6	<0.0001
Patients discharged on aspirin	97.6	96.5	98.5	<0.0001
Patients discharged on beta-blockers	93.0	91.7	94.0	<0.0001

All values listed are %.

ACE-I = angiotensin-converting enzyme inhibitor; AMI = acute myocardial infarction; ARB = angiotensin receptor blocker; CI = contraindication; CR = cardiac rehabilitation; LDL-C = low-density lipoprotein cholesterol; LVSD = left ventricular systolic dysfunction.

with significant reductions in both all-cause mortality (OR: 0.80, 95% CI: 0.68 to 0.93) and cardiac-specific mortality (OR: 0.74, 95% CI: 0.61 to 0.96) after a median follow-up of 15 months (22). Witt et al. (23) have subsequently extended these data by demonstrating that participation in contemporary, community-based cardiac rehabilitation programs, which enroll older and higher-risk patients not enrolled in clinical trials and provide more comprehensive approaches to secondary prevention in addition to exercise training, is associated with reduced mortality and fewer recurrent myocardial infarctions (23). Despite these proven benefits of cardiac rehabilitation, previous studies from

predominantly single centers within the U.S. report referral rates of less than 25% to 30% (11-15). In our study, only approximately one-half of eligible patients in this large, nationwide sample of 72,817 patients discharged from 156 GWTG participating hospitals were referred to cardiac rehabilitation. We also observed substantial variation among hospitals in the percentage of eligible patients referred to cardiac rehabilitation. In 35% of hospitals, fewer than 20% of eligible patients were referred, while only one-third referred over 60%. Given that GWTG hospitals in general (24,25), as well as the subset of those hospitals included in this analysis, report very good compliance with virtually all other secondary prevention performance measures, we believe our results represent a best-case scenario among hospitals with high adherence rates to core measures and sufficient resources for excellent data collection.

We also observed that individuals not referred to cardiac rehabilitation were less likely to have received other guideline-based therapies as defined by recent performance measures (20,21). Individuals not referred to cardiac rehabilitation were less likely to be discharged on aspirin, beta-blockers, angiotensin-converting enzyme inhibitors, or angiotensin receptor blockers if they had left ventricular systolic dysfunction, to receive lipid-lowering agents if they had a low-density lipoprotein cholesterol level >100 mg/dl, and to receive smoking cessation counseling. Most alarming, however, is that despite the wealth of data on the benefits of cardiac rehabilitation, the overall referral rate to cardiac rehabilitation (56%) was far lower than for any of the other performance measures studied (which ranged from 84% to 98%), suggesting that physician awareness about the benefits of cardiac rehabilitation is lower than for other interventions.

Cortes and Arthur (17) recently systematically reviewed 10 studies to examine referral patterns to cardiac rehabilitation. Major predictors of referral to cardiac rehabilitation included being English speaking, admitted to a hospital with a cardiac rehabilitation program, insurance status, and previous myocardial infarction. Other less significant predictors included younger age, cardiac catheterization, hypercholesterolemia, CABG surgery, hypertension, and

**Table 3** Predictors of Referral to Cardiac Rehabilitation in the Overall Population

Variable	Adjusted Odds Ratio (95% CI)
Age (yrs)	
≤50	Referent
51-65	1.00 (0.96-1.04)
66-80	0.92 (0.87-0.97)
>80	0.76 (0.71-0.82)
Admitting diagnosis	
STEMI	Referent
NSTEMI	0.84 (0.76-0.93)
Unspecified MI	0.73 (0.53-0.99)
CAD	0.71 (0.57-0.87)
Unstable angina	0.63 (0.50-0.78)
Comorbidities	
Dyslipidemia	1.15 (1.08-1.23)
Smoking	1.10 (1.05-1.15)
Hypertension	0.95 (0.91-0.99)
Prior MI	0.95 (0.90-0.99)
COPD	0.93 (0.89-0.97)
PAD	0.92 (0.87-0.98)
Stroke/TIA	0.91 (0.87-0.96)
Chronic dialysis	0.79 (0.70-0.90)
ICD	0.74 (0.55-1.00)
Procedures	
CABG	3.00 (2.32-3.87)
PCI	1.67 (1.50-1.85)

CI = confidence interval; other abbreviations as in Table 1.

smoking (17). Our results are similar. Younger age, ST-segment elevation myocardial infarction, and the performance of PCI or CABG surgery were associated with increased odds of cardiac rehabilitation referral. Individuals with dyslipidemia and those who smoked also had increased odds of cardiac rehabilitation referral; in contrast, most comorbidities were associated with lower odds of referral. Individuals with comorbidities may be perceived by physicians as less likely to benefit or less likely to participate in cardiac rehabilitation; however, in many instances, these individuals represent populations at significantly increased cardiovascular risk who may benefit from the more intense secondary prevention services provided in cardiac rehabilitation (23). A recent analysis examining patient and physician factors affecting cardiac rehabilitation referral concluded that among physicians, one of the most important factors influencing whether physicians refer patients to cardiac rehabilitation was the degree of the physician's perceived benefit of cardiac rehabilitation (26). Therefore, increased physician awareness about the benefits of cardiac rehabilitation in these higher-risk subgroups may increase referrals.

Obviously, individuals cannot enroll and participate in cardiac rehabilitation if they are not first referred. Mazzini et al. (27) report that, in a single center using a GWTG-based clinical pathway, 55% of patients with myocardial infarction were referred to cardiac rehabilitation at discharge. However, the subsequent enrollment rate in cardiac rehabilitation for these individuals who were referred before discharge was only 34% (27). With no more than one-half of eligible patients being referred and only one-third of those referred enrolling, it is not surprising that most studies, including a recent analysis of 267,427 Medicare beneficiaries with acute myocardial infarctions or CABG surgery, report overall enrollment rates in cardiac rehabilitation of only about 15% to 20% (9–15). One notable exception is Olmsted County, Minnesota, where 55% of individuals enrolled in cardiac rehabilitation after a myocardial infarction between 1982 and 1998 (2). Numerous barriers exist that prevent referred patients from enrolling in cardiac rehabilitation including cost, lack of insurance coverage, time commitment, and distance from a cardiac rehabilitation center. However, despite these barriers, simply increasing the proportion of eligible patients referred to cardiac rehabilitation, as recommended by numerous ACC/AHA guidelines (4–8) and a recently published American Association of Cardiovascular and Pulmonary Rehabilitation/ACC/AHA performance measure (18), could substantially increase cardiac rehabilitation enrollment.

Investigators in Canada have examined the effectiveness of computer-generated, automatic referral at discharge of eligible patients to cardiac rehabilitation (28–31). These authors report enrollment rates in cardiac rehabilitation ranging from as low as 43% to as high as 73% (28–31). These data support the hypothesis that failure to refer patients to cardiac rehabilitation represents one of the largest and, potentially, most easily overcome barrier to

participation in cardiac rehabilitation. By simply increasing referral rates, these authors report much higher rates of enrollment in cardiac rehabilitation than is reported using standard, physician-initiated referral practices. However, automatically referring patients to cardiac rehabilitation may not be enough. Mazzini et al. (27) report that among those who were referred to cardiac rehabilitation but did not enroll, 26% reported that they did not perceive that they had been referred, reflecting a gap between what is documented in the medical record and what the patient actually recalls after discharge. Additionally, multiple previous studies have demonstrated that physician endorsement of the benefits of cardiac rehabilitation is one of the most powerful predictors of attendance in cardiac rehabilitation among patients who are referred (13,14,16). Therefore, a computer-generated referral, although helpful in prompting physicians to initiate referrals, may not be sufficient to correct the current vast underutilization of cardiac rehabilitation.

**Study limitations.** First, hospital participation in the GWTG program is voluntary. Therefore, the overall proportion of eligible patients referred to cardiac rehabilitation and predictors of referral may not be the same in nonparticipating hospitals. Furthermore, even among participating hospitals, we are not able to determine the referral patterns of those hospitals with a large amount of missing data about cardiac rehabilitation referral. This does limit the generalizability of our findings; however, we are unlikely to have underestimated the proportion of eligible patients referred to cardiac rehabilitation in the U.S. Rather, we believe that our results represent a best-case scenario among hospitals with high adherence rates to core measures in general. Hospitals excluded from our analysis as well as non-GWTG participating institutions very likely refer a lower proportion of eligible CAD patients to cardiac rehabilitation. Therefore, the overall referral rate within the U.S. is likely lower than what we estimate in this analysis. Second, the GWTG program only collects in-hospital data and does not collect data on physician characteristics. Therefore, we are unable to assess what proportion of individuals are referred to cardiac rehabilitation after discharge from the hospital or what percentage of those referred actually attend cardiac rehabilitation. Furthermore, we are unable to determine what physician characteristics contribute to cardiac rehabilitation referral. Third, we lack detailed data on socioeconomic variables such as income and education levels of eligible patients, which prohibits us from assessing the impact that these variables have on referral rates. Fourth, we cannot exclude residual measured and unmeasured confounding variables that might account for these associations with cardiac rehabilitation referral.

## Conclusions

Overall, only approximately one-half of patients discharged from these GWTG participating hospitals who were eligible for cardiac rehabilitation after a myocardial infarction, PCI, or

CABG surgery were referred at hospital discharge. Older individuals as well as those with most of the comorbidities we studied had decreased odds of cardiac rehabilitation referral. Individuals with ST-segment elevation myocardial infarction or who underwent PCI or CABG surgery had increased odds for referral. More emphasis on increasing referral to cardiac rehabilitation is necessary to overcome the current underutilization of cardiac rehabilitation in the secondary prevention of cardiovascular disease.

**Reprint requests and correspondence:** Dr. Todd M. Brown, LHRB 306, 701 19th Street South, Birmingham, Alabama 35294. E-mail: [tmbrown@uab.edu](mailto:tmbrown@uab.edu).

## REFERENCES

1. Wenger NK. Current status of cardiac rehabilitation. *J Am Coll Cardiol* 2008;51:1619–31.
2. Witt BJ, Jacobsen SJ, Weston SA, et al. Cardiac rehabilitation after myocardial infarction in the community. *J Am Coll Cardiol* 2004;44:988–96.
3. Williams MA, Ades PA, Hamm LF, et al. Clinical evidence for a health benefit from cardiac rehabilitation: an update. *Am Heart J* 2006;152:835–41.
4. Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction). *J Am Coll Cardiol* 2004;44:e1–211.
5. Anderson JL, Adams CD, Antman EM, et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non-ST-Elevation Myocardial Infarction). *J Am Coll Cardiol* 2007;50:e1–157.
6. Gibbons RJ, Abrams J, Chatterjee K, et al. ACC/AHA 2002 guideline update for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for the Management of Patients with Chronic Stable Angina). *J Am Coll Cardiol* 2003;41:159–68.
7. Eagle KA, Guyton RA, Davidoff R, et al. ACC/AHA 2004 guideline update for coronary artery bypass graft surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for Coronary Artery Bypass Graft Surgery). *J Am Coll Cardiol* 2004;44:e213–310.
8. Smith SC Jr, Feldman TE, Hirshfeld JW Jr, et al. ACC/AHA/SCAI 2005 guideline update for percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/SCAI Writing Committee to Update the 2001 Guidelines for Percutaneous Coronary Intervention). *J Am Coll Cardiol* 2006;47:e1–121.
9. Suaya JA, Shepard DS, Normand S-L, Ades PA, Protas J, Stason WB. Use of cardiac rehabilitation by Medicare beneficiaries after myocardial infarction or coronary bypass surgery. *Circulation* 2007;116:1653–62.
10. Mochari H, Lee JR, Kligfield P, Mosca L. Ethnic differences in barriers and referral to cardiac rehabilitation among women hospitalized with coronary heart disease. *Prev Cardiol* 2006;9:8–13.
11. Blackburn GG, Foody JM, Spencer DL, Park E, Apperson-Hansen C, Pashkow FJ. Cardiac rehabilitation participation patterns in a large, tertiary care center: evidence for selection bias. *J Cardiopulm Rehabil* 2000;20:189–95.
12. Bittner V, Sanderson B, Breland J, Green D. Referral patterns to a university-based cardiac rehabilitation program. *Am J Cardiol* 1999;83:252–5.
13. Ades PA, Waldmann ML, Polk DM, Coflesky JT. Referral patterns and exercise response in the rehabilitation of female coronary patients aged  $\geq 62$  years. *Am J Cardiol* 1992;69:1422–5.
14. Barber K, Strommel M, Kroll J, Holmes-Rovner M, McIntosh B. Cardiac rehabilitation for community-based patients with myocardial infarction: factors predicting discharge recommendation and participation. *J Clin Epidemiol* 2001;54:1025–30.
15. Roblin D, Discker RA III, Orenstein D, Wilder M, Eley M. Delivery of outpatient cardiac rehabilitation in a managed care organization. *J Cardiopulm Rehabil* 2004;24:157–64.
16. Jackson L, Leclerc J, Erksine Y, Linden W. Getting the most out of cardiac rehabilitation: a review of referral and adherence predictors. *Heart* 2005;91:10–4.
17. Cortes O, Arthur HM. Determinants of referral to cardiac rehabilitation programs in patients with coronary artery disease: a systematic review. *Am Heart J* 2006;151:249–56.
18. Thomas RJ, King M, Lui K, Oldridge N, Pina IL, Spertus J. AACVPR/ACC/AHA 2007 performance measures on cardiac rehabilitation for referral to and delivery of cardiac rehabilitation/secondary prevention services. *J Am Coll Cardiol* 2007;50:1400–33.
19. Hong Y, LaBresh KA. Overview of the American Heart Association “Get With the Guidelines” programs: coronary heart disease, stroke, and heart failure. *Crit Pathways in Cardiol* 2006;5:179–86.
20. Krumholz HM, Anderson JL, Brooks NH, et al. ACC/AHA clinical performance measures for adults with ST-elevation and non-ST-elevation myocardial infarction: a report of the ACC/AHA Task Force on Performance Measures (ST-Elevation and Non-ST-Elevation Myocardial Infarction Performance Measures Writing Committee). *J Am Coll Cardiol* 2006;47:236–65.
21. Drozda JP, Messer JV, Spertus J. Clinical Performance Measure for Chronic Stable Coronary Artery Disease. Available at: <http://www.ama-assn.org/ama1/pub/upload/mm/370/cadminisetjune06.pdf>. Accessed January 12, 2009.
22. Taylor RS, Brown A, Ebrahim S, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004;116:682–92.
23. Witt BJ, Jacobsen SJ, Weston SA, et al. Cardiac rehabilitation after myocardial infarction in the community. *J Am Coll Cardiol* 2004;44:988–96.
24. LaBresh KA, Ellrodt AG, Gliklich R, Liljestrand J, Peto R. Get With the Guidelines for cardiovascular secondary prevention. *Arch Intern Med* 2004;164:203–9.
25. LaBresh KA, Fonarow GC, Smith SC Jr, et al. Improved treatment of hospitalized coronary artery disease patients with the Get With the Guidelines program. *Crit Pathways in Cardiol* 2007;6:98–105.
26. Grace SL, Gravelly-White S, Brual J, et al. Contribution of patient and physician factors to cardiac rehabilitation referral: a prospective multilevel study. *Nat Clin Pract Cardiovasc Med* 2008;10:653–62.
27. Mazzini MJ, Stevens GR, Whalen D, Ozonoff A, Balady GJ. Effect of an American Heart Association Get With the Guidelines Program-based clinical pathway on referral and enrollment into cardiac rehabilitation after acute myocardial infarction. *Am J Cardiol* 2008;101:1084–7.
28. Grace SL, Evindar A, Kung T, Scholey P, Stewart DE. Increasing access to cardiac rehabilitation: automatic referral to the program nearest home. *J Cardiopulm Rehabil* 2004;24:171–4.
29. Smith KM, Harkness K, Arthur HM. Predicting cardiac rehabilitation enrollment: the role of automatic physician referral. *Eur J Cardiovasc Prev Rehabil* 2006;13:60–6.
30. Grace SL, Evindar A, Kung T, Scholey PE, Stewart DE. Automatic referral to cardiac rehabilitation. *Med Care* 2004;42:661–9.
31. Grace SL, Scholey P, Suskin N, et al. A prospective comparison of cardiac rehabilitation enrollment following automatic vs. usual referral. *J Rehabil Med* 2007;39:239–45.

**Key Words:** coronary artery disease ■ exercise ■ prevention ■ cardiac rehabilitation.