

Surprisingly, in contrast to the large immobilizing effect observed upon the addition of *extraliposomal* inhibitor 792949, the inclusion of *intraliposomal* inhibitor caused only a minor spectral change. This observation indicates that binding of 792949 to CTP from the internal surface of the proteoliposomes (i.e., the *matrix-facing* conformation) occurs to a much lesser extent than does binding to CTP from the external surface (i.e., the *cytosolic-facing* conformation). We conclude that external 792949 affects spin-label mobility at both monomers within the functional homodimer suggesting a tight coordination of the two monomers. Supported by NIH grant GM-054642 to R.S.K.

### 3574-Pos

#### Characterization of Substrate Binding by the Bacterial Aspartate Transporter Glt<sub>Ph</sub> Through Equilibrium and Stopped-Flow Tryptophan Fluorescence Measurements

David Ewers, Toni Becher, Rabea Krefft, Patricia Hidalgo, Christoph Fahlke. Medizinische Hochschule Hannover, Hannover, Germany.

Excitatory amino acid transporters (EAATs) remove glutamate from the synaptic cleft to ensure low resting glutamate concentrations and to terminate glutamatergic synaptic transmission. We here study substrate binding to a bacterial EAAT paralogue with known structure, Glt<sub>Ph</sub> from *Pyrococcus horikoshii*, using fluorescence spectroscopy. We expressed mutant transporter with an inserted tryptophan in the TM3-TM4-linker, L130W Glt<sub>Ph</sub>, and studied tryptophan fluorescence of solubilized and purified L130W Glt<sub>Ph</sub>. In the presence of Na<sup>+</sup>, addition of aspartate causes changes in fluorescence intensities, as does addition of Na<sup>+</sup> in the presence of aspartate, allowing the construction of aspartate and Na<sup>+</sup> binding isotherms. Titrations of the mutant protein with aspartate at increasing temperatures resulted in increased apparent dissociation constants. Van't Hoff plots were linear over a measured range from 10 to 40°C, yielding a large negative aspartate binding enthalpy that is partly compensated by a negative binding entropy. Na<sup>+</sup> binding equilibria were less temperature dependent, but Na<sup>+</sup> binding showed to be enthalpy driven as well. Pre-equilibrium kinetics of substrate binding were monitored by measuring fluorescent changes after rapid application of substrates to L130W Glt<sub>Ph</sub>. Exponential fits to the binding transients required two time constants  $\tau_1$  and  $\tau_2$ , reflecting at least two underlying processes. Most of the change in fluorescence was associated with the fast process with  $\tau_1$  in the range of hundreds of ms.  $\tau_1^{-1}$  showed linear dependence on [Na<sup>+</sup>], suggesting that the fast process represents Na<sup>+</sup> binding. Aspartate uptake by Glt<sub>Ph</sub> was recently shown to exhibit a Q<sub>10</sub> of 3.7, indicating at least one significant conformational change during the transport cycle (Ryan et al. (2009) J Biol Chem 284, 17540-17548.).  $\tau_1$  was decreased by rising temperatures, but with lower Q<sub>10</sub> than the whole transport cycle.

### 3575-Pos

#### Detection of Substrate-Dependent Conformational Changes in HP1 of the Glutamate Transporter Glt<sub>PH</sub>

Derek Francis<sup>1</sup>, David S. Cafiso<sup>2</sup>, Joseph A. Mindell<sup>1</sup>.

<sup>1</sup>National Institutes of Health, Bethesda, MD, USA, <sup>2</sup>University of Virginia, Charlottesville, VA, USA.

Glutamate acts as the primary excitatory neurotransmitter in the mammalian central nervous system. Clearance of this neurotransmitter from the synapse is accomplished by a family of glutamate transporters known as EAATs, which move synaptic glutamate across the cell membrane into the cell against its concentration gradient. It has long been speculated that the mechanism for transport involves the movement of extracellular and intracellular gates, providing "alternating access" to a substrate binding site. Recently, the crystal structure of a related bacterial transporter, Glt<sub>Ph</sub>, was solved, revealing two helical hairpins (HP1 and HP2) which have been proposed to contribute to these gates. A number of studies have shown that HP2, which lies on the extracellular side of the protein, can adopt multiple conformations that either provide or restrict access to the substrate binding site. However, to date there is no structural information describing conformational changes involving HP1. Here we use the technique of site-directed spin labeling (SDSL) electron paramagnetic resonance (EPR) spectroscopy to explore the local structure and dynamics of residues within HP1 (residues 264-283) in purified Glt<sub>Ph</sub> reconstituted into proteoliposomes. The EPR spectra suggest that the protein exists in two conformational states under our purification conditions. Upon addition of substrate, we note changes in the relative abundance of these states. We are currently working to further characterize each of these conformational states in order to better understand the structural dynamics associated with substrate transport.

### 3576-Pos

#### Mechanism of Interaction of the Glutamate Transporter EAAC1 with K<sup>+</sup> Christof Grewer.

Binghamton University, Binghamton, NY, USA.

Forward glutamate transport by the excitatory amino acid carrier EAAC1 is coupled to the inward movement of three Na<sup>+</sup> and one H<sup>+</sup>, and the outward

movement of one K<sup>+</sup> ion. Internal K<sup>+</sup> is known to bind to the transporter after glutamate and Na<sup>+</sup> are unloaded to the cytosol, subsequently initiating relocation of the transporter binding sites to complete the transport cycle. However, parameters of K<sup>+</sup> interaction with EAAC1, such as affinity and voltage dependence, are currently unknown. Here, we determined the steady-state and pre-steady state kinetics of the interaction of K<sup>+</sup> with its extracellular binding site, and the subsequent K<sup>+</sup> transport step, by using transport current recording from EAAC1-transfected cells. Our results show that K<sup>+</sup> binds to its extracellular binding site with high affinity ( $K_m = 4.5$  mM). K<sup>+</sup> affinity is only weakly voltage dependent. However, transient transport currents were observed in response to steps of the transmembrane potential when K<sup>+</sup> was the only cation present. These currents were capacitive in nature and the charge movement followed a Boltzmann-like voltage dependence. Together, these results suggest that the cation binding process senses little of the transmembrane electric field, but that a subsequent K<sup>+</sup>-induced reaction step, possibly the K<sup>+</sup>-dependent transporter relocation, is electrogenic. The rate constant of the voltage dependent step was 70 s<sup>-1</sup>. This result is consistent with previous data that suggested the K<sup>+</sup>-induced relocation to be the rate-limiting step in the transport cycle. We propose a kinetic model, which is based on an alternating access mechanism, including a fast, voltage-independent K<sup>+</sup> binding step and a slow, electrogenic conformational change. Our model can be used to predict the kinetics of the K<sup>+</sup>-dependent half-cycle of the glutamate transport process. This work was supported by NIH grant 2R01NS049335-06A1.

### 3577-Pos

#### Sequence of Events in the Extracellular Half of the Transport Cycle in Glutamate Transporter

Zhijian Huang<sup>1</sup>, Emad Tajkhorshid<sup>2</sup>.

<sup>1</sup>Beckman Institute, Urbana, IL, USA, <sup>2</sup>Biochemistry Department, Beckman Institute, Urbana, IL, USA.

The crystal structure of Glt<sub>Ph</sub>, a bacterial homologue of glutamate transporter (Glt), revealed the structure of the outward-facing occluded state including the substrate and two Na<sup>+</sup> (Na1 and Na2). It has been well established, however, that substrate transport in Glt<sub>Ph</sub> is catalyzed by the co-transport of three Na<sup>+</sup> ions. However, the location of the third Na<sup>+</sup> (Na3) binding site remains unknown. Furthermore, only little is known regarding the sequence of binding events of the substrate and the co-transported ions to Glt<sub>Ph</sub>. In the present study, we investigate the binding sequence of substrate and Na<sup>+</sup> ions to their extracellular binding sites using molecular dynamics simulations of various bound states of the transporter characterizing the solvent accessibility of key residues involved in ion binding and identifying the resulting conformational changes in the transporter. The results show that extracellular water cannot access Asp312 (the putative Na3 binding site) in the apo and substrate-bound states, and that this residue becomes only accessible from the extracellular side upon Na<sup>+</sup> binding to the Na1 binding site. Based on the simulations, we propose that Na3 binds first to the Na1 binding site in the apo state, resulting in hydration of Asp312, and then moves into the Na3 binding site, the latter step likely being driven by membrane potential. The subsequent binding of a second Na<sup>+</sup> ion (Na1) and the substrate results in a partial closure of the extracellular gate and the formation of the Na2 binding site. Finally, Na2 enters its binding site and locks the extracellular gate resulting in formation of the occluded state. We also propose a putative Na3 binding site composed of three highly conserved residues, namely, Asp312, Thr92 and Asn310.

### 3578-Pos

#### Structural Transition Between the Ion-Releasing and Ion-Binding States of a Secondary Membrane Transporter

Jing Li, Emad Tajkhorshid.

University of Illinois at Urbana-Champaign, Urbana, IL, USA.

The crystal structure of Na<sup>+</sup>-coupled galactose symporter (vSGLT) reports the transporter in its substrate-bound state, with a Na<sup>+</sup> ion modeled in a binding site corresponding to that of a homologous protein, leucine transporter (LeuT). In molecular dynamics simulations, however, we find the Na<sup>+</sup> ion unstable, invariably and spontaneously diffusing out of the transporter through a pathway lined by D189, which appears to facilitate the diffusion of the ion toward the cytoplasm. Further analysis of the trajectories and close structural examination, in particular comparison of the Na<sup>+</sup> binding sites of vSGLT and LeuT, strongly indicates that the crystal structure of vSGLT actually represents an ion-releasing state of the transporter. The observed dynamics of the Na<sup>+</sup> ion, in contrast to the substrate, in a 200 ns equilibrium simulation, also suggests that the cytoplasmic release of the Na<sup>+</sup> ion precedes that of the substrate. Through comparison of the "open" conformation of the Na<sup>+</sup> binding site in vSGLT and the "close" conformation in LeuT, we used constrained simulation to develop a model for the ion-binding state in vSGLT. SMD simulations were then used to pull out substrate from the substrate-binding site both in

ion-releasing state and the modeled ion-binding state, confirming that it is more difficult to release the substrate in the presence of the Na<sup>+</sup> ion in its binding site.

Furthermore, local transition of the Na<sup>+</sup> binding site from an ion-releasing state to an ion-binding state in our constrained simulation induced significantly global conformational change in the protein, specifically, partial opening of the periplasmic side and closing of the cytoplasmic side, thus, capturing for the first time large-scale conformational changes between the inward-facing and outward-facing states in the transport cycle of this secondary transporter.

### 3579-Pos

#### Modeling of the Inward-Facing State of LeuT and Dynamics of the Outward-To-Inward Transition

Saher A. Shaikh, Emad Tajkhorshid.

University of Illinois Urbana Champaign, Urbana, IL, USA.

Leucine transporter (LeuT) is a bacterial aminoacid transporter belonging to the neurotransmitter:sodium symporter family. An 'alternating-access model' is proposed for LeuT function, wherein the transporter alternates between outward-facing and inward-facing states. While multiple crystal structures of LeuT bound with substrate and inhibitors have been reported, only the outward-facing state is known. The inward facing state, and the dynamics of the outward-to-inward transition remain undescribed.

Several transporters from different families report a structural fold similar to the basic 'LeuT fold' indicating the significance of this fold in transporter function. These structures include some in the inward-facing state. Exploiting this information, we have generated a model for the inward-facing state of LeuT. Since the inward-facing structure employed for modeling had very low sequence similarity, and moderate structural similarity to LeuT, a combination of several techniques was required for model generation. A detailed modeling approach was adopted, including sequence- and structure-based approaches, combined with molecular dynamics techniques, such as targeted MD. The final model retains the secondary structural features of LeuT and the substrate/ion binding sites, while adopting an inward-facing state.

We have also employed this model to study the dynamics of outward-to-inward transition of LeuT. The behavior of bound substrate and ions during this transition was recorded and shows interesting features relevant to the transport mechanism. Water permeation was monitored with the progress in transition. The main structural elements of LeuT involved in this transition are described. This study thus presents a model of the inward-facing state of LeuT and a description of a possibly general transport mechanism in transporters adopting the LeuT fold.

### 3580-Pos

#### Molecular Determinants of the Stoichiometry of Transport in GlpT

Giray Enkavi, Emad Tajkhorshid.

Department of Biochemistry, Center for Biophysics and Computational Biology, Beckman Institute for Advanced Science and Technology,

University of Illinois at Urbana-Champaign, Urbana, IL, USA.

The glycerol-3-phosphate/phosphate antiporter GlpT belongs to the Major facilitator superfamily (MFS), the largest group of secondary active membrane transporters. Stoichiometry of transport in organophosphate/phosphate antiporters has been a long-standing question. Experimental studies performed on hexose-6-phosphate transporter (UhpT), a close homolog of GlpT, suggested that the exchange stoichiometry might be regulated by pH in these antiporters. The crystal structure of GlpT, although devoid of a bound substrate, seemed to contradict with "variable stoichiometry" hypothesis, featuring a "single" putative binding site with two arginines (R45 and R269). The "putative" binding site also involves a histidine residue (H165) whose titration state have been suggested to play a key role in transport. We have previously identified one of the arginines (R45) as the binding site residue, but our simulations showed no indication of binding to the other arginine (R269). In order to examine the capacity of GlpT for binding two substrates simultaneously, and to investigate the molecular basis of the "variable stoichiometry" model, we have performed an exhaustive set of MD simulations in which binding of a second substrate to GlpT was simulated. The simulations, performed at different titration states of the substrates (Pi & G3P) and of H165, indicate that the GlpT binding site can, indeed, accommodate two substrates simultaneously upon protonation of H165 and R269 is the preferential binding residue for the second substrate. Moreover, combining of the trajectories of MD simulations with pKa calculations based on continuum electrostatics, we also analyzed the substrate-induced changes of the titration state in the binding site. Our results further indicate that H165 might act as a "pH sensor / stoichiometric switch" in addition to coordinating the substrate. Our findings might represent a general mechanism for transporters with "variable stoichiometry".

### 3581-Pos

#### Conformational Changes in the ApcT Amino Acid Transporter: Monte Carlo Normal Mode Following

Gennady V. Miloshevsky<sup>1,2</sup>, Ahmed Hassanein<sup>1</sup>, Peter C. Jordan<sup>2</sup>.

<sup>1</sup>Purdue University, West Lafayette, IN, USA, <sup>2</sup>Brandeis University, Waltham, MA, USA.

Amino acid/polyamine/organocation (APC) transporters belong to a large family (~250) of secondary transport proteins that catalyze bilayer translocation of a broad range of substrates. Monte Carlo Normal Mode Following [Miloshevsky & Jordan, *Structure* **14**, 1241 (2006); **15**, 1654 (2007)] is used to explore possible conformational change mechanisms in a proton-dependent APC transporter, ApcT, a bacterial homologue from *Methanocaldococcus jannaschii* [Shaffer *et al.*, *Science* **325**, 1010 (2009)]. ApcT was captured in an inward-facing apo state. Gating is initiated by global counter-torsions of the intracellular and extracellular domains of ApcT around the pore axis, with the extracellular half rotating clockwise and the intracellular half anticlockwise, and vice versa. The domain motions are highly concerted and cooperative. The stationary plane relative to which counter-torsion occurs passes through the center of ApcT parallel to the membrane. Intracellularly, overall rotation of the peripheral helices (TM7, TM5, TM8, TM3, TM4, TM9, TM10, TM11 and TM12) reconfigures TM6a significantly and TM1b slightly. These helices alternately approach and separate from the opposed peripheral TM10 and TM11, affecting the intracellular mouth. TM6a and TM1b move toward the protein's perimeter and become buried inside the protein. Loops and small helices on the intracellular surface of ApcT undergo large-scale rotations. Extracellular motion is similar. Overall peripheral helix rotation affects TM1a significantly and TM6b slightly, displacing them from and collapsing them onto TM8 and TM5. TM8 and TM5 alternately undergo large-scale bending near their midpoints. Normal mode following along the lowest-frequency eigenvector(s) reveals details of the gating transition in the ApcT transporter.

### 3582-Pos

#### The Allosteric Role of Ion Binding in the Functional Mechanisms of Transporters With LeuT Fold

Lei Shi, Harel Weinstein.

Weill Cornell Medical College, New York, NY, USA.

Recent crystallographic studies revealed that five transporter families without much sequence similarities among them have similar structure folds to LeuT, a bacterial neurotransmitter:sodium symporter (NSS). The LeuT fold is characterized by an internal two-fold structural pseudosymmetry. Interestingly, the transport cycle of at least some members of each of these families is dependent on a sodium gradient across the membrane. Remarkably, the role of sodium is mimicked by a proton in others. We report our computational findings focusing on the LeuT conformations with various combinations of bound substrate and ions, performed in the context of on-going collaborative studies utilizing electron paramagnetic resonance (EPR) spectroscopy and single-molecule fluorescence (smFRET) to identify dynamic details of the mechanism. The resulting mechanistic implications from the study of LeuT are generalized to two other transporter families, the sodium:solute symporter (SSS) and amino acid-polyamine-organocation (APC) transporter, using comparative molecular dynamics simulations. These comparative studies lead to the proposal of a set of common structure-function-dynamic elements recognizable in the conformational transitions of the transporters with LeuT-fold.

## Membrane Domains & Lipid Dynamics

### 3583-Pos

#### Neurologin-1 Oligomerization Induces Cell Morphology Changes Via Lipid Domain Nucleation

Nicole Mende, Sophie Pautot.

CRTD, Dresden, Germany.

Thus far, neurologin-1 (Nlg-1) has been known as a post-synaptic adhesion-signaling membrane protein involved in initiating synaptic contact, and in triggering presynaptic differentiation in a neurexin-expressing axon. Here, we are reporting that nlg-1 might also play a role in neuron morphology changes. Indeed, when nlg-1 was coexpressed in HEK-293 with psd-95, a scaffolding protein which binds nlg-1 PDZ domain, we have observed extensive cell morphology changes. Co-transfected cells exhibited long expansions resembling dendritic branches, as well as a significant increase in cell surface area. However, nlg-1 dimerization mutant did not lead to any major changes in morphology suggesting that nlg-1 multimerization was required.