



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

**Cognitive and anatomical data in a healthy cohort of adults**

P.D. Watson<sup>a,\*</sup>, E.J. Paul<sup>a</sup>, G.E. Cooke<sup>a</sup>, N. Ward<sup>a</sup>, J.M. Monti<sup>a</sup>,  
 K.M. Horecka<sup>a,h</sup>, C.M. Allen<sup>a</sup>, C.H. Hillman<sup>a,b</sup>, N.J. Cohen<sup>a,c</sup>,  
 A.F. Kramer<sup>a,c</sup>, A.K. Barbey<sup>a,c,d,e,f,g,h,\*</sup>

<sup>a</sup> Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana–Champaign, Urbana, IL, USA

<sup>b</sup> Department of Kinesiology and Community Health, University of Illinois at Urbana–Champaign, Urbana, IL, USA

<sup>c</sup> Department of Psychology, University of Illinois at Urbana–Champaign, Champaign, IL, USA

<sup>d</sup> Decision Neuroscience Laboratory, University of Illinois at Urbana–Champaign, Champaign, IL, USA

<sup>e</sup> Department of Bioengineering, University of Illinois at Urbana–Champaign, Urbana, IL, USA

<sup>f</sup> Department of Internal Medicine, University of Illinois at Urbana–Champaign, Champaign, IL, USA

<sup>g</sup> Department of Speech and Hearing Science, University of Illinois at Urbana–Champaign, Champaign, IL, USA

<sup>h</sup> Neuroscience Program, University of Illinois at Urbana–Champaign, Champaign, IL, USA

## ARTICLE INFO

*Article history:*

Received 18 January 2016

Received in revised form

30 March 2016

Accepted 30 March 2016

Available online 5 April 2016

*Keywords:*

Independent component analysis

Fluid intelligence

Neuroanatomy

Tractography

Individual differences

## ABSTRACT

We present data from a sample of 190 healthy adults including assessments of 4 cognitive factor scores, 12 cognitive tests, and 115 MRI-assessed neuroanatomical variables (cortical thicknesses, cortical and sub-cortical volumes, fractional anisotropy, and radial diffusivity). These data were used in estimating underlying sources of individual variation via independent component analysis (Watson et al., In press) [25].

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.neuroimage.2016.01.023>

\* Corresponding authors at: Decision Neuroscience Laboratory, Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana–Champaign, 405 North Mathews Avenue, Urbana, IL 61801, USA.

E-mail address: [pwatson1@illinois.edu](mailto:pwatson1@illinois.edu) (P.D. Watson).

URL: <http://DecisionNeuroscienceLab.org/> (P.D. Watson).

<http://dx.doi.org/10.1016/j.dib.2016.03.100>

2352–3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Specifications Table

Subject area	<i>Neuroscience</i>
More specific subject area	<i>Anatomical Neuroimaging</i>
Type of data	<i>Table of cognitive testing data and MRI assessed structural data.</i>
How data was acquired	<i>Cognitive testing, Freesurfer automated segmentation of T1 weighted 3D MPRAGE images on a Siemens Magnetom Trio 3T whole-body MRI</i>
Data format	<i>Analyzed</i>
Experimental factors	<i>Brief description of any pretreatment of samples</i>
Experimental features	<i>Multi-modal MRI collection prior to a large cognitive training intervention.</i>
Data source location	<i>Urbana, Illinois</i>
Data accessibility	<i>Public repository: Open Science framework INSIGHT project: <a href="https://osf.io/9ezwc/">https://osf.io/9ezwc/</a></i>

## Value of the data

- These data characterize individual variation across demographic, neuroanatomical, and cognitive factors.
- These provide a useful model of individual variation that can be used to control for individual differences.
- The relationship between these data and other neuroimaging (such as resting state) and cognitive data remains unexplored and would be a fruitful area of collaboration.
- These data can be used to estimate patterns of joint variance across and within different neuroimaging and behavioral methods.
- These patterns can be used to test specific cognitive–anatomical linkages.

## 1. Data

The data ([Supplementary Table 1](#)) includes cognitive and anatomical variables collected prior to a large, multi-modal cognitive training study [25]. They include:

- Demographic measures (i.e., age, sex, and education).
- Cardiovascular fitness measures.
- 4 cognitive factors estimated via structural equation modeling [15].
- Scores from the battery of 12 cognitive tests used to estimate these factors.
- 35 cortical thickness estimates and volume estimates for these same regions.
- 11 sub-cortical volumetric estimates.
- Total brain and total intracranial volume estimates.
- 7 estimates of ventricular size.
- 5 estimates of corpus callosum.
- 12 estimates of fractional anisotropy and in matter tracts.
- 12 estimates of radial diffusivity in white matter tracts.

## 2. Experimental design, materials and methods

### 2.1. Demographics

The 190 participants consisted of 85 females, and 105 males. The age range in our sample was 18–44 years, with a median of 22 years, and a mean of 24.3 years. The mean educational level of the

**Table 1**  
Included measures.

Data categories	Specific measures
Demographics & cardiovascular fitness	Age Years of education Sex VO <sub>2max</sub> percentile
Cognition	Fluid intelligence (fluid g) Working memory (wm) Executive function (ef) Episodic memory (em) BOMAT (correct trials) Number series (correct trials) Letter Sets (correct trials) Reading span Rotation span Symmetry span Garavan (inverse total errors) Keep Track Words Recalled Stroop (inverse cost) Immediate free recall Words Immediate free recall Pictures Immediate free recall Paired Associates
Cortical thicknesses	Superior parietal Postcentral Precuneus Lateral occipital Mean cortical thickness Superior temporal Inferior parietal Paracentral Precentral Middle temporal Banks of superior temporal sulcus Insula Superior frontal Supramarginal Transverse temporal Rostral middle frontal Caudal middle frontal Pars triangularis Pars opercularis Lateral orbitofrontal Pars orbitalis Frontal pole Posterior cingulate Inferior temporal Cuneus Peri calcarine Rostral anterior cingulate Medial orbitofrontal Caudal anterior cingulate Isthmus cingulate Fusiform Temporal pole Lingual Entorhinal Parahippocampal
Cortical volumes	Middle temporal Inferior parietal Inferior temporal Rostral anterior cingulate

**Table 1** (continued)

Data categories	Specific measures
Sub-cortical volumes	Posterior cingulate
	Rostral middle frontal
	Superior frontal
	Precentral
	Supra marginal
	Lateral orbitofrontal
	Fusiform
	Precuneus
	Insula
	Medial orbitofrontal
	Postcentral
	Superior temporal
	Caudal middle frontal
	Paracentral
	Superior parietal
	Isthmus cingulate
	Lateral occipital
	Transverse temporal
	Pars orbitalis
	Pars opercularis
	Caudal anterior cingulate
	Pars triangularis
	Entorhinal
	Temporal pole
	Parahippocampal
	Frontal pole
	Peri calcarine
	Cuneus
	Lingual
	Total Brain volume
	Total Intracranial Volume
	Hippocampus
	Ventral Diencephalon
	Cerebellum Cortex
	Cerebellum White Matter
	Thalamus
	Brain Stem
	Amygdala
	Putamen
	Accumbens area
	Pallidum
	Caudate
Ventricles	Surface Holes
	Lateral Ventricle
	Choroid plexus
	Third Ventricle
	Cerebrospinal fluid
	Inferior Lateral Ventricle
Corpus callosum	Fourth Ventricle
	CC Posterior
	CC Mid Posterior
	CC Central
	CC Mid Anterior
	CC Anterior
White matter tractography (Fractional Anisotropy)	Inferior fronto-occipital fasciculus
	Superior longitudinal fasciculus
	Temporal superior longitudinal fasciculus
	Inferior longitudinal fasciculus
	Anterior thalamic radiation
	Forceps minor Uncinate fasciculus

**Table 1** (continued)

Data categories	Specific measures
White matter tractography (Radial Diffusivity)	Cingulum bundle
	Corticospinal tract
	Forceps major
	Hippocampal cingulum bundle
	Inferior fronto-occipital fasciculus
	Superior longitudinal fasciculus
	Temporal superior longitudinal fasciculus
	Inferior longitudinal fasciculus
	Anterior thalamic radiation
	Forceps minor
	Uncinate fasciculus
	Cingulum bundle
	Corticospinal tract
Forceps major	
Hippocampal cingulum bundle	

participants was “some college” (i.e., median score 3, mean score 3.6) as reported on a scale from 1 to 5, where 1 denoted “less than a high school diploma”, 2 denoted “high school diploma or equivalent”, 3 denoted “some college”, 4 denoted “college degree”, and 5 denoted “post-graduate education.”

## 2.2. Aerobic fitness assessment

Maximal oxygen consumption ( $VO_{2max}$ ) was measured using a computerized indirect calorimetry system (ParvoMedics True Max 2400) and a modified Balke protocol [1] with averages for oxygen uptake ( $VO_2$ ) and respiratory exchange ratio (RER) assessed every 20 s. Participants ran on a motor-driven treadmill at a constant speed, with 2.0% increases in grade every two minutes until volitional exhaustion. The raw value was adjusted for body size, age, and gender to produce a  $VO_{2max}$  percentile score.

## 2.3. Cognitive tests and factor scores

Participants received a battery of 12 cognitive tests designed to estimate underlying latent variables corresponding to cognitive constructs (see Table 1). The four latent variables of interest were fluid intelligence (gf), working memory (wm), executive function (ef), and episodic memory (em). Each of these latent variables was measured with three cognitive tests as follows. Fluid intelligence (gf) was measured by the BOMAT, number series, and letter sets tests [3,4,7]. Working memory (wm) was measured by the reading, rotation, and symmetry span tests [8,23]. Executive function (ef) was measured by the Garavan, Keep Track, and Stroop tests [14,22,26]. Episodic memory (em) was measured by immediate free recall, words, pictures and paired associates tests [23,24,9]. Using a structural equation modeling approach [15], across the larger sample of 518 participants, we extracted estimates of the four cognitive construct latent variables (i.e., gf, wm, ef, em). Because Garavan and Stroop produce error scores, while all others are measures of accuracy, we inverted these two values (i.e., multiplied by  $-1$ ) in order to ensure all cognitive variables had the same sign.

## 2.4. Structural MRI protocol

High resolution T1-weighted brain images were acquired using a 3D MPRAGE (Magnetization Prepared Rapid Gradient Echo Imaging) protocol with 192 contiguous axial slices, collected in ascending fashion parallel to the anterior and posterior commissures, echo time (TE)=2.32 ms, repetition time (TR)=1900 ms, field of view (FOV)=230 mm, acquisition matrix 256 mm  $\times$  256 mm,

slice thickness=0.90 mm, and flip angle=9°. All images were collected on a Siemens Magnetom Trio 3T whole-body MRI scanner.

### 2.5. Automated volumetrics, cortical thickness estimates, and white-matter tractography

Automated brain tissue segmentation and reconstruction of the T1-weighted structural MRI images were performed using the standard recon-all processing pipeline in FreeSurfer, version 5.2.0 (Released May, 2013; <http://surfer-nmr.mgh.harvard.edu/>). This produced estimates of 1) cortical thickness, 2) cortical volumes, 3) sub-cortical volumes, 4) ventricles, and 5) corpus callosum [5,6,10–13]. Segmentations and tractography were manually checked for errors. Estimates in the left and right hemispheres were summed to produce bilateral estimates, and all values were converted to z-scores to control for differences in scale. A complete list of estimated structures appears in Table 1. FreeSurfer produced automated segmentation that closely approximates hand tracing, but like all segmentation procedures may introduce systematic bias.

The diffusion tensor imaging estimates for fractional anisotropy (FA) and radial diffusivity (RD) data was analyzed using tract-based spatial statistics in FSL [19–21]. This pipeline involves fitting a tensor model to the raw diffusion data using fMRIB's diffusion toolbox, and non-brain tissues were removed using FSL's brain extraction tool. All subjects' FA data were then aligned into a common space using the nonlinear registration tool FNIRT [18,2]. Next, the mean FA image was created and thinned to create a mean FA skeleton that represents the centers of all tracts common to the group. Each subject's aligned FA data was then projected onto this skeleton to create an estimate of the subject-level value associated with each tract.

### Acknowledgments

Research reported in this publication was supported by the Intelligence Advanced Research Projects Activity (IARPA), Contract no. 2014-13121700004 to the University of Illinois at Urbana-Champaign. The content is solely the responsibility of the authors and does not necessarily represent the views of IARPA.

University of Illinois at Urbana-Champaign, Institutional Review Board study approval number 14212.

### Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.03.100>.

### References

- [1] American College of Sports Medicine, ACSM's Guidelines for Exercise Testing and Prescription, 9th ed., Wolters Kluwer/Lippincott Williams & Wilkins, 2014.
- [2] J.L.R. Andersson, M. Jenkinson, S. Smith. Non-linear registration aka Spatial normalisation FMRIB Technical Report TR07JA2. In Practice. Retrieved from (<http://fmrib.medsci.ox.ac.uk/analysis/techrep/tr07ja2/tr07ja2.pdf>), 2007.
- [3] R.G. Bernreuter, C.H. Goodman, A study of the Thurstone primary mental abilities tests applied to freshman engineering students, *J. Educ. Psychol.* 32 (1) (1941) 55–60.
- [4] Bocumer Matrizentest, BOMAT-Advanced-Short Version, Hogrefe, Göttingen, 2009.
- [5] A.M. Dale, B. Fischl, M.I. Sereno, Cortical surface-based analysis. I. Segmentation and surface reconstruction, *NeuroImage* 9 (2) (1999) 179–194. <http://dx.doi.org/10.1006/nimg.1998.0395>.
- [6] R.S. Desikan, F. Ségonne, B. Fischl, B.T. Quinn, B.C. Dickerson, D. Blacker, R.J. Killiany, An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest, *NeuroImage* 31 (3) (2006) 968–980. <http://dx.doi.org/10.1016/j.neuroimage.2006.01.021>.
- [7] R. Ekstrom, J. French, H. Harman, D. Dermen, Manual for Kit of Factor-Referenced Cognitive Tests, Educational Testing Service, Princeton, NJ (1976) 117. <http://dx.doi.org/10.1073/pnas.0506897102>.
- [8] R.W. Engle, S.W. Tuholski, J.E. Laughlin, A.R.A. Conway, Working memory, short-term memory, and general fluid intelligence: a latent-variable approach, *J. Exp. Psychol. Gen.* (1999).

- [9] R.W. Engle, S.W. Tuholski, J.E. Laughlin, A.R.A. Conway, Working memory, short-term memory, and general fluid intelligence: a latent-variable approach, *J. Exp. Psychol. Gen.* 128 (3) (1999) 309–331.
- [10] B. Fischl, A.M. Dale, Measuring the thickness of the human cerebral cortex from magnetic resonance images, *Proc. Natl. Acad. Sci. USA* 97 (20) (2000) 11050–11055. <http://dx.doi.org/10.1073/pnas.200033797>.
- [11] B. Fischl, D.H. Salat, E. Busa, M. Albert, M. Dieterich, C. Haselgrove, A.M. Dale, Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain, *Neuron* 33 (3) (2002) 341–355. [http://dx.doi.org/10.1016/S0896-6273\(02\)00569-X](http://dx.doi.org/10.1016/S0896-6273(02)00569-X).
- [12] B. Fischl, D.H. Salat, A.J.W. Van Der Kouwe, N. Makris, F. Ségonne, B.T. Quinn, A.M. Dale, Sequence-independent segmentation of magnetic resonance images, *NeuroImage* (2004). <http://dx.doi.org/10.1016/j.neuroimage.2004.07.016>.
- [13] B. Fischl, A. Van Der Kouwe, C. Destrieux, E. Halgren, F. Ségonne, D.H. Salat, A.M. Dale, Automatically parcellating the human cerebral cortex, *Cereb. Cortex* 14 (1) (2004) 11–22. <http://dx.doi.org/10.1093/cercor/bhg087>.
- [14] H. Garavan, Serial attention within working memory, *Mem. Cognit.* 26 (2) (1998) 263–276. <http://dx.doi.org/10.3758/BF03201138>.
- [15] M.J. Kane, D.Z. Hambrick, S.W. Tuholski, O. Wilhelm, T.W. Payne, R.W. Engle, The generality of working memory capacity: a latent-variable approach to verbal and visuospatial memory span and reasoning, *J. Exp. Psychol. Gen.* 133 (2) (2004) 189–217. <http://dx.doi.org/10.1037/0096-3445.133.2.189>.
- [18] D. Rueckert, L.I. Sonoda, C. Hayes, D.L. Hill, M.O. Leach, D.J. Hawkes, Nonrigid registration using free-form deformations: application to breast MR images, *IEEE Trans. Med. Imaging* 18 (8) (1999) 712–721. <http://dx.doi.org/10.1109/42.796284>.
- [19] S.M. Smith, Fast robust automated brain extraction, *Hum. Brain Mapp.* 17 (3) (2002) 143–155. <http://dx.doi.org/10.1002/hbm.10062>.
- [20] S.M. Smith, M. Jenkinson, H. Johansen-Berg, D. Rueckert, T.E. Nichols, C.E. Mackay, T.E.J. Behrens, Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data, *NeuroImage* 31 (4) (2006) 1487–1505. <http://dx.doi.org/10.1016/j.neuroimage.2006.02.024>.
- [21] S.M. Smith, M. Jenkinson, M.W. Woolrich, C.F. Beckmann, T.E.J. Behrens, H. Johansen-Berg, P.M. Matthews, Advances in functional and structural MR image analysis and implementation as FSL, *NeuroImage* (2004). <http://dx.doi.org/10.1016/j.neuroimage.2004.07.051>.
- [22] J.R. Stroop, Studies of interference in serial verbal reactions, *J. Exp. Psychol.* (1935). <http://dx.doi.org/10.1037/h0054651>.
- [23] N. Unsworth, T.S. Redick, R.P. Heitz, J.M. Broadway, R.W. Engle, Complex working memory span tasks and higher-order cognition: a latent-variable analysis of the relationship between processing and storage, *Memory* 17 (6) (2009) 635–654. <http://dx.doi.org/10.1080/09658210902998047>.
- [24] B. Uttl, P. Graf, L.K. Richter, Verbal paired associates tests limits on validity and reliability, *Arch. Clin. Neuropsychol.* 17 (6) (2002) 567–581. [http://dx.doi.org/10.1016/S0887-6177\(01\)00135-4](http://dx.doi.org/10.1016/S0887-6177(01)00135-4).
- [25] P.D. Watson, E.J. Paul, G.E. Cooke, N. Ward, J.M. Monti, K.M. Horecka, C.M. Allen, C.H. Hillman, N.J. Cohen, A.F. Kramer, A. K. Barbey, Underlying sources of cognitive-anatomical variation in multi-modal neuroimaging and cognitive testing, *NeuroImage* 129 (2016) 439–449. <http://dx.doi.org/10.1016/j.neuroimage.2016.01.023>.
- [26] D.B. Yntema, Keeping track of several things at once, *Hum. Factors: J. Hum. Factors Ergon. Soc.* 5 (1) (1963) 7–17. <http://dx.doi.org/10.1177/001872086300500102>.