

# Depletion of Ascorbic Acid Restricts Angiogenesis and Retards Tumor Growth in a Mouse Model

Sucheta Telang, Amy L. Clem, John W. Eaton and Jason Chesney

Molecular Targets Program, Medical Oncology, James Graham Brown Cancer Center, University of Louisville, Louisville, KY, USA

## Abstract

**Angiogenesis requires the deposition of type IV collagen by endothelial cells into the basement membrane of new blood vessels. Stabilization of type IV collagen triple helix depends on the hydroxylation of proline, which is catalyzed by the iron-containing enzyme prolyl hydroxylase. This enzyme, in turn, requires ascorbic acid to maintain the enzyme-bound iron in its reduced state. We hypothesized that dietary ascorbic acid might be required for tumor angiogenesis and, therefore, tumor growth. Here, we show that, not surprisingly, ascorbic acid is necessary for the synthesis of collagen type IV by human endothelial cells and for their effective migration and tube formation on a basement membrane matrix. Furthermore, ascorbic acid depletion in mice incapable of synthesizing ascorbic acid (*Gulo*<sup>-/-</sup>) dramatically restricts the *in vivo* growth of implanted Lewis lung carcinoma tumors. Histopathological analyses of these tumors reveal poorly formed blood vessels, extensive hemorrhagic foci, and decreased collagen and von Willebrand factor expression. Our data indicate that ascorbic acid plays an essential role in tumor angiogenesis and growth, and that restriction of ascorbic acid or pharmacological inhibition of prolyl hydroxylase may prove to be novel therapeutic approaches to the treatment of cancer.**

*Neoplasia* (2007) 9, 47–56

**Keywords:** Ascorbic acid, tumor, angiogenesis, collagen, prolyl hydroxylase.

## Introduction

Ascorbic acid (vitamin C) is an essential dietary requirement in humans and other primates due to the genetic absence of a key enzyme in ascorbic acid synthesis, L-gulonolactone oxidase [1,2]. Humans deficient in ascorbic acid develop scurvy—a disease that came to prominence in the era of long sea voyages and was first systematically described by Lind [3] in 1753. Bleeding gums, hemorrhage, and ecchymoses, which are the major manifestations of this deficiency, are due to impaired collagen deposition leading to defective blood vessel formation and repair [4].

The clinical manifestations of scurvy reflect the crucial importance of ascorbic acid in angiogenesis and blood

vessel repair. Angiogenesis, in particular, is a prerequisite for tumor growth *in vivo*. Tumors are unable to grow beyond ~1 to 2 mm<sup>3</sup> without the formation of new blood vessels, which will deliver metabolic substrates and oxygen [5]. Effective angiogenesis requires the deposition of mature type IV collagen in the basement membrane of blood vessels by endothelial cells [6]. Type IV collagen folding, in turn, is dependent on the hydroxylation of proline by the iron-dependent enzyme prolyl hydroxylase, which requires ascorbic acid to reduce iron to its ferrous state for the reactivation of the enzyme [7]. These considerations provide general support for the idea that depletion of ascorbic acid might limit tumor growth by impairing blood vessel formation and repair.

However, there are reasons to suppose that ascorbic acid deficiency might have the opposite effect on tumor growth. For example, ascorbic acid is also important in maintaining normal immune function and in augmenting host immunity against cancer [8–10]. Although early studies attributed ascorbic acid-dependent augmentation of tumor immunity to improvement in cytolytic functions of lymphocytes and phagocytes, the precise mechanisms have not been clearly defined [11,12]. However, in the particular case of lymphocyte function, more recent data suggest that ascorbic acid may improve host immunity by inhibiting T-cell apoptosis [13].

Ascorbic acid deficiency may also affect the expression of hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) in such a way as to promote tumor growth. Neoplastic cells exposed to low oxygen respond by stabilizing HIF-1 $\alpha$ , which acts to induce the transcription of several mRNA transcripts that are essential for tumor growth, including glycolytic enzymes and vascular endothelial growth factor (VEGF). Ascorbic acid is a cofactor for the proline hydroxylation and subsequent proteosomal degradation of HIF-1 $\alpha$ . Depletion of ascorbic acid reportedly interrupts this degradation, causing high HIF-1 $\alpha$  expression and upregulation of several protumorigenic transcriptional targets [14]. Accordingly, dietary ascorbic acid restriction might

Abbreviations: HIF-1 $\alpha$ , hypoxia-inducible factor-1 $\alpha$ ; HUVECs, human umbilical vein endothelial cells; VEGF, vascular endothelial growth factor; *Gulo*, L-gulonolactone oxidase gene. Address all correspondence to: Jason Chesney, Molecular Targets Group, Medical Oncology, James Graham Brown Cancer Center, University of Louisville, 580 South Preston Street, Louisville, KY 40202. E-mail: jason.chesney@gmail.com

Received 11 October 2006; Revised 30 November 2006; Accepted 30 November 2006.

Copyright © 2007 Neoplasia Press, Inc. All rights reserved 1522-8002/07/\$25.00  
DOI 10.1593/neo.06664

enhance tumor growth by allowing increased stabilization of HIF-1 $\alpha$  and/or by impairing tumor immunity.

Thus, it is not clear whether ascorbic acid deficiency might increase or decrease tumor growth. A major impediment to testing the importance of ascorbic acid in tumor growth has been that mice, the most facile model for such studies, are able to synthesize ascorbic acid and do not require dietary vitamin C. Recently, mice bearing a homozygous deletion of the L-gulonolactone oxidase gene (*Gulo*<sup>-/-</sup>) have been generated; like humans, they require ascorbic acid supplementation to avoid manifestations of vitamin C deficiency [15]. Plasma ascorbic acid concentrations fall to approximately 10  $\mu$ M ( $\sim 1.7 \pm 0.2 \mu$ g/ml) in *Gulo*<sup>-/-</sup> mice kept on restricted dietary ascorbic acid for 3 to 5 weeks and, after  $\sim 6$  weeks of restriction, these animals demonstrate weight loss and hemarthroses [15].

Using this new animal model, we examined the requirement of dietary ascorbic acid for the growth of highly angiogenic Lewis lung carcinoma. Our results indicate that tumor growth is greatly diminished in vitamin C-deficient *Gulo*<sup>-/-</sup> mice. Tumors grown in these mice also show defective angiogenesis and intratumoral hemorrhagic foci. Contrary to what might be expected, however, there are no significant effects on HIF-1 $\alpha$  expression by tumor cells. The marked suppression of tumor growth in these vitamin C-deficient mice may provide a rationale for clinical trials in humans to evaluate the effects of dietary ascorbic acid restriction on cancer progression.

## Materials and Methods

### Cell Lines and Culture Conditions

Human umbilical vein endothelial cells (HUVECs) were obtained from Cambrex (Walkersville, MD) and were grown in culture with basal medium (EGM; Cambrex) supplemented with bovine brain extract, human endothelial cell growth factor, hydrocortisone, gentamicin, amphotericin B, and 2% fetal bovine serum (FBS) (EGM BulletKit; Cambrex). HUVECs used for experiments were between six and eight passages. Lewis lung carcinoma cells were obtained from ATCC (Manassas, VA) and cultured in Dulbecco's modified Eagle's medium (Invitrogen, Carlsbad, CA) supplemented with 10% FBS (HyClone Laboratories, Logan, UT) and gentamicin (Invitrogen). All cell lines were grown at 37°C in 5% CO<sub>2</sub>. For some experiments, 0 to 200  $\mu$ M L-ascorbic acid (Sigma, St. Louis, MO) was used to supplement the growth medium. Lewis lung carcinoma cells were exposed to hypoxia by incubation for 16 hours in 1% oxygen balanced with nitrogen and 5% CO<sub>2</sub> in a modular incubator chamber (Billups-Rothenburg, Del Mar, CA).

### Collagen Detection by Enzyme-Linked Immunosorbent Assay (ELISA)

HUVECs were plated at a density of  $4 \times 10^3$  cells/well on 96-well plates (Costar Corning, Corning, NY) in EGM containing 0 to 200  $\mu$ M L-ascorbic acid (Sigma). For these ELISA determinations, human placental collagen type IV (Rockland

Immunochemicals, Gilbertsville, PA) was used as a positive control. After 1 to 10 days in culture, HUVECs were detached with 0.25% trypsin-EDTA (Invitrogen), and the plates were washed with Tris-buffered saline (Tris/NaCl; 50 mM Tris, 150 mM NaCl, pH 8.5). The wells were then blocked with 1% bovine serum albumin (Sigma) in phosphate-buffered saline (PBS) for 1 hour at 37°C and washed with Tris/NaCl. The plates were incubated with rabbit anti-human type IV collagen (1:4000; Rockland Immunochemicals) for 2 hours at 37°C, washed thrice with Tris/NaCl/0.05% Tween-20, and then incubated with horseradish peroxidase (HRP)-conjugated donkey anti-rabbit immunoglobulin (1:8000; Rockland Immunochemicals) for 1 hour. After three washes with Tris/NaCl/Tween-20, color development in the presence of 2,2'-azino-bis [3-ethylbenzthiazoline-6-sulfonic acid] (ABTS; Rockland Immunochemicals) was read with a spectrophotometer at 420 nm. The assay was performed in quadruplicate and was repeated twice.

### Endothelial Tube Formation Assay

Twenty-four-well plates (Costar Corning) were coated with Matrigel basement membrane matrix (250  $\mu$ l/well; BD Biosciences, Bedford, MA) according to the manufacturer's instructions and allowed to polymerize at 37°C for 60 minutes. HUVECs grown in culture were detached, resuspended in EGM, and seeded on Matrigel-coated wells ( $8 \times 10^4$  well<sup>-1</sup>) in 0 to 200  $\mu$ M L-ascorbic acid. The plates were incubated at 37°C in 5% CO<sub>2</sub> and examined for tube formation through an inverted photomicroscope every 6 hours for 18 hours. At 18 hours, intact tubes formed were enumerated in three random fields, and representative photomicrographs were taken (original magnification,  $\times 40$ ). The assay was performed in triplicate and repeated twice.

### In Vivo Experiments

All animal experiments were conducted with the approval of the Institutional Animal Care and Use Committee at the University of Louisville (Louisville, KY). *Gulo*<sup>+/-</sup> heterozygous breeding pairs (C57Bl6/J mice B6.129P2-*Gulo*<sup>tm1Unc/Ucd</sup>) were obtained from the Mutant Mouse Regional Resource Center, and homozygous (*Gulo*<sup>-/-</sup>) mice were bred. Genotyping was performed using tail biopsy specimens with the REDExtract-N-Amp Tissue PCR Kit (Sigma) and primers as previously described [15]. All mice used in the experiments were < 3 months of age.

Mice were weaned at 3 weeks of age onto irradiated mouse chow (PicoLab Mouse Diet 5058; PMI Feeds, St. Louis, MO) containing 110 mg/kg ascorbic acid and were maintained on this diet during all experiments. Because this amount is not sufficient to maintain a physiological concentration of vitamin C in *Gulo*<sup>-/-</sup> animals [15], ascorbic acid was supplemented in drinking water (330 mg/l L-ascorbic acid + 0.01 mM EDTA), which was changed weekly.

For initial outgrowth experiments, *Gulo*<sup>-/-</sup> mice were depleted of ascorbic acid by withholding supplementation in drinking water for 4 weeks immediately after weaning and then were randomized to continued depletion or repletion by supplementation with 330 mg/l ascorbic acid in drinking

water. A third group of *Gulo*<sup>-/-</sup> mice was maintained on full supplementation with 330 mg/l ascorbic acid before and after tumor implantation. Each group contained 10 animals.

For studying the growth of established tumors, the *Gulo*<sup>-/-</sup> mice used were maintained on 165 mg/l ascorbic acid supplementation in water for 2 weeks before starting the study. They were then randomized either to depletion (by withholding supplementation) or to continued supplementation at 165 mg/l. Each group contained 10 animals.

For experiments on minimal supplementation, *Gulo*<sup>-/-</sup> mice were depleted by withholding ascorbic acid supplementation for 4 weeks immediately after weaning and then were randomized to either continued depletion, 10% of full supplementation (33 mg/l ascorbic acid), or full supplementation (330 mg/l ascorbic acid). Each group contained 10 animals.

Lewis lung carcinoma cells were collected from exponential growth phase culture, washed twice, and then resuspended in PBS ( $1 \times 10^7$  cells/ml). Groups of *Gulo*<sup>-/-</sup> mice were injected subcutaneously in the right flank with 0.1 ml of cell suspension ( $1 \times 10^6$  cells). Tumor measurements were started with the appearance of established tumors measuring 150 mg (between 5 and 7 days after implantation). After the initial appearance, tumors were measured with microcalipers every 2 days, and tumor masses were calculated using the formula:  $[(\text{width in mm})^2 \times (\text{length in mm})] / 2$  [16]. When tumor volume had reached 10% of body weight, animals were euthanized using carbon dioxide narcosis and tumors were removed and fixed in 10% formaldehyde for further analyses.

All data are expressed as the mean  $\pm$  SD of three experiments. Statistical significance was assessed by two-sample *t*-test (independent variable).

#### Ascorbic Acid Measurements

Ascorbic acid concentrations in plasma samples were measured by the 2,2'-dipyridyl method [17].

#### Protein Extraction and Western Blot Analysis

Cells were treated with 0.25% trypsin-EDTA, washed in PBS, and lysed in  $2 \times$  RIPA buffer. Protein samples were separated by 12.5% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred to a PVDF membrane. Membranes were blocked in TBS-Tween-20 (1%) containing 5% milk. Mouse monoclonal anti-HIF-1 antibody (1:500; Novus Biologicals, Littleton, CO) and mouse anti- $\beta$ -actin (1:5000; Sigma) were resuspended in 10 ml of TBS-Tween-20 (5% milk) and incubated with the membrane for 1 hour. The secondary antibodies used were rabbit anti-mouse or goat anti-mouse HRP-conjugated (1:8000; Pierce Biotechnology, Rockford, IL).

#### Histology and Immunohistochemistry

Five-micrometer sections of formalin-fixed and paraffin-embedded tumor tissues were treated with xylene to remove paraffin and then rehydrated. Hematoxylin-eosin staining and Masson's trichrome staining for collagen were performed using standard procedures. For immunohistochemical staining, deparaffinized and rehydrated sections were blocked by incubation with serum blocking buffer for 30 minutes

at room temperature. Tissue sections were incubated for 1 hour with rabbit anti-mouse von Willebrand factor (vWF) antibody (1:500; Dakocytomation, Carpinteria, CA) or mouse monoclonal anti-HIF-1 $\alpha$  (1:250; Novus Biologicals) for the detection of vWF and HIF-1 $\alpha$ , respectively. The sections were then incubated with biotinylated goat anti-rabbit (for vWF) or rabbit anti-mouse IgG (for HIF-1 $\alpha$ ) for 30 minutes (1:500; Vector Laboratories, Burlingame, CA) and developed with an avidin-biotin peroxidase reaction using 3,3'-diaminobenzidine tetrahydrochloride as chromogen. After counterstaining with Mayer's hematoxylin (Sigma), the sections were dehydrated and coverslips were attached with Permount (Fisher Scientific, Pittsburgh, PA). Appropriate negative controls (by omission of the primary antibody) were used. Anti-HIF-1 $\alpha$  antibody functionality and specificity were confirmed by Western blot analysis using Lewis lung carcinoma cells exposed to hypoxia for 16 hours as positive control.

Quantitative immunohistochemistry imaging analyses were performed using TIFFalyzer on images acquired by a digital RGB camera (DXM 1200F; Nikon, Tokyo, Japan) and saved in TIFF format [18,19]. All data are expressed as the mean  $\pm$  SD of three experiments. Statistical significance was assessed by two-sample *t*-test (independent variable).


## Results

### Type IV Collagen Secretion By Endothelial Cells Requires Ascorbic Acid

To test whether type IV collagen secretion by endothelial cells requires ascorbic acid, we pulsed cultured HUVECs with 0 to 200  $\mu$ M ascorbic acid and then examined the plates for the presence of immunoreactive type IV collagen using an ultrasensitive ELISA. We found that HUVECs produced type IV collagen in the absence of added ascorbic acid, but that the addition of ascorbic acid increased type IV collagen secretion. The culture medium (EGM) contains approximately 5  $\mu$ M ascorbic acid, which may be sufficient to allow for the observed type IV collagen secretion in the absence of added vitamin C. The concentration of added ascorbic acid that maximized collagen type IV accumulation (25  $\mu$ M) is approximately 50% below the physiological concentration of ascorbic acid in healthy humans and mice (Figure 1A). Finally, we should note that the collagen detected by this ELISA was not necessarily stabilized type IV collagen triple helix.

### Endothelial Cell Migration and Tube Formation Require Ascorbic Acid

Deposition of mature collagen type IV is essential for endothelial cell migration and forms a scaffold for new blood vessel formation. The importance of collagen type IV in angiogenesis is further emphasized by evidence that inhibitors of proline hydroxylation impair angiogenesis *in vitro* [20]. Migration and tube formation on the basement membrane matrix mimic blood vessel formation, and we speculated that restriction of ascorbic acid would limit migration and



**Figure 1.** Immunoreactive type IV collagen and intact tube formation by endothelial cells require the presence of a physiological concentration of ascorbic acid. Type IV collagen production by HUVECs in 0 to 200  $\mu\text{M}$  ascorbic acid was measured by ELISA. (A) Representative photomicrographs of tube formation on Matrigel basement membrane matrix in 0 to 200  $\mu\text{M}$  ascorbic acid (original magnification,  $\times 40$ ). (B) Intact tubes formed on Matrigel in 0 to 200  $\mu\text{M}$  ascorbic acid were enumerated (original magnification,  $\times 40$ ). (C).

tube formation. We cultured HUVECs on Matrigel in EGM containing 0 to 200  $\mu\text{M}$  added ascorbic acid and examined the cells for tube formation by light microscopy (Figure 1B). Alignment of HUVECs into tube-like structures began within 2 to 4 hours and was completed by 18 hours. Very few intact tubes were formed in 0 to 12.5  $\mu\text{M}$  added ascorbic acid (despite the estimated 5- $\mu\text{M}$  background ascorbic acid in the medium). As the concentration of ascorbic acid was increased, greater numbers of intact tubes were observed, reaching a maximum at 50 to 100  $\mu\text{M}$  ( $23 \pm 1.5$  and  $27.6 \pm 2$  tubes, respectively, per  $\times 40$  high-power field). Interestingly and for unknown reasons, we observed decreased numbers of intact tubes at supraphysiologic concentrations of ascorbic acid (Figure 1C).

#### Restriction of Dietary Ascorbic Acid Depletes Plasma Ascorbic Acid in *Gulo*<sup>-/-</sup> Mice

Restriction of vitamin C in the diet of *Gulo*<sup>-/-</sup> mice has previously been demonstrated to deplete plasma ascorbic acid [15]. We sought to confirm this finding in our animals before embarking on experiments with tumor growth. We measured plasma ascorbic acid concentrations in *Gulo*<sup>-/-</sup> mice before limiting dietary ascorbic acid and at weekly intervals after starting restriction until the animals were euthanized. We found a gradual decrease in the concentration of plasma ascorbic acid over the 4 weeks of restriction from a baseline concentration of  $7.35 \pm 0.6 \mu\text{g/ml}$  (approximately 42  $\mu\text{M}$ ) to  $0.73 \pm 0.1 \mu\text{g/ml}$  by 4 weeks when the animals were fed only a diet containing 110 mg/kg ascorbic acid without further supplementation. We observed only a small further decline in the concentration of plasma ascorbic acid in restricted *Gulo*<sup>-/-</sup> mice over the course of tumor growth ( $0.47 \pm 0.12 \mu\text{g/ml}$  by 6 weeks) (Figure 2A).

#### Restriction of Dietary Ascorbic Acid Suppresses Tumor Growth in *Gulo*<sup>-/-</sup> Mice In Vivo

Our findings that endothelial cell migration and tube formation require the presence of physiological concentrations of ascorbic acid supported the generally accepted idea that vitamin C is required for effective blood vessel formation *in vivo*. To determine whether suppression of angiogenesis by a scorbutic state would limit tumor growth, we used the highly angiogenic tumor Lewis lung carcinoma in *Gulo*<sup>-/-</sup> mice. Before starting our experiments, we first sought to examine the survival and proliferation of Lewis lung carcinoma cells *in vitro* in the absence of added ascorbic acid in the medium. We found that the cells survived and demonstrated no significant differences in growth from cells cultured in the presence of added ascorbic acid (data not shown).

To study the effect of ascorbic acid restriction on initial tumor outgrowth, animals were depleted of ascorbate for 4 weeks and then implanted subcutaneously with Lewis lung carcinoma cells. After implantation, depleted animals were either: 1) continued on a depleted diet (D), or 2) made replete with ascorbic acid (D/R). We also included a group of *Gulo*<sup>-/-</sup> animals maintained on full supplementation before and after tumor implantation (R) ( $n = 10$  in all groups). We followed tumor volumes from appearance until the point at which tumor burden necessitated euthanization. At 12 days of growth after tumor appearance, tumors in the depleted animals were significantly smaller than either in depleted/replete groups or in fully supplemented groups (D,  $332.4 \pm 67.3 \text{ mm}^3$ ; D/R,  $2992.6 \pm 248 \text{ mm}^3$ ; R,  $1703 \pm 316.4 \text{ mm}^3$ ;  $P < .005$  in D versus either D/R or R at 12 days;  $n = 10$  for all groups) (Figure 2B). Interestingly, tumors in the depleted/replete mice were significantly larger than those in animals maintained on ascorbic acid before and after tumor implantation.

We then examined the effects of ascorbic acid depletion on the growth of previously established tumors. For this, we implanted Lewis lung carcinoma cells in *Gulo*<sup>-/-</sup> animals maintained on 165 mg/l ascorbic acid in drinking water for 2 weeks before the start of the study. When tumors were


palpable, we randomized animals to ascorbic acid depletion (D) or continued ascorbic acid supplementation of 165 mg/l (R) ( $n = 10$  in both groups). At 12 days of growth, we found that tumors in the depleted group were significantly smaller than those in the replete group (D,  $744.3 \pm 198.4 \text{ mm}^3$ ; R,  $1934.7 \pm 384 \text{ mm}^3$ ;  $P < .005$ ;  $n = 10$ ) (Figure 2C).

As observed in previous studies, depletion of ascorbic acid beyond approximately 6 weeks in homozygous *Gulo*<sup>-/-</sup> mice led to weight loss and decreased activity with the onset of hemarthroses and petechial bleeding. We therefore sought to test whether minimal supplementation of ascorbic acid would delay the onset of scorbutic symptoms while still retarding tumor growth. Following depletion of ascorbic acid for 4 weeks in *Gulo*<sup>-/-</sup> mice, we randomized animals to either continued depletion (D), 10% of the full supplementation dose (33 mg/l ascorbic acid) (10%), or full supplementation (330 mg/l ascorbic acid) (R), and implanted Lewis lung carcinoma cells subcutaneously in all the ani-


mals. We followed tumor growth from the time of appearance up to 10 days of growth and found that tumors in the D group and in the 10% supplemented group were both significantly smaller than those in the fully supplemented group (D,  $800 \pm 92.9 \text{ mm}^3$ ; 10% of full supplementation,  $776 \pm 270 \text{ mm}^3$ ; R,  $2529 \pm 562 \text{ mm}^3$ ;  $P < .05$  for D and 10% of full supplementation *versus* R;  $n = 10$ ) (Figure 2D). Importantly, the mice on the 10% ascorbic acid diet did not exhibit weight loss or bleeding (data not shown).

#### Tumors from Animals on Low-Ascorbic-Acid Diets Are Hemorrhagic and Show Decreased Capillary Density

We speculated that the differences we observed in tumor size between ascorbic acid-restricted and supplemented animals were due to inadequate delivery of essential nutrients as a result of poor blood vessel formation. To examine this, we first stained sections of representative tumors from depleted and replete mice with Masson's trichrome to



**Figure 2.** Tumor growth is significantly reduced in *Gulo*<sup>-/-</sup> mice on restricted ascorbic acid. Plasma ascorbic acid concentrations in *Gulo*<sup>-/-</sup> mice were measured before and at weekly intervals following depletion of ascorbic acid and during tumor growth for 6 weeks ( $n = 7$  in all groups). (A) Lewis lung carcinoma cells were injected subcutaneously into *Gulo*<sup>-/-</sup> mice depleted of ascorbic acid for 4 weeks following which animals were either continued on a depleted diet (D) or a replete diet (R). A control group of *Gulo*<sup>-/-</sup> mice was fully supplemented with ascorbic acid before and after injection with Lewis lung carcinoma cells (R). Tumors were measured with microcalipers on indicated days ( $n = 10$  per group). (B) *Gulo*<sup>-/-</sup> mice implanted with Lewis lung carcinoma cells were randomized to ascorbic acid depletion (D) or full repletion (R) when tumors were visible, and tumors were measured with microcalipers on indicated days ( $n = 10$  per group). (C) *Gulo*<sup>-/-</sup> mice depleted of ascorbic acid were implanted with Lewis lung carcinoma cells and, when tumors were palpable, were randomized to depletion (D), 10% repletion (10%), or 100% (R) repletion, and tumors were followed for 10 days ( $n = 10$  per group) (D).



**Figure 3.** Tumors in ascorbic acid–restricted *Gulo*<sup>-/-</sup> mice exhibit poorly formed vasculature with multiple hemorrhagic foci. Tissue sections from representative tumors in replete (A, C, E, and G) and depleted (B, D, F, and H) *Gulo*<sup>-/-</sup> mice were examined for vasculature after staining with Masson's trichrome (MT) stain (A and C), anti-vWF polyclonal antibody (vWF; E and F), or preimmune rabbit serum (ISO; G and H). Endothelial cells stained with  $\alpha$ -vWF are in brown and shown by arrows (A and B: original magnification,  $\times 200$ ; C–H: original magnification,  $\times 400$ ).

determine the character of the tumor vasculature. The vasculature in tumors excised from the depleted/replete (D/R) group was well defined (Figure 3, A and C). In contrast, tumors in depleted animals showed multiple areas of hemorrhage and poorly formed blood vessels (Figure 3, B and D). Furthermore, the absolute number of blood vessels was markedly reduced in depleted tumors in comparison with that in tumors from

fully supplemented animals (blood vessels per  $\times 100$  high-power field: D,  $5 \pm 2$ ; D/R,  $18.6 \pm 3$ ; R,  $25 \pm 5$ ;  $P < .005$ ).

To further evaluate blood vessel development and to enumerate capillaries, we stained tumors from depleted and replete animals with an antibody specific for vWF—a marker of mature endothelial cells [21,22]. We observed increased vWF staining indicative of endothelial cells in

tumor sections from the depleted/replete group of mice (Figure 3E) in comparison to minimal staining in depleted tumors (Figure 3F).

#### Ascorbic Acid Restriction Limits Collagen Synthesis in Tumors

Because ascorbic acid is required for the action of prolyl hydroxylase and, hence, collagen stabilization, we expected that collagen staining in tumors excised from mice on restricted ascorbic acid would be decreased. We stained representative tissue sections from all groups with Masson's trichrome and found decreased collagen staining in the fibrotic sheath of the depleted group in comparison with that in the replete group (Figure 4).

#### Ascorbic Acid Restriction Does Not Affect HIF-1 $\alpha$ Protein Expression


As mentioned in the Introduction, ascorbic acid acts as a cofactor in HIF proline hydroxylation by prolyl hydroxylase, which promotes proteosomal destruction of HIF. *In vitro* studies have demonstrated that physiological concentrations of ascorbic acid suppress HIF-1 $\alpha$  levels in transformed cells [14]. Extrapolating from these results, it would be possible that lack of HIF hydroxylation (secondary to ascorbic acid deficiency) might slow its proteosomal destruction and increase its intracellular levels. However, staining of tissue sections of tumors with an anti-HIF-1 $\alpha$  antibody did not show differences in HIF-1 $\alpha$  expression. The function and specificity of this antibody were confirmed by a band at 120 kDa by Western blot analysis in Lewis lung carcinoma cells exposed to hypoxia (data not shown). In addition, quantitative immunohistochemical analyses of tumor sec-

tions in depleted and replete groups, in comparison with isotype control sections, did not reveal a significant difference in HIF-1 $\alpha$  expression (HIF-1 $\alpha$  in replete tumors,  $490 \pm 118.0$  vs  $487 \pm 28.3$  energy units/pixel in ascorbic acid-depleted tumors). Therefore, at least in the *in vivo* environment, ascorbic acid depletion may selectively suppress tumor blood vessel formation/stability without affecting HIF levels (Figure 5).


#### Discussion

The importance of vitamin C in carcinogenesis and tumor growth has been the subject of controversy for years. Early observational clinical studies demonstrated a 10% regression rate [23] and up to a 20-fold increase in survival in patients with advanced cancer treated with high-dose ascorbic acid (10 g/day intravenously for 10 days, followed by oral administration of the same dose) in comparison to controls retrospectively analyzed from a population of similar terminal cancer patients [24,25]. The results of these studies were widely disputed, in part due to lack of randomization. Two subsequent randomized placebo-controlled double-blinded trials using oral vitamin C [26,27] demonstrated no improvement in survival in patients treated with high doses of ascorbic acid. Perhaps as a result, the use of megadoses of vitamin C as a possible therapy for cancer fell by the wayside and has not been revisited until recently [28,29].

The present work was based on precisely the opposite hypothesis: that restriction of the availability of ascorbic acid would suppress tumor growth, perhaps through diminishing angiogenesis. Ascorbic acid plays an essential role in angiogenesis as a cofactor for prolyl hydroxylase in catalyzing hydroxyproline synthesis [7]. Hydroxyproline is required for



**Figure 4.** Collagen synthesis is decreased in subcutaneous tumors formed in *Gulo*<sup>-/-</sup> mice on an ascorbic acid-restricted diet. Sections from representative tumors harvested from replete (A and C) and depleted (B and D) *Gulo*<sup>-/-</sup> mice were evaluated for collagen after staining with Masson's trichrome. Collagen appears as blue-stained fibrillar material indicated by arrows (A and B: original magnification,  $\times 200$ ; C and D: original magnification,  $\times 400$ ).



**Figure 5.** HIF-1 $\alpha$  protein expression is similar in tumors from replete *Gulo*<sup>-/-</sup> animals and tumors from depleted *Gulo*<sup>-/-</sup> animals. Tissue sections from representative tumors in replete (A and C) and depleted (B and D) *Gulo*<sup>-/-</sup> mice were examined by immunohistochemistry using an anti-HIF-1 $\alpha$  mouse monoclonal antibody (A and B) or preimmune mouse serum (C and D) (original magnification,  $\times 400$ ). Arrows indicate representative areas of HIF-1 $\alpha$  expression.

effective type IV collagen production by endothelial cells. Collagen type IV forms a critical part of the vascular basement membrane and is vital for new blood vessel formation.

Tumor growth and metastasis are dependent on angiogenesis. One early study demonstrated that tumor growth in the avascular anterior chamber of the rabbit eye was arrested at a small size ( $\sim 1 \text{ mm}^3$ ) due to the absence of neovascularization, whereas tumors grew readily on the surface of the iris where new vasculature could develop [30]. Tumors secrete angiogenic growth factors such as VEGF, which promote endothelial cell proliferation and new blood vessel development [31]. That anti-VEGF antibodies and a dominant interfering form of the VEGF receptor 2 (flk-1) have proven to be successful in impairing neovascularization and in causing tumor regression demonstrates the importance of angiogenesis in tumor growth [32,33].

An early study on the importance of ascorbic acid in angiogenesis was carried out in guinea pigs, which, like primates and *Gulo*<sup>-/-</sup> mice, are incapable of vitamin C synthesis. In subcutaneously implanted polyvinyl sponges (which promote angiogenic activity), investigators found dilated blood vessels, interstitial hemorrhage, and disintegration of connective tissues in scorbutic guinea pigs [34,35]. A later investigation demonstrated decreased type IV collagen mRNA in blood vessels in the presence of restricted ascorbic acid [35]. In addition, inhibitors of prolyl hydroxylase (e.g., ciclopirox and 2,2'-dipyridyl) cause inhibition of both tube formation on Matrigel and aortic arch sprouting *in vitro* [20]. Similarly, proline analogs such as *cis*-hydroxyproline, which impair  $\alpha$  helix formation and collagen secretion, lead to the formation of large and dilated blood vessels in a plasma clot culture model [36] and inhibit angiogenesis in chick chorioallantoic membranes [37]. These earlier observations are in accord with our *in vitro* data from cultured HUVECs showing

a requirement for ascorbic acid for both type IV collagen synthesis and endothelial tube formation on a basement membrane matrix. Our data indicate that although tube formation is maximal at concentrations of ascorbic acid that correspond to physiological concentrations, supraphysiologic concentrations appear to retard endothelial cell migration and tube formation. In contrast, type IV collagen was not affected by supraphysiologic ascorbic acid.

Our experiments in homozygous *Gulo*<sup>-/-</sup> mice indicate that dietary ascorbic acid restriction in these mice retards both initial tumor outgrowth and the growth of previously established tumors. Histopathological data from tumors in ascorbic acid-restricted mice demonstrate a paucity of blood vessels (from both microscopic examination and staining for vWF) and areas of hemorrhage. In contrast, tumors grown in mice supplemented with ascorbic acid have numerous and well-defined blood vessels and few areas of hemorrhage. Collagen staining of tumors in ascorbic acid-deficient and ascorbic acid-replete mice also suggests that collagen synthesis in these tumors is affected by dietary restriction of vitamin C. The possibility of using ascorbic acid restriction clinically is supported by our observation that restricting ascorbic acid to 10% of the full supplementation dose delays the onset of scorbutic symptoms in *Gulo*<sup>-/-</sup> mice while still significantly retarding tumor growth. Interestingly, we observed a trend toward increased tumor growth in mice that had been depleted of and then made replete with ascorbic acid. We postulate that host support for tumor growth may be increased secondary to overcompensation after ascorbic acid depletion. For example, tumor-associated endothelial cells may have upregulated surface transporters in response to low extracellular ascorbic acid, and this overexpression may confer the capacity for increased angiogenesis. However, the precise mechanism(s) for this



observation is not obvious, and we intend to examine this phenomenon in a future study.

This is, by no means, the first attempt to modify tumor growth through depletion of ascorbic acid. One early study reported the effects of ascorbic acid depletion on the growth and biology of sarcomas initiated by 20-methylcholanthrene in guinea pigs [38]. These investigators found that animals given high doses of ascorbic acid showed accelerated tumor growth, whereas tumors grew slowly (or even regressed) in guinea pigs fed a scorbutic diet. However, other studies in guinea pigs have provided conflicting results regarding the requirements for ascorbic acid in collagen synthesis [35,39], making interpretation of the effects of ascorbic acid depletion on tumor growth in guinea pigs difficult.

A recent study by Parsons et al. [40] evaluated tumor growth in a mammary tumor model in *Gulo*<sup>-/-</sup> mice bred in a Balb/c background. In these experiments, mice were ascorbic acid-depleted for 2 weeks before tumor implantation and did not show any differences in tumor growth in comparison with control animals on full ascorbic acid supplementation. We cannot, with certainty, explain why these results are so divergent from those reported here, but differences in tumor type and/or regimen used to deplete ascorbic acid might be involved. In our experiments, we restricted dietary ascorbic acid for 4 weeks so that the serum ascorbic acid concentration decreased to approximately five-fold below the normal physiological range. Furthermore, we chose to study Lewis lung carcinoma because this tumor relies heavily on angiogenesis [41,42]. Finally, strain differences in mice used in each study may also have affected the outcomes reported.

Tumors have been found to have an unusual metabolism of ascorbic acid, actively concentrating ascorbic acid relative to surrounding normal fluids and tissues [43,44]. This occurs across a concentration gradient, and uptake appears to involve the oxidized form dehydroascorbic acid, which enters through facilitative glucose transporters (GLUTs) [45]. In certain malignant cell types (e.g., melanoma), dehydroascorbic acid uptake may be up to 10 times greater than that exhibited by normal melanocytes [44]. However, despite evidence demonstrating that ascorbic acid is concentrated by cancer cells, the precise function performed by vitamin C in tumor cells is far from clear. In U937 lymphoma cells, ascorbic acid was reported to decrease Fas pathway-mediated apoptosis [46], and similar findings have been made on apoptosis engendered by the NF- $\kappa$ B pathway in several other tumor cell lines [47]. Furthermore, pretreatment of cells with the readily transported form of ascorbic acid, dehydroascorbic acid, has been shown to decrease radiation-induced apoptosis in HL60 promyelocytic leukemia cells [48]. These data indicate that ascorbic acid, in addition to supporting angiogenic activities, may also promote tumor cell survival and protect against oxidative stress.

Ascorbic acid depletion may lead to stabilization of HIF-1 $\alpha$  and thus contribute to tumor growth through factors downstream of HIF such as VEGF and GLUT-1, thus increasing angiogenesis and glycolysis [14]. Surprisingly, our immunohistochemical analyses of tumors demonstrated that HIF-1 $\alpha$  expression was not significantly altered by the absence of

ascorbic acid. These data indicate that ascorbic acid depletion may preferentially affect collagen formation—and, thus, new vessel formation—without affecting HIF-1 $\alpha$  stability.

Our results, although obtained with a single mouse model and a single type of implantable tumor, support the idea that ascorbic acid depletion powerfully restricts tumor growth *in vivo*, likely through interference with angiogenesis. Recent reports continue to advocate an increase in ascorbic acid intake over the recommended daily allowance for optimal reduction of the risk of cancer and chronic disease [49]. At the least, our results suggest that ascorbic acid supplementation in patients with cancer should be reevaluated. Additionally, we believe that decreasing dietary ascorbic acid or disrupting prolyl hydroxylases may provide novel therapeutic approaches to the treatment of cancer.

### Acknowledgements

We gratefully acknowledge helpful discussions with Richard Bucala and Otto Grubraw.

### References

- [1] Chatterjee I (1970). Biosynthesis of ascorbic acid in animals. *Methods Enzymol* **18A**, 28–34.
- [2] Grollman AP and Lehninger AL (1957). Enzymic synthesis of L-ascorbic acid in different animal species. *Arch Biochem Biophys* **69**, 458–467.
- [3] Lind J (1753). A Treatise on the Scurvy.
- [4] Hodges RE, Hood J, Canham JE, Sauberlich HE, and Baker EM (1971). Clinical manifestations of ascorbic acid deficiency in man. *Am J Clin Nutr* **24**, 432–443.
- [5] Folkman J (1971). Tumor angiogenesis: therapeutic implications. *N Engl J Med* **285**, 1182–1186.
- [6] Bonanno E, Lurlaro M, Madri JA, and Nicosia RF (2000). Type IV collagen modulates angiogenesis and neovessel survival in the rat aorta model. *In Vitro Cell Dev Biol Anim* **36**, 336–340.
- [7] Myllylä Raii E-RK-SaKIK (1978). The role of ascorbate in the prolyl hydroxylase reaction. *Biochem Biophys Res Commun* **83**, 441–448.
- [8] De la Fuente M and Victor VM (2001). Ascorbic acid and N-acetylcysteine improve *in vitro* the function of lymphocytes from mice with endotoxin-induced oxidative stress. *Free Radic Res* **35**, 73–84.
- [9] Jeng KC, Yang CS, Siu WY, Tsai YS, Liao WJ, and Kuo JS (1996). Supplementation with vitamins C and E enhances cytokine production by peripheral blood mononuclear cells in healthy adults. *Am J Clin Nutr* **64**, 960–965.
- [10] Naidu KA (2003). Vitamin C in human health and disease is still a mystery? An overview. *Nutr J* **2**, 7.
- [11] Cameron E and Pauling L (1974). The orthomolecular treatment of cancer: I. The role of ascorbic acid in host resistance. *Chem Biol Interact* **9**, 273–283.
- [12] Cameron E, Pauling L, and Leibovitz B (1979). Ascorbic acid and cancer: a review. *Cancer Res* **39**, 663–681.
- [13] Campbell JD, Cole M, Bunditratavorn B, and Vella AT (1999). Ascorbic acid is a potent inhibitor of various forms of T cell apoptosis. *Cell Immunol* **194**, 1–5.
- [14] Knowles HJ, Raval RR, Harris AL, and Ratcliffe PJ (2003). Effect of ascorbate on the activity of hypoxia-inducible factor in cancer cells. *Cancer Res* **63**, 1764–1768.
- [15] Maeda N, Hagihara H, Nakata Y, Hiller S, Wilder J, and Reddick R (2000). Aortic wall damage in mice unable to synthesize ascorbic acid. *Proc Natl Acad Sci USA* **97**, 841–846.
- [16] Taetle R, Rosen F, Abramson I, Venditti J, and Howell S (1987). Use of nude mouse xenografts as preclinical drug screens: *in vivo* activity of established chemotherapeutic agents against melanoma and ovarian carcinoma xenografts. *Cancer Treat Rep* **71**, 297–304.
- [17] Omaye ST, Turnbull JD, and Sauberlich HE (1979). Selected methods for the determination of ascorbic acid in animal cells, tissues, and fluids. *Methods Enzymol* **62**, 3–11.

- [18] Matkowskyj KA, Cox R, Jensen RT, and Benya RV (2003). Quantitative immunohistochemistry by measuring cumulative signal strength accurately measures receptor number. *J Histochem Cytochem* **51**, 205–214.
- [19] Matkowskyj KA, Schonfeld D, and Benya RV (2000). Quantitative immunohistochemistry by measuring cumulative signal strength using commercially available software Photoshop and MATLAB. *J Histochem Cytochem* **48**, 303–312.
- [20] Clement PM, Hanauske-Abel HM, Wolff EC, Kleinman HK, and Park MH (2002). The antifungal drug ciclopirox inhibits deoxyhypusine and proline hydroxylation, endothelial cell growth and angiogenesis *in vitro*. *Int J Cancer* **100**, 491–498.
- [21] Jaffe EA, Hoyer LW, and Nachman RL (1973). Synthesis of antihemophilic factor antigen by cultured human endothelial cells. *J Clin Invest* **52**, 2757–2764.
- [22] Sadler JE (1991). von Willebrand factor. *J Biol Chem* **266**, 22777–22780.
- [23] Cameron E and Campbell A (1974). The orthomolecular treatment of cancer: II. Clinical trial of high-dose ascorbic acid supplements in advanced human cancer. *Chem Biol Interact* **9**, 285–315.
- [24] Cameron E and Pauling L (1976). Supplemental ascorbate in the supportive treatment of cancer: prolongation of survival times in terminal human cancer. *Proc Natl Acad Sci USA* **73**, 3685–3689.
- [25] Cameron E and Pauling L (1978). Supplemental ascorbate in the supportive treatment of cancer: reevaluation of prolongation of survival times in terminal human cancer. *Proc Natl Acad Sci USA* **75**, 4538–4542.
- [26] Creagan ET, Moertel CG, O'Fallon JR, Schutt AJ, O'Connell MJ, Rubin J, and Frytak S (1979). Failure of high-dose vitamin C (ascorbic acid) therapy to benefit patients with advanced cancer. A controlled trial. *N Engl J Med* **301**, 687–690.
- [27] Moertel CG, Fleming TR, Creagan ET, Rubin J, O'Connell MJ, and Ames MM (1985). High-dose vitamin C *versus* placebo in the treatment of patients with advanced cancer who have had no prior chemotherapy. A randomized double-blind comparison. *N Engl J Med* **312**, 137–141.
- [28] Padayatty SJ and Levine M (2000). Reevaluation of ascorbate in cancer treatment: emerging evidence, open minds and serendipity. *J Am Coll Nutr* **19**, 423–425.
- [29] Padayatty SJ and Levine M (2001). New insights into the physiology and pharmacology of vitamin C. *CMAJ* **164**, 353–355.
- [30] Gimbrone MA Jr, Leapman SB, Cotran RS, and Folkman J (1972). Tumor dormancy *in vivo* by prevention of neovascularization. *J Exp Med* **136**, 261–276.
- [31] Ferrara N and Davis-Smyth T (1997). The biology of vascular endothelial growth factor. *Endocr Rev* **18**, 4–25.
- [32] Kim KJ, Li B, Winer J, Armanini M, Gillett N, Phillips HS, and Ferrara N (1993). Inhibition of vascular endothelial growth factor–induced angiogenesis suppresses tumour growth *in vivo*. *Nature* **362**, 841–844.
- [33] Millauer B, Shawver LK, Plate KH, Risau W, and Ullrich A (1994). Glioblastoma growth inhibited *in vivo* by a dominant-negative *Flk-1* mutant. *Nature* **367**, 576–579.
- [34] Stolman JMGHM and Gould BS (1961). Ascorbic acid and blood vessels. *Arch Pathol* **72**, 59–69.
- [35] Mahmoodian F and Peterkofsky B (1999). Vitamin C deficiency in guinea pigs differentially affects the expression of type IV collagen, laminin, and elastin in blood vessels. *J Nutr* **129**, 83–91.
- [36] Nicosia RF, Belser P, Bonanno E, and Diven J (1991). Regulation of angiogenesis *in vitro* by collagen metabolism. *In Vitro Cell Dev Biol* **27A**, 961–966.
- [37] Ingber D and Folkman J (1988). Inhibition of angiogenesis through modulation of collagen metabolism. *Lab Invest* **59**, 44–51.
- [38] Migliozi JA (1977). Effect of ascorbic acid on tumour growth. *Br J Cancer* **35**, 448–453.
- [39] Spanheimer RG and Peterkofsky B (1985). A specific decrease in collagen synthesis in acutely fasted, vitamin C–supplemented, guinea pigs. *J Biol Chem* **260**, 3955–3962.
- [40] Parsons KK, Maeda N, Yamauchi M, Banes AJ, and Koller BH (2006). Ascorbic acid–independent synthesis of collagen in mice. *Am J Physiol Endocrinol Metab* **290**, E1131–E1139.
- [41] Lamagna C, Hodiuala-Dilke KM, Imhof BA, and Aurrand-Lions M (2005). Antibody against junctional adhesion molecule-C inhibits angiogenesis and tumor growth. *Cancer Res* **65**, 5703–5710.
- [42] Watanabe K, Hasegawa Y, Yamashita H, Shimizu K, Ding Y, Abe M, Ohta H, Imagawa K, Hojo K, Maki H, et al. (2004). Vasohibin as an endothelium-derived negative feedback regulator of angiogenesis. *J Clin Invest* **114**, 898–907.
- [43] Langemann H, Torhorst J, Kabiersch A, Krenger W, and Honegger CG (1989). Quantitative determination of water- and lipid-soluble antioxidants in neoplastic and non-neoplastic human breast tissue. *Int J Cancer* **43**, 1169–1173.
- [44] Spielholz C, Golde DW, Houghton AN, Nualart F, and Vera JC (1997). Increased facilitated transport of dehydroascorbic acid without changes in sodium-dependent ascorbate transport in human melanoma cells. *Cancer Res* **57**, 2529–2537.
- [45] Vera JC, Rivas CI, Fischbarg J, and Golde DW (1993). Mammalian facilitative hexose transporters mediate the transport of dehydroascorbic acid. *Nature* **364**, 79–82.
- [46] Perez-Cruz I, Carcamo JM, and Golde DW (2003). Vitamin C inhibits FAS-induced apoptosis in monocytes and U937 cells. *Blood* **102**, 336–343.
- [47] Carcamo JM, Pedraza A, Borquez-Ojeda O, and Golde DW (2002). Vitamin C suppresses TNF alpha–induced NF kappa B activation by inhibiting I kappa B alpha phosphorylation. *Biochemistry* **41**, 12995–13002.
- [48] Witenberg B, Kletter Y, Kalir HH, Raviv Z, Fenig E, Nagler A, Halperin D, and Fabian I (1999). Ascorbic acid inhibits apoptosis induced by X irradiation in HL60 myeloid leukemia cells. *Radiat Res* **152**, 468–478.
- [49] Carr AC and Frei B (1999). Toward a new recommended dietary allowance for vitamin C based on antioxidant and health effects in humans. *Am J Clin Nutr* **69**, 1086–1107.