



Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb

Exotic $\mu\tau jj$ events from heavy ISS neutrinos at the LHCE. Arganda^a, M.J. Herrero^b, X. Marcano^{b,*}, C. Weiland^{b,c}^a Departamento de Física Teórica, Facultad de Ciencias, Universidad de Zaragoza, E-50009 Zaragoza, Spain^b Departamento de Física Teórica and Instituto de Física Teórica, IFT-UAM/CSIC, Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain^c Graduate School of Science and Engineering, Shimane University, Matsue, 690-8504, Japan

ARTICLE INFO

Article history:

Received 7 September 2015

Received in revised form 13 October 2015

Accepted 8 November 2015

Available online 11 November 2015

Editor: B. Grinstein

ABSTRACT

In this letter we study new relevant phenomenological consequences of the right-handed heavy neutrinos with masses at the $\mathcal{O}(1)$ TeV energy scale, working within the context of the Inverse Seesaw Model that includes three pairs of quasi-degenerate pseudo-Dirac heavy neutrinos. We propose a new exotic signal of these heavy neutrinos at the CERN Large Hadron Collider containing a muon, a tau lepton, and two jets in the final state, which is based on the interesting fact that this model can incorporate large Lepton Flavor Violation for specific choices of the relevant parameters, particularly, the neutrino Yukawa couplings. We will show here that an observable number of $\mu\tau jj$ exotic events, without missing energy, can be produced at this ongoing run of the LHC.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

One of the most interesting properties of the Inverse Seesaw Model (ISS) is that it allows for the existence of heavy Majorana neutrinos with masses within the reach of the CERN Large Hadron Collider (LHC), and with couplings to the Standard Model (SM) particles that can be sizable, therefore leading to observable production and decay rates. The version of the ISS that incorporates three pairs of SM-singlet heavy neutrinos, one pair per generation, is especially appealing because, on the one hand, it can explain the observed small neutrino masses and neutrino oscillation data and, on the other hand in connection with the LHC physics, it can lead to new interesting phenomenology with experimental signatures distinct from those of the SM particles. The most frequently studied signatures of heavy neutrinos are those related to their Majorana nature [1–3] and, in particular, the most characteristic signal is the same-sign dilepton plus two jets events which is being searched for at the LHC. The rates of this type of events within the ISS are directly related to the Majorana mass matrix μ_X that introduces the Lepton Number Violation in the model, which in turn is built in as to precisely fit the observed light neutrino masses via a low scale realization of the Seesaw Mechanism where $M_{\text{light}} \simeq m_D M_R^{T-1} \mu_X M_R^{-1} m_D^T$ and m_D and M_R are

* Corresponding author.

E-mail addresses: ernesto.arganda@unizar.es (E. Arganda), maria.herrero@uam.es (M.J. Herrero), xabier.marcano@uam.es (X. Marcano), cedric.weiland@uam.es (C. Weiland).

<http://dx.doi.org/10.1016/j.physletb.2015.11.013>

0370-2693/© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

the Dirac and extended right handed (RH) neutrino sector mass matrices respectively. The usually assumed hierarchy among the three input scales, $\mu_X \ll m_D \ll M_R$, then produces the interesting mentioned features of the ISS. Furthermore, due to the smallness of μ_X , the heavy neutrinos combine in nearly degenerate pairs to form pseudo-Dirac fermions with masses close to M_R and with a splitting of order $\mathcal{O}(\mu_X)$. This pseudo-Dirac character of the heavy neutrinos has also been explored in the literature in connection with the appearance of other interesting signals at the LHC, like the trilepton final state [4,5], and other multi-lepton signals [6].

We propose here a new exotic signal of the ISS heavy neutrinos at the LHC that is based on another interesting feature of the ISS, the fact that it can incorporate large Lepton Flavor Violation (LFV) for specific choices of the model parameters, particularly the Yukawa couplings Y_ν , and lead to a new rich phenomenology, including the appearance of rare processes like the radiative decays, $l_i \rightarrow l_j \gamma$ ($i \neq j$) and others. Moreover, it has been shown recently [7] that sizable branching ratios for the LFV Higgs boson decays, like $H \rightarrow \mu\tau$, can also be produced within this ISS context, that can be enhanced in its supersymmetric version [8] for specific choices of the model parameters, reaching values which are compatible with the present excess observed by CMS and ATLAS in $H \rightarrow \mu\tau$ [9,10], while respecting the present upper bounds on the related radiative decay $\tau \rightarrow \mu\gamma$ [11]. Our specific proposal here is to look at rare LHC events with one muon, one tau lepton and two jets in the final state that are produced in these ISS scenarios with large LFV, and that presumably will have a very small SM background. This letter then summarizes our computation of the rates

for these exotic $\mu\tau jj$ events due to the production and decays of the heavy quasi-Dirac neutrinos at the LHC within the ISS.

2. ISS scenarios with large LFV

We consider the ISS model that extends the SM with two kinds of fermionic gauge singlets: RH neutrinos, denoted here by ν_R and coupled to the ν_L via Yukawa interactions, and X , with opposite lepton number. We assume one pair of them per generation. The relevant ISS Lagrangian for the mass terms is:

$$\mathcal{L}_{\text{ISS}} = -Y_\nu^{ij} \bar{L}_i \tilde{\Phi} \nu_{Rj} - M_R^{ij} \bar{\nu}_{Ri}^c X_j - \frac{1}{2} \mu_X^{ij} \bar{X}_i^c X_j + \text{h.c.}, \quad (1)$$

where L is the SM lepton doublet, Φ is the SM Higgs doublet with $\langle \Phi \rangle = v = 174 \text{ GeV}$, $\tilde{\Phi} = i\sigma_2 \Phi^*$, Y_ν is the 3×3 neutrino Yukawa coupling matrix, M_R is a lepton number conserving complex 3×3 mass matrix, and μ_X is a Majorana complex 3×3 symmetric mass matrix that violates lepton number conservation by two units. The diagonalization of the relevant 9×9 neutrino mass matrix, containing the Dirac mass matrix, $m_D = Y_\nu \langle \Phi \rangle$, and M_R ,

$$M_{\text{ISS}} = \begin{pmatrix} 0 & m_D & 0 \\ m_D^T & 0 & M_R \\ 0 & M_R^T & \mu_X \end{pmatrix}, \quad (2)$$

provides the 9 physical neutrino states $n_j (j = 1, \dots, 9)$: three light neutrinos $\nu_{1,2,3}$ that are identified with the ones experimentally observed, and six heavy neutrinos $N_{1..6}$ which are the object of our interest here. In order to match the predictions of this ISS model with the low energy neutrino data, one considers small μ_X given by the following parameterization in terms of the light neutrino mixing matrix U_{PMNS} [12]:

$$\mu_X = M_R^T m_D^{-1} U_{\text{PMNS}}^* m_\nu U_{\text{PMNS}}^\dagger m_D^{T-1} M_R. \quad (3)$$

This parametrization, which works quite well in the $\mu_X \ll m_D \ll M_R$ limit, was introduced in Ref. [7] and allows us to use Y_ν and M_R as input parameters of the model. For the rest of this letter, we focus on the simplest case with diagonal M_R matrix and degenerate entries, $M_{R1,2,3} \equiv M_R$. Within this mentioned small μ_X limit, the heavy neutrinos can be grouped into pseudo-Dirac pairs with nearly degenerate masses. Both the light neutrino masses and the small differences among the quasi-degenerate heavy neutrino masses are governed by the small size of μ_X , basically following the same pattern as in the one generation case where:

$$m_\nu = \frac{m_D^2}{m_D^2 + M_R^2} \mu_X, \quad (4)$$

$$m_{N_{a/b}} = \pm \sqrt{M_R^2 + m_D^2} + \frac{M_R^2 \mu_X}{2(m_D^2 + M_R^2)}. \quad (5)$$

For the present work, the corresponding pairings of the six heavy neutrinos are given by: $N_{1/2}$, $N_{3/4}$ and $N_{5/6}$, with $m_{N_{1/2}} \leq m_{N_{3/4}} \leq m_{N_{5/6}}$. Their corresponding masses in the $M_R \gg m_D$ limit are close to M_R which will be taken here in the range accessible to the LHC, namely $\leq \mathcal{O}(1 \text{ TeV})$.

Regarding the heavy neutrino interactions, we give here just the relevant ones, namely, those to W gauge bosons, which enter both in the heavy neutrino production and in the decays of our interest, and those to Z gauge and Higgs bosons, which enter in the computation of the total heavy neutrino width:

$$\mathcal{L}_W = -\frac{g}{\sqrt{2}} W^{\mu-} \bar{l}_i B_{li n_j} \gamma_\mu P_L n_j + \text{h.c.}, \quad (6)$$

$$\mathcal{L}_Z = -\frac{g}{4c_W} Z^\mu \bar{n}_i \gamma_\mu [C_{n_i n_j} P_L - C_{n_i n_j}^* P_R] n_j, \quad (7)$$

$$\mathcal{L}_H = -\frac{g}{2m_W} H \bar{n}_i C_{n_i n_j} [m_{n_i} P_L + m_{n_j} P_R] n_j, \quad (8)$$




Fig. 1. Heavy neutrino flavor mixing, according to eq. (14), for the three benchmark scenarios in (11). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

where, assuming the charged leptons to be diagonal,

$$B_{li n_j} = U_{ij}^{\nu*}, \quad (9)$$

$$C_{n_i n_j} = \sum_{k=1}^3 U_{ki}^\nu U_{kj}^{\nu*}, \quad (10)$$

and U^ν is the 9×9 neutrino rotation matrix that diagonalizes the mass matrix (2) according to $U^{\nu T} M_{\text{ISS}} U^\nu = \text{diag}(m_{n_1}, \dots, m_{n_9})$.

Since we are interested in the ISS with large LFV, and more specifically producing large τ - μ transitions, we will work with some benchmark scenarios that produce these wanted features. In particular we will use the three benchmark scenarios proposed in [7] that are defined by the following input Yukawa matrices, Y_ν :

$$Y_{\tau\mu}^{(1)} = f \begin{pmatrix} 0 & 1 & -1 \\ 0.9 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad (11a)$$

$$Y_{\tau\mu}^{(2)} = f \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & -1 \\ -1 & 1 & -1 \end{pmatrix}, \quad (11b)$$

$$Y_{\tau\mu}^{(3)} = f \begin{pmatrix} 0 & -1 & 1 \\ -1 & 1 & 1 \\ 0.8 & 0.5 & 0.5 \end{pmatrix}. \quad (11c)$$

These scenarios have been explored fully in [7] and display large LFV ratios for processes involving τ - μ transitions (this motivates the given name, $Y_\nu = Y_{\tau\mu}^{(1,2,3)}$), whereas they strongly suppress processes involving τ - e and μ - e transitions, which can be easily understood with the use of the approximate formulas presented also in [7]:

$$\text{BR}_{\tau \rightarrow \mu \gamma}^{\text{approx}} = 8 \times 10^{-17} \frac{m_\tau^5 (\text{GeV}^5)}{\Gamma_\tau (\text{GeV})} \left| \frac{v^2}{2M_R^2} (Y_\nu Y_\nu^\dagger)_{23} \right|^2, \quad (12)$$




Fig. 2. The two main processes, Drell-Yan and γW fusion, producing exotic $\tau\mu jj$ events via heavy neutrino production and decay at the LHC.

$$\text{BR}_{H \rightarrow \mu\bar{\tau}}^{\text{approx}} = 10^{-7} \frac{v^4}{M_R^4} \left| (Y_\nu Y_\nu^\dagger)_{23} - 5.7 (Y_\nu Y_\nu^\dagger Y_\nu Y_\nu^\dagger)_{23} \right|^2. \quad (13)$$

For f values close to its perturbativity limit, $f = \sqrt{6\pi}$, and $M_R \sim \mathcal{O}(1 \text{ TeV})$ these lead to rates of up to $\text{BR}(H \rightarrow \mu\bar{\tau}) \sim 10^{-5}$ while respecting the present bounds on the radiative decays, $\text{BR}(\mu \rightarrow e\gamma) \leq 5.7 \times 10^{-13}$ [13], $\text{BR}(\tau \rightarrow \mu\gamma) \leq 4.4 \times 10^{-8}$ and $\text{BR}(\tau \rightarrow e\gamma) \leq 3.3 \times 10^{-8}$ [11]. Moreover, in the case of the SUSY version of this ISS model, the Higgs decay rates induced by the SUSY loop contributions have been found in [8] to be as large as the present LHC sensitivity of about $\text{BR}(H \rightarrow \mu\tau) \sim 10^{-2}$ [9,10].

As we see from eq. (12), the constrains from LFV radiative decays set upper bounds on the combination $|(Y_\nu Y_\nu^\dagger)_{ij}|$, with $i \neq j$. In terms of the heavy neutrino flavor mixing matrix, B_{iN} , they constrain the combination $|(B_{iN} B_{iN}^\dagger)_{ij}|$, but not the B_{iN_j} itself, which is the relevant element that controls the flavor pattern of each of the heavy neutrinos.

In Fig. 1 we show the flavor content of each heavy neutrino for the three benchmark scenarios in (11), where the length of the colored bars is calculated as

$$S_{iN_i} = \frac{|B_{iN_i}|^2}{\sum_{l=e,\mu,\tau} |B_{iN_i}|^2}, \quad (14)$$

and, therefore, represents the relative mixing of the heavy neutrino N_i with a given flavor l . We learn from Fig. 1 that, although the three scenarios share the property of suppressing the LFV μ - e and τ - e rates while maximizing the τ - μ ones, the heavy neutrino flavor mixing pattern is different in each scenario. We also see that some heavy neutrinos carry both μ and τ flavors, specially in the first two scenarios, pointing towards signals with both μ and τ leptons. It should be further noticed, that the values of these B_{iN} mixing parameters are determined within the ISS in terms of the input m_D and M_R mass matrices and, therefore, in the limit $m_D \ll M_R$ they are suppressed as $B_{iN} \sim \mathcal{O}(m_D M_R^{-1})$. In addition, they are constrained from ElectroWeak Precision Observables (EWPO) [14] by:

$$|B_{eN}|^2 < 3.0 \times 10^{-3}, \quad (15)$$

$$|B_{\mu N}|^2 < 3.2 \times 10^{-3}, \quad (16)$$

$$|B_{\tau N}|^2 < 6.2 \times 10^{-3}. \quad (17)$$

3. Numerical results

Heavy neutrinos with masses of the TeV order can be produced at present and future colliders, in particular in the second run of the LHC. The dominant production mechanism in this case is the Drell-Yan (DY) process, Fig. 2 left, where the heavy neutrino is produced in association with a charged lepton. The γW fusion,

Fig. 2 right, also produces the same signal with two extra jets and, in fact, can be relevant especially for large neutrino masses in the $\mathcal{O}(1 \text{ TeV})$ energy range [15,16].

In order to estimate the heavy neutrino production at the LHC, we have implemented the model in MadGraph 5 [17] using FeynRules [18]. Following Ref. [16], we have used a K -factor of 1.2 for the DY-process and split the γW process into three regimes characterized by the virtuality of the photon¹: elastic, inelastic and deep inelastic scattering (DIS) regimes. In particular, we have set the boundaries between these three regimes to $\Lambda_\gamma^{\text{Elastic}} = 1.22 \text{ GeV}$ and $\Lambda_\gamma^{\text{DIS}} = 15 \text{ GeV}$. In order to regularize possible collinear singularities, we have also imposed the following cuts to the transverse momentum and pseudorapidity of the outgoing leptons:

$$p_T^l > 10 \text{ GeV}, \quad |\eta^l| < 2.4. \quad (18)$$

The results for the three benchmark scenarios in (11) are shown in Fig. 3, where the dominant DY production cross sections normalized by f^2 are plotted as functions of the mass parameter M_R . We see that the production cross sections can be of the fb order, reachable at the LHC, for masses $M_R \lesssim 600 \text{ GeV}$. Notice that the results are always equal for the pseudo-Dirac pairs, since their Majorana character plays a subleading role in their production.

We can also learn that the flavor of the associated charged lepton is different depending on the heavy neutrino produced and the scenario considered, and that this pattern can be understood looking at the mixing in Fig. 1. Thus, $N_{3/4}$ are mainly electronic neutrinos in the first and third scenarios, therefore, they are basically produced exclusively with electrons. $N_{1/2}$ are equally produced with muons and taus in the first scenario, dominantly produced with electrons in the second scenario, and mainly produced with taus in the third scenario. $N_{5/6}$ are equally produced with muons and taus in the first and second scenarios, and mainly produced with muons in the third scenario.

Once the heavy neutrinos are produced, they will decay inside the detector. As mentioned above, in the limit $M_R \gg m_D$ the heavy neutrino masses are close to M_R , with small differences of $\mathcal{O}(m_D^2 M_R^{-1})$ between the different pseudo-Dirac pairs and, therefore, assuming they are practically degenerate, their decay into each other should be suppressed, with the dominant channels being $N_j \rightarrow Z\nu_i, H\nu_i, W^\pm l_i^\mp$. The relevant decays into $W^+ l_i^-$ or $W^- l_i^+$ have a partial width given by:

$$\Gamma_{N_j \rightarrow W l_i} = \frac{\sqrt{(M_j^2 - m_i^2 - m_W^2)^2 - 4m_i^2 m_W^2}}{16\pi M_j^3} |\overline{F_W}|^2, \quad (19)$$

with

¹ We warmly thank Richard Ruiz, Tao Han and Daniel Alva for their generous help and clarifications in the implementation of the γW process.




Fig. 3. Heavy neutrino DY-production at the LHC for the three scenarios. The cross sections are normalized by the Yukawa coupling intensity f in (11). Processes not shown are negligible. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

$$|\overline{F}_W|^2 = \frac{g^2}{4M_W^2} |B_{iN_j}|^2 \times \left\{ (M_j^2 - m_i^2)^2 + M_W^2 (M_j^2 + m_i^2) - 2M_W^4 \right\}. \quad (20)$$

We have also seen that, in agreement with [3,4], in the large M_j limit, the heavy neutrinos decay with equal branching ratios to all the bosons in all the scenarios if we sum over all the flavors, namely,

$$\begin{aligned} \text{BR}(N_j \rightarrow h\nu) &= \text{BR}(N_j \rightarrow Z\nu) = \\ \text{BR}(N_j \rightarrow W^+l^-) &= \text{BR}(N_j \rightarrow W^-l^+) = 25\%. \end{aligned} \quad (21)$$

It is, however, interesting to study the rich flavor structure of the decay products, which depends on the decaying heavy neutrino and the scenario we are considering. Like in the production, the flavor preference of the decays to $W^\pm l^\mp$, which are the ones relevant to this study and are given in Fig. 2, also follows the same pattern as in Fig. 1. Therefore, we can expect the production and decay of the heavy neutrinos to lead to exotic $\mu\tau jj$ events with no missing energy and $M_{jj} \sim M_W$, with M_{jj} the invariant mass of the two jets.

Using the narrow width approximation, the total cross section of the exotic events we are interested in is given by:

$$\sigma(pp \rightarrow \mu\tau jj) =$$




Fig. 4. Number of exotic $\mu\tau jj$ events at the LHC for the three scenarios and for three values of f . For each f , the bottom solid line is the prediction of $\mu\tau jj$ events from DY and the upper lines on top of each of the three shadowed regions are the predictions after adding the $\mu\tau jj$ events from γW , for $p_T^{\max} = 10, 20, 40$ GeV from bottom to top, respectively. The p_T^{\max} cut is only applied to the two extra jets in the events from γW . The upper red shadowed areas are excluded by $\tau \rightarrow \mu\gamma$. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

$$\begin{aligned} \sum_{i=1}^6 \left\{ \sigma(pp \rightarrow N_i \mu^\pm) \text{BR}(N_i \rightarrow W^\pm \tau^\mp) \right. \\ \left. + \sigma(pp \rightarrow N_i \tau^\pm) \text{BR}(N_i \rightarrow W^\pm \mu^\mp) \right\} \\ \times \text{BR}(W^\pm \rightarrow jj). \end{aligned} \quad (22)$$

Fig. 4 shows the expected number of exotic events $\mu\tau jj$ at the LHC for an integrated luminosity of $\mathcal{L} = 300 \text{ fb}^{-1}$ at $\sqrt{s} = 14 \text{ TeV}$. The lower solid lines for each choice of f are the number of events considering only the DY-production.

Moreover, γW fusion processes can also contribute to this kind of exotic events if the p_T of the extra jets are below a maximum value p_T^{\max} and, therefore, they can be considered as soft or collinear jets which can escape detection. In this case the predicted total number of exotic events are the sum of the events produced by DY and γW channels. These total contributions for different values of $p_T^{\max} = 10, 20$ and 40 GeV are shown as the border lines on top of the shadowed areas with gradual decreasing intensity above each solid line. In addition we have included in the plots red shadowed areas that represent the regions excluded by the experimental upper bound on $\text{BR}(\tau \rightarrow \mu\gamma)$. We can see that, after considering all the constraints, the three scenarios lead

to an interesting number of $\mathcal{O}(10\text{--}100)$ total $\mu\tau jj$ exotic events for the range of M_R studied here of [200 GeV, 1 TeV].

The SM backgrounds for events with two leptons of different flavor have been studied in Ref. [19]. However, a high efficiency in the τ -tagging and a good reconstruction of the W boson invariant mass from the two leading jets would help in reducing the background. In that case, the main background would come from processes with photons or jets misidentified as leptons, mainly from $W/Z + \gamma^*$, $W/Z + \text{jets}$ and multijet events with at least four jets with one of them misidentified as a muon and another one as a tau; and from $Z/\gamma^* + \text{jets} \rightarrow \mu^+\mu^- + \text{jets}$ if one of the muons is misidentified as a τ candidate. Nevertheless, a dedicated background study for these particular $\mu\tau jj$ exotic events is beyond the scope of this letter and it will be done in a future work.

4. Conclusions

In this letter we have proposed a new interesting way to study the production and decay of the heavy neutrinos of the ISS in connection with LFV. We have presented the computation of the predicted number of exotic $\mu\tau jj$ events which can be produced in these ISS scenarios with large LFV by the production of heavy pseudo-Dirac neutrinos together with a lepton of flavor ℓ , both via DY and γW fusion processes, and their subsequent decay into W plus a lepton of different flavor. We have concluded that, for the three benchmark scenarios studied here, a number of $\mathcal{O}(10\text{--}100)$ total $\mu\tau jj$ exotic events without missing energy can be produced at this ongoing run of the LHC, for values of M_R from 200 GeV to 1 TeV. Similarly, other rare processes like $\tau e jj$ or $\mu e jj$ could be produced within other ISS scenarios with large LFV, although for the latter ones, while being interesting since they could provide in addition observable CP asymmetries [20], the number of events would be strongly limited by the $\mu \rightarrow e\gamma$ upper bound. Of course, a more realistic study of these exotic events, including detector simulation, together with a full background study should be done in order to reach a definitive conclusion, but this is beyond the scope of this letter and will be addressed in a future work.

Acknowledgements

This work is supported by the European Union Grant No. FP7 ITN INVISIBLES (Marie Curie Actions, Grant No. PITN-GA-2011-289442), by the CICYT through Grant No. FPA2012-31880, by the Spanish Consolider-Ingenio 2010 Programme CPAN (Grant No. CSD2007-00042), and by the Spanish MINECO's "Centro de Exce-

lencia Severo Ochoa" Programme under Grant No. SEV-2012-0249. E.A. is financially supported by the Spanish DGIID-DGA Grant No. 2013-E24/2 and the Spanish MICINN Grants No. FPA2012-35453 and No. CPAN-CSD2007-00042. X.M. is supported through the FPU Grant No. AP-2012-6708. C.W. receives financial support as an International Research Fellow of the Japan Society for the Promotion of Science.

References

- [1] A. Pilaftsis, *Z. Phys. C* 55 (1992) 275, arXiv:hep-ph/9901206; A. Datta, M. Guchait, A. Pilaftsis, *Phys. Rev. D* 50 (1994) 3195, arXiv:hep-ph/9311257.
- [2] F. del Aguila, J.A. Aguilar-Saavedra, R. Pittau, *J. High Energy Phys.* 0710 (2007) 047, arXiv:hep-ph/0703261.
- [3] A. Atre, T. Han, S. Pascoli, B. Zhang, *J. High Energy Phys.* 0905 (2009) 030, arXiv:0901.3589 [hep-ph].
- [4] A. Das, N. Okada, *Phys. Rev. D* 88 (2013) 113001, arXiv:1207.3734 [hep-ph]; A. Das, P.S. Bhupal Dev, N. Okada, *Phys. Lett. B* 735 (2014) 364, arXiv:1405.0177 [hep-ph].
- [5] G. Bambhaniya, S. Goswami, S. Khan, P. Konar, T. Mondal, *Phys. Rev. D* 91 (2015) 075007, arXiv:1410.5687 [hep-ph]; G. Bambhaniya, S. Khan, P. Konar, T. Mondal, *Phys. Rev. D* 91 (9) (2015) 095007, arXiv:1411.6866 [hep-ph].
- [6] F. del Aguila, J.A. Aguilar-Saavedra, *Nucl. Phys. B* 813 (2009) 22, arXiv:0808.2468 [hep-ph].
- [7] E. Arganda, M.J. Herrero, X. Marcano, C. Weiland, *Phys. Rev. D* 91 (1) (2015) 015001, arXiv:1405.4300 [hep-ph].
- [8] E. Arganda, M.J. Herrero, X. Marcano, C. Weiland, arXiv:1508.04623 [hep-ph].
- [9] V. Khachatryan, et al., CMS Collaboration, *Phys. Lett. B* 749 (2015) 337, arXiv:1502.07400 [hep-ex].
- [10] G. Aad, et al., ATLAS Collaboration, arXiv:1508.03372 [hep-ex].
- [11] B. Aubert, et al., BaBar Collaboration, *Phys. Rev. Lett.* 104 (2010) 021802, arXiv:0908.2381 [hep-ex].
- [12] B. Pontecorvo, *Sov. Phys. JETP* 6 (1957) 429; B. Pontecorvo, *Zh. Eksp. Teor. Fiz.* 33 (1957) 549; Z. Maki, M. Nakagawa, S. Sakata, *Prog. Theor. Phys.* 28 (1962) 870.
- [13] J. Adam, et al., MEG Collaboration, *Phys. Rev. Lett.* 110 (2013) 201801, arXiv:1303.0754 [hep-ex].
- [14] F. del Aguila, J. de Blas, M. Perez-Victoria, *Phys. Rev. D* 78 (2008) 013010, arXiv:0803.4008 [hep-ph].
- [15] P.S.B. Dev, A. Pilaftsis, U.k. Yang, *Phys. Rev. Lett.* 112 (8) (2014) 081801, arXiv:1308.2209 [hep-ph].
- [16] D. Alva, T. Han, R. Ruiz, *J. High Energy Phys.* 1502 (2015) 072, arXiv:1411.7305 [hep-ph].
- [17] J. Alwall, et al., *J. High Energy Phys.* 1407 (2014) 079, arXiv:1405.0301 [hep-ph].
- [18] A. Alloul, N.D. Christensen, C. Degrande, C. Duhr, B. Fuks, *Comput. Phys. Commun.* 185 (2014) 2250, arXiv:1310.1921 [hep-ph].
- [19] G. Aad, et al., ATLAS Collaboration, *Phys. Rev. Lett.* 115 (3) (2015) 031801, arXiv:1503.04430 [hep-ex].
- [20] S. Bray, J.S. Lee, A. Pilaftsis, *Nucl. Phys. B* 786 (2007) 95, arXiv:hep-ph/0702294.