

# Ouabain- and $\text{Ca}^{2+}$ -sensitive ATPase activity of chimeric Na- and Ca-pump molecules

Douglas B. Luckie, Kimberly L. Boyd and Kunio Takeyasu

Department of Physiology, University of Virginia School of Medicine, Charlottesville, VA 22908, USA

Received 31 December 1990; revised version received 11 February 1991

Chimeric ion-pumps, consisting of the N-terminal 2/3 of the  $\alpha_1$ -subunit of the ouabain-sensitive chicken  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase and the C-terminal 1/3 of the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase, were expressed in ouabain-insensitive mouse L cells. These chimeric molecules exhibited ouabain-sensitive ATPase activity very similar to that of the wild-type chicken  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase. This ATPase activity could be stimulated by adding  $\text{Ca}^{2+}$  to the assay system. These results suggest that the sites for ouabain-inhibition are restricted to the N-terminal 2/3 of the Na-pump, and the C-terminal 1/3 of the Ca-pump interacts with  $\text{Ca}^{2+}$ .

$\text{Na}^+$ ,  $\text{K}^+$ -ATPase;  $\text{Ca}^{2+}$ -ATPase; Ouabain; Chimeric molecule; Chicken

## 1. INTRODUCTION

Recent cDNA cloning and sequencing studies have shown that a variety of E1-E2 type ATPases (ion pumps) may form very similar higher order structures, even though the primary amino acid sequences of these ATPases are rather different [1]. On the other hand, these ATPases show distinct specificities for particular ions, inhibitors and subunits. One approach to identifying the functional domains that determine selectivity of ion transport is to construct chimeric molecules between distinct functional proteins that have very similar structures. This type of approach has been successfully applied to structure/function analysis of several membrane proteins [2-6].

Proposed models of the higher order structures of E1-E2 ATPases predict multiple (7-10) transmembrane segments, and conserved amino acid sequences for the ATP-binding- and phosphorylation-domains between the fourth and the fifth potential transmembrane domains ( $M_4$  and  $M_5$ ) (Fig. 1). The region just before  $M_5$  is also very well conserved among these distinct ion-pumps [1]. We have taken advantage of this, and constructed a set of chicken sodium pump/calcium pump chimeric cDNAs that would preserve the overall topological structure of the encoded chimeric proteins as ion pumps. Mouse L cells that express the chicken  $\beta$ -

subunit [7] were transfected with these chimeric cDNAs. Because the  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase of mouse L cells is relatively ouabain-resistant, this avian/mouse hybrid system allows us to monitor enzymatic functions of the relatively ouabain-sensitive chimeric molecules.

## 2. MATERIALS AND METHODS

### 2.1. Construction of mutant cDNAs encoding chicken Na-pump/Ca-pump chimeric molecules

Wild-type cDNAs encoding the chicken sodium pump  $\alpha_1$ -subunit [8] and the chicken SR calcium-pump [9] were used for construction of chimeric cDNAs. A unique restriction site, *Eco*NI, which is endogenous to both the SR calcium pump cDNA and the PM sodium pump  $\alpha_1$ -subunit cDNA, was used as an exchange point for recombination. This exchange occurred at nucleotide 2134 on the  $\alpha_1$ -subunit cDNA and at nucleotide 2170 on the calcium pump cDNA, both of which encode lysine<sup>\*</sup> residues within the evolutionarily conserved FSBA-binding domain [1,10] (-Pro-Ala-Leu-Lys<sup>\*</sup>-Lys-Ala-) just before  $M_5$  of the two proteins. These chimeric cDNAs as well as wild type cDNAs were cloned into an expression vector [7] derived from a plasmid pSV<sub>2</sub>CAT [11].

### 2.2. Transfection and cell culture

The detailed methods for cell culture were described previously [7,8]. Mouse L cells expressing the  $\beta_1$ -subunit of the chicken  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase [7] were co-transfected by the calcium phosphate precipitation method, as described by Small and Scangos [12], with 0.1  $\mu\text{g}$  of pSVneo DNA and about 1  $\mu\text{g}$  of chimeric cDNA/100 mm diameter tissue culture plate. The transfected cells were selected in a medium containing G418 sulfate (Gibco) (~500  $\mu\text{g}/\text{ml}$ ), and then screened for expression of the chicken chimeric molecules using chicken specific monoclonal antibodies (see below).

### 2.3. Immunofluorescence microscopy

Monoclonal antibodies 7C (specific for the chicken  $\alpha_1$ -subunit [8]) and 5D2 (specific for the chicken SR-calcium pump [13]), and TRITC-labeled goat-anti-mouse IgG were used to identify the chicken proteins in mouse cells. Fluorescence images were obtained with a laser scanning confocal microscope (BioRad MRC 600). Further details of immuno-fluorescent staining were described in previous reports [7,8].

Correspondence address: K. Takeyasu, Department of Physiology, University of Virginia School of Medicine, Charlottesville, VA 22908, USA

Abbreviations: ATPase, adenosine-tri-phosphatase; FITC, fluorescein-isothiocyanate; FSBA, 5'-(*p*-fluorosulfonyl)benzoyl-adenosine; PM, plasma membrane; SR, sarcoplasmic reticulum; TRITC, tetramethylrhodamine




Fig. 1. Construction of chicken chimeric ion-pumps. Schematic representation of a possible structure of chimera I. Chimera II (not shown) consists of the N-terminal 2/3 of the SR calcium-pump and the C-terminal 1/3 of the sodium pump. Phosphorylation site (Pi) and FITC- and FSBA-binding domains are shown, and potential transmembrane segments are numbered.

#### 2.4. ATPase activity assays

The rate of ATP hydrolysis was determined with an enzyme coupled system, for each ADP produced in the assay, a molecule of NADH was oxidized to  $\text{NAD}^+$  and this was monitored spectroscopically by a decrease in absorbance at 340 nm. [14-16]. The ATPase assay medium (1 ml) contained 5  $\mu\text{l}$  of cell homogenate (300-500  $\mu\text{g}$  total protein), 120 mM NaCl, 15 mM KCl, 30 mM triethanolamine, pH 7.4, 4 mM  $\text{MgCl}_2$ , 3 mM  $\text{Na}_2\text{ATP}$ , 0.5 mM EGTA, 2 mM  $\text{NaN}_3$ , 2.5 mM phosphoenolpyruvate, 0.5 mM NADH, 10 U/ml pyruvate kinase, and 30 U/ml lactate dehydrogenase.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity was obtained by calculating the difference in activity in the presence and absence of 1 mM ouabain. Protein concentration was determined by the method of Bradford [17].

### 3. RESULTS AND DISCUSSION

Utilizing an avian/murine hybrid expression system [7,8,18], relatively ouabain-sensitive chicken sodium-pump functions can be studied in relatively ouabain-

insensitive mouse L cells. When a chicken  $\beta_1$ -subunit cDNA was stably transfected into mouse L cells, the chicken  $\beta_1$ -subunit protein was expressed on mouse cell surfaces associated with the mouse  $\alpha$ -subunit. These cells are designated as  $\text{Ltk}^+ \beta$  cells [7]. When an additional chicken  $\alpha_1$ -subunit cDNA was introduced into these mouse  $\text{Ltk}^+ \beta$  cells, the chicken  $\alpha_1$ -subunit was expressed both on the mouse cell surface and inside the cells. These cells are designated as  $\text{Ltk}^+ \beta \text{neo}^+ \alpha$  cells [18]. Normally, mouse cells transfected with the chicken  $\alpha_1$ -subunit cDNA express  $10^5$ - $10^6$  ouabain-sensitive sites per cell during 10 mM butyrate treatment [8]. When mouse cells are transfected with a cDNA encoding chicken SR calcium pump, the avian calcium pump protein was always expressed inside the cells [9]. These cell lines were used as control cells in this study.

Fig. 2 shows confocal fluorescence micrographs of the cloned mouse cells expressing chicken chimeric ion pumps ( $\text{Ltk}^+ \beta \text{neo}^+ \text{ChiI}$  and  $\text{Ltk}^+ \beta \text{neo}^+ \text{ChiII}$ ). Chimera I (containing the N-terminal 2/3 of the sodium pump  $\alpha_1$ -subunit) was found only internally, while chimera II (containing the N-terminal 2/3 of the calcium pump) was found both on the plasma membrane and inside the cell. Extensive studies of the expression of sodium-pump subunits in a variety of systems have demonstrated that the  $\alpha$ -subunit contains exclusively the domains conferring ouabain-sensitivity and catalytic activity to the sodium pump [5,8,19-21], while the  $\beta$ -subunit is critically involved in the stabilization and transport of the  $\alpha$ -subunit to the PM [18,19,22,23]. A recent study of expression of sodium/calcium pump chimeric molecules in *Xenopus* oocytes demonstrated that the C-terminal 1/3 of the  $\alpha$ -subunit contains the domain(s) that assemble with the  $\beta$ -subunit [24]. Subcellular distributions of chimeras I and II (Fig. 2) are consistent with these results. Our in-




Fig. 2. Laser confocal micrographs of cloned mouse L cells that express the chicken Na-pump/Ca-pump chimeric molecules. (Left) Cells transfected with chimera I cDNAs were fixed, permeabilized, and stained with monoclonal IgG 7C specific to the chicken  $\alpha_1$ -subunit [8]. (Right) Cells transfected with chimera II cDNAs were fixed, permeabilized, and stained with monoclonal IgG 5D2 specific to the chicken SR calcium pump [13]. Scale bar = 10  $\mu\text{m}$ .

interpretation is that chimera I lacks the domains that assemble with the  $\beta$ -subunit, and is, therefore, not transported to the plasma membrane, while chimera II acquires the assembling domains and is transported to the cell surface.

Fig. 3 illustrates the effects of varying concentrations of ouabain on the ATPase activity (measured in the presence of  $\text{Na}^+$  and  $\text{K}^+$  and absence of  $\text{Ca}^{2+}$ ) of cloned mouse L cells transfected with various chicken cDNAs. The inhibition of the ATPase activity of both  $\text{Ltk}^+ \beta\text{neo}^+ \alpha$  and  $\text{Ltk}^+ \beta\text{neo}^+ \text{ChII}$  cells by ouabain is biphasic;  $\sim 60\%$  shows  $\text{IC}_{50}$  of  $< 10 \mu\text{M}$  and the rest  $\text{IC}_{50}$  of  $\sim 300 \mu\text{M}$ . On the other hand, ouabain has a monophasic inhibitory effect ( $\text{IC}_{50}$ :  $\sim 300 \mu\text{M}$ ) in untransfected mouse  $\text{Ltk}^-$  and  $\text{Ltk}^+ \beta\text{neo}^+ \text{ChII}$  cells. At present, functions of chimera II distinct from those of the endogenous mouse ATPase cannot be detected in our system (see also below). However, it has been known that the calcium-pump does not bind ouabain, and that the endogenous mouse sodium pump does not have high affinity sites for ouabain. Thus, this new high affinity site for ouabain in transfected cells must come from the chicken wild-type  $\alpha_1$ -subunit and chimera I. Recently, using chimeras and site-directed mutagenesis, Price and Lingrel [5] have demonstrated that the extracellular domain between M1 and M2 consisting of 11 amino acids (especially Arg-116 and Asp-127) is the major ouabain binding domain. However, whether ouabain binds only to this site or to additional site(s) in order to exert its action remains to be determined. The




Fig. 3. Dose/response relationships for ouabain-inhibition of the ATPase activity of homogenate from untransfected ( $\text{Ltk}^-$ ) and transfected ( $\text{Ltk}^+ \beta\text{neo}^+ \text{ChII}$ ,  $\text{Ltk}^+ \beta\text{neo}^+ \text{ChIII}$ , and  $\text{Ltk}^+ \beta\text{neo}^+ \alpha$ ) mouse cells. ATPase activities of cell homogenate were measured in the presence of 0.2% Triton X-100 which partially solubilizes and clears turbid sample and allows chemicals access to both sides of the membrane. At appropriate concentrations, detergents have been found to activate the Na,K-ATPase [27]. In our system, 0.2% Triton X-100 increases ATPase activity on the average 5-fold in the presence of  $\text{Ca}^{2+}$ - and mitochondrial ATPase inhibitors (0.5 mM EGTA, 2 mM  $\text{NaN}_3$ ). The ATPase activity (60–80 nmol ADP cleaved/mg protein/min) that can be inhibited by 1 mM ouabain was defined as 100%. Ouabain-sensitive ATPase activity was  $\sim 64\%$  of overall ATPase activity measured in the presence of 0.5 mM EGTA and 2 mM  $\text{NaN}_3$ .

results from the inhibitory effect of ouabain on chimera I suggest that ouabain does not have to bind to the ectodomains after  $\text{M}_2$ .

Some of the potential transmembrane segments, including  $\text{M}_4$ ,  $\text{M}_5$ ,  $\text{M}_6$ , of both sodium- and calcium-pumps contain glutamic and aspartic acid residues. Mutagenesis experiments have demonstrated that these charged amino acids are critical for both sodium pump [25] and calcium pump activities [26]. In our chimeric molecules, these amino acids are preserved. Therefore, it is interesting to identify the ion specificity of these chimeric molecules. Fig. 4 shows the effect of  $\text{Ca}^{2+}$  ions on the chimeric molecules, illustrating that  $\text{Ca}^{2+}$  ions added to the assay system stimulated the ATPase activity of chimera I. Under the same conditions, neither the chicken wild-type nor the mouse endogenous ATPase showed this level of stimulation by  $\text{Ca}^{2+}$ . Thus, when combined with the N-terminal 2/3 of the sodium pump  $\alpha_1$ -subunit, the C-terminal 1/3 of the calcium pump appears to undergo the proper protein folding required for  $\text{Ca}^{2+}$ -stimulation of the ATPase activity.

We have established many mouse L cell lines stably transfected with the chicken wild-type  $\text{Ca}^{2+}$ -ATPase cDNA as well as chimera II cDNA. However, we have not been able to detect chicken calcium-pump specific




Fig. 4. Effect of calcium ions on the ATPase activity of homogenate from transfected cells. The same ATPase assay system as described in Fig. 3 (in the presence of Na and K ions) was activated at time 0, and then 1  $\mu\text{l}$  of 0.5 M  $\text{CaCl}_2$  was added to the 1 ml assay system after a few minutes (shown by arrows in the figure). This created a final free  $\text{Ca}^{2+}$  level of  $\sim 4 \mu\text{M}$  after EGTA chelation. The addition of the same level of free Ca ions to the assay system leads to maximal stimulation of the  $\text{Ca}^{2+}$ -ATPase activity of biochemically purified chicken skeletal SR  $\text{Ca}^{2+}$ -ATPase. (A)  $\text{Ltk}^+ \beta\text{neo}^+ \text{ChII}$  cells. Sample contained 450  $\mu\text{g}$  protein and calibration (adding 50 nmol of ADP to sample after the assay's completion) determined 0.222 AU to equal 50 nmol ADP. ATPase activity before and after  $\text{Ca}^{2+}$ -addition was 92.6 and 211 nmol ADP cleaved/mg protein/min, respectively. (B)  $\text{Ltk}^+ \beta$  cells. Sample contained 296  $\mu\text{g}$  protein and calibration determined 0.355 AU to equal 50 nmol ADP in this assay. ATPase activity before and after  $\text{Ca}^{2+}$  addition was 86.7 and 94.1 nmol ADP cleaved/mg protein/min, respectively.

ion-stimulated ATPase activity for unknown reasons. The only method currently available for detecting the chicken  $\text{Ca}^{2+}$ -ATPase activity (suggested to us by Dr. G. Inesi, Univ. Maryland) is to express them in COS-1 cells [26]. Probably expression of the chimeric and the wild type molecules in COS-1 cells will be useful for further characterization of ion specificity.

The  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase is supposed to exist as an  $\alpha$ -/ $\beta$ -subunit complex in order to exert its full activity [1], while the SR  $\text{Ca}^{2+}$ -ATPase does not require any additional subunits for its full functioning. In this sense, it is intriguing that chimera I (N-terminal sodium-/C-terminal calcium-pump) exhibits ouabain-sensitive and  $\text{Ca}^{2+}$ -stimulated ATPase activity. Further detailed analysis of chimera I, as well as careful characterization of chimera II as a ' $\text{Ca}^{2+}$ -ATPase' and a ' $\text{Na}^+$ ,  $\text{K}^+$ -ATPase', should provide critical information about the structure, conformational changes, and functional relationships of the E1-E2 ATPases.

*Acknowledgements:* We wish to thank Drs. D.M. Fambrough and N.J. Karlin for a gift of a cDNA encoding the SR  $\text{Ca}^{2+}$ -ATPase, Dr. Z. Kaprielian for a monoclonal antibody, 5D2, Dr. K.J. Renaud for a monoclonal antibody, 7C, Dr. H.C. Kutchai for ATPase assays. Drs. A.P. Somlyo and Z. Shao for laser-scanning confocal microscopy and for critical discussion. This study was supported by a grant from NIH (RO1-GM44373 to K.T.). D.B.L. and K.L.B. were supported by NIH Training Grant T32 HL07284. K.T. is an Established Investigator of the American Heart Association (1991-1996) and a member of the Cancer Center and the Diabetes Center at the University of Virginia.

## REFERENCES

- [1] Jorgensen, P.L. and Andersen, J.P. (1988) *J. Membr. Biol.* 103, 95-120.
- [2] Kobilka, B.K., Kobilka, T.-S., Daniel, K., Regan, J.W., Caron, M.G. and Lefkowitz, R.J. (1988) *Science* 240, 1310-1316.
- [3] Fukuda, K., Kubo, T., Maeda, A., Akiba, I., Bujo, H., Nakai, J., Mishina, M., Higashida, H., Neher, E., Marty, A. and Numa, S. (1989) *Trends Pharmacol. Sci. (Suppl.)*, 4-10.
- [4] Lax, I., Bellot, F., Hawk, R., Ulrich, A., Clivet, D. and Schlessinger, J. (1989) *EMBO J.* 8, 421-427.
- [5] Price, E.M. and Lingrel, J.B. (1988) *Biochemistry* 27, 8400-8408.
- [6] Noguchi, S., Ohta, T., Takeda, K., Ohtsubo, M. and Kawamura, M. (1988) *Biochem Biophys. Res. Commun.* 155, 1237-1243.
- [7] Takeyasu, K., Tamkun, M.M., Siegel, N.R. and Fambrough, D.M. (1987) *J. Biol. Chem.* 262, 10733-10740.
- [8] Takeyasu, K., Tamkun, M.M., Renaud, K.J. and Fambrough, D.M. (1988) *J. Biol. Chem.* 263, 4347-4354.
- [9] Karlin, N.J., Kaprielian, Z. and Fambrough, D.M. (1989) *Mol. Cell. Biol.* 9, 1978-1986.
- [10] Ohta, T., Nagano, K. and Yoshida, M. (1986) *Proc. Natl. Acad. Sci. USA* 83, 2071-2075.
- [11] Gorman, C.M., Moffat, L. and Howard, B.H. (1982) *Mol. Cell. Biol.* 2, 1044-1051.
- [12] Small, J. and Scangos, G. (1984) *Gene Anal. Techn.* 1, 13-17.
- [13] Kaprielian, Z. and Fambrough, D.M. (1987) *Dev. Biol.* 124, 490-503.
- [14] Kutchai, H. and Campbell, K.P. (1989) *Biochemistry* 28, 4830-4839.
- [15] Madden, T.D., Chapman, D. and Quinn, P.J. (1979) *Nature* 279, 538-541.
- [16] Schwartz, Z., Allen, J.C. and Harigaya, S. (1969) *J. Pharmacol. Exp. Ther.* 168, 31-40.
- [17] Bradford, M. (1976) *Anal. Biochem.* 72, 248-254.
- [18] Takeyasu, K., Renaud, K., Taormino, J., Wolitsky, B.A., Barnstein, A., Tamkun, M.M. and Fambrough, D.M. (1989) *Curr. Top. Membr. Transp.* 34, 143-165.
- [19] Noguchi, S., Mishina, M., Kawamura, M. and Numa, S. (1987) *FEBS Lett.* 225, 27-32.
- [20] Kent, R.B., Emanuel, J.R., Neria, Y.B., Levenson, R. and Housman, D.E. (1987) *Science* 237, 901-903.
- [21] Emanuel, J.R., Schulz, J., Zhou, X.-M., Kent, R.B., Housman, D., Cantley, L. and Levenson, R. (1988) *J. Biol. Chem.* 263, 7726-7733.
- [22] McDonough, A.A., Geering, K. and Farley, R.A. (1990) *FASEB J.* 4, 1598-1605.
- [23] Geering, K. (1990) *J. Membr. Biol.* 115, 109-121.
- [24] Kawamura, M. (1991) in: *The Sodium Pump*, Rockefeller Univ. Press (in press).
- [25] Lingrel, J.B., Orłowski, J., Price, E., Jewell, E., Szamraj, O., Pathak, B. and Tyson, P. (1991) in: *The Sodium Pump*, Rockefeller Univ. Press (in press).
- [26] Clarke, D.M., Loo, T.W., Inesi, G. and MacLennan, D.H. (1989) *Nature* 339, 476-478.
- [27] Forbush, B. (1983) *Anal. Biochem.* 128, 159-163.