

inhibition and drug washout, were not determined. Here, these issues are examined using Chinese hamster ovary (CHO) cells, transiently transfected with mouse HCN1 DNA, measurement of resulting currents using the whole-cell patch clamp technique, and a fast perfusion system for the application of lidocaine. We found that the onset of lidocaine action is fast (~6 seconds) with maximum inhibition occurring at approximately 20 seconds, and that its effects are partially reversible. With 200 μM lidocaine perfused directly onto the cell under study, If elicited in response to -85mV and -100mV test pulses was reduced by $18.4 \pm 3.6\%$ ($n=6$ cells) and $8.3 \pm 1.0\%$ ($n=7$ cells), respectively, with average current return of $49.1 \pm 10.9\%$ and $73.9 \pm 9.4\%$ to the level prior to lidocaine addition. The fact that currents did not return completely may be attributed to a phenomenon called current rundown, or to incomplete washout of lidocaine. The significant difference in current reduction between -85mV and -100mV suggests that the effect of lidocaine on HCN1 channels depends on voltage, with greater If inhibition at less negative voltages.

2460-Pos Board B430

Ligand Binding and Gating in HCN2 Channels

Jana Kusch¹, Christoph Biskup¹, Susanne Thon¹, Eckhard Schulz², Klaus Benndorf¹.

¹Universitätsklinikum Jena, Jena, Germany, ²Fachhochschule Schmalkalden, Schmalkalden, Germany.

HCN channels are nonselective tetrameric cation channels that are activated by hyperpolarizing voltages and modulated by the ligand cAMP. They generate spontaneous rhythmic activity in heart and brain. Ligand binding to the intracellular cyclic nucleotide-binding site accelerates activation kinetics, shifts the steady-state activation to more positive voltages and increases the open probability. Though it is relatively simple to determine an apparent affinity for the ligand action, it is not so simple to determine the true ligand affinity during channel activation because, according to the principle of reciprocity, ligand binding and efficacy depend on each other, i.e. the affinity of the binding sites must increase when the channel opens. It is therefore important to determine the binding of the ligands to the channels and channel activation simultaneously.

Activation of homotetrameric HCN2 channels was studied in inside-out macro-patches simultaneously with ligand binding by means of patch-clamp fluorometry, using a fluorescent cAMP (fcAMP) that activates the channels in a similar manner as cAMP. With 1 μM fcAMP the binding of the ligand to the open channels exceeded that to closed channels. The slowness of the activation time course of HCN2 channels allowed us to monitor the ligand binding during the activation process. As predicted, the slow activation time course was accompanied by an increase of ligand binding. Moreover, the increase of binding was exponential whereas activation obeyed the typical sigmoidal time course. Hence, in the superimposed normalized time courses, the initial binding preceded activation whereas at later times activation preceded binding. These results show that activation gating indeed increases the binding affinity for the ligands, quite as predicted by the principle of reciprocity, and that the gating of the first of the four subunits, which does not lead to channel opening, is associated with ligand binding.

2461-Pos Board B431

Electrophysiological Evaluation of Novel Blockers of If Current

Martina Del Lungo¹, Michele Melchiorre², Laura Sartiani¹, Martin Biel³, Andras Varro⁴, Maria N. Romanelli², Elisabetta Cerbai¹.

¹Center of Molecular Medicine University of Florence, Florence, Italy,

²Department of Pharmaceutical Sciences, University of Florence, Florence, Italy,

³Department of Pharmacy and Pharmacology, University of Munich, Munich, Germany, ⁴Department of Pharmacology and Pharmacotherapy, University of Szeged, Szeged, Hungary.

In the sino-atrial node (SAN) a major role in rhythm generation is played by f-channels. They mediate a sodium-potassium inward current (If) activated upon hyperpolarization and encoded by HCN genes (HCN1-4). If is overexpressed in cardiac diseases at ventricular level, where it may contribute to the increased propensity for arrhythmias. Selective f-channel blockers have a potential therapeutic use as bradycardic and antiarrhythmic agents. Zatebradine and ivabradine act as f-channel blockers but they lack cardiac selectivity, blocking the neuronal HCN isoforms. As a consequence an unmet need exists to develop new blockers selective for the mammalian SAN channel isoform, HCN4. Zatebradine analogues (C1-C5) were synthesized and the effect on If was measured on patch-clamped HEK293 cells expressing mHCN1, mHCN2 and hHCN4 and native guinea-pig and rabbit SAN cells. At 10 μM concentration all compounds reduced maximal If amplitude; however, potencies (defined by EC50) differed considerably. Ivabradine, taken as reference compound, showed no isoform selectivity; C1 and C4 were more potent on HCN1, the ratio being 4.3 (HCN2/HCN1) and 7.5 (HCN4/HCN1). C2 was more potent on HCN4, the ratio being 6 and 17 vs. HCN1 and HCN2, respectively. C3 was equipotent on HCN1 and HCN4 and C5 had low activity on all isoforms. Blockade was concentration-dependent, did not reverse upon drug removal and did not change current activa-

tion properties. Finally, data obtained in SAN cells show that effects on native If resemble those obtained on HCN4 isoform, in accordance with the hypothesis that HCN4 has a major contribution in SAN cells. Present results indicate that drug interaction with different HCN isoforms has diverse structural requirements. Current investigations are aimed to characterize the pharmacological profile of the new f-channel blockers and improve their isoform selectivity (Supported by EU - LSH M/CT/2006/018676, Normacor).

2462-Pos Board B432

Functional Consequences Of Disease-associated Mutations In The Pore Region Of Human Cone Photoreceptor CNG Channels

Katja Koeppen, Peggy Reuter, Thomas Ladewig, Bernd Wissinger.

Institute for Ophthalmic Research, Tuebingen, Germany.

CNGA3 encodes the A-subunit of the cone photoreceptor cyclic nucleotide-gated (CNG) channel. Mutations in the CNGA3 gene have been associated with achromatopsia, a congenital, autosomal-recessively inherited retinal disorder characterized by lack of color vision, severely reduced visual acuity, photophobia and nystagmus. The aim of this study was the functional characterization of five mutant CNGA3 channels with amino acid substitutions in the pore region (S341P, L363P, G367V, P372S and E376K), which had been identified in achromatopsia patients. Mutant channels were heterologously expressed in HEK293 cells and their functional properties were assessed by calcium imaging and patch-clamp measurements. For patch-clamp recordings mutant CNGA3 was co-expressed with the wild-type B3 subunit present in native channels and transfected cells were incubated at 27°C in order to enhance folding and trafficking of the channel mutants. Furthermore, immunocytochemical experiments were performed after incubation at either 27°C or 37°C to determine the extent of co-localization of mutant channels with the cell membrane.

All five pore mutations rendered the channel non-functional in calcium imaging experiments, indicating severely reduced calcium influx through the mutant channel pore. Interestingly, cGMP-induced potassium currents could be recorded from patches containing channels with the mutations S341P, G367V and E376K. Even though macroscopic currents were small compared to wild type channels, these three pore mutants have been shown to possess residual potassium conductivity. While channels with the mutation G367V, P372S or E376K showed normal co-localization with the plasma membrane after incubation at 37°C, reduced surface expression was observed for channel mutants S341P and L363P, suggesting impaired folding and/or trafficking of the mutant proteins.

2463-Pos Board B433

Bimodal Agonism In A Cyclic Nucleotide-Gated Channel Is Coordinated By Two Adjacent Binding Domains

Kerry Chan, Edgar Young.

Simon Fraser University, Burnaby, BC, Canada.

Bimodal agonism is a cGMP-dependent desensitization found in the catfish CNGA2 subtype of the cyclic nucleotide-gated channel. In bimodal agonism, initial cGMP binding events at lower concentrations (<3 mM) increase channel open probability (P_o); however, additional cGMP binding events at higher concentrations (>3 mM) decrease P_o . Understanding bimodal agonism could lead to building cyclic nucleotide analogues that can specifically open or close the channel. The C-terminal region (β 7-C helix) of the ligand-binding domain (BD) was previously shown to determine ligand selectivity and efficacy, so we tested its role in bimodal agonism. The corresponding region from the normal (non-bimodal) CNGA4 BD was substituted into the bimodal CNGA2 BD to form a chimeric channel subunit. We expressed the chimera as homomers and tested its activation in the excised patch-clamp. Steady-state currents were measured at high and low concentrations of cGMP. The chimera is bimodal ($I_{10\text{mM cGMP}}/I_{3\text{mM cGMP}}$ is 0.72) suggesting that the C-terminal region of the BD is not essential in bimodal agonism. The roles of intersubunit interactions in bimodal agonism were studied through fusing together two bimodal and two normal pseudo-subunits into tandem tetramers. Tetramers with the two bimodal pseudo-subunits arranged either adjacent (*cis*) or diagonally opposite (*trans*) to each other were tested. The *cis* tetramer is bimodal ($I_{30\text{mM cGMP}}/I_{3\text{mM cGMP}}$ is 0.87) but the *trans* tetramer is not ($I_{30\text{mM cGMP}}/I_{3\text{mM cGMP}}$ is 1.05). The *cis* and *trans* tetramers have an identical number of bimodal subunits yet their capability for bimodal agonism (and by extension their P_o at high cGMP concentration) is different. This suggests that the bimodal subunits could be coupled in a dimer within a "dimer-of-dimers" mechanism. This also suggests a putative additional cGMP binding pocket may be located in the regions between two adjacent bimodal BDs.

2464-Pos Board B434

The Voltage Sensor Of Cng1 Channels Becomes Functional When The Hydrophobic Bond Between Phe380 And Leu356 Is Impaired

Monica Mazzolini.

SISSA, Trieste, Italy.