

Available online at www.sciencedirect.com**ScienceDirect**

Agriculture and Agricultural Science Procedia 10 (2016) 135 – 139

Agriculture and Agricultural Science

Procedia

5th International Conference "Agriculture for Life, Life for Agriculture"

Soils zones in Romania and pedogenetic processes

Anca-Luiza STĂNILĂ*, Mihail DUMITRU

*National Research Institute for Soil Science, Agrochemistry and Environment – ICPA Bucharest,
61 Mărăști Blvd, District 1, 011464, Bucharest, Romania*

Abstract

Orohydrographic conditions, climate and vegetation that are specific to our country's territory they have made a small area to develop most soils globe. Romania presents a diversity of soils, from the semi-arid zones in which specific zones encountered the wet and cold. Most soils in our country are distributed in a horizontal zonality plains and low hills are represented by cernisols (kastanozems, chernozems, phaeozems) and luvisols (preluvosols, luvosols) and a vertical zonality of high hills and mountain regions including cambisols zone (eutricambosols, districambosols), spodosols zone (prepodzols, podzols) and umbrisols zone (nigrosols, humosiosols).

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the University of Agronomic Sciences and Veterinary Medicine Bucharest

Keywords: soils zones, pedogenetic processes, climate, vegetation, taxonomy.

1. Introduction

Soil zone represents a territory characterized by the predominance of a soil type. The concept of zonality of the soil has been issued, as it is known by V. V. Dokuceaev (1898), in our country is evidenced by Gh. M. Murgoci (1911) and was widely used in soil geography.

Soil zonality is determined by climate zonality, natural law of nature, and the earth's rotation sphericity, and influence on climate and mountainous regions and the maritime currents.

* Corresponding author. Tel.: +4021-318-4463; Fax: +4021-318-4348.
E-mail address: luizastanila2011@yahoo.com

Typically, a soil zone is established by pedogenetic processes relative to climate, vegetation and course in relief (Posea and Cruceru, 2005). Based on these considerations, our land soils are judiciously disposed in the plains, plateaus and low hills in a horizontal zonality gradually passing into regions of high hills and mountains in a vertical zonality (Murgoci, 1911).

Of course, that vertical zonality, is identical, but not analogue, with horizontal zonality; there are specific zones that are missing in mountainous horizontal zonality as, for example, cambisols zone and the humosols specific alpine meadows.

2. Materials and Methods

The field studies consisted of mapping and spatial reambulating the studied area on maps at 1:10.000 and 1:25.000 scale, with collection of numerous soil and groundwater samples, with observations on relief, micro-relief, parent material, climate, etc.

The basic research and mapping unit of the areas with soil zone was the soil profile, thus allowing the study of morphological characteristics of the soils. As a result, soils were classified based on intrinsic properties, namely the soil profile, taking into account diagnostic horizons and characteristics (Soil survey methodology, 3 Volume, 1987).

Soil profiles were located on the ground so that to form a network of studied points. The method of parallel routes, located almost at equal distances has been used, to cover more or less uniformly the whole working area.

The morphological description of soil profiles was done according to the Romanian System of Soil Taxonomy (SRTS, 2003, 2012), ICPA Bucharest.

In order to establish the soils diagnosis, their morphological features have been taken into account, namely the thickness of morphological horizons, color, texture, structure, composition, adhesion, etc. Soil samples were taken from genetic horizons both in modified and unchanged settings.

In modified settings, soil samples of 20 cm thickness were taken in bags, for the chemical characterization to be carried.

In natural (unchanged) settings, soil samples were taken using a metal cylinder of known volume (200 cm³), to characterize the physical and hydro-physical features, as well as the momentary soil moisture.

3. Results and Discussions

Horizontal zonality start with the *cernisols zone*, represented by types kastanozems, chernozems and phaeozems, to which pedogenetic processes dominant are bioaccumulation with the formation of the mollic A horizon (Am); followed by clay migration with the development of the horizon B argic (Bt) and argillisation which leads to the appearance of cambic horizon (Bv) (Figure 1, 2, 3, Table 1) (Florea et al., 2012; Ispas and Stănilă, 2015).

Cernisols zone overlaps plains, plateaus and low hills (0-550 m), in hot dry climates sub-humid moderate heat, until cool and humid (Suceava Plateau) under the vegetation of the steppe and forest-steppe. This zone occupies a quarter of the country (26.7%).

The next zone is that of the *luvisols*, including as soil types preluvosols and luvosols (Figure 4). Of pedogenetic processes it should be mentioned clay migration with the formation of eluvial horizons (El, Ea) and iluvial horizon (Bt argic). This zone overlaps the plains and high plateaus, Sub-Carpathian hills and piedmont (200-800 m).

The climate remains relatively warm and semi-humid with average annual temperatures of 7-10°C and 550-800 mm rainfall, under the forests of quercinee or quercinee mixture with *Fagus sylvatica*. Occupies a different quarter of the country's 25.5 %.

Layout of vertical soil starts with *cambisols zone* which includes eutricambosols associated with regions of high hills districambosols and only districambosols low mountainous regions. Argillisation is pedogenetic processes that led to the formation of cambic horizon (Bv), diagnostic for cambisols.

Cambisols zone overlaps the lower mountain stage (500-1450 m), with cool climate - wet, average annual temperatures of 3-6°C and 6-8°C and annual rainfall between 700-1300 mm under beech and mixed forests (beech-spruce).

Spodisols zone with the main soil types prepodzols and podzols that forms as a result of the podzolization runs in alpine East (1450-1900 m) (2200 m) (Figure 5).

The climate is very humid and cold with average annual temperature of 2-6°C and average annual rainfall of 800-1400 mm and a characteristic vegetation consists of forests of spruce, fir and spruce that are thinned and then gradually disappear, their place being taken by *Pinus mugo* and *Juniperus communis*.




Fig. 1. Map soils zones in Romania.

Table 1. The main features of soils zones.

No.	Soils zones	Pedogenetic processes main	Types of soil	Horizons pedogenetic features	Major relief units	Climate	Vegetation
1.	Cernisols	Bioaccumulation Clay migration Argillisation	Kastanozems Chernozems Phaeozems	Am Bv Bt	Plains, plateaus and low hills (0-550 m)	Hot dry - subhumid moderate heat, until cool and humid	Steppe and forest steppe zone
2.	Luvisols	Clay migration	Preluvsols Luvosols	Ao El, Ea Bt	Plains and high plateaux Sub-Carpathian hills and piedmont (200 -800 m)	Heat-semi-wet	Forest zone (oak and beech subzone)
3.	Cambisols	Argillisation	Eutricambosols Districambosols	Ao Bv	High hills, mountains (500-1450 m)	Cool - wet	Forest zone (sublevel of beech and mixed forest)
4.	Spodosols	Podzolization	Prepodzols Podzols	O, Ao (Au) Es Bs, Bhs	Mountains- (1450-1900 m) (2200 m)	Very cold-wet	Subalpine floor
5.	Umbrisols	Organic matter humification Debasification- alteration	Nigrisols Humosiosols	O, Au Bv	Mountains- (2000-2500 m)	Very cool - very wet	Alpine meadows

The last area that corresponds to the upper alpine stage, with altitudes between 2000-2500 m the climate is very cold and very humid with average annual temperatures of -2°C and reserves 1300-1400 mm of water and vegetation is represented only by alpine meadows, it is a *umbrisols*. It is represented by two types of soil: nigrosols and humosiosols.

Pedogenetic processes mention that formation of organic matter humification horizon umbric (Au), organic horizon (Of) then debasification and alteration formation horizon cambic (Bv).



Fig. 2. Cambic chernozem in the Moldavian Plain.



Fig. 3. Argic phaeozem in the Burnas Plain.

Cambisols, spodosols and umbrisols represents another quarter of the country (25.5%), but the share is held by cambisols a rate of over 19%.

That, essentially, any change in climate and vegetation entail a change in the soil cover. Such climate and vegetation, soils main land mass formed at different zones (strips).

Making a comparison between natural of soils zones and the climate and vegetation, we find that they do not coincide, nor should coincide, because the soil formation also attended other factors besides climate and vegetation.



Fig. 4. Albic luvisol in the Candesti Piedmont.



Fig. 5. Podzol in the Bucegi Mountains.

4. Conclusions

Arguably if we go in every region of the country from the side the lowest relief to the higher parts of its soils zones are located in plains, plateaus and low hills in a horizontal zonality passing gradually in regions high hills and mountains in a vertical zonality.

Regarding the direction of the areas of soil, they are oriented along parallels geographic only in Oltenia and Muntenia, while in Moldova, Banat and Crisana, they have about the direction of the meridians in Transylvania there is a disposition somewhat concentric soils zones.

All these are determined by the direction of the mountain range, namely because regular orography of the country.

References

- Florea, N., Munteanu, I., Rusu, C., Dumitru, M., Ianoș, Gh., Răducu, D., Rogobete, Gh., Țărău, D., 2012. Romanian System of Soil Taxonomy. Sitech Publishing House, Bucharest.
- Ispas, Șt., Stănilă, A.-L., 2015. Soils of Romania. Publisher Wallachia University Press, Târgoviște.
- Murgoci, Gh., 1911. Natural soils zones in Romania. Extract from the Yearbook of the Geological Institute of Romania, year IV, 1910, Fascicule I.
- Posea, Gr., Cruceru, N., 2005. Geomorphology Romania. Romania for Tomorrow Publishing House, Bucharest.
- ***, 1987. Soil survey methodology (3 Volume). ICPA, Bucharest.