



Exactly solvable quantum mechanics and infinite families of multi-indexed orthogonal polynomials

Satoru Odake^{a,*}, Ryu Sasaki^b

^a Department of Physics, Shinshu University, Matsumoto 390-8621, Japan

^b Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

ARTICLE INFO

Article history:

Received 3 May 2011

Received in revised form 11 June 2011

Accepted 25 June 2011

Available online 1 July 2011

Editor: T. Yanagida

Keywords:

Shape invariance

Orthogonal polynomials

ABSTRACT

Infinite families of *multi-indexed* orthogonal polynomials are discovered as the solutions of exactly solvable one-dimensional quantum mechanical systems. The simplest examples, the one-indexed orthogonal polynomials, are the infinite families of the *exceptional* Laguerre and Jacobi polynomials of types I and II constructed by the present authors. The totality of the integer indices of the new polynomials are finite and they correspond to the degrees of the 'virtual state wavefunctions' which are 'deleted' by the generalisation of Crum–Adler theorem. Each polynomial has another integer n which counts the nodes.

© 2011 Elsevier B.V. Open access under CC BY license.

1. Introduction

We will report on the discovery of infinite families of *multi-indexed* orthogonal polynomials, which form the main parts of the eigenfunctions of exactly solvable one-dimensional quantum mechanical systems. This will add a huge number of new members to the inventory of exactly solvable quantum mechanics [1,2], which had seen rapid increase due to the recent discovery of various kinds of infinite families of *exceptional* orthogonal polynomials [3–8]. The method for deriving the multi-indexed orthogonal polynomials is simple generalisation of the well-known method of Crum [9] and its modification by Krein and Adler [10]. Starting from an exactly solvable Hamiltonian system, Crum's method provides another exactly solvable system by successively deleting the lowest lying eigenstate. The modification due to Krein and Adler uses deletion of a finite number of higher eigenstates satisfying certain conditions. The present method is based on the same spirit of 'deleting' a finite number of *virtual states* instead of the eigenstates. Here the virtual state wavefunctions are the solutions of the Schrödinger equation of the original system and they have *negative energies*, with respect to the groundstate eigenvalue. The virtual state wavefunctions have *no node in the interior* and they are *square-non-integrable*. Their reciprocals are *square-non-integrable*, too. We apply this method to the Hamiltonian systems of the radial oscillator potential and the Darboux–Pöschl–Teller potential, which are the best known examples of exactly solvable quantum mechanics

[1,2]. Their eigenfunctions consist of the Laguerre and the Jacobi polynomials, respectively.

When only one virtual state of degree $\ell (\geq 1)$ is deleted, the resulting system is the exceptional X_ℓ Laguerre or Jacobi polynomials of type I or II [3,5]. In general M distinct virtual states of degree $v_j (\geq 1)$ are deleted and the resulting eigenfunctions consist of orthogonal polynomials indexed by the integers (v_1, \dots, v_M) . The multi-indexed orthogonal polynomials are natural generalisation of the exceptional orthogonal polynomials. They start at degree $\ell \geq 1$ instead of a degree 0 constant term, thus avoiding the constraints by Bochner's theorem [11,12]. As solutions of exactly solvable quantum mechanical systems they constitute a complete system of orthogonal functions. The first examples of the exceptional Laguerre and Jacobi polynomials are constructed by Gómez-Ullate et al. [12] in the framework of Sturm–Liouville theory and by Quesne [13] within the shape invariant [14] and exactly solvable quantum mechanics. They are equivalent to the lowest members ($\ell = 1$) of the X_ℓ Laguerre and Jacobi polynomials derived in [3] by the present authors. The present method could be considered as the generalisation of the Darboux–Crum transformations [15] applied for the derivation of the exceptional orthogonal polynomials [16–20].

It is expected that these new polynomials would play important roles in physics, mathematics and related disciplines. The multi-indexed Jacobi polynomials provide infinitely many global solutions of second order Fuchsian differential equations with $3 + \ell$ regular singularities [21]. Other theoretical developments are: non-linear identities underlying the shape invariance [22] are clarified, the structure of the extra zeros of the new orthogonal polynomials [23] are exemplified, two-step Darboux transformations [24]

* Corresponding author.

E-mail address: odake@azusa.shinshu-u.ac.jp (S. Odake).

are explicitly constructed for the type II exceptional Laguerre polynomials, various use of non-physical states [25,26] for generating the exceptional polynomials are suggested and several applications [27,28] are reported. In this connection, let us mention references of multi-step Darboux–Crum transformations, higher order susy transformations, \mathcal{N} -fold susy transformations, etc. [29]. Some of them utilise non-physical solutions.

This Letter is organised as follows. The general scheme of virtual states deletion is presented in section two. The explicit forms of the new polynomials are derived in section three, together with the forward–backward shift operators and the second order differential operators. The final section is for a summary and comments.

2. Adler–Crum scheme for virtual states deletion

Here we present the method of virtual states deletion in its generic and formal form. For rigorous treatments of possible singularities at the boundaries, we need to specify the details of the Hamiltonian systems, which will be done in the next section. Throughout this Letter we choose all the solutions of the Schrödinger equations to be real. We start with an exactly solvable one-dimensional quantum mechanical system defined in an interval $x_1 < x < x_2$,

$$\mathcal{H}\phi_n(x) = \mathcal{E}_n\phi_n(x) \quad (n \in \mathbb{Z}_{\geq 0}), \quad (1)$$

$$(\phi_m, \phi_n) \stackrel{\text{def}}{=} \int_{x_1}^{x_2} dx \phi_m(x)^* \phi_n(x) = h_n \delta_{mn}. \quad (2)$$

For simplicity, we assume that it has discrete eigenvalues only with vanishing groundstate energy: $0 = \mathcal{E}_0 < \mathcal{E}_1 < \mathcal{E}_2 < \dots$, which allows us to express the potential in terms of the groundstate wavefunction $\phi_0(x)$ which has no node:

$$\mathcal{H} = -\frac{d^2}{dx^2} + U(x), \quad U(x) = \frac{\partial_x^2 \phi_0(x)}{\phi_0(x)}. \quad (3)$$

Let us suppose that the Schrödinger equation has a finite or infinite number of *virtual state* solutions with *negative energies*:

$$\mathcal{H}\tilde{\phi}_v(x) = \tilde{\mathcal{E}}_v \tilde{\phi}_v(x), \quad \tilde{\mathcal{E}}_v < 0 \quad (v \in \mathcal{V}), \quad (4)$$

which *have no node* in the interior $x_1 < x < x_2$. Here \mathcal{V} denotes the index set of the virtual states. We also require the infinite norm conditions, $(\tilde{\phi}_v, \tilde{\phi}_v) = (1/\tilde{\phi}_v, 1/\tilde{\phi}_v) = \infty$ and this means that the virtual state wavefunctions vanish at one boundary and diverge at the other:

$$\begin{aligned} \lim_{x \rightarrow x_j} \tilde{\phi}_v(x) &= \lim_{x \rightarrow x_j} \partial_x \tilde{\phi}_v(x) = 0, \\ \lim_{x \rightarrow x_k} \tilde{\phi}_v(x) &= \lim_{x \rightarrow x_k} \partial_x \tilde{\phi}_v(x) = \infty, \end{aligned} \quad (5)$$

where $j \neq k \in \{1, 2\}$ and $\forall v \in \mathcal{V}$. The set of eigenfunctions $\{\phi_n(x)\}$ forms a complete set in an appropriate Hilbert space \mathbb{H} , for example:

$$\mathbb{H} \stackrel{\text{def}}{=} \left\{ \psi(x) \in L^2[x_1, x_2] \mid \lim_{x \downarrow x_1} (x - x_1)^{\frac{1}{2}} \partial_x \psi(x) = 0, \right. \\ \left. \lim_{x \uparrow x_2} (x_2 - x)^{\frac{1}{2}} \partial_x \psi(x) = 0 \right\}.$$

(For $x_1 = -\infty$ and $x_2 = \infty$, replace $(x - x_1)^{\frac{1}{2}} \rightarrow (-x)^{\frac{1}{2}}$ and $(x_2 - x)^{\frac{1}{2}} \rightarrow x^{\frac{1}{2}}$.) That is any function in \mathbb{H} which is orthogonal to all the eigenfunctions has a zero norm:

$$(\Psi, \phi_n) = 0 \quad (\forall n \in \mathbb{Z}_{\geq 0}) \quad \Rightarrow \quad (\Psi, \Psi) = 0.$$

After M -steps of deleting the lowest lying eigenstates by Crum's method [9], the resulting Hamiltonian system is iso-spectral to the original one given by

$$\mathcal{H}^{[M]} \phi_n^{[M]}(x) = \mathcal{E}_n \phi_n^{[M]}(x) \quad (n = M, M+1, \dots),$$

$$\phi_n^{[M]}(x) \stackrel{\text{def}}{=} \frac{W[\phi_0, \phi_1, \dots, \phi_{M-1}, \phi_n](x)}{W[\phi_0, \phi_1, \dots, \phi_{M-1}](x)},$$

$$(\phi_m^{[M]}, \phi_n^{[M]}) = \prod_{j=0}^{M-1} (\mathcal{E}_n - \mathcal{E}_j) \cdot h_n \delta_{mn},$$

$$U^{[M]}(x) \stackrel{\text{def}}{=} U(x) - 2\partial_x^2 \log |W[\phi_0, \phi_1, \dots, \phi_{M-1}](x)|,$$

in which $W[f_1, \dots, f_n](x) \stackrel{\text{def}}{=} \det(\partial_x^{j-1} f_k(x))_{1 \leq j, k \leq n}$ is the Wronskian. These formulas have been repeatedly rediscovered under various names, higher derivative susy, \mathcal{N} -fold susy, etc. [29]. The corresponding Adler–Crum formulas for deleting the set of M distinct eigenstates specified by $\mathcal{D} \stackrel{\text{def}}{=} \{d_1, d_2, \dots, d_M\}$, $d_j \geq 0$, are

$$\mathcal{H}^{[M]} \phi_n^{[M]}(x) = \mathcal{E}_n \phi_n^{[M]}(x) \quad (n \in \mathbb{Z}_{\geq 0} \setminus \mathcal{D}),$$

$$\phi_n^{[M]}(x) \stackrel{\text{def}}{=} \frac{W[\phi_{d_1}, \phi_{d_2}, \dots, \phi_{d_M}, \phi_n](x)}{W[\phi_{d_1}, \phi_{d_2}, \dots, \phi_{d_M}](x)},$$

$$(\phi_m^{[M]}, \phi_n^{[M]}) = \prod_{j=1}^M (\mathcal{E}_n - \mathcal{E}_{d_j}) \cdot h_n \delta_{mn},$$

$$U^{[M]}(x) \stackrel{\text{def}}{=} U(x) - 2\partial_x^2 \log |W[\phi_{d_1}, \phi_{d_2}, \dots, \phi_{d_M}](x)|.$$

The set \mathcal{D} must satisfy the conditions $\prod_{j=1}^M (m - d_j) \geq 0$, $\forall m \in \mathbb{Z}_{\geq 0}$ [10,30]. The exactly solvable Hamiltonian system obtained by deleting M distinct *virtual states* $\mathcal{D} = \{d_1, \dots, d_M\}$, $d_j \in \mathcal{V}$ looks almost the same:

$$\mathcal{H}^{[M]} \phi_n^{[M]}(x) = \mathcal{E}_n \phi_n^{[M]}(x) \quad (n \in \mathbb{Z}_{\geq 0}), \quad (6)$$

$$\mathcal{H}^{[M]} \tilde{\phi}_v^{[M]}(x) = \tilde{\mathcal{E}}_v \tilde{\phi}_v^{[M]}(x) \quad (v \in \mathcal{V} \setminus \mathcal{D}), \quad (7)$$

$$\phi_n^{[M]}(x) \stackrel{\text{def}}{=} \frac{W[\tilde{\phi}_{d_1}, \tilde{\phi}_{d_2}, \dots, \tilde{\phi}_{d_M}, \phi_n](x)}{W[\tilde{\phi}_{d_1}, \tilde{\phi}_{d_2}, \dots, \tilde{\phi}_{d_M}](x)}, \quad (8)$$

$$(\phi_m^{[M]}, \phi_n^{[M]}) = \prod_{j=1}^M (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_j}) \cdot h_n \delta_{mn}, \quad (9)$$

$$\tilde{\phi}_v^{[M]}(x) \stackrel{\text{def}}{=} \frac{W[\tilde{\phi}_{d_1}, \tilde{\phi}_{d_2}, \dots, \tilde{\phi}_{d_M}, \tilde{\phi}_v](x)}{W[\tilde{\phi}_{d_1}, \tilde{\phi}_{d_2}, \dots, \tilde{\phi}_{d_M}](x)}, \quad (10)$$

$$U^{[M]}(x) \stackrel{\text{def}}{=} U(x) - 2\partial_x^2 \log |W[\tilde{\phi}_{d_1}, \tilde{\phi}_{d_2}, \dots, \tilde{\phi}_{d_M}](x)|. \quad (11)$$

It should be stressed that the final results after M deletions are independent of the orders of deletions. The algebraic side of the derivation of (6)–(11) is basically the same as that for the Adler–Crum case. The essential difference is the verification of the inheritance at each step of the properties of the virtual states; non-nodeness and the proper boundary behaviours. This is guaranteed by choosing the parameters of the theory g and h large enough, see (23) and (24).

Here we show the process $\mathcal{H} \rightarrow \mathcal{H}^{[1]}$. It is elementary to show

$$\mathcal{H} = \hat{\mathcal{A}}_d^\dagger \hat{\mathcal{A}}_d + \tilde{\mathcal{E}}_d, \quad \hat{\mathcal{A}}_d \stackrel{\text{def}}{=} \frac{d}{dx} - \frac{\partial_x \tilde{\phi}_d(x)}{\tilde{\phi}_d(x)}, \quad \forall d \in \mathcal{V}. \quad (12)$$

Since $\tilde{\phi}_d(x)$ has no node, the operator $\hat{\mathcal{A}}_d$ and $\hat{\mathcal{A}}_d^\dagger$ are regular in the interior. Next we define $\mathcal{H}^{[1]}$ by interchanging the order of these two operators:

$$\mathcal{H}^{[1]} \stackrel{\text{def}}{=} \hat{\mathcal{A}}_d \hat{\mathcal{A}}_d^\dagger + \tilde{\mathcal{E}}_d, \quad U^{[1]}(x) = U(x) - 2\partial_x^2 \log|\tilde{\phi}_d(x)|. \quad (13)$$

It is straightforward to show that \mathcal{H} and $\mathcal{H}^{[1]}$ are iso-spectral

$$\mathcal{H}^{[1]} \phi_n^{[1]}(x) = \mathcal{E}_n \phi_n^{[1]}(x),$$

$$\phi_n^{[1]}(x) \stackrel{\text{def}}{=} \hat{\mathcal{A}}_d \phi_n(x) = \frac{W[\tilde{\phi}_d, \phi_n](x)}{\tilde{\phi}_d(x)} \quad (n \in \mathbb{Z}_{\geq 0}),$$

$$\mathcal{H}^{[1]} \tilde{\phi}_v^{[1]}(x) = \tilde{\mathcal{E}}_v \tilde{\phi}_v^{[1]}(x),$$

$$\tilde{\phi}_v^{[1]}(x) \stackrel{\text{def}}{=} \hat{\mathcal{A}}_d \tilde{\phi}_v(x) = \frac{W[\tilde{\phi}_d, \tilde{\phi}_v](x)}{\tilde{\phi}_d(x)} \quad (v \in \mathcal{V} \setminus \{d\}),$$

$$\phi_n(x) = \frac{\hat{\mathcal{A}}_d^\dagger}{\mathcal{E}_n - \tilde{\mathcal{E}}_d} \phi_n^{[1]}(x),$$

$$(\phi_m^{[1]}, \phi_n^{[1]}) = (\mathcal{E}_m - \tilde{\mathcal{E}}_d)(\phi_m, \phi_n) = (\mathcal{E}_m - \tilde{\mathcal{E}}_d)h_n \delta_{mn}.$$

Obviously the state d is deleted from the virtual state spectrum of $\mathcal{H}^{[1]}$, for $\hat{\mathcal{A}}_d \tilde{\phi}_d(x) = 0$. Since the zero-modes of $\hat{\mathcal{A}}_d$ and $\hat{\mathcal{A}}_d^\dagger$ are $\tilde{\phi}_d(x)$ and $1/\tilde{\phi}_d(x)$, which have infinite norms, the sets of eigenfunctions $\{\phi_n(x)\}$ and $\{\phi_n^{[1]}(x)\}$ are iso-spectral and in one to one correspondence. Thus the new set $\{\phi_n^{[1]}(x)\}$ is complete. If not, there must exist a finite norm vector Ψ in the Hilbert space which is orthogonal to all the basis: $(\Psi, \phi_n^{[1]}) = 0, \forall n \in \mathbb{Z}_{\geq 0}$. This would lead to a contradiction $(\hat{\mathcal{A}}_d^\dagger \Psi, \phi_n) = 0, \forall n \in \mathbb{Z}_{\geq 0}$, since $\hat{\mathcal{A}}_d^\dagger \Psi \neq 0$. Next we show the logic for demonstrating that the virtual state solutions $\{\tilde{\phi}_v^{[1]}(x)\}$ have no node in the interior. By using the Schrödinger equations for them we obtain

$$\partial_x W[\tilde{\phi}_d, \tilde{\phi}_v](x) = (\tilde{\mathcal{E}}_d - \tilde{\mathcal{E}}_v) \tilde{\phi}_d(x) \tilde{\phi}_v(x), \quad (14)$$

which has no node. Since we can verify that $W[\tilde{\phi}_d, \tilde{\phi}_v](x)$ vanishes at one boundary, no-nodeness of $W[\tilde{\phi}_d, \tilde{\phi}_v](x)$ in the interior follows. When the virtual states d and v belong to different types (I and II) the Wronskians might not vanish at both boundaries. In that case we can show that they have the same sign at both boundaries $W[\tilde{\phi}_d, \tilde{\phi}_v](x_1)W[\tilde{\phi}_d, \tilde{\phi}_v](x_2) > 0$. Likewise we can show, for all the explicit examples in the next section that $\tilde{\phi}_v^{[1]}$ and $1/\tilde{\phi}_v^{[1]}$ have infinite norms. The steps going from $\mathcal{H}^{[1]} \rightarrow \mathcal{H}^{[2]}$ and further are essentially the same.

3. Multi-indexed Laguerre and Jacobi polynomials

Here we will apply the method of virtual states deletion to two well-known systems of exactly solvable quantum mechanics with the radial oscillator and Darboux–Pöschl–Teller potentials:

$$U(x) = \begin{cases} x^2 + \frac{g(g-1)}{x^2} - (1+2g), & : \text{L}, \\ x_1 = 0, x_2 = \infty, g > \frac{1}{2} \\ \frac{g(g-1)}{\sin^2 x} + \frac{h(h-1)}{\cos^2 x} - (g+h)^2, & \\ x_1 = 0, x_2 = \frac{\pi}{2}, g, h > \frac{1}{2} & : \text{J}, \end{cases} \quad (15)$$

in which L and J stand for the names of their eigenfunctions, the Laguerre and Jacobi polynomials. For the actual parameter ranges consistent with specific deletions, see (23)–(24). The eigenfunctions are factorised into the groundstate eigenfunction and the polynomial in the sinusoidal coordinate $\eta = \eta(x)$ [31]:

$$\phi_n(x; \lambda) = \phi_0(x; \lambda) P_n(\eta(x); \lambda), \quad (16)$$

in which λ stands for the parameters, g for L and (g, h) for J. Their explicit forms are

$$\text{L: } \phi_0(x; g) \stackrel{\text{def}}{=} e^{-\frac{1}{2}x^2} x^g, \quad \eta(x) \stackrel{\text{def}}{=} x^2,$$

$$P_n(\eta; g) \stackrel{\text{def}}{=} L_n^{(g-\frac{1}{2})}(\eta),$$

$$\mathcal{E}_n(g) \stackrel{\text{def}}{=} 4n, \quad h_n(g) \stackrel{\text{def}}{=} \frac{1}{2n!} \Gamma\left(n + g + \frac{1}{2}\right); \quad (17)$$

$$\text{J: } \phi_0(x; g, h) \stackrel{\text{def}}{=} (\sin x)^g (\cos x)^h, \quad \eta(x) \stackrel{\text{def}}{=} \cos 2x,$$

$$P_n(\eta; g, h) \stackrel{\text{def}}{=} P_n^{(g-\frac{1}{2}, h-\frac{1}{2})}(\eta),$$

$$\mathcal{E}_n(g, h) \stackrel{\text{def}}{=} 4n(n + g + h),$$

$$h_n(g, h) \stackrel{\text{def}}{=} \frac{\Gamma(n + g + \frac{1}{2}) \Gamma(n + h + \frac{1}{2})}{2n!(2n + g + h) \Gamma(n + g + h)}. \quad (18)$$

They have two types of virtual states, which are again polynomial solutions. The total number is finite depending on the parameter values g and h , except for L1 which has infinite. The virtual states wavefunctions for L are:

$$\text{L1: } \tilde{\phi}_v^{\text{I}}(x) \stackrel{\text{def}}{=} e^{\frac{1}{2}x^2} x^g \xi_v^{\text{I}}(\eta(x); g),$$

$$\xi_v^{\text{I}}(\eta; g) \stackrel{\text{def}}{=} P_v(-\eta; g), \quad v \in \mathbb{Z}_{\geq 0},$$

$$\tilde{\mathcal{E}}_v^{\text{I}} \stackrel{\text{def}}{=} -4\left(g + v + \frac{1}{2}\right), \quad \tilde{\delta}^{\text{I}} \stackrel{\text{def}}{=} -1; \quad (19)$$

$$\text{L2: } \tilde{\phi}_v^{\text{II}}(x) \stackrel{\text{def}}{=} e^{-\frac{1}{2}x^2} x^{1-g} \xi_v^{\text{II}}(\eta(x); g),$$

$$\xi_v^{\text{II}}(\eta; g) \stackrel{\text{def}}{=} P_v(\eta; 1 - g), \quad v = 0, 1, \dots, \left[g - \frac{1}{2} \right]',$$

$$\tilde{\mathcal{E}}_v^{\text{II}} \stackrel{\text{def}}{=} -4\left(g - v - \frac{1}{2}\right), \quad \tilde{\delta}^{\text{II}} \stackrel{\text{def}}{=} 1, \quad (20)$$

in which $[a]'$ denotes the greatest integer less than a and $\tilde{\delta}^{\text{I,II}}$ will be used later. For no-nodeness of $\xi_v^{\text{I,II}}$, see (2.39) of [22]. The virtual states wavefunctions for J are:

$$\text{J1: } \tilde{\phi}_v^{\text{I}}(x) \stackrel{\text{def}}{=} (\sin x)^g (\cos x)^{1-h} \xi_v^{\text{I}}(\eta(x); g, h),$$

$$\xi_v^{\text{I}}(\eta; g, h) \stackrel{\text{def}}{=} P_v(\eta; g, 1 - h), \quad v = 0, 1, \dots, \left[h - \frac{1}{2} \right]',$$

$$\tilde{\mathcal{E}}_v^{\text{I}} \stackrel{\text{def}}{=} -4\left(g + v + \frac{1}{2}\right)\left(h - v - \frac{1}{2}\right), \quad \tilde{\delta}^{\text{I}} \stackrel{\text{def}}{=} (-1, 1); \quad (21)$$

$$\text{J2: } \tilde{\phi}_v^{\text{II}}(x) \stackrel{\text{def}}{=} (\sin x)^{1-g} (\cos x)^h \xi_v^{\text{II}}(\eta(x); g, h),$$

$$\xi_v^{\text{II}}(\eta; g, h) \stackrel{\text{def}}{=} P_v(\eta; 1 - g, h), \quad v = 0, 1, \dots, \left[g - \frac{1}{2} \right]',$$

$$\tilde{\mathcal{E}}_v^{\text{II}} \stackrel{\text{def}}{=} -4\left(g - v - \frac{1}{2}\right)\left(h + v + \frac{1}{2}\right), \quad \tilde{\delta}^{\text{II}} \stackrel{\text{def}}{=} (1, -1). \quad (22)$$

The larger the parameter g and/or h become, the more the virtual states are ‘created’. The L1 system is obtained from the J1 in the limit $h \rightarrow \infty$ [5]. This explains the infinitely many virtual states of L1. Let us denote by $\mathcal{V}^{\text{I,II}}$ the index sets of the virtual states of types I and II for L and J. Due to the parity property of the Jacobi polynomial $P_n^{(\alpha, \beta)}(-x) = (-1)^n P_n^{(\beta, \alpha)}(x)$, the two virtual state polynomials ξ_v^{I} and ξ_v^{II} for J are related by $\xi_v^{\text{II}}(-\eta; g, h) = (-1)^v \xi_v^{\text{I}}(\eta; g, h)$. For no-nodeness of $\xi_v^{\text{I,II}}$, see (2.40) of [22] and (3.2) of [6]. It is also obvious that the virtual state wavefunctions $\tilde{\phi}_v$ and their reciprocals $1/\tilde{\phi}_v$ are not square integrable in all four types. There is a third type of negative energy




Fig. 1. Schematic picture of virtual states deletion. The black circles denote eigenstates. The down and up triangles denote virtual states of types I and II. The deleted virtual states are denoted by white triangles.

polynomial solutions for both L and J. For J, the degree n solution is $(\sin x)^{1-g}(\cos x)^{1-h}P_n^{(-g+\frac{1}{2}, -h+\frac{1}{2})}(\eta)$. Even when they have no node in the interior, their reciprocals are square integrable and they cannot be used for the virtual states deletion. The label 0 is special in that the wavefunctions satisfy $\tilde{\phi}_0^I(x; \lambda)\tilde{\phi}_0^{II}(x; \lambda) = c_{\mathcal{F}}^{-1} \frac{d\eta(x)}{dx}$ since $\xi_0^{I,II} = 1$. Here the constant $c_{\mathcal{F}} = 2$ for L and $c_{\mathcal{F}} = -4$ for J. We will not use the label 0 states for deletion.

Since there are two types of virtual states available, the general deletion is specified by the set of $M + N$ positive integers $\mathcal{D} \stackrel{\text{def}}{=} \{d_1^I, \dots, d_M^I, d_1^{II}, \dots, d_N^{II}\}$, $d_j^{I,II} \geq 1$, which are the degrees of the deleted virtual state wavefunctions. Note that the subcases of either $M = 0$ or $N = 0$ are meaningful. In order to accommodate all these virtual states, the parameters g and h must be larger than certain bounds:

$$\text{L: } g > \max\left\{N + \frac{3}{2}, d_j^{II} + \frac{1}{2}\right\}, \quad (23)$$

$$\text{J: } g > \max\left\{N + 2, d_j^{II} + \frac{1}{2}\right\},$$

$$h > \max\left\{M + 2, d_j^I + \frac{1}{2}\right\}. \quad (24)$$

See Fig. 1 for the schematic structure of virtual states deletion.

Like the original eigenfunctions (16) the n -th eigenfunction $\phi_{\mathcal{D},n}(x; \lambda) \equiv \phi_n^{[M,N]}(x)$ after the deletion (9) can be clearly factorised into an x -dependent part, the common denominator polynomial $\mathcal{E}_{\mathcal{D}}$ in η and the multi-indexed polynomial $P_{\mathcal{D},n}$ in η :

$$\begin{aligned} \phi_n^{[M,N]}(x) &\equiv \phi_{\mathcal{D},n}(x; \lambda) = c_{\mathcal{F}}^{M+N} \psi_{\mathcal{D}}(x; \lambda) P_{\mathcal{D},n}(\eta(x); \lambda), \\ \psi_{\mathcal{D}}(x; \lambda) &\stackrel{\text{def}}{=} \frac{\phi_0(x; \lambda^{[M,N]})}{\mathcal{E}_{\mathcal{D}}(\eta(x); \lambda)}, \end{aligned} \quad (25)$$

in which $\phi_0(x; \lambda)$ is the groundstate wavefunction (17)–(18). Here the shifted parameters $\lambda^{[M,N]}$ after the $[M, N]$ deletion are $\lambda^{[M,N]} \stackrel{\text{def}}{=} \lambda - M\delta^I - N\delta^{II}$, explicitly they are

$$\lambda^{[M,N]} = \begin{cases} g + M - N & : \text{L}, \\ (g + M - N, h - M + N) & : \text{J}. \end{cases} \quad (26)$$

Needless to say, the denominator polynomial $\mathcal{E}_{\mathcal{D}}$ has no node in the interior. The polynomials $P_{\mathcal{D},n}$ and $\mathcal{E}_{\mathcal{D}}$ are expressed in terms of Wronskians of the variable η :

$$\begin{aligned} P_{\mathcal{D},n}(\eta; \lambda) &\stackrel{\text{def}}{=} W[\mu_1, \dots, \mu_M, \nu_1, \dots, \nu_N, P_n](\eta) \\ &\times \begin{cases} e^{-M\eta} \eta^{(M+g+\frac{1}{2})N} & : \text{L}, \\ \left(\frac{1-\eta}{2}\right)^{(M+g+\frac{1}{2})N} \left(\frac{1+\eta}{2}\right)^{(N+h+\frac{1}{2})M} & : \text{J}, \end{cases} \end{aligned} \quad (27)$$

$$\begin{aligned} \mathcal{E}_{\mathcal{D}}(\eta; \lambda) &\stackrel{\text{def}}{=} W[\mu_1, \dots, \mu_M, \nu_1, \dots, \nu_N](\eta) \\ &\times \begin{cases} e^{-M\eta} \eta^{(M+g-\frac{1}{2})N} & : \text{L}, \\ \left(\frac{1-\eta}{2}\right)^{(M+g-\frac{1}{2})N} \left(\frac{1+\eta}{2}\right)^{(N+h-\frac{1}{2})M} & : \text{J}, \end{cases} \end{aligned} \quad (28)$$

$$\begin{aligned} \mu_j &= \begin{cases} e^{\eta} \xi_{d_j^I}^I(\eta; g) & : \text{L}, \\ \left(\frac{1+\eta}{2}\right)^{\frac{1}{2}-h} \xi_{d_j^I}^I(\eta; g, h) & : \text{J}, \end{cases} \\ \nu_j &= \begin{cases} \eta^{\frac{1}{2}-g} \xi_{d_j^{II}}^{II}(\eta; g) & : \text{L}, \\ \left(\frac{1-\eta}{2}\right)^{\frac{1}{2}-g} \xi_{d_j^{II}}^{II}(\eta; g, h) & : \text{J}, \end{cases} \end{aligned} \quad (29)$$

in which P_n in (27) denotes the original polynomial, $P_n(\eta; g)$ for L and $P_n(\eta; g, h)$ for J. These polynomials depend on the order of deletions only through the sign of permutation. The multi-indexed polynomial $P_{\mathcal{D},n}$ is of degree $\ell + n$ and the denominator polynomial $\mathcal{E}_{\mathcal{D}}$ is of degree ℓ in η , in which ℓ is given by

$$\ell \stackrel{\text{def}}{=} \sum_{j=1}^M d_j^I + \sum_{j=1}^N d_j^{II} - \frac{1}{2}M(M-1) - \frac{1}{2}N(N-1) + MN \geq 1. \quad (30)$$

Here the label n specifies the energy eigenvalue \mathcal{E}_n of $\phi_{\mathcal{D},n}$. Hence it also counts the nodes due to the oscillation theorem. Although they do not satisfy the three term recurrence relations, the multi-indexed polynomials $\{P_{\mathcal{D},n}\}$ form a complete set of orthogonal polynomials with the orthogonality relations:

$$\begin{aligned} \int d\eta \frac{W(\eta; \lambda^{[M,N]})}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)^2} P_{\mathcal{D},m}(\eta; \lambda) P_{\mathcal{D},n}(\eta; \lambda) \\ = h_n(\lambda) \delta_{nm} \\ \times \begin{cases} \prod_{j=1}^M (n+g+d_j^I + \frac{1}{2}) \cdot \prod_{j=1}^N (n+g-d_j^{II} - \frac{1}{2}) & : \text{L}, \\ 4^{-M-N} \prod_{j=1}^M (n+g+d_j^I + \frac{1}{2})(n+h-d_j^I - \frac{1}{2}) \\ \times \prod_{j=1}^N (n+g-d_j^{II} - \frac{1}{2})(n+h+d_j^{II} + \frac{1}{2}) & : \text{J}, \end{cases} \end{aligned} \quad (31)$$

where the weight function of the original polynomials $W(\eta; \lambda) d\eta = \phi_0(x; \lambda)^2 dx$ reads explicitly

$$W(\eta; \lambda) \stackrel{\text{def}}{=} \begin{cases} \frac{1}{2} e^{-\eta} \eta^{g-\frac{1}{2}} & : \text{L}, \\ \frac{1}{2^{g+h+1}} (1-\eta)^{g-\frac{1}{2}} (1+\eta)^{h-\frac{1}{2}} & : \text{J}. \end{cases} \quad (32)$$

This is obtained by rewriting the orthogonality relations of the eigenfunctions (9) after the $[M, N]$ deletion.

In the rest of this section we explore various properties of the new multi-indexed polynomials $\{P_{\mathcal{D},n}\}$. As for the exceptional orthogonal polynomials [3–5, 21, 20, 7], the lowest degree polynomial $P_{\mathcal{D},0}(\eta; \lambda)$ is related to the denominator polynomial $\mathcal{E}_{\mathcal{D}}(\eta; \lambda)$ by the parameter shift $\lambda \rightarrow \lambda + \delta$ ($\delta = 1$ for L and $\delta = (1, 1)$ for J):

$$\begin{aligned} P_{\mathcal{D},0}(\eta; \lambda) &= \mathcal{E}_{\mathcal{D}}(\eta; \lambda + \delta) \\ &\times \begin{cases} (-1)^M \prod_{j=1}^N (g-d_j^{II} - \frac{1}{2}) & : \text{L}, \\ 2^{-M} \prod_{j=1}^M (h-d_j^I - \frac{1}{2}) \\ \times (-2)^{-N} \prod_{j=1}^N (g-d_j^{II} - \frac{1}{2}) & : \text{J}. \end{cases} \end{aligned} \quad (33)$$

The virtual state wavefunction with the index $v \in \mathcal{D}$ leading to the final $[M, N]$ deletion is $\tilde{\phi}_v^{[M', N']}$ (10) with the index set $\mathcal{D}' \stackrel{\text{def}}{=} \mathcal{D} \setminus \{v\}$, in which $[M', N'] = [M - 1, N]$ if $v \in \mathcal{V}^I$, and $[M, N - 1]$ if $v \in \mathcal{V}^{II}$. It has the following form:

$$\begin{aligned} &\tilde{\phi}_v^{[M', N']}(x; \lambda) \\ &= c_{\mathcal{F}}^{M+N-1} \frac{\mathcal{E}_{\mathcal{D}}(\eta(x); \lambda)}{\mathcal{E}_{\mathcal{D}'}(\eta(x); \lambda)} \\ &\times \begin{cases} (-1)^N \phi_0(ix; \lambda^{[M-1, N]}) i^{-(g+M-N-1)}, & v \in \mathcal{V}^I : \text{L}, \\ \phi_0(x; t^{II}(\lambda^{[M, N-1]})), & v \in \mathcal{V}^{II} : \text{L}, \\ (-1)^N \phi_0(x; t^I(\lambda^{[M-1, N]})), & v \in \mathcal{V}^I : \text{J}, \\ \phi_0(x; t^{II}(\lambda^{[M, N-1]})), & v \in \mathcal{V}^{II} : \text{J}. \end{cases} \quad (34) \end{aligned}$$

Here the twist operator t acting on the parameters is defined as $t^{II}(g) = 1 - g$ for L, $t^I(g, h) = (g, 1 - h)$ and $t^{II}(g, h) = (1 - g, h)$ for J. Note that the virtual states (20)–(22) have the form $\phi_v(x; \lambda) = \phi_v(x; t(\lambda))$. These relations are essential for the determination of the lower bounds of the parameters given in (23)–(24), which guarantee that the virtual state properties are correctly inherited at each step of deletion.

The Hamiltonian $\mathcal{H}_{\mathcal{D}}(\lambda) \equiv \mathcal{H}^{[M, N]}$ of the $[M, N]$ deleted system can be expressed in terms of its groundstate eigenfunction $\phi_{\mathcal{D}, 0}(x; \lambda) \equiv \phi_0^{[M, N]}(x; \lambda)$ with the help of (33):

$$\mathcal{H}_{\mathcal{D}}(\lambda) = \mathcal{A}_{\mathcal{D}}(\lambda)^\dagger \mathcal{A}_{\mathcal{D}}(\lambda), \quad (35)$$

$$\mathcal{A}_{\mathcal{D}}(\lambda) \stackrel{\text{def}}{=} \frac{d}{dx} - \frac{\partial_x \phi_{\mathcal{D}, 0}(x; \lambda)}{\phi_{\mathcal{D}, 0}(x; \lambda)},$$

$$\phi_{\mathcal{D}, 0}(x; \lambda) \propto \phi_0(x; \lambda^{[M, N]}) \frac{\mathcal{E}_{\mathcal{D}}(\eta(x); \lambda + \delta)}{\mathcal{E}_{\mathcal{D}}(\eta(x); \lambda)}. \quad (36)$$

This has the same form as the various Hamiltonians of the exceptional orthogonal polynomials including those of the discrete quantum mechanics [3–5,7]. Reflecting the construction [19,6] it is shape invariant [14,3,22]

$$\mathcal{A}_{\mathcal{D}}(\lambda) \mathcal{A}_{\mathcal{D}}(\lambda)^\dagger = \mathcal{A}_{\mathcal{D}}(\lambda + \delta)^\dagger \mathcal{A}_{\mathcal{D}}(\lambda + \delta) + \mathcal{E}_1(\lambda). \quad (37)$$

This means that the operators $\mathcal{A}_{\mathcal{D}}(\lambda)$ and $\mathcal{A}_{\mathcal{D}}(\lambda)^\dagger$ relate the eigenfunctions of neighbouring degrees and parameters:

$$\mathcal{A}_{\mathcal{D}}(\lambda) \phi_{\mathcal{D}, n}(x; \lambda) = f_n(\lambda) \phi_{\mathcal{D}, n-1}(x; \lambda + \delta), \quad (38)$$

$$\mathcal{A}_{\mathcal{D}}(\lambda)^\dagger \phi_{\mathcal{D}, n-1}(x; \lambda + \delta) = b_{n-1}(\lambda) \phi_{\mathcal{D}, n}(x; \lambda), \quad (39)$$

in which the constants $f_n(\lambda)$ and $b_{n-1}(\lambda)$ are the factors of the eigenvalue $f_n(\lambda) b_{n-1}(\lambda) = \mathcal{E}_n(\lambda)$:

$$f_n(\lambda) = \begin{cases} -2 & : \text{L}, \\ -2(n + g + h) & : \text{J}, \end{cases} \quad b_{n-1}(\lambda) = -2n \quad : \text{L, J}.$$

The forward and backward shift operators are defined by

$$\mathcal{F}_{\mathcal{D}}(\lambda) \stackrel{\text{def}}{=} \psi_{\mathcal{D}}(x; \lambda + \delta)^{-1} \circ \mathcal{A}_{\mathcal{D}}(\lambda) \circ \psi_{\mathcal{D}}(x; \lambda) \quad (40)$$

$$= c_{\mathcal{F}} \frac{\mathcal{E}_{\mathcal{D}}(\eta; \lambda + \delta)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)} \left(\frac{d}{d\eta} - \frac{\partial_{\eta} \mathcal{E}_{\mathcal{D}}(\eta; \lambda + \delta)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda + \delta)} \right), \quad (41)$$

$$\mathcal{B}_{\mathcal{D}}(\lambda) \stackrel{\text{def}}{=} \psi_{\mathcal{D}}(x; \lambda)^{-1} \circ \mathcal{A}_{\mathcal{D}}(\lambda)^\dagger \circ \psi_{\mathcal{D}}(x; \lambda + \delta) \quad (42)$$

$$= -4c_{\mathcal{F}}^{-1} c_2(\eta) \frac{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda + \delta)} \times \left(\frac{d}{d\eta} + \frac{c_1(\eta, \lambda^{[M, N]})}{c_2(\eta)} - \frac{\partial_{\eta} \mathcal{E}_{\mathcal{D}}(\eta; \lambda)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)} \right), \quad (43)$$

in which the functions $c_1(\eta; \lambda)$ and $c_2(\eta)$ are those appearing in the (confluent) hypergeometric equations for the Laguerre and Jacobi polynomials

$$\text{L: } c_1(\eta, \lambda) \stackrel{\text{def}}{=} g + \frac{1}{2} - \eta, \quad c_2(\eta) \stackrel{\text{def}}{=} \eta,$$

$$\text{J: } c_1(\eta, \lambda) \stackrel{\text{def}}{=} h - g - (g + h + 1)\eta, \quad c_2(\eta) \stackrel{\text{def}}{=} 1 - \eta^2.$$

Their action on the multi-indexed polynomials $P_{\mathcal{D}, n}(\eta; \lambda)$ is

$$\mathcal{F}_{\mathcal{D}}(\lambda) P_{\mathcal{D}, n}(\eta; \lambda) = f_n(\lambda) P_{\mathcal{D}, n-1}(\eta; \lambda + \delta), \quad (44)$$

$$\mathcal{B}_{\mathcal{D}}(\lambda) P_{\mathcal{D}, n-1}(\eta; \lambda + \delta) = b_{n-1}(\lambda) P_{\mathcal{D}, n}(\eta; \lambda). \quad (45)$$

The second order differential operator $\tilde{\mathcal{H}}_{\mathcal{D}}(\lambda)$ governing the multi-indexed polynomials is:

$$\begin{aligned} \tilde{\mathcal{H}}_{\mathcal{D}}(\lambda) &\stackrel{\text{def}}{=} \psi_{\mathcal{D}}(x; \lambda)^{-1} \circ \mathcal{H}_{\mathcal{D}}(\lambda) \circ \psi_{\mathcal{D}}(x; \lambda) = \mathcal{B}_{\mathcal{D}}(\lambda) \mathcal{F}_{\mathcal{D}}(\lambda) \\ &= -4 \left(c_2(\eta) \frac{d^2}{d\eta^2} + \left(c_1(\eta, \lambda^{[M, N]}) \right. \right. \\ &\quad \left. \left. - 2c_2(\eta) \frac{\partial_{\eta} \mathcal{E}_{\mathcal{D}}(\eta; \lambda)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)} \right) \frac{d}{d\eta} \right. \\ &\quad \left. + c_2(\eta) \frac{\partial_{\eta}^2 \mathcal{E}_{\mathcal{D}}(\eta; \lambda)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)} \right. \\ &\quad \left. - c_1(\eta, \lambda^{[M, N]} - \delta) \frac{\partial_{\eta} \mathcal{E}_{\mathcal{D}}(\eta; \lambda)}{\mathcal{E}_{\mathcal{D}}(\eta; \lambda)} \right), \quad (46) \end{aligned}$$

$$\tilde{\mathcal{H}}_{\mathcal{D}}(\lambda) P_{\mathcal{D}, n}(\eta; \lambda) = \mathcal{E}_n(\lambda) P_{\mathcal{D}, n}(\eta; \lambda). \quad (47)$$

Since all the zeros of $\mathcal{E}_{\mathcal{D}}(\eta; \lambda)$ are simple, (47) is a Fuchsian differential equation for the J case. The characteristic exponents at the zeros of $\mathcal{E}_{\mathcal{D}}(\eta; \lambda)$ are the same everywhere, 0 and 3. The multi-indexed polynomials $\{P_{\mathcal{D}, n}(\eta; \lambda)\}$ provide infinitely many global solutions of the above Fuchsian equation (47) with $3 + \ell$ regular singularities for the J case [21]. The L case is obtained in a confluent limit. These situations are basically the same as those of the exceptional polynomials.

Although we have restricted $d_j^{I, II} \geq 1$, there is no obstruction for deletion of $d_j^{I, II} = 0$. In terms of the multi-indexed polynomial (27), the level 0 deletions imply the following:

$$\begin{aligned} P_{\mathcal{D}, n}(\eta; \lambda) |_{d_M^I=0} &= P_{\mathcal{D}', n}(\eta; \lambda - \delta^I) \times A, \\ \mathcal{D}' &= \{d_1^I - 1, \dots, d_{M-1}^I - 1, d_1^{II} + 1, \dots, d_N^{II} + 1\}, \quad (48) \end{aligned}$$

$$\begin{aligned} P_{\mathcal{D}, n}(\eta; \lambda) |_{d_M^{II}=0} &= P_{\mathcal{D}', n}(\eta; \lambda - \delta^{II}) \times B, \\ \mathcal{D}' &= \{d_1^I + 1, \dots, d_M^I + 1, d_1^{II} - 1, \dots, d_{N-1}^{II} - 1\}, \quad (49) \end{aligned}$$

where the multiplicative factors A and B are

$$A = \begin{cases} (-1)^M \prod_{j=1}^N (d_j^{II} + 1) & : \text{L}, \\ -(-2)^{-M} \prod_{j=1}^{M-1} (g - h + d_j^I + 1) \\ \quad \times (-2)^{-N} \prod_{j=1}^N (d_j^{II} + 1) \cdot (n + h - \frac{1}{2}) & : \text{J}, \end{cases}$$

$$B = \begin{cases} (-1)^M \prod_{j=1}^M (d_j^I + 1) \cdot (n + g - \frac{1}{2}) & : \text{L}, \\ 2^{-M} \prod_{j=1}^M (d_j^I + 1) \cdot (-2)^{-N} \\ \quad \times \prod_{j=1}^{N-1} (h - g + d_j^{II} + 1) \cdot (n + g - \frac{1}{2}) & : \text{J}. \end{cases}$$

From (33), $\mathcal{E}_{\mathcal{D}}$ behaves similarly. Therefore including the level 0 deletion corresponds to $M + N - 1$ virtual states deletions. This is why we have restricted $d_j^{I, II} \geq 1$.

These relations (48)–(49) can be used for studying the equivalence of $\mathcal{H}_{\mathcal{D}}$. It should be stressed that the same polynomials can have different but equivalent sets of multi-indices \mathcal{D} , which defines the denominator polynomial $\mathcal{E}_{\mathcal{D}}$. Since the overall scale of $\mathcal{E}_{\mathcal{D}}$ is irrelevant (see (36)) for the multi-indexed polynomial $\{P_{\mathcal{D}, n}\}$, we consider the equivalence class of $\mathcal{E}_{\mathcal{D}}$. For example, it

is straightforward to verify the following equivalence and their duals $I \leftrightarrow II$:

$$\mathcal{E}_{\{1^1, \dots, k^1\}}(\eta; \lambda - \tilde{\delta}^{II}) \sim \mathcal{E}_{\{k^{II}\}}(\eta; \lambda - k\tilde{\delta}^I) \quad (k \geq 1), \quad (50)$$

$$\begin{aligned} & \mathcal{E}_{\{m^I, \dots, (k+m)^I\}}(\eta; \lambda - m\tilde{\delta}^{II}) \\ & \sim \mathcal{E}_{\{(k+1)^{II}, \dots, (k+m)^{II}\}}(\eta; \lambda - (k+1)\tilde{\delta}^I) \quad (k \geq 1). \end{aligned} \quad (51)$$

The simplest, $k = 1$, of (50) corresponds to the well-known fact that the X_1 types I and II polynomials are identical for L and J [5]. Deleting the first k consecutive excited states, like the l.h.s. of (50) has led to many interesting phenomena in the ordinary and discrete quantum mechanics [30,33,32]. Classification of the equivalent classes leading to the same polynomials is a challenging future problem.

The exceptional X_ℓ orthogonal polynomials of types I and II [12, 13,3,5,21,19] correspond to the simplest cases of one virtual state deletion of that type, $\mathcal{D} = \{\ell^I\}$ or $\{\ell^{II}\}$, $\ell \geq 1$:

$$\xi_\ell(\eta; \lambda) = \mathcal{E}_{\mathcal{D}}(\eta; \lambda + \ell\tilde{\delta} + \tilde{\delta}), \quad (52)$$

$$P_{\ell,n}(\eta; \lambda) = P_{\mathcal{D},n}(\eta; \lambda + \ell\tilde{\delta} + \tilde{\delta}) \times A, \quad (53)$$

where $\tilde{\delta} = \tilde{\delta}^{I,II}$ and the multiplicative factor A is $A = -1$ for XL1, $(n+g+\frac{1}{2})^{-1}$ for XL2, $2(n+h+\frac{1}{2})^{-1}$ for XJ1 and $-2(n+g+\frac{1}{2})^{-1}$ for XJ2. Most formulas between (35) and (47) look almost the same as those appearing in the theory of the exceptional orthogonal polynomials [3–5,21,19,20,6,7].

In a recent paper [24] Gómez-Ullate et al. discussed “two-step Darboux transformations”, which corresponds to the examples of $\mathcal{D} = \{m_1^{II}, m_2^{II}\}$ for L in our scheme. This paper stimulated the present work.

4. Summary and comments

This is a first short report on the discovery of infinite families of multi-indexed orthogonal polynomials. They are obtained as the main part of exactly solvable quantum mechanical systems, which are deformations of the radial oscillator and the Darboux–Pöschl–Teller potentials. Although these new polynomials start at degree ℓ (≥ 1), they form a complete set of orthogonal polynomials. The exactly solvable modified Hamiltonian and the eigenfunctions are obtained by applying the modified Crum’s theorem due to Adler to the original system to delete a finitely many virtual state solutions of types I and II. The simplest case of one virtual state deletion reproduces the exceptional orthogonal polynomials [12,13,3,5,21]. For the Jacobi case, the new polynomials provide infinitely many examples of global solutions of second order Fuchsian differential equations with $3+\ell$ regular singularities [21]. Like the undeformed theory, the Laguerre results can be obtained from the Jacobi results in certain confluence limits. But we presented all results in parallel, for better understanding of the structure. The parameter ranges (23) and (24) are conservative sufficient conditions with which the proper boundary behaviours and the completeness are guaranteed.

As a first report, only the basic results are presented. Some important issues cannot be included due to space restrictions, for example, an alternative proof of shape invariance, the action of various intertwining operators, etc. Many important problems are remaining to be clarified. For example, the bispectrality [19], generating functions [21], properties of the zeros of the denominator polynomial $\mathcal{E}_{\mathcal{D}}$ and the extra zeros of the multi-indexed polynomials $\{P_{\mathcal{D},n}\}$ [23], classification of the equivalent classes leading to the same new polynomials.

The exceptional Wilson, Askey–Wilson, Racah and q -Racah polynomials were constructed in the framework of discrete quantum mechanics [4,20,7,8,34,35], and the Crum–Adler theorem for

discrete quantum mechanics were also presented [33,30,32]. The method of virtual states deletion presented in this Letter is applicable to discrete quantum mechanics and it is easy to write down general formulas like (6)–(11). This gives multi-indexed orthogonal polynomials corresponding to the deformations of the Wilson, Askey–Wilson, Racah and q -Racah polynomials. In concrete examples, more than two types of virtual state solutions are available and much richer structures are expected. We will report on these topics elsewhere.

Before closing this Letter, let us make two small comments. The original Schrödinger equation (1) has a general solution for an arbitrary energy \mathcal{E} , $\mathcal{H}\psi(x) = \mathcal{E}\psi(x)$. If we define the corresponding $[M, N]$ deletion solution

$$\psi^{[M,N]}(x) \stackrel{\text{def}}{=} \frac{W[\tilde{\phi}_{d_1}^I, \dots, \tilde{\phi}_{d_M}^I, \tilde{\phi}_{d_1}^{II}, \dots, \tilde{\phi}_{d_N}^{II}, \psi](x)}{W[\tilde{\phi}_{d_1}^I, \dots, \tilde{\phi}_{d_M}^I, \tilde{\phi}_{d_1}^{II}, \dots, \tilde{\phi}_{d_N}^{II}](x)},$$

it solves the deformed Schrödinger equation with the same energy, too

$$\mathcal{H}^{[M,N]}\psi^{[M,N]}(x) = \mathcal{E}\psi^{[M,N]}(x).$$

Among the multi-indices, we can include a finite number of continuous real parameters as labelling the type I virtual solutions as described in [6,16,17]. The resulting orthogonal functions are no longer polynomials, but most properties of the multi-indexed polynomials are shared by these new orthogonal functions.

Acknowledgements

We thank T. Oshima and Y. Haraoka for useful discussion on Fuchsian differential equations. R.S. is supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), No. 22540186.

References

- [1] L. Infeld, T.E. Hull, Rev. Mod. Phys. 23 (1951) 21.
- [2] F. Cooper, A. Khare, U. Sukhatme, Phys. Rep. 251 (1995) 267.
- [3] S. Odake, R. Sasaki, Phys. Lett. B 679 (2009) 414.
- [4] S. Odake, R. Sasaki, Phys. Lett. B 682 (2009) 130.
- [5] S. Odake, R. Sasaki, Phys. Lett. B 684 (2010) 173. (Remark: J1(J2) in this reference corresponds to J2(J1) in later references.)
- [6] S. Odake, R. Sasaki, J. Phys. A 44 (2011) 195203 (14 pp.).
- [7] S. Odake, R. Sasaki, Prog. Theor. Phys. 125 (2011) 851.
- [8] S. Odake, R. Sasaki, arXiv:1104.0473 [math-ph].
- [9] M.M. Crum, Quart. J. Math. Oxford Ser. (2) 6 (1955) 121.
- [10] M.G. Krein, Dokl. Acad. Nauk SSSR 113 (1957) 970; V.É. Adler, Theor. Math. Phys. 101 (1994) 1381.
- [11] S. Bochner, Math. Zeit. 29 (1929) 730.
- [12] D. Gómez-Ullate, N. Kamran, R. Milson, J. Approx Theory 162 (2010) 987; D. Gómez-Ullate, N. Kamran, R. Milson, J. Math. Anal. Appl. 359 (2009) 352.
- [13] C. Quesne, J. Phys. A 41 (2008) 392001; B. Bagchi, C. Quesne, R. Roychoudhury, Pramana J. Phys. 73 (2009) 337; C. Quesne, SIGMA 5 (2009) 084.
- [14] L.E. Gendenshtein, JETP Lett. 38 (1983) 356.
- [15] G. Darboux, Théorie générale des surfaces, vol. 2, Gauthier-Villars, Paris, 1888.
- [16] G. Junker, P. Roy, Phys. Lett. A 232 (1997) 155; G. Junker, P. Roy, Ann. Phys. 270 (1998) 155.
- [17] D. Dutta, P. Roy, J. Math. Phys. 51 (2010) 042101 (9 pp.).
- [18] D. Gómez-Ullate, N. Kamran, R. Milson, J. Phys. A 43 (2010) 434016 (16 pp.); arXiv:1101.5584 [math-ph].
- [19] R. Sasaki, S. Tsujimoto, A. Zhedanov, J. Phys. A 43 (2010) 315204 (20 pp.).
- [20] S. Odake, R. Sasaki, J. Phys. A 43 (2010) 335201 (18 pp.).
- [21] C.-L. Ho, S. Odake, R. Sasaki, arXiv:0912.5447 [math-ph].
- [22] S. Odake, R. Sasaki, J. Math. Phys. 51 (2010) 053513 (9 pp.).
- [23] C.-L. Ho, R. Sasaki, arXiv:1102.5669 [math-ph].
- [24] D. Gómez-Ullate, N. Kamran, R. Milson, arXiv:1103.5724 [math-ph].
- [25] C.-L. Ho, arXiv:1104.3511 [math-ph].
- [26] Y. Grandati, arXiv:1101.0055 [math-ph]; Y. Grandati, arXiv:1103.5023 [math-ph].

- [27] B. Midya, B. Roy, *Phys. Lett. A* 373 (2009) 4117.
- [28] C.-L. Ho, *Ann. Phys.* 326 (2011) 797.
- [29] A.A. Andrianov, M.V. Ioffe, V.P. Spiridonov, *Phys. Lett. A* 174 (1993) 273;
H. Aoyama, M. Sato, T. Tanaka, *Phys. Lett. B* 503 (2001) 423;
C.D.J. Fernández, V. Hussin, *J. Phys. A* 32 (1999) 3603;
V.G. Bagrov, B.F. Samsonov, *Phys. Lett. A* 210 (1996) 60;
F. Cannata, M.V. Ioffe, A.I. Neelov, D.N. Nishnianidze, *J. Phys. A* 37 (2004) 10339.
- [30] L. García-Gutiérrez, S. Odake, R. Sasaki, *Prog. Theor. Phys.* 124 (2010) 1.
- [31] S. Odake, R. Sasaki, *J. Math. Phys.* 47 (2006) 102102 (33 pp.);
S. Odake, R. Sasaki, *Phys. Lett. B* 641 (2006) 112.
- [32] S. Odake, R. Sasaki, *Prog. Theor. Phys.* 126 (2011) 1, doi:10.1143/PTP.126.1, arXiv:1101.5468 [math-ph].
- [33] S. Odake, R. Sasaki, *Prog. Theor. Phys.* 122 (2009) 1067.
- [34] S. Odake, R. Sasaki, *J. Math. Phys.* 49 (2008) 053503 (43 pp.).
- [35] S. Odake, R. Sasaki, *Prog. Theor. Phys.* 119 (2008) 663.