



# Mitochondrial d-loop variation, coat colour and sex identification of Late Iron Age horses in Switzerland



Julia Elsner<sup>a,\*</sup>, Sabine Deschler-Erb<sup>a</sup>, Barbara Stopp<sup>a</sup>, Michael Hofreiter<sup>b</sup>, Jörg Schibler<sup>a</sup>, Angela Schlumbaum<sup>a</sup>

<sup>a</sup> Integrative Prehistory and Archaeological Science, Department of Environmental Sciences, Basel University, Spalenring 145, 4055 Basel, Switzerland

<sup>b</sup> Institute for Biochemistry and Biology, University of Potsdam, Karl-Liebknecht-Str. 24–25, D-14476 Potsdam, Germany

## ARTICLE INFO

### Article history:

Received 26 November 2015

Accepted 7 March 2016

Available online 14 March 2016

### Keywords:

Genetic diversity

Ancient DNA

Iron Age

Horse

Coat colour

Withers height

Greying

Sex identification

## ABSTRACT

In the Celtic world, horses enjoyed a prominent position as status symbols and objects of veneration, yet little is known about these Celtic horses except that they were rather small. The Late Iron Age was a time defined by increasing inter-cultural contact between Celtic peoples and the Romans. This is, amongst other features, observable in the phenotypes of domestic livestock such as horses. Amongst the usually small animals, larger ones are rarely but regularly encountered in the archaeological record. We have investigated mitochondrial (mt) DNA d-loop diversity, sex and coat colour using bones from 34 horses of different size from three Swiss sites (Mormont, Basel-Gasfabrik, Aventicum) most of them dating from 150 to 50 BCE. The aim was to characterise the diversity of matrilineages and coat colourations of Iron Age horses, and to identify molecular sex. We detected eleven mt haplotypes clustering into six haplogroups (B, D, F, I, X2, X3) in the ancient dataset ( $n = 19$ ). Large individuals were all male, but smaller stallions were also identified; molecular sexing confirmed and augmented to morphological results. The horses were bay, chestnut and black in colour, and spottings or dilutions were absent in all animals. With a simplified primer system to detect premature greying, white coats can be excluded as well. The limited colour range proposes selection for monochrome animals. Additionally, ancient matrilineages were compared to modern horses from regions appertaining to the Late Roman Republic and to European pony breeds. Based on Principal Component Analysis (haplotype frequencies) and  $F_{ST}$ -values (genetic distances) the mtDNA variation of the Iron Age horses investigated here has survived in modern European breeds, particularly in northern European ponies.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Several Late Iron Age sites have been identified in Switzerland, including the eponymous sacrificial site in La Tène, canton Neuchâtel. In recent years comprehensive interdisciplinary investigations were carried out at Basel-Gasfabrik (Pichler et al., in preparation) and on the hill of Mormont (Brunetti et al., 2014) to gain detailed insight into the socio-economy of Celtic people in the region. In the Late Iron Age, contacts between Celtic groups and Romans intensified. Mainly the Celtic nobility were susceptible for Roman influence and long distance trade increased in the alpine foreland (Brem and Haldimann, 1999) to import prestige items like wine and tableware, and, according to Caesar, also horses (bell. gall. 4,2,2: Möller, 2013). Several written sources describe ancient horse and mule trade across Europe, Asia and Africa (see e.g. Peters, 1998: 140f), yet the scientific evidence for this is scarce (Paulus and Uerpmann, 2007; Nuviala et al., 2014). Horses were venerated status symbols in ancient times (Green, 1992); they were significant for transportation and warfare; however, horse meat was also

eaten on a regular basis by the Celts. Horses played an important part in Celtic symbolism as well, apparent in Epona, patroness of horses and riders, and in the ritual life as indicated for example from perforated horse skulls from La Tène (Méniel, 2007) or carcass depositions in Gournay-sur-Aronde, France (Rapin et al., 1980). However, most Iron Age sites in Switzerland do not contain more than 5% horse remains (Schibler et al., 1999: 121). Celtic horses are described as small (110–130 cm withers height), whereas the Romans also bred larger animals (>140 cm) (Müller-Lhotska, 1984: 114; Peters, 1998: 149f; Arbogast et al., 2002: 44f). Therefore, when encountering small and large horses in Celtic contexts, archaeozoologists often consider the small ones to be local, while larger ones are regarded as imported, endowed or looted. Some authors argue that the reason for the small size of the Celtic horse was poor pasture and low breeding interest (Boessneck et al., 1971: 106), but this interpretation contradicts the Celts' reputation as stock breeders and mounted warriors (Hyland, 1990: 173; Green, 1992: 66; Sidnell, 2006: 162). As small horses are more versatile and quick on the battlefield (Junkelmann, 1990: 44), it can be assumed that the Celts intentionally bred small horses. However, by the Late Iron Age, small and large horses were found to be coexisting in Celtic sites for example in southern Germany (Manching) or Luxembourg (Titelberg)

\* Corresponding author.

E-mail address: [julia.elsner@email.de](mailto:julia.elsner@email.de) (J. Elsner).

(Boessneck et al., 1971; Méniel, 2014: 18). Little is known about the genetic diversity of Celtic and Roman horses (Di Bernardo et al., 2004; Bower et al., 2013), yet it is speculated that the Romans introduced new female lineages to conquered regions (Bower et al., 2013).

In ancient DNA (aDNA) studies, the analysis of the non-recombining maternally inherited mitochondrial (mt) DNA has been widely applied due to its high copy number in the cell and the high variability within the non-coding displacement-loop (d-loop). Studies of modern horse mtDNA diversity have revealed an intricate phylogenetic structure. Based on 616 base pairs six major haplogroups were defined (Vila et al., 2001; Jansen et al., 2002). The inclusion of additional aDNA data (Cieslak et al., 2010) and whole mitochondrial genomes (Lippold et al., 2011b; Achilli et al., 2012) led to revisions of the initial nomenclature, resulting in now 18 or 19 haplogroups. Apparently, geographic structuring is low today, and was so in wild horse populations, but there is also evidence that some matrilineages were regionally confined (Cieslak et al., 2010; Lira et al., 2010). Studies featuring ancient horse mtDNA suggest multiple domestications and an increasing diversity of female lineages from the Bronze Age onwards (Vila et al., 2001; Cieslak et al., 2010).

Animals with unique colour variants may have been specially valued. Individual phenotypic information can be retrieved from nuclear DNA, e.g. from single nucleotide polymorphisms (SNPs). An increasing number of genetic loci associated with coat colour has been detected so far (for reviews see Rieder, 2009; Cieslak et al., 2011) including the pleiotropic effect some of them have on behaviour and health (Reissmann, 2009: 18f; Trut et al., 2009; Bellone, 2010).

Morphological sex identification in archaeological horse remains is feasible when determinant skeletal elements are preserved; these include the canines, which dominantly prevail in males, and the pelvis (Montane et al., 1913). Molecular sex identification is possible using the amelogenin gene (Iwase et al., 2003) and the zinc-finger gene on the X and Y chromosomes (Senese et al., 1999; Han et al., 2010).

Here we investigate horse remains, mostly with known withers heights, from three key Late Iron Age sites in Switzerland where Roman cultural influence is noticeable in the archaeological record. We explore mitochondrial d-loop variation, molecular sex and coat

colour markers of 32 Late La Tène and two Roman time horses. The diversity of the ancient female lineages is compared to modern indigenous horse and pony breeds. This is the first study focusing on variation in horses from a defined region inhabited by Celtic peoples, zooming in on a narrow time frame.

## 2. Material and methods

### 2.1. Sites and material

We have selected 32 horse remains from Late Iron Age contexts (La Tène D, 150 BCE–30 CE), the sites of Mormont, Basel-Gasfabrik and pre-Roman Aventicum in Switzerland (Fig. 1). Two samples dating to the late 1st to early 2nd century AD from Aventicum have been added to the analysis. Sample details are given in Table 1. All sites comprised more horse remains than investigated here.

#### 2.1.1. Mormont

The site on the hill of Mormont near Eclépens and La Sarraz, canton Vaud, was first discovered in 2006 during gravel works and addressed as a Helvetic sanctuary (Dietrich et al., 2007). After examining the faunal remains, a second scenario was proposed: the site might have been a temporal camp of refugees or migrants (Méniel, 2014: 195f). Since the investigations by other archaeological disciplines are not finished yet, both interpretations have to be regarded as preliminary. According to typological dating the main occupation time of the site was La Tène D1b (130/120–90/80 BCE), which is supported by absolute dendrochronological dates between 107 and 102 BCE (Kaenel, 2014). During this time c. 300 structures were constructed including 260 pits, filled with various items ranging from mill stones, ceramic and metal pots, and decorative objects to an abundance of faunal remains and even several human bones. Horses (n = 40) are the third most common animal based on minimum number of individuals (MNI), after cattle (n = 174) and pigs (n = 59) (Méniel, 2014: 12). Given the short occupation time, they might stem from contemporaneous populations. The dead animals had been treated in different ways, some were deposited as articulated (partial) skeletons or cadavers, other bones show cut marks



Fig. 1. Location of investigated sites in Switzerland. 1 – Mormont, canton Vaud; 2 – Avenches/Aventicum, canton Vaud; 3 – Basel-Gasfabrik, canton Basel-City.

**Table 1**  
 Details of archaeological sites and samples, and summary of results. Given are name, location and date of sites, and laboratory and archaeological code, morphological sex and age identification if determinant elements (pelvis, teeth) are present, skeletal element for withers height estimation and genetic analyses, as well as mitochondrial haplotype, GenBank accession code of d-loop sequences; molecular sex and coat colour. The measurement for greatest length (GL) and smallest breadth of diaphysis (SD) are given in mm, withers height (WH) was estimated after May (1985) and is given in cm. For some of the samples from Mormont, radius GL and SD from the same individual are given in parenthesis (Méniel, 2014: 264). Details on the mitochondrial haplotypes are given in Table 5, on molecular sexing in Table 6 and nuclear loci for the coat colour analysis in Table 7. Nomenclature of mitochondrial haplotypes follows Cieslak et al. (2010).

Site, location (altitude, latitude, longitude), date	Samples										
	Lab code	Arch code	Morph. sex, age	Skeletal element	GL	SD	WH	Mitochondrial haplotype	GenBank ID	Molecular sex	Coat colour
Mormont, 605 m, 46.66/6.54, Late La Tène, c. 100 BCE	MO1	Id 4 fosse 293	Female, 18 y	Metacarpus sinister	191 (285)	27 (31)	116.5 (117)	I	KC893855	–	–
	MO2	Id 35 fosse 288	–	Metacarpus sinister	181	27	110.5	D3	KC893856	Female	–
	MO3	Id 3 fosse 42	Female, 11.5 y	Metacarpus sinister	191 (281)	25.5 (29.3)	116.5 (115.5)	–	–	–	–
	MO4	Id 30 fosse 143	–	Metacarpus sinister	199	29	121.5	D2	KC893857	–	Chestnut
	MO5	Id 21 fosse 94	Female, 15 y	Metacarpus sinister	209	30	127.5	X2b	KC893858	Female	Black/not grey
	MO6	Id 44 fosse 196	–	Metacarpus sinister	202	30.5	123	B1	KC893859	Male	Black
	MO7	Id 2 fosse 131	Female, 10 y	Metacarpus sinister	189 (281)	27 (30.5)	115 (115.5)	–	–	–	–
	MO8	Id 27 fosse 146	–	Metacarpus sinister	205	28	125	–	–	–	–
	MO9	Id 42 fosse 275	–	Metacarpus sinister	194	29	118	X3c1 + 15,615G	KC893860	Male	Black/not grey
	MO10	Id 01 fosse 45	Male, 10 y	Metacarpus dexter	241 (365)	36 (41)	147 (150)	A/B/C/D/F?	KC893861	–	–
	MO11	Id 10 fosse 146	–	Metacarpus sinister	Incomplete	–	–	–	–	–	–
	MO12	Id 32 fosse 146	–	Metacarpus sinister	190	28	116	F	KC893862	–	–
	MO13	Id 53 fosse 210	Male, 5.5 y	Metacarpus sinister	191 (290)	29.7 (32)	116.5 (119)	–	–	–	–
	MO14	Id 58 fosse 542	–	Metacarpus sinister	Incomplete	–	–	K?	KC893863	–	–
	MO15	Id 54 fosse 210	Male, 5 y	Metacarpus sinister	208	29.7	127	X2b	KC893864	–	–
	MO16	Id 16 fosse 205	Male, 4 y	Metacarpus sinister	234 (340)	35.5 (39.2)	143 (140)	F	KC893865	Male	Bay/not grey
	MO17	Id 57 fosse 542	–	Metacarpus sinister	194	30	118	X2b	KC893866	–	Chestnut/not grey
	MO18	Id 55 fosse 169	Male, 1.5 y	Metacarpus sinister	Incomplete (subadult)	–	–	X2	KC893867	–	–
Basel-Gasfabrik, 255 m, 47.57/7.58, Late La Tène, 150–80 BCE	BGF1	Fk22519 pit44 excavation 1990/42	–	Metacarpus	201.5	30.4	123	F	KC893850	–	Chestnut
	BGF2	Fk22576 pit284 excavation 1990/32	–	Metacarpus	222.8	35.5	136	X2b	KC893851	–	–
	BGF3	Fk"2" pit114 excavation 1941/4	–	Metacarpus	196.9	31.4	120	X3c1 + 631G	KC893852	–	–
	BGF4	Fk"2" pit114 excavation 1941/4	–	Metacarpus	178.1	24.9	109	–	–	–	Chestnut
	BGF5	Fk 18796 pit256 excavation 1989/5	–	Radius/Ulna	276	33	113.5	F	KC893854	–	–
	BGF6	Fk(18943)18714 pit255 excavation 1989/5	–	Upper (pre)molar	–	–	–	–	–	–	–
Avenches, 630 m, 46.87/7.05, *Late La Tène, c. 50 BCE, **Roman, late 1st to early 2nd century CE	AV1**	K10656, Sk4	–	Radius	330	35	135.5	–	–	–	–
	AV2*	K10574	–	Radius/ulna	315	36	129.5	X3c1 + 15,615G	KC893849	–	–
	AV3**	K10656, Sk3	–	Radius	358	39.7	147	X2	KC893848	–	–

\* refers to AV2 and Late La Tène. \*\* refers to AV1, AV3 and Roman.

and traces of burning. We have examined 18 randomly chosen equid metacarpi in this study.

### 2.1.2. Basel-Gasfabrik

In Basel, a settlement of the Rauricii was discovered in 1911 at the left border of the river Rhine (Hecht and Niederhäuser, 2011: 6), today within the area of chemical industry (Gasfabrik, GF). It was excavated in several campaigns during the following 100 years. The

unfortified proto-urban site was inhabited between 150 and 80 BCE (La Tène D1) as indicated for example by fibula types and coins. It comprised houses, gardens and pens as well as craft zones, and two cemeteries. The majority of archaeological and faunal remains stem from the c. 585 pits which had been filled with "waste" after they had become unusable as e.g. storage cellars (Hecht and Niederhäuser, 2011: 47). Five horse limb bones and one tooth were chosen for the analysis based on calculability of the horses' withers height (except for the tooth).

### 2.1.3. Aventicum

Avenches, canton Vaud, was once the capital of the Helvetii known as Aventicum (Castella and Meylan, 2008). Celtic temples had been built in the first century BC, and under the reign of the Roman Emperor Augustus, who forced the urbanisation of the Celtic provinces, Roman town structures were added in the first decade CE which persisted about three hundred years (Blanc and Frei-Stolba, 2001). Aventicum became a Roman colony in 71 CE and was incorporated into *Germania Superior* when the province was formed in 85 CE (Blanc et al., 2001). The three radii studied here stem from sacrificial pits nearby the temples *au Lavoëx* and were chosen due to their large size (Deschler-Erb, 2015).

### 2.2. Modern horse populations

The matrilineages of the ancient horses were compared to modern horse mtDNA variability. The comparison of modern animal sequences to ancient genetic data has to be treated with caution. In ancient times, there were no breeds in today's sense (Johnstone, 2004: 88f) yet different morphotypes were described, for example by Vegetius (mulom. 3,6,2: Lommatzsch, 1903). We selected breeds that come from regions belonging to the Late Roman Republic in c. 100 BCE (Crook et al., 1994: add. mat. 1), contemporaneous to the occupation of both Mormont and Basel-GF. They were grouped into: Spain (*Hispania Citerior* and *Ulterior*); Italy (*Italia*, *Sardinia*, *Gallia Cisalpina*), and Croatia (*Illyricum*). In addition, breeds from Switzerland, southern Germany and eastern France represent the home region of the Celtic horses investigated here. Breeds were chosen due to their "indigenous" status according to the Domestic Animal Diversity Information System (DAD-IS; dad.fao.org), to availability on GenBank (ncbi.nlm.nih.gov) in April 2015 and when the authors sampled material from maternally unrelated individuals. As Celtic horses are ponies by today's standards (withers heights <148 cm) (Fédération Equestre Internationale, 2015: 60) we compared them also to European pony breeds from Spain, Italy, the Carpathians, European Russia, and Northern Europe (including the British Isles). In total, we assembled sequences from 35 museum and 726 modern specimens representing 30 breeds (Table S1).

### 2.3. Metrics

Measurements of greatest length (GL) and smallest breadth of diaphysis (SD) of metacarpi and/or radii from Mormont were taken from Méniel (2014: 263f); and from Aventicum and Basel-GF measured according to von den Driesch (1976). Withers heights were calculated from GL of all bones according to May (1985); note that this results in an estimation disregarding sexual dimorphism or robustness.

Metacarpus :  $GL \text{ (mm)} * 0.6102 = WH \text{ (cm)}$

Radius :  $GL \text{ (mm)} * 0.4111 = WH \text{ (cm)}$

### 2.4. Sample preparation, DNA extraction and PCR analysis

Sample preparation and DNA extraction were performed as previously described in Elsner et al. (2014) following the User Developed Protocol: „Purification of total DNA from compact animal bone using the DNeasy® Blood & Tissue Kit“ (Qiagen, Basel, Switzerland) in dedicated aDNA facilities physically separated from post-PCR laboratories. Standard precautions and measures in aDNA processing were adhered to (Shapiro and Hofreiter, 2012).

PCR for mtDNA and the nuclear (nc) loci (grey, *AMELX/Y*, *ZFX/Y*) was set up in 25 µl volumes containing 1.5 U AmpliTaq Gold, 1 × GeneAmp 10 × PCR Gold Buffer (150 mM Tris-HCl, 500 mM KCl, pH 8.0) and 2 mM MgCl<sub>2</sub> (all Applied Biosystems, Hombrechtikon, Switzerland); 0.4 mM dNTP Mix (Promega, Dübendorf, Switzerland); 0.2 µM of

**Table 2**

Primer sequences for mitochondrial haplotyping, and amplification strategy. Target lengths are given excluding primers. Reference sequence from Xu and Arnason (1994).

Name	Sequence 5'-3'	Position of target in reference sequence NC_001640	Annealing temperature (°C)	Length of target (bp, colour)
Ec1_f	TTCTTCCCCTAAACGACAACA	15492-15563	52	72 (green)
Ec1_r	GACGTACATAGGCCATTCATAAGA			
Ec4_f	GAATGGCCTATGTACGCTGCTG	15591-15669	55	79 (4f+4r, pink)
Ec4_r	GACTTGGATGGGGTATGCAC			
Ec4b_r	ATATTATGTACATGCTTATTATTCA	15591-15609	50	19 (4f+4br, red)
Ec2_f	ACATAACACCATACCCACCTGACA	15557-15659	55	103 (violet)
Ec2_r	GATGGGGTATGCACGATCAATAAT			
Ec5_f	ACCCATCCCAAGTCAATCA	15696-15758	55	63 (5f+5r, light blue)
Ec5_r	TAGTTGGGAGGGTTGCTG			
Eca1_r	GGCTGGTGATTAAGCTCGT	15696-15730	52	35 (5f+1r, blue)

each primer; 20 µg/µl BSA (bovine serum albumin, Roche, Basel, Switzerland), and up to 10 µl template DNA on a Mastercycler ProS (Eppendorf, Allschwil, Switzerland). The cycling conditions were: 12 min initial denaturation, followed by either 50 (for mtDNA) or 70 (for ncDNA) cycles of denaturation at 95 °C for 40 s, annealing at 52 °C to 58 °C (see Tables 2–4) for 30 s, and extension at 72 °C for 30 s, with a final extension of 60 s at 72 °C. Non-template controls were performed alongside all amplifications. Amplification products were visualised on 2% agarose gels. To authenticate sequence patterns and to exclude potential contamination, at least two PCR products from two independent extractions were obtained.

PCR products of mtDNA amplifications were cloned with the TOPO TA Cloning Kit (Invitrogen, Zug, Switzerland) following the manufacturer's protocol, except that the reaction volume was halved. Two clones of each PCR product were Sanger sequenced by Microsynth (Balgach, Switzerland). The products of the ncDNA PCR were premixed with tailed sequencing primers (Binladen et al., 2007) and also Sanger sequenced by Microsynth.

The coat colour SNPs were amplified in multiplex PCR; the first step in 20 µl volumes containing 2.5 U AmpliTaq Gold, 1 × GeneAmp 10 × Gold Buffer, and 4 mM MgCl<sub>2</sub>, 0.1 µM of each primer, 20 µg/µl BSA, and 4 µl template DNA. After 30 cycles of amplification as described above, PCR products were diluted 1:20; 5 µl were used in the second, singleplex amplification step (40 cycles) including 0.5 U AmpliTaq Gold, 1 × GeneAmp 10 × Gold Buffer, and 4 mM MgCl<sub>2</sub>, 0.2 µM of each primer, and 20 µg/µl BSA. Again, non-template controls were performed and PCR products visualised on 2% agarose gels.

Pyrosequencing of coat colour SNPs of the Basel-GF and Aventicum samples was performed in the Institute for Animal Sciences, Humboldt University Berlin and of Mormont specimens at the Life Science Training Facilities in the Biozentrum, University of Basel according to manufacturer's instructions and Ludwig et al. (2009) on PSQ™ 96MA (Biotage, Uppsala, Sweden).

### 2.5. Mitochondrial haplotyping

To investigate the matrilineages six partially overlapping and interspersed targets of the mtDNA d-loop with different lengths covering nucleotide positions 15,492–15,669 and 15,696–15,758 (Table 2) (Xu and Arnason, 1994) were amplified. Primers were taken and/or modified from Weber (2005), Cieslak et al. (2010) and Elsner et al. (2014) except for Ec4b\_r (this study).

### 2.6. Molecular sex identification

For the molecular sex identification primer sets Ecab\_AMELX/Y from Lippold et al. (2011a) and ZFX/Y\_Equus (this study, Table 3) were applied.



**Table 3**  
Primer sequences to identify the molecular sex of horse remains. Reference sequences from Han et al. (2010).

Name	Sequence 5'-3'	Reference sequence	Position of target in reference sequence	Annealing temperature (°C)	Length of target (bp)
ZFX_Equus_F	ATTATATCTGGCCAGGACT	DQ179230	326–382	55	58
ZFX_Equus_R	TGCCTAGCTTCCAAATCTAA				
ZFY_Equus_F	GAATTCCTACATGCCATA	DQ179229	214–263	55	50
ZFY_Equus_R	ATAAAGTCATGAGCCGGATA				

### 2.7. Coat colour analysis


The investigation of nuclear SNPs in the genes associated with coat colour phenotypes was performed as described in Ludwig et al. (2009). Samples were tested for the basic colours bay, chestnut and black via the *MC1R*- and *ASIP*-loci (Marklund et al., 1996; Rieder et al., 2001). Epistatic dilutions Cream (Mariat et al., 2003) and Silver (Reissmann et al., 2007) and spottings Overo (Santschi et al., 1998), Tobiano (Brooks et al., 2002) and Sabino (Brooks and Bailey, 2005) were examined. In addition, the grey-locus, resulting in a complete white coat in adult horses regardless of original colour, was investigated. This phenotype is caused by a duplication in intron 6 of the *STX17* gene (Rosengren Pielberg et al., 2008). We designed primer pairs to target short fragments of c. 100 bp, first, for the region that spans the end and the beginning of the 4.6 kb duplication (Table 4, red) to detect the mutated allele as well as second, a primer pair to target the end of the fragment (Table 4, blue) which would be present in both wildtype (not grey) and mutated genotype (grey). The suitability of the primer systems was tested on five randomly chosen Freiberger (bay or chestnut) and Camargue (grey) horses each (Fig. S1) at the Institute of Genetics, University of Bern.

### 2.8. Sequence data analysis

D-loop and *STX17* sequences were examined with BioEdit (Hall, 1999). Mitochondrial haplotypes were determined by comparison with the reference sequence NC\_001640 (Xu and Arnason, 1994). We used the nomenclature of Cieslak et al. (2010) to determine the haplotype (Table S2). Haplotype and nucleotide diversity as well as pairwise difference  $F_{ST}$  values were computed with Arlequin 3.5 (Excoffier and Lischer, 2010), excluding five ancient samples with missing data (see Table 5). The comparison of Iron Age and modern sequences was done on the basis of haplotype frequencies and pairwise differences for each breed, geographic origin and type (horse, pony). Sequences were pruned to positions 15,494–15,740 to include as many samples as possible.  $F_{ST}$  values were interpreted as follows: values from 0 to 0.05 indicate non to little genetic differentiation, 0.05–0.15 moderate differentiation, 0.15–0.25 large differentiation, and values above 0.25 very large differentiation. Negative  $F_{ST}$  values equate to zero (Hartl and Clark, 2007: 288). Based on relative haplotype frequencies, Principal Component Analysis (PCA) was computed with PAST (Hammer et al., 2001). Singleton haplotypes were removed from this analysis. Median Joining Networks (MJN) (Bandelt et al., 1999) were constructed with Network 4.6.1.2 (fluxus-engineering.com) using the following parameters: insertions and deletions were double weighted (default 50),

**Table 4**  
Primer sequences to detect i) the duplication causing premature greying (grey-locus) and ii) the wildtype (no duplication). Figure shows position of target sequences. Reference sequences from Rosengren Pielberg et al. (2008).

Name	Sequence 5'-3'	Position of target in reference sequence		Annealing temperature (°C)	Length of target (bp, colour)
		EU606026	EU606027		
Both_f	GCACCACCTGGGAAGTCA				
Both_r	CTGGAGTGTGCCACGAAATC	5830–5900	10407–10477	52	71 (blue)
Dup_r	GAGAAGTTGGCCAAAGCAG	-	5830–5889	58	61 (red)



transition; transversion ratio was set to 1:10, and variable sites were down weighted according to the number of deviations from the majority at each nucleotide position excluding haplotype defining sites (Tables S2 & S3). The dataset was reduced to 360 modern (one representative per haplotype per breed) and 14 ancient sequences. The probabilities  $P$  of allelic dropout for the SNP loci were calculated as in Gagneux et al. (1997):  $P = K * (K/2)^{n-1}$ , where  $K$  is the number of observed allelic dropouts divided by the number of all positive amplifications of heterozygous individuals, and  $n$  is the number of PCR replications.

## 3. Results

### 3.1. Metrics


The measurements of SD and GL as well as the estimation of withers height of the La Tène and Roman time horses revealed variation at all archaeological sites. Furthermore, based on metacarpi the individuals cluster into a smaller and into a larger group, based on the radii into three groups (Fig. 2). Note that not all individuals are represented by both skeletal elements. There is a linear correlation between length and breadth ( $R^2 = 0.81$  for metacarpi,  $R^2 = 0.9$  for radii) indicating isometric growth and no robustness differences.

The archaeozoological defined standard withers height of Celtic horses ranges between c. 110 cm to 130 cm (Peters, 1998; Arbogast et al., 2002); the majority of animals investigated here accord to this (Fig. 3). Five horses were larger than 130 cm: three from the Late Iron

**Table 5**

Differences to the reference sequence NC\_001640 between nucleotide positions 15,492–15,669 and 15,696–15,758. Samples indicated by an asterisk have missing data at nucleotide positions: MO10, MO14, MO18: 15,564–15,669; BGF2, AV3: 15,564–15,690 and 15,610–15,669. Cieslak et al. (2010) disregard hotspot nucleotide positions 15,585; 15,597; 15,604 and 15,650 for haplotype assignment, they are mentioned here for completeness.

Sample	Differences to NC_001640 (Xu and Arnason, 1994)	Haplotype after Cieslak et al. (2010)
MO1	495C; 538G; 602T; 650G; 709T; 720A	I
MO2	495C; 650G; 666A; 720A	D3
MO4	495C	D2
MO5	494C; 495C; 496G; 534T; 585A; 603C; 649G; 720A	X2b
MO6	495C; 602T; 617C; 659C; 720A	B1
MO9	495C; 542T; 585A; 597G; 602T; 615G; 635T; 650G; 666A; 703C; 720A	X3c1 + 615
MO10*	495C; 720A	Haplogroup A/B/C/D/F?
MO12	495C; 585A; 601C; 602T; 720A	F
MO14*	495C; 546T; 703C; 720A	Haplogroup K?
MO15	494C; 495C; 496G; 534T; 603C; 649G; 720A	X2b
MO16	495C; 585A; 601C; 602T; 720A	F
MO17	494C; 495C; 496G; 534T; 603C; 649G; 720A	X2b
MO18*	494C; 495C; 496G; 534T; 720A	Haplogroup X2
BGF1	495C; 585A; 601C; 602T; 720A	F
BGF2*	494C; 495C; 496G; 534T; 603C; 649N; 720A	X2b
BGF3	495C; 542T; 597G; 602T; 631G; 635T; 650G; 666A; 703C; 720A	X3c1 + 631
BGF5	495C; 585A; 601C; 602T; 720A	F
AV2	495C; 542T; 585A; 597G; 602T; 615G; 635T; 650G; 666N; 703C; 720A	X3c1 + 615
AV3*	494C; 495C; 496G; 534T; 602T; 603C; 604A; 720A	Haplogroup X2



**Fig. 2.** Greatest length (GL) against smallest breadth of diaphysis (SD) of metacarpi (left panel) and radii (right panel). Bones from Mormont (black), Basel-GF (grey) and Aventicum (light grey). Mormont individuals with data from both skeletal elements indicated by symbol; morphological and/or molecular sex determination: f = female, m = male.

Age: two from Mormont (MO10: 149 cm, MO16: 142 cm) and one from Basel-GF (BGF2: 136 cm), and both Roman time horses (AV1: 136 cm, AV3: 147 cm) from Aventicum. The average withers height of the Mormont horses was, based on 15 measurable metacarpi and six radii, 124 cm (119 cm without the two large animals). The average in Basel-GF was 120 cm and in Aventicum 137 cm (141 cm for the Roman time individuals).

**3.2. Mitochondrial haplotyping**


In Mormont ten metacarpi were genetically well preserved, three only allowed for a partial amplification of the target region, and five had no amplifiable mtDNA preserved. It was possible to amplify mtDNA from four out of six samples from Basel-GF and from two out of three radii (one Iron Age, one Roman) from Aventicum. This results in an overall amplification success of 56% (19/34).

We identified eleven haplotypes belonging to six haplogroups (hg). Most prevalent is hg X2 (n = 6), followed by hg F (n = 4), hg X3 (n = 3), hg D (n = 2), hg B (n = 1) and hg I (n = 1). Additionally, one sample each might be associated with hg K and hgs A/B/C/D/F, respectively

(Table 5). Haplotype sharing within and between archaeological sites was observed: haplotype F is shared by MO12, MO16, BGF1 and BGF5. MO15 and MO17 have the same haplotype X2b, MO5 and BGF2 also share variants of X2b, and despite some missing data MO18 and AV3 can be assigned to haplogroup X2 as well. MO9, BGF3 and AV2 possess variants of haplotype X3c1, one of which previously undescribed (BGF3). Horses with withers height below 130 cm carried haplotypes B1, D2, D3, F, I, X2b, and X3c1; they share haplotype F and haplogroup X2 with the large horses. Haplotype (0.92 ± 0.06) and nucleotide (0.3 ± 0.02) diversity is very high in the investigated Iron Age horses (Table S4). This result is mirrored by the horses from Mormont: haplotype diversity = 0.97 ± 0.06, nucleotide diversity = 0.3 ± 0.02.

**3.3. Molecular sex identification**

Molecular sex could be identified for five specimens from Mormont, the other samples were not preserved well enough to enable the amplification of targets in the amelogenin and/or zinc-finger genes (Table 6). This confirms the results of the morphological sex determination for MO5 and MO16; MO2, MO6 and MO9 were previously undetermined.



**Fig. 3.** Estimated withers height (May, 1985) in cm from metacarpi and/or radii from Mormont (black), Basel-GF (grey) and Aventicum (light grey). When measurements of both bones were available, the mean was used.

**Table 6**  
Results for molecular sex identification with AMELX/Y and ZFX/Y.

Sample	AMELX	AMELY	ZFX	ZFY	Sex
MO2	1/3	0/3	1/2	0/3	Female
MO5	1/3	0/2	2/2	0/4	Female
MO6	0/2	1/2	2/2	1/3	Male
MO9	0/2	1/1	2/2	2/3	Male
MO16	0/1	0/1	2/2	2/2	Male

### 3.4. Coat colour

For six of the Mormont and two of the Basel-GF samples DNA preservation allowed for the amplification of SNPs in genes associated with coat colour (Table 7). The specimens from Aventicum were not preserved well enough for SNP typing. All positive samples revealed basic colourations. One large horse (MO16) was bay, of the smaller horses (<130 cm) three had a black coat and four were chestnut. We did not detect any mutations in the dilution and spotting loci. The Mormont specimens were also tested for the duplication that causes premature greying. For five individuals the target at the end of the 4.6 kb fragment present in both wildtype and mutated genotype could be amplified twice (once for MO6, see Table 7), but the shorter PCR products for the grey-locus were negative four times for each individual suggesting all tested animals were not grey. The probability of allelic dropout for the only heterozygous SNP, *MCR1R*, was  $p = 0.0015$  with four PCR replications.

### 3.5. Comparison with modern horses

The Median Joining Network shows the widespread distribution of Late Iron Age horse mtDNA d-loop sequences within the variation of southern European horse and European pony breeds (Fig. 4). The ancient horses are represented in the major haplogroups (B, D, F, I, X2, X3) except for hg K (see above).

The fixation index based on the number of pairwise differences between each breed and the ancient horses indicated substantial structuring ( $F_{ST} = 0.12$ ,  $p < 0.001$ ) (see Table S5 for pairwise population  $F_{ST}$ ). The genetic differentiation between the geographic regions is moderate ( $F_{ST} = 0.06$ ,  $p < 0.001$ ).  $F_{ST}$ -values for the regional groups indicate close relationships between all European horse and pony breeds and the Celtic horses, except for the Spanish breeds which are moderately different (Table S6). This outcome is supported by PCA based on the frequencies of haplotypes with loadings >10% (Fig. 5; frequencies of all haplotypes in Fig. S2). The Iron Age horses stand out from the modern breeds because of their unique variability but tend to be closer to the northern European ponies which share all haplotypes present in the ancient dataset.

**Table 7**  
Results from the coat colour analysis. In addition to the phenotype, the genotypes for the eight SNPs investigated are shown; differences from the wildtype are indicated in bold and only detected at the *ASIP*- and *MCR1*-loci. The duplication causing premature greying was not detected in any of the Mormont samples (Avenches and Basel-GF samples were not tested). Unreproducible results are parenthesised.

Sample	Phenotype	Nuclear Genes								GREY
		<i>ASIP</i>	<i>MCR1R</i>	<i>EDNRB</i> Overo spotting	<i>KIT13</i> Tobiano spotting	<i>KIT16</i> Sabino spotting	<i>MATP</i> Cream dilution	<i>SILV9</i> Silver dilution	<i>SILV11</i>	
MO4	(Chestnut)	<b>(a/a)</b>	<b>(e/e)</b>	ov/ov	KM0/KM0	(sb1/sb1)	C/C	(z/z)	z/z	–
MO5	Black	<b>a/a</b>	<b>E/e</b>	ov/ov	KM0/KM0	sb1/sb1	C/C	z/z	z/z	No
MO6	Black	<b>a/a</b>	<b>E/e</b>	ov/ov	KM0/KM0	sb1/sb1	C/C	z/z	z/z	(No)
MO9	Black	<b>a/a</b>	<b>E/e</b>	ov/ov	KM0/KM0	sb1/sb1	C/C	z/z	z/z	No
MO16	Bay	A/A	E/E	ov/ov	KM0/KM0	sb1/sb1	C/C	z/z	z/z	No
MO17	Chestnut	<b>a/a</b>	<b>e/e</b>	ov/ov	KM0/KM0	sb1/sb1	C/C	z/z	(z/z)	No
BGF1	Chestnut	<b>a/a</b>	<b>e/e</b>	ov/ov	KM0/KM0	sb1/sb1	C/C	(z/z)	z/z	–
BGF4	Chestnut	–	<b>e/e</b>	–	KM0/KM0	–	C/C	z/z	z/z	–

Differences from the wildtype are indicated in bold; the wildtype is given in capital letters in the genetic nomenclature.

## 4. Discussion

This study provides first insight into mtDNA d-loop diversity, molecular sex and coat colour of Late La Tène horses in Switzerland. DNA preservation was as expected for the region and time frame explored, however, the results for the nuclear markers were better for the recently excavated site on the hill of Mormont (Pruvost et al., 2007; Bollongino et al., 2008). The withers height of the investigated horses fell mostly within the variation of standard Iron Age horses, yet five individuals were larger than 130 cm reaching up to 149 cm. Two of these were identified as male, but smaller stallions were also present, thus sexual dimorphism does not explain the height differences. Moreover, the relation between length and breadth (robustness) is linear, indicating that smaller and larger horses did not belong to different morphotypes (Brooks et al., 2010; Dzierzecka and Komosa, 2013). Only one horse exceeded the defined pony range of maximal 148 cm. The investigation of the maternal lineages revealed high nucleotide and haplotype diversity within the upper range of variance in modern European breeds (see Table S4). The horses belonged to six haplogroups (B, D, F, I, X2, X3) which are rare in wild horses (Weinstock et al., 2005; Cieslak et al., 2010; Lorenzen et al., 2011; Orlando et al., 2013). They were first described from domestic horses from sites in Novosibirsk, Russia, dating to c. 2000 BCE, and Moldova and Romania around 1250 BCE (Cieslak et al., 2010). A haplogroup-package dominated by hgs D, K, X2 and X3 is subsequently found in archaeological sites throughout Eurasia (Vila et al., 2001; Di Bernardo et al., 2004; Keyser-Tracqui et al., 2005; McGahern et al., 2006; Cai et al., 2007; Cai et al., 2009; Lira et al., 2010; Cieslak et al., 2010; Priskin et al., 2010; Bower et al., 2013).

The large matrilineage diversity is striking, particularly given that the majority of horses, namely those from Mormont, belonged to a potentially contemporaneous local population which should include directly related individuals. The occurrence of shared lineages in the sites Mormont and Basel-GF might indicate horse exchange between Helvetii and Rauricii. The ancient matrilineages show affinities to modern horse and pony breeds from most regions investigated. They tend to have bequeathed more to northern European and British Isles pony breeds and less to Spanish breeds. However, considering that domestic horses were first introduced to Switzerland in the Late Neolithic or Early Bronze Age, i.e. around 2500–2000 BCE (Schibler and Studer, 1998: 177), it is remarkable that most major haplogroups are present in the Iron Age archaeological record, notably in view of the rareness of archaeological horse remains.

With morphological and molecular methods, five mares and seven stallions were identified, all from the hill site of Mormont. This corresponds to the male:female ratio of 1:1.5 amongst all morphologically sex determined horses in Mormont (Méniel, 2014: 17).

Coat colouration seems to be rather uniform. We would expect more variety: Ludwig et al. (2009) documented a rapid and substantial increase in coat colourations already in the Bronze Age, at least in Asia




Fig. 4. Median Joining Network of 14 ancient and 355 modern mtDNA d-loop sequences. Circle size is proportional to haplotype frequency. Nomenclature follows Cieslak et al. (2010).

and eastern Europe, and in northern Europe spotted Iron Age horses were detected (Svensson et al., 2012) in comparable datasets. However, monochrome horses were favoured over skewbalds and white markings on face and leg according to written Roman sources (e. g. Corp.

Hippiatr. Gr., 1, 115; Oder and Hoppe, 1924–1927; Geop. 16,2; Wappmann, 1985). Breeders probably had noticed hearing, visual, and neurological impairment and even stillbirths coming along predominantly with spottings, especially in homozygous individuals




Fig. 5. PCA plot based on relative mtDNA haplotype frequencies of the haplotypes present in the Iron Age horses and additional haplotypes with loadings over 10% (see Fig. S2). The two dimensions display 60% of the total variance.



(Reissmann, 2009: 20; Bellone, 2010) and thus may have selected against those phenotypes. Notably premature greying resulting in a white coat regardless of original colour was also not detected. Again, we expected a different result, as white horses played an important role in Roman and also in Germanic ceremonies (Tacitus, Germ. 1,10; Kretschmer, 1986; Johnstone, 2004: 61). However, of the horses investigated, only the specimens from Aventicum stem from an unambiguous ritual context and those were not preserved well enough for genetic coat colour identification. Assuming that ceremonial activities were also performed on the hill of Mormont, we find no evidence that specially coloured horses were preferred in ritual, which apparently also applies for sex as both females and males were present.

Some important archaeozoological questions concerning horse exchange, breeding aims and selection criteria e.g. for ceremony have to remain open yet, due to both the small sample size and insufficient data for comparison. Further complicating is the fact that Roman horses, too, were not standardised and the workaday animals were usually smaller than military or circus breedings (Junkelmann, 1990: 39). It appears that mtDNA d-loop variation offers too little structuring, but because it is highly conservative the investigation of the paternal inherited Y-chromosome cannot contribute to the question of horse provenience either (Lippold et al., 2011a; Wallner et al., 2013). The exploration of nuclear markers beyond sex and colouration (Schubert et al., 2014) might give insight into possibly distinct characteristics of horses bred by Celtic and Roman stud farmers. Further approaches like stable isotope analysis, particularly of strontium (Slovak and Paytan, 2011), seem even more promising to solve these important issues. It might thus be necessary to dig deeper into the past and to include archaeological material from both Rome and its provinces, and regions outside the Roman Empire to trace the origin and breeding history of small and large horses.

## 5. Conclusion

The horse remains excavated from the archaeological sites Mormont, Basel-GF and Aventicum, dating to 150–50 BCE (La Tène D) and to c. 100 CE, revealed a high diversity of matrilineages. The most frequent haplogroups F and X2 were shared between the sites, and between small and large horses. While all determinable large individuals (> 140 cm withers height) were male, both sexes were present amongst horses with withers heights between 110 and 130 cm. The detection of solely monochrome coats and the absence of white colourations might indicate selection. The simplified approach to detect premature greying developed in this study provides a relevant tool for aDNA research concerning the phenotypes of (pre)historic horses. Compared to modern indigenous breeds, Late Iron Age horses seem to have bequeathed more to northern European ponies than to Spanish breeds. By broadening the methodological spectrum and by complementing the dataset both spatially and temporally, we hope to address archaeozoological questions on horse exchange, breeding aims, and selection criteria e.g. in ritual contexts more comprehensively.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.jasrep.2016.03.007>.

## Acknowledgements

Samples were kindly provided by Archéologie Vaudoise (Mormont), Site et Musée Romain d'Avenches (Aventicum) and Archäologische Bodenforschung Basel-Stadt (Basel-Gasfabrik). The authors thank Tosso Leeb and Michaela Drögemüller (Institute for Genetics, University of Bern), Andreas Papassotiropoulos and Kim Dung Huynh (Biozentrum, University of Basel), Arne Ludwig (Institute for Zoo- and Wildlife Research) and Mélanie Pruvost (then Institute for Animal Sciences, Humboldt University Berlin), Patrice Méniel and Pauline Nuviala (University of Burgundy, Dijon), co-editor Andy Howard, University of Durham, and one anonymous reviewer. Julia Elsner was funded by Swiss National Science Foundation (K-31K1\_120528/1), Freiwillige

Akademische Gesellschaft Basel, Nikolaus und Bertha Burckhardt-Bürgin-Foundation.

## References

- Achilli, A., Olivieri, A., Soares, P., Lancioni, H., Hooshar Kashani, B., Perego, U.A., Nergadze, S.G., Carossa, V., Santagostino, M., Capomaccio, S., Felicetti, M., Al-Achkar, W., Penedo, M.C., Verini-Supplizi, A., Houshmand, M., Woodward, S.R., Semino, O., Silvestrelli, M., Giullotto, E., Pereira, L., Bandelt, H.J., Torroni, A., 2012. Mitochondrial genomes from modern horses reveal the major haplogroups that underwent domestication. *PNAS* 109, 2449–2454. <http://dx.doi.org/10.1073/pnas.1111637109>.
- Arbogast, R.M., Benoit, C., Lepetz, S., Méniel, P., Yvinec, J.-H., 2002. *Archéologie du cheval. Des origines à la période moderne en France*. Hespérides, Paris.
- Bandelt, H.J., Forster, P., Rohl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16, 37–48.
- Bellone, R.R., 2010. Pleiotropic effects of pigmentation genes in horses. *Anim. Genet.* 41, 100–110. <http://dx.doi.org/10.1111/j.1365-2052.2010.02116.x>.
- Binladen, J., Gilbert, M.T., Campos, P.F., Willerslev, E., 2007. 5'-tailed sequencing primers improve sequencing quality of PCR products. *Biotechniques* 42, 174–176. <http://dx.doi.org/10.2144/000112316>.
- Blanc, P., Frei-Stolba, R., 2001. Die Stadtentwicklung. *Archäol. Schweiz.* 24. <http://dx.doi.org/10.5169/seals-18959>.
- Blanc, P., Frey-Kupper, S., Frei-Stolba, R., 2001. Die Anfänge der Hauptstadt der Helvetier. *Archäol. Schweiz.* 24. <http://dx.doi.org/10.5169/seals-18956>.
- Boessneck, J., von den Driesch, A., Meyer-Lemppenau, U., Wechsler-von Ohlen, E., 1971. Die Tierknochenfunde aus dem Oppidum von Manching. Franz Steiner, Wiesbaden.
- Bollongino, R., Tresset, A., Vigne, J.D., 2008. Environment and excavation: pre-lab impacts on ancient DNA analyses. *C.R. Palevol* 7, 91–98. <http://dx.doi.org/10.1016/j.crpv.2008.02.002>.
- Bower, M.A., Edwards, C.J., Evans, C., Downes, C.S., 2013. Horse husbandry in Iron Age and Romano-British East Anglia: ancient DNA analysis of horses at Langdale Hale, Cambridgeshire. In: Evans, C., Appleby, G., Lucy, S., Regan, R. (Eds.), *Process and History: Romano-British Communities at Colne Fen, Earith: An Inland Port and Supply Farm*. University Press, Cambridge, pp. 140–143.
- Brem, H., Haldimann, M.-A., 1999. Handel und Verkehr: Die Latènezeit. In: Müller, F., Kaenel, G., Lüscher, G. (Eds.), *SPM IV: Eisenzeit*. Verlag Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel, pp. 215–219.
- Brooks, S.A., Bailey, E., 2005. Exon skipping in the KIT gene causes a Sabino spotting pattern in horses. *Mamm. Genome* 16, 893–902. <http://dx.doi.org/10.1007/s00335-005-2472-y>.
- Brooks, S.A., Makvandi-Nejad, S., Chu, E., Allen, J.J., Streeter, C., Gu, E., McCleery, B., Murphy, B.A., Bellone, R., Sutter, N.B., 2010. Morphological variation in the horse: defining complex traits of body size and shape. *Anim. Genet.* 41, 159–165. <http://dx.doi.org/10.1111/j.1365-2052.2010.02127.x>.
- Brooks, S.A., Terry, R.B., Bailey, E., 2002. A PCR-RFLP for KIT associated with tobiano spotting pattern in horses. *Anim. Genet.* 33, 301–303. <http://dx.doi.org/10.1046/j.1365-2052.2002.00854.x>.
- Brunetti, C., Kaenel, G., Méniel, P. (Eds.), 2014. *Les Helvètes au Mormont. Une énigme dans le monde celtique*. SAS Archeodunum, Chaponnay.
- Cai, D.W., Han, L., Xie, C.Z., Li, S.N., Zhou, H., Zhu, H., 2007. Mitochondrial DNA analysis of Bronze Age horses recovered from Chifeng region, Inner Mongolia, China. *Prog. Nat. Sci.* 17, 544–550.
- Cai, D.W., Tang, Z.W., Han, L., Speller, C.F., Yang, D.Y.Y., Ma, X.L., Cao, J.E., Zhu, H., Zhou, H., 2009. Ancient DNA provides new insights into the origin of the Chinese domestic horse. *J. Archaeol. Sci.* 36, 835–842. <http://dx.doi.org/10.1016/j.jas.2008.11.006>.
- Castella, D., Meylan, M.F., 2008. *Topographie sacrée et rituels: le cas d'Aventicum, capital des Helvètes: Actes du colloque international d'Avenches, 2–4 Novembre 2006*. Antiqua.
- Cieslak, M., Pruvost, M., Benecke, N., Hofreiter, M., Morales, A., Reissmann, M., Ludwig, A., 2010. Origin and history of mitochondrial DNA lineages in domestic horses. *PLoS One* 5, e15311. <http://dx.doi.org/10.1371/journal.pone.0015311>.
- Cieslak, M., Reissmann, M., Hofreiter, M., Ludwig, A., 2011. Colours of domestication. *Biol. Rev. Camb. Philos. Soc.* 86, 885–899. <http://dx.doi.org/10.1111/j.1469-185X.2011.00177.x>.
- Crook, J.A., Lintott, A., Rawson, E. (Eds.), 1994. *The Cambridge Ancient History Volume 9: The Last Age of the Roman Republic, 146–43 BC*. University Press, Cambridge.
- Deschler-Erb, S., 2015. Tier und Kult - Spezielle Tierknocheneponierungen der Spätlatène- und Römerzeit aus Aventicum/Avenches (CH) im nordalpinen Vergleich. *Musée Romain d'Avenches, Avenches*.
- Di Bernardo, G., Galderisi, U., Del Gaudio, S., D'Aniello, A., Lanave, C., De Robertis, M.T., Cascino, A., Cipollaro, M., 2004. Genetic characterization of Pompeii and Herculaneum Equidae buried by Vesuvius in 79 AD. *J. Cell. Physiol.* 199, 200–205. <http://dx.doi.org/10.1002/jcp.10461>.
- Dietrich, E., Kaenel, G., Weidmann, D., Jud, P., Méniel, P., Moinat, P., 2007. Le sanctuaire helvète du Mormont. *Archäol. Schweiz.* 30, 2–13. <http://dx.doi.org/10.5169/seals-109261>.
- Dzierzecka, M., Komosa, M., 2013. Variability of the proximal phalanx in warmblood and coldblood horses - morphological and structural analyses. *Belg. J. Zool.* 143, 119–130.
- Elsner, J., Schibler, J., Hofreiter, M., Schlumbaum, A., 2014. Burial condition is the most important factor for mtDNA PCR amplification success in Palaeolithic equid remains from the Alpine foreland. *Archaeol. Anthropol. Sci.* 1–11. <http://dx.doi.org/10.1007/s12520-014-0213-4>.
- Excoffier, L., Lischer, H.E., 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10, 564–567. <http://dx.doi.org/10.1111/j.1755-0998.2010.02847.x>.

- Fédération Equestre Internationale FEI, 2015. 2015 Veterinary Regulations. Fédération Equestre Internationale, Lausanne.
- Gagneux, P., Boesch, C., Woodruff, D.S., 1997. Microsatellite scoring errors associated with noninvasive genotyping based on nuclear DNA amplified from shed hair. *Mol. Ecol.* 6, 861–868.
- Green, M.J., 1992. *Animals in Celtic life and Myth*. Routledge, London New York.
- Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41, 95–98 (doi: citeulike-article-id:691774).
- Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontol. Electron.* 4, 1–9.
- Han, S.-H., Yang, B.-C., Ko, M.-S., Oh, H.-S., Lee, S.-S., 2010. Length difference between equine ZFX and ZFY genes and its application for molecular sex determination. *J. Assist. Reprod. Genet.* 27, 725–728. <http://dx.doi.org/10.1007/s10815-010-9467-7>.
- Hartl, D.L., Clark, A.G., 2007. *Principles of Population Genetics*. Sinauer Associates, Inc. Publishers, Sunderland, Mass.
- Hecht, Y., Niederhäuser, A., 2011. Alltagskultur und Totenrituale der Kelten. Ein Siedlungszentrum am Oberrhein um 100 v. Chr. – The everyday culture and funerary rituals of the Celts. A central settlement on the Upper Rhine around 100 BC. Christoph Merian Verlag, Basel.
- Hyland, A., 1990. *Equus: The Horse in the Roman World*. Yale University Press, New Haven, London.
- Iwase, M., Satta, Y., Hirai, Y., Hirai, H., Imai, H., Takahata, N., 2003. The amelogenin loci span an ancient pseudoautosomal boundary in diverse mammalian species. *PNAS* 100, 5258–5263. <http://dx.doi.org/10.1073/pnas.0635848100>.
- Jansen, T., Forster, P., Levine, M.A., Oelke, H., Hurler, M., Renfrew, C., Weber, J., Olek, K., 2002. Mitochondrial DNA and the origins of the domestic horse. *PNAS* 99, 10905–10910. <http://dx.doi.org/10.1073/pnas.152330099>.
- Johnstone CJ (2004) A Biometric Study of Equids in the Roman World. (PhD thesis Archaeology department). York: University of York.
- Junkelmann, M., 1990. *Die Reiter Roms. Teil I: Reise, Jagd, Triumph und Circusrennen*. Philipp von Zabern, Mainz.
- Kaenel, G., 2014. De quand date l'occupation du Mormont? In: Brunetti, C., Kaenel, G., Méniel, P. (Eds.), *Les Helvètes au Mormont Une énigme dans le monde celtique*. SAS Archeodunum, Chaponnay, pp. 74–75.
- Keyser-Tracqui, C., Blandin-Frappin, P., Francfort, H.P., Ricaut, F.X., Lepetz, S., Crubezy, E., Samashev, Z., Ludes, B., 2005. Mitochondrial DNA analysis of horses recovered from a frozen tomb (Berel site, Kazakhstan, 3rd Century BC). *Anim. Genet.* 36, 203–209. <http://dx.doi.org/10.1111/j.1365-2052.2005.01316.x>.
- Kretschmer, M. (Ed.), 1986. *Tacitus: Germania*. Verlag Aschendorff, Münster.
- Lippold, S., Knapp, M., Kuznetsova, T., Leonard, J.A., Benecke, N., Ludwig, A., Rasmussen, M., Cooper, A., Weinstock, J., Willerslev, E., Shapiro, B., Hofreiter, M., 2011a. Discovery of lost diversity of paternal horse lineages using ancient DNA. *Nat. Commun.* 2, 450. <http://dx.doi.org/10.1038/ncomms1447>.
- Lippold, S., Matzke, N.J., Reissmann, M., Hofreiter, M., 2011b. Whole mitochondrial genome sequencing of domestic horses reveals incorporation of extensive wild horse diversity during domestication. *BMC Evol. Biol.* 11, 328. <http://dx.doi.org/10.1186/1471-2148-11-328>.
- Lira, J., Linderholm, A., Olaria, C., Brandstrom Durling, M., Gilbert, M.T., Ellegren, H., Willerslev, E., Liden, K., Arsuaga, J.L., Gotherstrom, A., 2010. Ancient DNA reveals traces of Iberian Neolithic and Bronze Age lineages in modern Iberian horses. *Mol. Ecol.* 19, 64–78. <http://dx.doi.org/10.1111/j.1365-294X.2009.04430.x>.
- Lommatzsch, E. (Ed.), 1903. *P. Vegeti Renati digestorum artis mulomedicinae libri*. B. G. Teubner, Leipzig.
- Lorenzen, E.D., Nogues-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K.A., Ugan, A., Borregaard, M.K., Gilbert, M.T., Nielsen, R., Ho, S.Y., Goebel, T., Graf, K.E., Byers, D., Stenderup, J.T., Rasmussen, M., Campos, P.F., Leonard, J.A., Koepfli, K.P., Froese, D., Zazula, G., Stafford Jr., T.W., Aaris-Sorensen, K., Batra, P., Haywood, A.M., Singarayer, J.S., Valdes, P.J., Boeskorov, G., Burns, J.A., Davydov, S.P., Haile, J., Jenkins, D.L., Kosintsev, P., Kuznetsova, T., Lai, X., Martin, L.D., McDonald, H.G., Mol, D., Meldgaard, M., Munch, K., Stephan, E., Sablin, M., Sommer, R.S., Sipko, T., Scott, E., Suchard, M.A., Tikhonov, A., Willerslev, R., Wayne, R.K., Cooper, A., Hofreiter, M., Sher, A., Shapiro, B., Rahbek, C., Willerslev, E., 2011. Species-specific responses of Late Quaternary megafauna to climate and humans. *Nature* 479, 359–364. <http://dx.doi.org/10.1038/nature10574>.
- Ludwig, A., Pruvost, M., Reissmann, M., Benecke, N., Brockmann, G.A., Castanos, P., Cieslak, M., Lippold, S., Llorente, L., Malaspina, A.S., Slatkin, M., Hofreiter, M., 2009. Coat color variation at the beginning of horse domestication. *Science* 324, 485. <http://dx.doi.org/10.1126/science.1172750>.
- Mariot, D., Taourit, S., Guerin, G., 2003. A mutation in the MTP gene causes the cream coat colour in the horse. *Genet. Sel. Evol.* 35, 119–133. <http://dx.doi.org/10.1051/gse:2002039>.
- Marklund, L., Moller, M.J., Sandberg, K., Andersson, L., 1996. A missense mutation in the gene for melanocyte-stimulating hormone receptor (MC1R) is associated with the chestnut coat color in horses. *Mamm. Genome* 7, 895–899. <http://dx.doi.org/10.1007/s003359900264>.
- May, E., 1985. *Widerristhöhe und Langknochenmaße bei Pferden - ein immer noch aktuelles Problem*. *Z. Säugetierkunde* 50, 368–382.
- McGahern, A.M., Edwards, C.J., Bower, M.A., Heffernan, A., Park, S.D., Brophy, P.O., Bradley, D.G., MacHugh, D.E., Hill, E.W., 2006. Mitochondrial DNA sequence diversity in extant Irish horse populations and in ancient horses. *Anim. Genet.* 37, 498–502. <http://dx.doi.org/10.1111/j.1365-2052.2006.01506.x>.
- Méniel, P., 2007. *Les animaux de La Tène*. In: Musée Schwab Bienne MSB and Landesmuseum Zürich LZ (eds) *La Tène La recherche, les questions, les réponses*. Verlag Museum Schwab, Biel, pp. 61–65.
- Méniel, P., 2014. *Les restes animaux du site du Mormont. Eclépens et La Sarraz, canton de Vaud, vers 100 avant J.-C.* Cahiers d'Archéologie Romande, Lausanne.
- Möller, L. (Ed.), 2013. *Der gallische Krieg/Gaius Julius Caesar*. Marixverlag, Wiesbaden.
- Montane, L., Bourdelle, E., Bressou, C., 1913. *Anatomie régionale des animaux domestiques, I: Cheval*. Baillière, Paris.
- Müller-Lhotska, U.A., 1984. *Das Pferd in der Schweiz. Von der Prähistorie bis zum ausgehenden Mittelalter* (PhD thesis History department) University of Zürich, Zürich.
- Nuviala, P., Nowell, G., Ughetto, J., Balasse, M., 2014. *L'origine des chevaux du site du Mormont (canton de Vaud, Suisse): L'apport des isotopes du carbone, de l'oxygène et du strontium*. In: Barral, P., Guillaumet, M.-J., Roulière-Lambert, M.-J., Saracino, M., Vitali, D. (Eds.), *Les Celtes et le nord de l'Italie Premier et second âges du fer*. Société archéologique de l'Est, Vérone, pp. 603–610.
- Oder, E., Hoppe, K. (Eds.), 1924–1927. *Corpus hippiatricorum graecorum*. B. G. Teubner, Leipzig.
- Orlando, L., Ginolhac, A., Zhang, G., Froese, D., Albrechtsen, A., Stiller, M., Schubert, M., Cappellini, E., Petersen, B., Moltke, I., Johnson, P.L., Fumagalli, M., Vilstrup, J.T., Raghavan, M., Korneliusen, T., Malaspina, A.S., Vogt, J., Szklarczak, D., Kelstrup, C.D., Vinther, J., Dolocan, A., Stenderup, J., Velazquez, A.M., Cahill, J., Rasmussen, M., Wang, X., Min, J., Zazula, G.D., Seguin-Orlando, A., Mortensen, C., Magnussen, K., Thompson, J.F., Weinstock, J., Gregersen, K., Røed, K.H., Eisenmann, V., Rubin, C.J., Miller, D.C., Antczak, D.F., Bertelsen, M.F., Brunak, S., Al-Rasheid, K.A., Ryder, O., Andersson, L., Mundy, J., Krogh, A., Gilbert, M.T., Kjaer, K., Sicheritz-Ponten, T., Jensen, L.J., Olsen, J.V., Hofreiter, M., Nielsen, R., Shapiro, B., Wang, J., Willerslev, E., 2013. Recalibrating *Equus* evolution using the genome sequence of an early Middle Pleistocene horse. *Nature* 499, 74–78. <http://dx.doi.org/10.1038/nature12323>.
- Paulus, S., Uerpmann, H.P., 2007. *Physikalisch-chemische Untersuchungen an Equidenzähnen aus Kalkriebe zur Bestimmung von Knochenhaltung und Jahreszeit der Schlacht*. In: Wilbers-Rost, S., Uerpmann, H.P., Uerpmann, M., Grosskopf, B., Tolksdorf-Lienemann, E. (Eds.), *Kalkriebe 3: Interdisziplinäre Untersuchungen auf dem Oberes in Kalkriebe, Archäologische Befunde und naturwissenschaftliche Begleituntersuchungen*. Verlag Philipp von Zabern, Mainz, pp. 145–156.
- Peters, J., 1998. *Römische Tierhaltung und Tierzucht*. Marie Leidorf GmbH, Rahden/Westf.
- Pichler, S.L., Lassau, G., Alt, K.W., Röder, B. and Schibler, J. (eds) (in preparation). *Über die Toten zu den Lebenden. Interdisziplinäre Synthese*. Basel: ABBIS.
- Priskin, K., Szabo, K., Tomory, G., Bogacsi-Szabo, E., Csanyi, B., Eordogh, R., Downes, C.S., Rasko, I., 2010. Mitochondrial sequence variation in ancient horses from the Carpathian Basin and possible modern relatives. *Genetica* 138, 211–218. <http://dx.doi.org/10.1007/s10709-009-9411-x>.
- Pruvost, M., Schwarz, R., Correia, V.B., Champlot, S., Braguier, S., Morel, N., Fernandez-Jalvo, Y., Grange, T., Geigl, E.M., 2007. Freshly excavated fossil bones are best for amplification of ancient DNA. *PNAS* 104, 739–744. <http://dx.doi.org/10.1073/pnas.0610257104>.
- Rapin, A., Méniel, P., Brunaux, J.-L., 1980. *Un sanctuaire gaulois à Gournay-sur-Aronde (Oise) Gallia* pp. 1–25.
- Reissmann, M., 2009. *Die Farben der Pferde*. Genetik, Klassifizierung, Charakteristik. Cadmos, Brunsbek.
- Reissmann, M., Bierwolf, J., Brockmann, G.A., 2007. Two SNPs in the SILV gene are associated with silver coat colour in ponies. *Anim. Genet.* 38, 1–6. <http://dx.doi.org/10.1111/j.1365-2052.2006.01553.x>.
- Rieder, S., 2009. Molecular tests for coat colour in horses. *J. Anim. Breed. Genet.* 126, 415–424. <http://dx.doi.org/10.1111/j.1439-0388.2009.00832.x>.
- Rieder, S., Taourit, S., Mariot, D., Langlois, B., Guerin, G., 2001. Mutations in the agouti (ASIP), the extension (MC1R), and the brown (TYRP1) loci and their association to coat color phenotypes in horses (*Equus caballus*). *Mamm. Genome* 12, 450–455. <http://dx.doi.org/10.1007/s003350020017>.
- Rosengren Pielberg, G., Golovko, A., Sundstrom, E., Curik, I., Lennartsson, J., Seltenhammer, M.H., Druml, T., Binns, M., Fitzsimmons, C., Lindgren, G., Sandberg, K., Baumung, R., Vetterlein, M., Stromberg, S., Grabherr, M., Wade, C., Lindblad-Toh, K., Ponten, F., Heldin, C.H., Solkner, J., Andersson, L., 2008. A cis-acting regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nat. Genet.* 40, 1004–1009. <http://dx.doi.org/10.1038/ng.185>.
- Santschi, E.M., Purdy, A.K., Valberg, S.J., Vrotsos, P.D., Kaese, H., Mickelson, J.R., 1998. Endothelin receptor B polymorphism associated with lethal white foal syndrome in horses. *Mamm. Genome* 9, 306–309.
- Schibler, J., Stopp, B., Studer, J., 1999. *Haustierhaltung und Jagd*. In: Müller, F., Kaenel, G., Lüscher, G. (Eds.), *SPM IV: Eisenzeit*. Schweizerische Gesellschaft für Ur- und Frühgeschichte, Basel, pp. 116–136.
- Schibler, J., Studer, J., 1998. *Haustierhaltung und Jagd während der Bronzezeit der Schweiz*. In: Hochuli, S., Niffeler, U., Rychener, J. (Eds.), *SPM III: Bronzezeit*. Gesellschaft für Ur- und Frühgeschichte der Schweiz, Basel, pp. 171–192.
- Schubert, M., Jonsson, H., Chang, D., Der Sarkissian, C., Ermini, L., Ginolhac, A., Albrechtsen, A., Dupanloup, I., Foucal, A., Petersen, B., Fumagalli, M., Raghavan, M., Seguin-Orlando, A., Korneliusen, T.S., Velazquez, A.M., Stenderup, J., Hoover, C.A., Rubin, C.J., Alfathan, A.H., Alquraishi, S.A., Al-Rasheid, K.A., MacHugh, D.E., Kalbfleisch, T., MacLeod, J.N., Rubin, E.M., Sicheritz-Ponten, T., Andersson, L., Hofreiter, M., Marques-Bonet, T., Gilbert, M.T., Nielsen, R., Excoffier, L., Willerslev, E., Shapiro, B., Orlando, L., 2014. Pre-historic genomes reveal the genetic foundation and cost of horse domestication. *PNAS* 111, E5661–E5669. <http://dx.doi.org/10.1073/pnas.1416991111>.
- Senese, C., Penedo, M.C.T., Shiue, Y.L., Bowling, A.T., Millon, L.V., 1999. A HaellII PCR-RFLP in the ZFY/ZFX genes of horses. *Anim. Genet.* 30, 390–391. <http://dx.doi.org/10.1046/j.1365-2052.1999.00526-10.x>.
- Shapiro, B., Hofreiter, M. (Eds.), 2012. *Ancient DNA - Methods and Protocols*. Springer, New York.
- Sidnell, P., 2006. *Warhorse. Cavalry in Ancient Warfare*. Continuum Books, London, New York.

- Slovak, N.M., Paytan, A., 2011. Applications of Sr isotopes in archaeology. In: Baskaran, M. (Ed.), *Handbook of Environmental Isotope Geochemistry*. Springer, Berlin, Heidelberg, pp. 743–768.
- Svensson, E.M., Telldahl, Y., Sjöling, E., Sundkvist, A., Hulth, H., Sjövald, T., Gotherstrom, A., 2012. Coat colour and sex identification in horses from Iron Age Sweden. *Ann. Anat.* 194, 82–87. <http://dx.doi.org/10.1016/j.aanat.2011.11.001>.
- Trut, L., Oskina, I., Kharlamova, A., 2009. Animal evolution during domestication: the domesticated fox as a model. *BioEssays* 31, 349–360. <http://dx.doi.org/10.1002/bies.200800070>.
- Vila, C., Leonard, J.A., Gotherstrom, A., Marklund, S., Sandberg, K., Liden, K., Wayne, R.K., Ellegren, H., 2001. Widespread origins of domestic horse lineages. *Science* 291, 474–477. <http://dx.doi.org/10.1126/science.291.5503.474>.
- von den Driesch, A., 1976. *A Guide to the Measurement of Animal Bones From Archaeological Sites*. Harvard University Press, Cambridge, Mass.
- Wallner, B., Vogl, C., Shukla, P., Burgstaller, J.P., Druml, T., Brem, G., 2013. Identification of genetic variation on the horse y chromosome and the tracing of male founder lineages in modern breeds. *PLoS One* 8, e60015. <http://dx.doi.org/10.1371/journal.pone.0060015>.
- Wappmann, U., 1985. *Buch 16 und 17 der Geoponica. Übersetzung und Besprechung* (PhD thesis Veterinary Medicine. München).
- Weber, C., 2005. *Molekulargenetische Typisierung einer skythenzeitlichen Pferdepopulation* (Master thesis Paleogenomics Mainz: Johannes Gutenberg-Universität).
- Weinstock, J., Willerslev, E., Sher, A., Tong, W., Ho, S.Y., Rubenstein, D., Storer, J., Burns, J., Martin, L., Bravi, C., Prieto, A., Froese, D., Scott, E., Xulong, L., Cooper, A., 2005. Evolution, systematics, and phylogeography of Pleistocene horses in the new world: a molecular perspective. *PLoS Biol.* 3, e241. <http://dx.doi.org/10.1371/journal.pbio.0030241>.
- Xu, X., Arnason, U., 1994. The complete mitochondrial DNA sequence of the horse, *Equus caballus*: extensive heteroplasmy of the control region. *Gene* 148, 357–362.