

content in WZ ( $4.09 \pm 1.13 \times 10^3 \mu\text{g}/\text{mL}/\text{cm}^3$ ) measured by DMMB assay was significantly higher than that in RZ ( $1.57 \pm 0.42 \times 10^3 \mu\text{g}/\text{mL}/\text{cm}^3$ ;  $P=0.004$ ). There was a significant correlation ( $R^2=0.62$ ) between the proteoglycan estimated by sugar area of FT-IRIS and the proteoglycan content measured by DMMB assay.

**Conclusions:** Collagen was low and proteoglycan was high in WZ, whereas collagen was high and proteoglycan was low in RZ. The results indicate that proportion of collagen and proteoglycan is not uniform but different in WZ and in RZ. Correlations between FT-IRIS analysis and biochemical assays for collagen and for proteoglycan indicate that FT-IRIS may be a useful tool for quantitative analysis of extracellular matrix of meniscus.

#### 443 CELLULAR AND BIOMECHANICAL SEGMENTAL CHARACTERIZATION OF HUMAN MENISCUS

H. Pereira<sup>1,2</sup>, A.M. Frias<sup>1</sup>, S.G. Caridade<sup>1</sup>, J.F. Mano<sup>1</sup>, J.M. Oliveira<sup>1</sup>, J. Espregueira-Mendes<sup>1,3</sup>, R.L. Reis<sup>1</sup>. <sup>1</sup>3Bs Res. Group – Univ. of Minho; ICVS/3B's – PT Government Associate Lab., Guimarães, Portugal; <sup>2</sup>Orthopedic Dept. Centro Hosp.ar Póvoa de Varzim, Vila da Conde, Portugal; <sup>3</sup>Saúde Atlântica Sports Ctr. – F.C. Porto Stadium; Porto Univ. Res. Ctr., Porto, Portugal

**Purpose:** The regeneration of meniscus lesions using tissue engineering strategies has recently been attracting great deal of attention. Tissue engineering driven approaches however, require a better knowledge on the segmental composition of meniscus from both biological and biomechanical point of view. With this work we aim to contribute to the knowledge of this tissue aiming future clinical applications in particular, the aspects dealing with the segmental analysis of cellular phenotype and distribution, mechanical properties and extracellular matrix composition.

**Methods:** Human tissue was obtained from local hospitals by means of surgery or biopsy, in accordance with local ethical committee guidelines. For this study we evaluated menisci from 30 donors, 26 lateral and 23 medial menisci were enrolled for study. Only morphologically intact menisci, not submitted to previous surgery were included. Each meniscus was divided into anterior, middle and posterior segments prior to mechanical, biological or histological evaluation. We isolated human meniscus cells (HMC's) using explants or using an enzymatic standard protocol. Flow cytometry analysis was performed in order to characterize meniscus cells population. Micro-computed tomography (Micro-CT) of freeze-dried meniscus was also carried out. Histomorphometric analysis of menisci stained sections (haematoxylin and eosin – H&E, safranin O and collagen I) was performed for segmental characterization of ECM and cells distribution. Dynamic mechanical analysis (DMA) was carried out for anterior, middle and posterior segments of fresh menisci. Within the region of interest, samples were cut in cylindrical shapes with 4mm diameter and 4mm thickness using a biopsy punch and were stored in PBS solution. The viscoelastic measurements were performed, at 37°C in PBS (pH 7.4), using a TRITEC8000B DMA from Triton Technology (UK), equipped with the compressive mode.

**Results:** Micro-CT analysis revealed that meniscus (freeze-dried) possessed a mean porosity of 53%, a mean pore size and trabeculae thickness of 85  $\mu\text{m}$  and 80  $\mu\text{m}$ , respectively. DMA analysis has shown, as expected, variability within samples due to their human nature, however we could observe a trend of increasing menisci stiffness: medial anterior (0.25 MPa at 1 Hz) < lateral anterior < lateral middle < medial posterior < lateral posterior < medial middle (0.9 MPa at 1 Hz). Cells isolated from the different samples are a mixed population of cells, i.e. fibrochondrocyte-like and MSCs (cells are positive for CD105, CD73 and CD90, and lack CD34 and CD45). Figure 1 illustrates the histological evaluation and it shows that meniscus ECM is composed of collagen-type I. Moreover, this fibrocartilagenous tissue has higher cell density in the periphery as compared to meniscus core. Cellular density among the different segments (anterior, middle, posterior) of meniscus was quantified using the H&E 2-D histological images.

**Conclusions:** To our knowledge, this is the first study of segmental characterization of fresh human menisci, without changes due to freezing or cryopreservation, in respect to biomechanical properties, further considering cells phenotype and distribution. This study provides deeper insights on human meniscus properties, contributing for the

development of adequate acellular and cellular tissue engineering strategies for the regeneration of meniscus.




Fig. 1. (a) Menisci segments considered in the different evaluation (anterior, middle and posterior). Evaluation of the (b) presence of collagen type I in the ECM; (c) safranin O staining demonstrating the presence of fibrocartilage distribution; (d) H&E stained sections at different magnification [low (d.1) and high (d.2) magnification].

## Pain: Clinical and Pathophysiology

#### 444 EXPERIMENTAL KNEE JOINT PAIN DURING STRENGTH TRAINING INCREASES MUSCLE STRENGTH GAIN IN HEALTHY SUBJECTS: A RANDOMISED CONTROLLED TRIAL

T.J. Sørensen<sup>1</sup>, H. Langberg<sup>2</sup>, H. Bliddal<sup>3</sup>, M. Henriksen<sup>3</sup>. <sup>1</sup>Unit for Musculoskeletal Function and Physiotherapy, Inst. of Sports Sci. and Clinical Biomechanics, Univ. of Southern Denmark, Odense, Denmark; <sup>2</sup>Inst. of Sports Med. – Copenhagen, Copenhagen Univ. Hosp. – Bispebjerg, Copenhagen, Denmark; <sup>3</sup>Clinical Motor Function Lab., The Parker Inst., Copenhagen Univ. Hosp. – Frederiksberg, Frederiksberg, Denmark

**Purpose:** Knee joint pain and reduced quadriceps strength are cardinal symptoms in many knee pathologies such as knee osteoarthritis. In people with painful knee pathologies quadriceps exercise reduce pain, improves physical function and increases muscle strength. A general assumption is that pain compromises muscle function and thus may prevent effective rehabilitation. The aim of the study was to evaluate the effects of experimental knee joint pain during quadriceps strength training on muscle strength gain in healthy individuals.

**Methods:** Twenty-seven healthy, untrained volunteers participated in a randomized controlled trial of unilateral quadriceps strengthening (8 weeks/3 times per week). Participants were randomized to perform the resistance training during pain induced by injections of either painful hypertonic saline (pain group, N=13) or a control condition with injection of non-painful isotonic saline (control group, N=14) into the infra-patellar fat pad. The resistance training consisted of two quadriceps strengthening exercises (leg press and knee extension machine exercises). All participants performed 3 sets of each exercise with loads corresponding to 80% of 1 repetition maximum (RM). Each set was performed to the point of muscular fatigue (inability to maintain the target load; approx. 8–12 repetitions). The primary outcome measure was change in maximal isokinetic muscle strength in knee extension/flexion (60, 120 and 180 deg/s.).