



ELSEVIER



<http://www.elsevier.com/locate/jiph>

SHORT REPORT

# First microbiota assessments of children's paddling pool waters evaluated using 16S rRNA gene-based metagenome analysis



CrossMark

Toko Sawabe<sup>a,\*</sup>, Wataru Suda<sup>b,c</sup>, Kenshiro Ohshima<sup>b</sup>,  
Masahira Hattori<sup>b,d</sup>, Tomoo Sawabe<sup>e</sup>

<sup>a</sup> Department of Food and Nutrition, Hakodate Junior College, 52-1, Takaoka-cho, Hakodate 042-0955, Japan

<sup>b</sup> Laboratory of Metagenomics, Graduate School of Frontier Sciences, The University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa, Chiba 277-8561, Japan

<sup>c</sup> Department of Microbiology and Immunology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

<sup>d</sup> Cooperative Major in Advanced Health Science, Graduate School of Advanced Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

<sup>e</sup> Laboratory of Microbiology, Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate 041-0821, Japan

Received 2 July 2015; received in revised form 29 September 2015; accepted 1 November 2015

## KEYWORDS

Children's paddling pool;  
Microbiota;  
Contamination;  
Metagenome;  
Small subunit rRNA gene

**Summary** Insufficient chloric sterilization of children's paddling pool waters increases the risk of diarrheal illness. Therefore, we investigated the microbiota changes after children use pools. First, we applied 16S rRNA gene-based metagenome analysis to understand the dynamics of microbiota in pool water, especially with respect to the bio-contamination by potential pathogens. *Proteobacteria* were major taxa detected in every pool water sample after children spent time in the pool. In more detail, *Gammaproteobacteria* comprised the dominant class, which was followed by *Betaproteobacteria*. Five phyla, *Bacteroidetes*, *Firmicutes*, *Actinobacteria* and *Deinococcus-Thermus* phyla were minor groups. The pool water microbiota are likely to be a consortium of intestinal and skin microbiota from humans. Interestingly, the ratio of *Gammaproteobacteria* and *Betaproteobacteria* differed according to the age of the children who used the pool, which means the pool water was additionally contaminated by soil microbiota as a result of the children's behavior. Furthermore, potential pathogens, such as *Campylobacter* spp.,

\* Corresponding author.

E-mail address: [sawabe@hakodate-jc.ac.jp](mailto:sawabe@hakodate-jc.ac.jp) (T. Sawabe).

*Comamonas testosteroni* and *Burkholderia pseudomallei*, were also found. Considering the standard plate counts, the abundances of these human pathogens are unlikely to be a sufficiently infectious dose. We suggest the importance of sanitary measures in paddling pool waters to reduce bio-contamination from both humans and the environment.

© 2015 King Saud Bin Abdulaziz University for Health Sciences. Published by Elsevier Limited. All rights reserved.

Children's paddling pools serve as a focal point in the transmission of infection and have caused a verotoxin-producing *Escherichia coli* O157 outbreak [1]. The risk of spreading diarrheal illness is increased by the insufficient chloric sterilization of pools [2]. O157 outbreaks in nursery facilities in Japan have been increasing annually; there were seven cases in 2010, four cases in 2011, nine cases in 2012 and 23 cases in 2013 [3,4]. Therefore, the Ministry of Health, Labor and Welfare in Japan published "Infection measure guideline in child day-care facilities, 2014", recommending thorough chloric sterilization water quality management for children's pools including a density of free residual chlorine that ranges from 0.4 to 1.0 ppm. On the other hand, chlorine may increase the risk of respiratory organ disease in infants [5]. Additionally, small inflatable and plastic pools for infants are typically filled with tap water that lacks chloric sterilization. In this study, 16S rRNA gene-based metagenome analysis was performed to identify the dynamics of the microbiota in pool water.


Ten water samples (T1 to 7 and F1 to 3) were obtained from children's paddling pools (ca. 270 L) at two nursery facilities (the facility T and F) from July 15 to August 1, 2014. These pools were filled with tap water (chloride density supplied: approximately 1.0 ppm) that had a lower density than 1.0 ppm by volatilization of chlorine. The numbers of children in the pool ranged from four to 16, and the age ranged between zero to six years. In the facility T samples, children from zero to six years old were present in every sample. In the F facility, the pool was used group-by-group; the groups were divided according to the age of the children, e.g., F1, F2 and F3 consisted of one- to two-year-old children three- to five-year-old children and only two-year-old children, respectively.

Indicator bacteria were evaluated in each sample based on standard protocols. *E. coli* counts were estimated using a sheet medium "Sanita-kun" (JNC Co., Ltd, Japan). The standard plate counts (SPC), coliforms and *E. coli* were  $9.8 \times 10^2$  to  $4.6 \times 10^3$  CFU/mL,  $2.0\text{--}1.5 \times 10^2$  CFU/mL (except

T4) and  $0\text{--}9.4 \times 10^2$  CFU/mL, respectively. The counts after the children spent time in the pools were beyond the criteria for public health in Japan, including SPC <200 CFU/mL and *E. coli* non-detectable. In previous studies, there was no correlation between the number of bacteria and number of users observed [6,7]. The indicator bacteria were below the undetectable limit in the tap water used in the children's paddling pools.

Five liters of water samples were filtered using a Sterivex filter (Merck Millipore Co.) together with pressure vessels DV-5 (Advantec, Tokyo, Japan) and filter-sterilized nitrogen gas. The microbes that were collected by the filter were treated with lysostaphin (L7386, Sigma-Aldrich) and lysozyme (L7651, Sigma-Aldrich) at 37 °C overnight; then, the bacterial genomic DNA was extracted from the lysate using the Wizard genomic DNA purification system (Promega, Madison, WI). Genomic DNA was used for 16S rRNA gene-based metagenome analysis after measuring the concentration with the Quantus™ and QuantiFluor™ ONE dsDNA System (Promega). Sequencing of the V1-V2 region of the 16S rRNA gene was performed using 454 GS Junior (Roche Applied Science), and 3000 reads per sample were used for the bacterial community comparison with a previously reported method [8]. The sequence data for this study are deposited in DDBJ/GenBank/EMBL under PRJDB4242.


The metagenome analyses revealed three interesting points. First, five phyla (*Proteobacteria*, *Bacteroidetes*, *Firmicutes*, *Actinobacteria* and *Deinococcus-Thermus*) were detected in the pool water microbiota after the children had been in the pool, accounting for 99.8–100% of the sample (Fig. 1A). *Proteobacteria* was the major phylum in the pool water microbiota, ranging from 76.2 to 100%. In more detail, *Gammaproteobacteria* ranged from 50% to 97%, which was followed by *Betaproteobacteria*, *Alphaproteobacteria*, *Deltaproteobacteria* and *Epsilonproteobacteria*. The phyla, *Bacteroidetes*, *Firmicutes* and *Actinobacteria*, were minor groups that occurred



**Figure 1** Microbiota of pool water samples. (A) Phylum composition of the pool water samples. ■ *Proteobacteria*, □ *Bacteroidetes*, ■ *Firmicutes*, ■ *Actinobacteria*, ■ *Deinococcus-Thermus*, □ *others*. (B) PCoA plot of the Weighted UniFrac analysis for the pool water microbiota. black: T facility, white: F facility.

at a frequency of less than 20% in the pool water microbiota. *Deinococcus-Thermus* was also detected in every sample. *Deinococcus-Thermus* is the dominant phyla on hand and/or human skin microbiota [9]. The water microbiota likely includes intestinal and skin microbiota from humans [10]. All samples were also analyzed using a PCoA plot of Weighted UniFrac. The pool water microbiota is unlikely to be similar in every sample, which means the microbiota may be affected by the status of children's activities, conditions and behaviors (Fig. 1B).

Second, comparisons between the microbiota in samples F1 and F2 further revealed a sharp contrast in how children's behavior affects the microbiota in the pool water when we look at the high abundance of reads for *Betaproteobacteria* in sample F2 (Fig. 2). The F2 sample pool was used by three-to five-year-old children. According to observation records of these children's behavior during pool time, they frequently got in and out of the pool and played on the bare ground surrounding the pool. As a result, the *Betaproteobacteria* may be from soil contamination.



**Figure 2** Difference in the class level bacterial composition between samples F1 and F2. ■ *Gammaproteobacteria*, □ *Betaproteobacteria*, ■ *Alphaproteobacteria*, □ *Deltaproteobacteria*, ■ *Epsilonproteobacteria*.

Third, the microbiota analysis also provides interesting insights into bio-contamination from human feces. Reads affiliated with *E. coli* (0.03–2.8%) were detected in almost all samples, except for the T6 sample, and *Campylobacter hominis* (0.03%) was detected in the T4 sample, which means the pool water may be bio-contaminated by feces from healthy humans. Potential pathogens, such as *Campylobacter* spp. (*C. coli* and *C. ureolyticus*), were also identified, but they were below 1% in samples T4, T6 and F2. *Comamonas testosteroni* (9.4%) and *Burkholderia pseudomallei* (0.03%) were present in sample F2, which suggests contamination of potent human pathogens when children played in the pool. *C. testosteroni* is found in a wide range of natural habitats, and it infects patients with underlying disease at a hospital [11]. Considering the SPC, the abundances of these human pathogens are unlikely sufficient infectious doses in this study. In conclusion, the results suggest the importance of sanitary measures in paddling pool waters to reduce bio-contamination from both humans and the environment. Washing the bodies of children before they enter the pool using clear tap water is recommended to maintain the conditions of "the SPC <200 CFU/mL and *E. coli* non-detectable".

## Funding

No funding sources.

## Competing interests

None declared.

## Ethical approval

Not required.

## Acknowledgment

This work was supported by JSPS KAKENHI Grant Number 25560031.

## References

- [1] Brewster DH, Brown MI, Robertson D, Houghton GL, Bimson J, Sharp JCM. An outbreak of *Escherichia coli* O157 associated with a children's paddling pool. *Epidemiol Infect* 1994;112:441–7.
- [2] Friedman MS, Roels T, Koehler JE, Feldman L, Bibb WF, Blake P. *Escherichia coli* O157:H7 outbreak associated with an improperly chlorinated swimming pool. *Clin Infect Dis* 1999;29:298–303.
- [3] Infectious Diseases Weekly Report (IDWR) 2013 week50, National Institute of Infectious Diseases, Japan.
- [4] Infectious Agents Surveillance Report (IASR), Enterohemorrhagic *Escherichia coli* infection in Japan as of April 2013, The Topic of This Month, vol. 34 (5) (No. 399).
- [5] Voisin C, Sardella A, Marcucci BA. Infant swimming in chlorinated pools and the risks of bronchiolitis, asthma and allergy. *Eur Respir J* 2010;36:41–7.
- [6] Sawabe T, Kimura M, Sasaki S. Investigation of cleanliness for water inflatable swimming pools: a case study of Hakodate junior college related facilities. *Bull Hakodate Jr Coll* 2013;39:49–52.
- [7] Sawabe T, Yamada M. Sanitary risk assessment of inflatable swimming pool water. *Bull Hakodate Jr Coll* 2014;40:21–6.
- [8] Said HS, Suda W, Nakagome S, Chinen H, Oshima K, Kim S, et al. Dysbiosis of salivary microbiota in inflammatory bowel disease and its association with oral immunological biomarkers. *DNA Res* 2014;21:15–25.
- [9] Zongxin L, Xia L, Yueqiu L, Li Y, Karen EN, Yuezhu W, et al. Pyrosequencing analysis of the human microbiota of healthy Chinese undergraduates. *BMC Genomics* 2013;14:390.
- [10] Khanna S, Tosh PK. A clinician's primer on the role of the microbiome in human health and disease. *Mayo Clin Proc* 2014;89:107–14.
- [11] Kim HJ, Lee Y, Oh K, Choi SH, Huh JW. Septic shock due to unusual pathogens, *Comamonas testosteroni* and *Acinetobacter guillouiae* in an immune competent patient. *Korean J Crit Care Med* 2015;3:180–3.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

**ScienceDirect**