

Contents lists available at SciVerse ScienceDirect

Physica A

journal homepage: www.elsevier.com/locate/physa

Exact field-driven interface dynamics in the two-dimensional stochastic Ising model with helicoidal boundary conditions

J. Ricardo G. Mendonça*

Instituto de Física, Universidade de São Paulo – Caixa Postal 66318, 05314-970 São Paulo, SP, Brazil

Instituto de Física, Universidade Federal de Uberlândia – Caixa Postal 593, 38400-902 Uberlândia, MG, Brazil

ARTICLE INFO

Article history:

Received 28 May 2012

Received in revised form 2 July 2012

Available online 31 July 2012

This paper is dedicated to Professor Sílvio R. A. Salinas (IF/USP, Brazil) on the occasion of his 70th birthday.

Keywords:

Stochastic Ising model
Bethe ansatz
RSOS growth model
Zero range process
Exclusion process
XXZ quantum chain
KPZ universality class

ABSTRACT

We investigate the interface dynamics of the two-dimensional stochastic Ising model in an external field under helicoidal boundary conditions. At sufficiently low temperatures and fields, the dynamics of the interface is described by an exactly solvable high-spin asymmetric quantum Hamiltonian that is the infinitesimal generator of the zero range process. Generally, the critical dynamics of the interface fluctuations is in the Kardar–Parisi–Zhang universality class of critical behavior. We remark that a whole family of RSOS interface models similar to the Ising interface model investigated here can be described by exactly solvable restricted high-spin quantum XXZ-type Hamiltonians.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Mappings from two-dimensional (2D) Ising model interface configurations to diffusion processes are known at least since the work of Rost [1], and have been explored many times since then [2–11]. In some cases [2–4], the interface dynamics of the model at zero temperature in the absence of an external field was mapped into the one-dimensional symmetric simple exclusion process, with the main result being the solution of a first-passage time problem showing that the mean lifetime of a shrinking domain is proportional to its initial area, providing a microscopic derivation for this well known experimental fact. It was also recognized that the resulting exclusion process can be recast as a probabilistic cellular automaton with a transition matrix equivalent to the transfer matrix (in a diagonal direction) of the symmetric six-vertex model in one of its critical lines [3,4].

The relationship between interfaces, exclusion processes, and vertex models was explored further [12–15], and it was realized that a Heisenberg Hamiltonian with pure imaginary Dzyaloshinsky–Moriya interaction that commutes with the transfer matrix of a six-vertex model describes the single-step surface growth model [16,17], as well as a discrete-velocity version of the noisy Burgers equation, which in turn is equivalent to the Kardar–Parisi–Zhang equation [18]. In the interacting particle system scenario, the above mentioned Hamiltonian is but the infinitesimal generator of the asymmetric simple exclusion process [12,14]. Conversely, a host of results concerning symmetric and asymmetric simple exclusion processes have been translated into the Ising interface problem and, in particular, the investigation of the motion of tagged particles,

* Correspondence to: Instituto de Física, Universidade de São Paulo – Caixa Postal 66318, 05314-970 São Paulo, SP, Brazil. Tel.: +55 11 9937 3760.
E-mail addresses: jricardo@if.usp.br, jricardo@infis.ufu.br.

as first introduced in the study of the hydrodynamic behavior of exclusion-type processes [19,20], has provided a partial explanation for the relationship between the characteristics of different asymptotic growth regimes in some $(1 + 1)$ -dimensional stochastic growth models [6,7,9].

In this article we show that under suitable generalized, but otherwise quite natural periodic boundary conditions the dynamics of an interface in the 2D stochastic Ising model in the presence of an external driving field can be mapped via a particle-height transformation into the dynamics of hopping particles without exclusion known as the zero range process [21–27]. The infinitesimal generator of the zero range process is equivalent to a high-spin, in general asymmetric quantum Hamiltonian that is exactly solvable by the Bethe ansatz [28–30]. We argue that the critical behavior of a generalized particle-height model must be on the Kardar–Parisi–Zhang universality class of critical behavior, since this is the critical behavior of the corresponding generalized quantum chains. This may have implications in the study of related models such as the dynamics of k -mers and other Ising-type lattice configurations.

The article is organized as follows. In Section 2 we introduce the 2D Ising model in an external field and the single-spin flip rates in terms of which the dynamics of the Ising contours will be analyzed, and in Section 3 we show that it can be described by diffusing particles without exclusion and exhibit the infinitesimal generator of the process. Section 4 contains a brief exposition of the exact solution of the zero range process by the Bethe ansatz and a discussion on its dynamical critical exponent. In Section 5, we show that a whole class of interface models similar to the Ising interface model can be described by exactly solvable, generalized restricted XXZ-type Hamiltonians with many modeling possibilities. Finally, in Section 6 we summarize our results and indicate some directions for further investigation.

2. The 2D stochastic Ising model in a field

The 2D Ising model in an external field is described by the Hamiltonian

$$H(\mathbf{S}) = -J \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} S_{\mathbf{r}} S_{\mathbf{r}'} - B \sum_{\mathbf{r}} S_{\mathbf{r}}, \quad (1)$$

where $\mathbf{S} = \{S_{\mathbf{r}} : \mathbf{r} \in \Lambda_L^N\}$ with $S_{\mathbf{r}} \in \{-1, +1\}$ are Ising spins, $\Lambda_L^N \subset \mathbb{Z}^2$ is a member of a family of semi-infinite lattices of $|\Lambda_L^N| = L \times \infty$ sites, and $\langle \mathbf{r}, \mathbf{r}' \rangle$ denotes pairs of nearest neighbor sites on Λ_L^N . The integer index N in Λ_L^N refers to the boundary conditions, that are free in the infinite direction and helicoidal with pitch N in the finite direction, i. e., $\mathbf{r} + L\mathbf{x} + N\mathbf{y} \equiv \mathbf{r}$ for all $\mathbf{r} \in \Lambda_L^N$. When $N = 0$ we recover the usual periodic boundary condition, which is however uninteresting for our purposes, as we will see later. In the above Hamiltonian we take $J > 0$, making the model ferromagnetic, and for definiteness we take $B \geq 0$.

We introduce a dynamics on the Ising spins through the master equation

$$\frac{d}{dt} P(\mathbf{S}, t) = \sum_{\tilde{\mathbf{S}} \in \Omega(\Lambda_L^N)} \left[W(\tilde{\mathbf{S}} \rightarrow \mathbf{S}) P(\tilde{\mathbf{S}}, t) - W(\mathbf{S} \rightarrow \tilde{\mathbf{S}}) P(\mathbf{S}, t) \right] \quad (2)$$

for the probability $P(\mathbf{S}, t)$ of observing the configuration $\mathbf{S} \in \Omega(\Lambda_L^N) = \{-1, +1\}^{\Lambda_L^N}$ at instant t , where $W(\mathbf{S} \rightarrow \tilde{\mathbf{S}})$ is the rate at which configuration $\tilde{\mathbf{S}}$ is reached from configuration \mathbf{S} per unit time. The rates $W(\mathbf{S} \rightarrow \tilde{\mathbf{S}})$ should be translation invariant and verify the condition of detailed balance $W(\tilde{\mathbf{S}} \rightarrow \mathbf{S}) P(\tilde{\mathbf{S}}) = W(\mathbf{S} \rightarrow \tilde{\mathbf{S}}) P(\mathbf{S})$, with $P(\mathbf{S}) \propto \exp[-\beta H(\mathbf{S})]$ the Gibbs equilibrium probability distribution and for the sake of notational economy we omitted the dependence of $P(\mathbf{S})$ and $W(\mathbf{S} \rightarrow \tilde{\mathbf{S}})$ on J , B , and the inverse temperature $\beta = 1/k_B T$. In this work we consider heat-bath single-spin flip transition rates given by

$$W(\mathbf{S} \rightarrow \tilde{\mathbf{S}}) \equiv W(S_{\mathbf{r}} \rightarrow \tilde{S}_{\mathbf{r}}) = \frac{1}{Z_{\mathbf{r}}} \exp[-\beta H(S_{\mathbf{r}})], \quad (3)$$

with

$$H(S_{\mathbf{r}}) = -J \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} S_{\mathbf{r}} S_{\mathbf{r}'} - B S_{\mathbf{r}} \quad \text{and} \quad Z_{\mathbf{r}} = \sum_{S_{\mathbf{r}} \in \{-1, +1\}} \exp[-\beta H(S_{\mathbf{r}})], \quad (4)$$

where $\langle \mathbf{r}' : \mathbf{r} \rangle = \{\mathbf{r}' \in \Lambda_L^N : |\mathbf{r}' - \mathbf{r}| = 1\}$.

Let $w(S_{\mathbf{r}}) = \frac{1}{2} \sum_{\langle \mathbf{r}', \mathbf{r} \rangle} |S_{\mathbf{r}'} - S_{\mathbf{r}}|$ be the number of spins neighboring $S_{\mathbf{r}}$ that have the sign opposite to it. In terms of this quantity, the single-spin flip rates read

$$W(S_{\mathbf{r}} \rightarrow \tilde{S}_{\mathbf{r}}) = \frac{1}{1 + \exp[4\beta J(2 - w(S_{\mathbf{r}})) + 2\beta B S_{\mathbf{r}}]}. \quad (5)$$

At sufficiently low temperatures, as long as $B < 2J$ spins with $w(S_{\mathbf{r}}) = 0, 1$ will hardly flip, because their transition rates become exponentially small when compared with the other rates, of the order of $\exp[-2\beta(2J - B)]$ at maximum. Processes with $w(S_{\mathbf{r}}) = 3, 4$ correspond to fast processes, since at sufficiently low temperatures and again in the range $B < 2J$




Fig. 1. Staircase-like 2D Ising interface separating the “+” and the “-” phases. In this figure, $L = 18$ and $N = 15$. When $B > 0$ ($B < 0$), the “+” (“-”) phase invades the “-” (“+”) phase, while for $B = 0$ the interface only fluctuates about its initial shape. When $\beta \rightarrow \infty$ and $|B| < 2J$, the number of bonds N is conserved.

their rates become close to unity, $W(S_r \rightarrow \tilde{S}_r) \geq 1 - \exp[-2\beta(2J - B)]$. In the low-temperature limit and in the range $B < 2J$, thus, the heat-bath single-spin flip rates (5) define a process in which only spins with $w(S_r) \geq 2$ have an appreciable flipping rate, and henceforth we ignore the flipping of spins with $w(S_r) < 2$.¹ Spins with $w(S_r) = 3, 4$, in turn, can be avoided by choosing initial configurations in which the “+” phase is separated from the “-” phase by a single-valued, non-self-intersecting staircase-like interface as in Fig. 1. With initial configurations of this type and within the regime of low temperatures and fields, we are left with a process in which only spins with $w(S_r) = 2$ flip. Since spins with $w(S_r) = 2$ lie at the interface, the above-defined spin flip dynamics actually defines an interface dynamics. In the next section we map this dynamics into an interacting particle system on the integers.

3. Mapping to the zero range process

Various possibilities exist to map the dynamics of 2D Ising interfaces into a system of interacting particles on the line. One possible map is obtained by associating with every vertical bond in the interface a particle and with every horizontal bond a hole [3]. In this way we end up with a simple exclusion process in which particles hop in an augmented one-dimensional lattice. Another possibility was given in Refs. [6,7]. In this case, one considers a set of L particles on a one-dimensional lattice occupying the positions x_ℓ , $1 \leq \ell \leq L$. If one associates the particle labels ℓ with a horizontal coordinate, and the particle positions x_ℓ with the heights of an interface, then one has a one-to-one map between the set of particles and an interface. With the additional constraint $x_{\ell+1} - x_\ell \geq 1$, the resulting model was called the particle-height model. The particle-height model thus establishes a map between the low-temperature dynamics of an Ising interface and the simple exclusion process, although the constraint on the particles’ positions seems a bit artificial in the interface scenario.

Our mapping of the Ising interface to a set of hopping particles on the integers is equivalent to the particle-height mapping with $x_{\ell+1} - x_\ell \geq 0$. Let Γ_L^N be the set of all single-valued, non-self-intersecting staircase-like Ising interface configurations γ_L^N of length $|\gamma_L^N| = L + N$, $N \geq 0$, in the infinite strip A_L^N of width L with helicoidal boundary condition $\mathbf{r} + L\mathbf{x} + N\mathbf{y} \equiv \mathbf{r}$. The dynamics of the γ_L^N interfaces under the action of the flipping rates $W(S_r)$ in the regime where $\beta \rightarrow \infty$ and $B < 2J$ preserves the length of the interfaces, i.e., given an initial configuration $\gamma_L^N(t = 0) \in \Gamma_L^N$, all subsequent configurations $\gamma_L^N(t \geq 0) \in \Gamma_L^N$. The number N of vertical bonds in the interface thus decomposes the state space of interface configurations into an infinite number of disjoint sectors, $\Gamma_L = \bigcup_{N \geq 0} \Gamma_L^N$.

The interface configurations γ_L^N are single-valued functions with respect to the horizontal coordinate. Let $h_\ell \in \mathbb{Z}$, $1 \leq \ell \leq L$ denote the height of the Ising interface at site ℓ . Then the height differences $n_\ell = h_{\ell+1} - h_\ell$ are nonnegative due to the special form of γ_L^N , and their dynamics relates to the dynamics of the interface as follows. Let us consider the case

¹ In realistic pseudo-two-dimensional $S = \frac{1}{2}$ Ising-like materials, e.g. in the antiferromagnetic compounds K_2XF_4 with $X = \text{Mn, Fe, Co, or Ni}$, $J/k_B \sim 1-100$ K, such that $\beta J \gg 1$ implies $T < 1$ K [31]. The values for which $B < 2J$ thus lie in the range $B \lesssim 1.5$ T, of the order of half the magnetic field strength of a typical medical MRI system.

$0 < B < 2J$, and that the “+” phase lies below the “−” phase, as in Fig. 1. In this way, every “−” (“+”) spin that flips contributes to the growth of the “+” (“−”) phase, increasing (decreasing) the height variable associated with its horizontal position by one unit. The single-spin flipping rates are given by

$$- \rightarrow + : p = \frac{1}{1 + e^{-2\beta B}} \quad \text{and} \quad + \rightarrow - : q = \frac{1}{1 + e^{2\beta B}} = 1 - p. \tag{6}$$

When $B = 0$, we have a spin (time) reversal invariant system, and the interface does not move bodily. Otherwise, when $0 < B < 2J$ and $\beta \rightarrow \infty$, $p = 1$ and $q = 0$, and the surface can only grow. The general situation $0 < q < p < 1$ is obtained by taking $\beta \rightarrow \infty$ and $B \rightarrow 0$ with $\beta B = \text{constant}$. The height differences moves corresponding to the flipping processes are $(n_{\ell-1}, n_{\ell}) \rightarrow (n_{\ell-1} - 1, n_{\ell} + 1)$ when $h_{\ell} \rightarrow h_{\ell} + 1$, and $(n_{\ell-1}, n_{\ell}) \rightarrow (n_{\ell-1} + 1, n_{\ell} - 1)$ when $h_{\ell} \rightarrow h_{\ell} - 1$. The maximum possible height difference in Λ_L^N is $n_{\ell} = N$, in which case $h_1 = h_2 = \dots = h_{\ell} = 0$ and $h_{\ell+1} = h_{\ell+2} = \dots = h_L = N$ for some ℓ . More generally, we have $\sum_{\ell=1}^{L-1} n_{\ell} = h_L - h_1 = N$. Now one appreciates the role of the helicoidal boundary conditions on Λ_L^N : the pitch N gives the total number of interacting particles in the height differences scenario. The boundary condition in the height differences scenario is simply periodic.

The dynamics of the variables n_{ℓ} is but the zero range dynamics, in which particles hop on the lattice without exclusion [21–27]. Except for the (immaterial) absolute values of the heights, the dynamics of the n_{ℓ} variables contains all the information about the evolving Ising interface. The elementary processes for the n_{ℓ} variables are

$$(n_{\ell-1} + 1, n_{\ell}) \xrightleftharpoons[p]{q} (n_{\ell-1}, n_{\ell} + 1), \quad 0 \leq n_{\ell-1}, n_{\ell} \leq N - 1, \quad 2 \leq \ell \leq L. \tag{7}$$

As is well known [32,33], we may write the master equation for reaction–diffusion processes on the lattice as a Schrödinger-like equation in Euclidean time, the infinitesimal generator of the Markov semigroup playing the role of the quantum Hamiltonian. In this scenario, the infinitesimal generator of the above zero range process is given by the $N^L \times N^L$ operator

$$H_N = \sum_{\ell=1}^L \sum_{m+n=0}^{N-1} \left[p \left(E_{\ell}^{m+1, m+1} E_{\ell+1}^{n, n} - E_{\ell}^{m, m+1} E_{\ell+1}^{n+1, n} \right) + q \left(E_{\ell}^{m, m} E_{\ell+1}^{n+1, n+1} - E_{\ell}^{m+1, m} E_{\ell+1}^{n, n+1} \right) \right], \tag{8}$$

where $E_{\ell}^{m, n} = \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes E^{m, n} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1}$, with $\mathbf{1}$ the $N \times N$ identity matrix and $(E^{m, n})_{i, j} = \delta_{i, m} \delta_{j, n}$ the $N \times N$ matrix with a single unit element in row m and column n occupying the ℓ -th position in the direct product. The operator H_N can be diagonalized by the coordinate Bethe ansatz and, indeed, it has been diagonalized by this and related methods many times in the literature [28–30,34]. We will thus not reproduce a complete resolution of (8) here. Instead, we just outline the technique and quote the main results regarding the process defined by H_N of interest to us.

4. Bethe ansatz solution

4.1. Bethe ansatz equations

We are interested in the solutions of the eigenvalue equation

$$H_N |\Psi_N\rangle = E_N |\Psi_N\rangle, \tag{9}$$

where H_N is given in (8) and

$$|\Psi_N\rangle = \sum_{x_1 \leq x_2 \leq \dots \leq x_N} \Phi(x_1, x_2, \dots, x_N) |x_1, x_2, \dots, x_N\rangle \tag{10}$$

is the eigenfunction written in the basis that specifies the positions of the N particles in the system, with $\Phi(x_1, x_2, \dots, x_N)$ the coefficient for the configuration $|x_1, x_2, \dots, x_N\rangle$. Notice that since there is no exclusion, particle positions can coincide.

If the positions of the particles obey $x_{j+1} > x_j$, $1 \leq j \leq N$, the eigenvalue equation (9) is satisfied by the ansatz (10) with coefficients

$$\Phi(x_1, x_2, \dots, x_N) = \sum_P A_{P(1)P(2)\dots P(N)} \exp \left[i \sum_{j=1}^N k_{P(j)} x_j \right] \tag{11}$$

and eigenvalue

$$E_N(k_1, k_2, \dots, k_N) = \sum_{j=1}^N \epsilon(k_j), \tag{12}$$

where $\epsilon(k) = 1 - pe^{-ik} - qe^{ik}$ is the “single particle energy,” the first summation in (11) is over all the $N!$ permutations P of the indices $(1, 2, \dots, N)$ used to label the positions of the particles, and the “wave numbers” k_1, k_2, \dots, k_N are chosen so that $|\Psi_N\rangle$ satisfies (9). We see that when the particles are far apart, in the case being just not on the same site, they behave

as if they were free, and the total “energy” of the system is the sum of the “energies” of single particles. When a pair of particles sit on the same site, $x_{j+1} = x_j$, we obtain from (9)–(12) that the amplitudes $A_{P(1)P(2)\dots P(N)}$ should satisfy

$$\frac{A_{P(1)P(2)\dots P(j)P(j+1)\dots P(N)}}{A_{P(1)P(2)\dots P(j+1)P(j)\dots P(N)}} = -\frac{e^{ik_{P(j+1)}}}{e^{ik_{P(j)}}} e^{i\Theta_{P(j)P(j+1)}}, \tag{13}$$

where the “two-particle scattering phase” $\Theta_{j\ell}$ is defined by

$$e^{i\Theta_{j\ell}} = \frac{p + qe^{i(k_j+k_\ell)} - e^{ik_j}}{p + qe^{i(k_j+k_\ell)} - e^{ik_\ell}}. \tag{14}$$

The boundary condition $\Phi(x_2, x_3, \dots, x_{1+L}) = \Phi(x_1, x_2, \dots, x_N)$ furnishes the additional relation

$$A_{P(1)P(2)\dots P(N)} = e^{ik_{P(1)}L} A_{P(2)P(3)\dots P(1)}. \tag{15}$$

Iterating relation (13) N times, (15) gives us the Bethe ansatz equations for the “wave numbers” k_j in the N -particle sector,

$$e^{ik_jL} = (-1)^{N-1} \prod_{\ell=1}^N \left(\frac{e^{ik_\ell}}{e^{ik_j}} \right) e^{i\Theta_{j\ell}} = (-1)^{N-1} \prod_{\ell=1}^N \left(\frac{e^{ik_\ell}}{e^{ik_j}} \right) \frac{p + qe^{i(k_j+k_\ell)} - e^{ik_j}}{p + qe^{i(k_j+k_\ell)} - e^{ik_\ell}}, \quad 1 \leq j \leq N. \tag{16}$$

The solutions k_1, k_2, \dots, k_N of these equations give through (12) the eigenvalues of (8). Notice that since H_N is in general nonhermitian, the k_j are in general complex numbers.

The eigenfunctions (10) with the coefficients (11) should also be eigenfunctions of the translation operator T that shifts the positions of the particles to the left by one site, since $[H, T] = 0$. The eigenvalues e^{iP} of T are given by

$$T|\Psi_N\rangle = e^{iP}|\Psi_N\rangle = \left(\prod_{j=1}^N e^{ik_j} \right) |\Psi_N\rangle, \tag{17}$$

where we have defined the total momentum-like P by

$$P = \sum_{j=1}^N k_j \pmod{2\pi} = \frac{2\pi\ell}{L}, \quad 0 \leq \ell \leq L-1. \tag{18}$$

With P defined above, Eq. (16) can be rewritten as

$$e^{ik_j(L+N)} = (-1)^{N-1} e^{i\sum_{\ell=1}^N k_\ell} \prod_{\ell=1}^N e^{i\Theta_{j\ell}} = (-1)^{N-1} e^{iP} \prod_{\ell=1}^N \frac{p + qe^{i(k_j+k_\ell)} - e^{ik_j}}{p + qe^{i(k_j+k_\ell)} - e^{ik_\ell}}, \quad 1 \leq j \leq N. \tag{19}$$

The learned reader will recognize in (19) the Bethe ansatz equations for the asymmetric simple exclusion process of N particles in a lattice of $L + N$ sites with twisted boundary conditions, the angle of twist being given by the total momentum P of the system. For stochastic processes, the relevant momentum sector is the $P = 0$ sector, since the coefficients $\Phi(x_1, x_2, \dots, x_N)$ have to be all real and positive. In this sector, the correspondence between the zero range process and the asymmetric simple exclusion process is exact.

4.2. The dynamical critical exponent

Numerical simulations together with theoretical arguments and explicit calculations indicate that the critical behavior of the interface is independent of the particular values of p and q as long as $p \neq q$ [13–15,17]. It has then become usual to investigate the Bethe ansatz equations (16) with $p = 1, q = 0$, since this facilitates the analysis considerably. The more general cases $0 < q < p < 1$ were investigated in Refs. [15,35]. The $p = 1, q = 0$ case corresponds to a 2D Ising interface evolving in a finite field $B > 0$ but at zero temperature. However, at least for very low (but nonzero) temperatures, one has the same kind of critical behavior as observed at zero temperature [36]. In the totally asymmetric simple exclusion process, another simplification of the Bethe ansatz equations comes with the choice of the half-filled sector $2N = L$. The analogous choice for the zero range process is to consider the sector with $N = L$, which corresponds in the interface scenario to an interface with average slope $\pi/4$.

The dynamical critical exponent z , that measures the degree of anisotropy between the spatial and temporal correlation lengths, can be determined from the asymptotic behavior of the gap $E_N^{(1)}(L)$ of H_N through $\text{Re}\{E_N^{(1)}(L)\} \sim L^{-z}$. For the asymmetric exclusion process with arbitrary $p \neq q$ and $\varrho = N/L$, the large L asymptotic value of $E_N^{(1)}(L)$ is given by

$$E_N^{(1)}(L) = -2C|p - q|\sqrt{\varrho(1 - \varrho)}L^{-3/2} \pm 2\pi i|(p - q)(1 - 2\varrho)|L^{-1}, \tag{20}$$

with an exact (numerically evaluated) $C = 6.509\,189 \dots$ [35]. The dynamical critical exponent of the asymmetric exclusion process is then $z = 3/2$, indicating that it belongs to the Kardar–Parisi–Zhang universality class of critical behavior [18].

For the zero range process, the available calculations of $E_N^{(1)}(L)$ are based on the analysis of slightly generalized models, with non-uniform hopping rates or in which particles can hop together [26,27]. The totally asymmetric zero-range process with uniform rates in the $N = L$ sector was solved in Refs. [29,30], with the result that the first gap behaves like

$$E_N^{(1)}(L) \sim a_0 L^{-3/2} - i \frac{\pi}{2} L^{-1}, \quad (21)$$

with an exact (numerically evaluated) $a_0 = 2.301\,345 \dots$, first obtained in Ref. [14]; see also [15].² We thus see that, in either case, the gap behaves asymptotically as $E_N^{(1)}(L) \sim L^{-3/2}$, the dynamical critical exponent $z = 3/2$, and both processes – the asymmetric simple exclusion process and the asymmetric zero-range process – belong to the Kardar–Parisi–Zhang universality class of critical behavior, and so does the driven interface dynamics of the 2D stochastic Ising model in the regime of low temperatures and fields.

5. Generalized particle-height model and interface dynamics

The particle-height model mentioned in Section 3 can be generalized to processes in which the particle positions observe the constraint $x_{\ell+1} - x_\ell \geq s$, generating a process in which particles move only if the next particle is far apart by at least s sites. Identifying a particle with the up spin state and a hole with the down spin state in the σ^z basis, the time evolution of this restricted exclusion processes is governed by the infinitesimal generator [29,30]

$$H_s = - \sum_{\ell=1}^L P_s \left[p \sigma_\ell^- \sigma_{\ell+1}^+ + q \sigma_\ell^+ \sigma_{\ell+1}^- + \frac{1}{4} (\sigma_\ell^z \sigma_{\ell+1}^z - \mathbf{1}) \right] P_s, \quad (22)$$

where p and $q = 1 - p$ are the rates at which particles hop respectively to the right and to the left,

$$P_s = \prod_{\ell=1}^L \left[\frac{1}{2} (\mathbf{1} - \sigma_\ell^z) + \frac{1}{2} (\mathbf{1} + \sigma_\ell^z) \prod_{j=1}^{s-1} \frac{1}{2} (\mathbf{1} - \sigma_{\ell+j}^z) \right] \quad (23)$$

is the operator that projects out configurations in which particles are closer than by s sites, $\sigma^\pm = \frac{1}{2} (\sigma^x \pm i \sigma^y)$ and σ^z are the usual Pauli spin- $\frac{1}{2}$ matrices, and $\mathbf{1}$ is the 2×2 identity matrix.

The infinitesimal generator H_s is exactly solvable by the Bethe ansatz for any choice of the integer s [29,30]. The $s = 1$ case recovers the generator of the asymmetric simple exclusion process, while the generic $s > 1$ case describes, e.g., the driven diffusion of s -mers on the lattice [37]. In fact, it has been shown that an arbitrary mixture of s -mers of different sizes, diffusing with the same rates but ruled by evolution operators H_s with different s , can be integrated exactly, with the eigenspectrum depending only on the average size \tilde{s} of the s -mers [29,30].

For the generalized particle-height model, either with a single type of particle or with an admixture of particles of different sizes, the simplifying filling fraction analogous to the condition $N = L$ in the zero-range process is given by $(1 + \tilde{s})N = L$, where \tilde{s} is the average size of the particles in the system, not necessarily a semipositive integer. Clearly, the larger the average particle size \tilde{s} , the smaller the average interface slope $N/L = 1/1 + \tilde{s}$. In this case, the spectral gap of the process has been found to scale like [29,30]

$$E_s^{(1)}(L) \sim a_0 L^{-3/2} + i \left(\frac{\tilde{s} - 1}{\tilde{s} + 1} \right) \frac{\pi}{2} L^{-1}, \quad (24)$$

with the same $a_0 = 2.301\,345 \dots$ as before; compare with (21). We can then predict that the interface dynamics obtained from the generalized particle-height model with $x_{\ell+1} - x_\ell \geq s$ also belongs to the Kardar–Parisi–Zhang universality class of critical behavior.

A most interesting thing would be to visualize how the Ising interface configurations evolve in the case of a particle-height model that includes besides particles of positive sizes also particles with negative sizes, since in this case handles and loops could develop. More generally, it would be of interest to find physical applications of the operator (22) for negative values of s or \tilde{s} , since they are all exactly solvable and display the same type of critical behavior.

6. Summary and conclusions

We showed that it is possible to map the interface dynamics of the 2D stochastic Ising model in the regime of low temperatures and fields into an exactly solvable interacting particle system of hopping particles without exclusion. The infinitesimal generator of the process is exactly solvable by the Bethe ansatz, with an spectral gap in the asymmetric case scaling like $L^{-3/2}$ with the system size. The 2D Ising interface in the presence of a driving field then grows and fluctuates according to the Kardar–Parisi–Zhang universality class of critical behavior. We remark that most studies (e.g., in the realm

² Notice that the constants C in (20) and a_0 in (21) are related by $C = 2\sqrt{2}a_0$.

of nucleation dynamics) of 2D Ising interfaces are carried out in zero temperature, and several results were obtained in the absence of external fields. In our study, following [5,36], we allow for finite temperatures and fields, as long as the conditions stated in Section 2 are met. When $\beta = \infty$ ($T = 0^+$) or $B = 0$, the process becomes symmetric and the Bethe ansatz analysis reduces to a simple spin-wave analysis. In this case the gap becomes $E_N^{(1)}(L) = 2 \sin^2(\pi/L)$ (independent of N as long as $N < L$), with asymptotic behavior $E_N^{(1)}(L \gg 1) = 2\pi^2 L^{-2}$, and the interface only fluctuates, without moving or growing, according to the $z = 2$ Edwards–Wilkinson universality class of critical behavior [38].

It would be desirable to explore the mapping of the Ising interface to the zero range process to investigate step-step correlation functions by tagged-particle methods within the context of exact Bethe solutions [6,7,9,19,20,23,39], as well as the dynamics of special configurations like semi-infinite strips of the minority phase (“Ising fingers”) [40] using some of the ideas exposed here. We hope to return to these subjects soon.

Acknowledgments

The author thanks Professor Francisco C. Alcaraz (IFSC/USP) for having called his attention to operator (22) and for helpful conversations. This work was partially supported by the Brazilian agency CNPq through grant PDS 151999/2010-4.

References

- [1] H. Rost, Nonequilibrium behaviour of many-particle processes: density profile and local equilibrium, *Z. Wahrscheinlichkeitstheor. Verwandte Geb.* 58 (1981) 41–53.
- [2] J.-P. Marchand, Ph.A. Martin, A microscopic derivation of the classical nucleation equation, *Physica A* 127 (1984) 681–691; Exclusion process and droplet shape, *J. Stat. Phys.* 44 (1986) 491–504; *J. Stat. Phys.* 50 (1988) 469–471 (erratum).
- [3] D. Kandel, E. Domany, Rigorous derivation of domain growth kinetics without conservation laws, *J. Stat. Phys.* 58 (1990) 685–706.
- [4] D. Kandel, E. Domany, B. Nienhuis, A six-vertex model as a diffusion problem: derivation of correlation functions, *J. Phys. A: Math. Gen.* 23 (1990) L755–L762.
- [5] A.L.C. Ferreira, S.K. Mendiratta, Simulation of domain-wall dynamics in the SOS approximation of the 2D anisotropic Ising model: field driven case, *J. Phys. A: Math. Gen.* 24 (1991) 4397–4406.
- [6] S.N. Majumdar, M. Barma, Tag diffusion in driven systems, growing interfaces, and anomalous diffusion, *Phys. Rev. B* 44 (1991) 5306–5308; Two-tag correlation functions in one-dimensional lattice gases, *Physica A* 177 (1991) 366–372.
- [7] M. Barma, Dynamics of field-driven interfaces in the two-dimensional Ising model, *J. Phys. A: Math. Gen.* 25 (1992) L693–L699.
- [8] H. Spohn, Interface motion in models with stochastic dynamics, *J. Stat. Phys.* 71 (1993) 1081–1132.
- [9] P.M. Binder, M. Paczuski, M. Barma, Scaling of fluctuations in one-dimensional interface and hopping models, *Phys. Rev. E* 49 (1994) 1174–1181.
- [10] D.B. Abraham, T.J. Newman, G.M. Schütz, Nonequilibrium dynamics of finite interfaces, *Phys. Rev. Lett.* 72 (1994) 3266–3269.
- [11] T. Strobel, Interface motion in a planar spin-flip model derived from exclusion on the line, *J. Stat. Phys.* 79 (1995) 923–950.
- [12] D. Dhar, An exactly solved model for interfacial growth, *Phase Transit.* 9 (1987) 51–51.
- [13] J. Krug, H. Spohn, Kinetic roughening of growing interfaces, in: C. Godrèche (Ed.), *Solids Far from Equilibrium: Growth, Morphology, and Defects*, Cambridge University Press, Cambridge, 1992, pp. 479–582.
- [14] L.-H. Gwa, H. Spohn, Six-vertex model, roughened surfaces, and an asymmetric spin Hamiltonian, *Phys. Rev. Lett.* 68 (1992) 725–728; Bethe solution for the dynamical-scaling exponent of the noisy Burgers equation, *Phys. Rev. A* 46 (1992) 844–854.
- [15] D. Kim, Bethe ansatz solution for crossover scaling functions of the asymmetric XXZ chain and the Kardar–Parisi–Zhang-type growth model, *Phys. Rev. E* 52 (1995) 3512–3524.
- [16] P. Meakin, P. Ramanlal, L.M. Sander, R.C. Ball, Ballistic deposition on surfaces, *Phys. Rev. A* 34 (1986) 5091–5103.
- [17] M. Plischke, Z. Rácz, D. Liu, Time-reversal invariance and universality of two-dimensional growth models, *Phys. Rev. B* 35 (1987) 3485–3495.
- [18] M. Kardar, G. Parisi, Y.-C. Zhang, Dynamic scaling of growing interfaces, *Phys. Rev. Lett.* 56 (1986) 889–892.
- [19] A. De Masi, P.A. Ferrari, Self-diffusion in one-dimensional lattice gases in the presence of an external field, *J. Stat. Phys.* 38 (1985) 603–613.
- [20] P.A. Ferrari, The simple exclusion process as seen from a tagged particle, *Ann. Probab.* 14 (1986) 1277–1290.
- [21] F. Spitzer, Interaction of Markov processes, *Adv. Math.* 5 (1970) 246–290.
- [22] T.M. Liggett, Infinite particle systems with zero range interactions, *Ann. Probab.* 1 (1973) 240–253.
- [23] E.D. Andjel, Invariant measures for the zero range process, *Ann. Probab.* 10 (1982) 525–547.
- [24] M.R. Evans, Phase transitions in one-dimensional nonequilibrium systems, *Braz. J. Phys.* 30 (2000) 42–57; M.R. Evans, R.A. Blythe, Nonequilibrium dynamics in low-dimensional systems, *Physica A* 313 (2002) 110–152; M.R. Evans, T. Hanney, Nonequilibrium statistical mechanics of the zero-range process and related models, *J. Phys. A: Math. Gen.* 38 (2005) R195–R239.
- [25] M.R. Evans, E. Levine, P.K. Mohanty, D. Mukamel, Modelling one-dimensional driven diffusive systems by the zero-range process, *Eur. Phys. J. B* 41 (2004) 223–230.
- [26] T. Sasamoto, M. Wadati, One-dimensional asymmetric diffusion model without exclusion, *Phys. Rev. E* 58 (1998) 4181–4190.
- [27] A.M. Povolotsky, Bethe ansatz solution of zero-range process with non-uniform stationary state, *Phys. Rev. E* 69 (2004) 061109.
- [28] F.C. Alcaraz, R.Z. Bariev, Exact solution of a vertex model with unlimited number of states per bond, *J. Phys. A: Math. Gen.* 32 (1999) L25–L31.
- [29] F.C. Alcaraz, R.Z. Bariev, Exact solution of the asymmetric exclusion model with particles of arbitrary size, *Phys. Rev. E* 60 (1999) 79–88.
- [30] F.C. Alcaraz, R.Z. Bariev, An exactly solvable constrained XXZ chain, in: M.T. Batchelor, L.T. Wille (Eds.), *Statistical Physics on the Eve of the 21st Century—In Honour of J.B. McGuire on the Occasion of his 65th Birthday*, in: Series on Advances in Statistical Mechanics, vol. 14, World Scientific, Singapore, 1999, pp. 412–424.
- [31] L.J. de Jongh, A.R. Miedema, Experiments on simple magnetic model systems, *Adv. Phys.* 23 (1973) 1–260; W.B. Wolf, The Ising model and real magnetic materials, *Braz. J. Phys.* 30 (2000) 794–810.
- [32] F.C. Alcaraz, M. Droz, M. Henkel, V. Rittenberg, Reaction–diffusion processes, critical dynamics, and quantum chains, *Ann. Phys. NY* 230 (1994) 250–302.
- [33] G.M. Schütz, Exactly solvable models for many-body systems far from equilibrium, in: C. Domb, J. Lebowitz (Eds.), *Phase Transitions and Critical Phenomena*, Vol. 19, Academic Press, London, 2000, pp. 1–251.
- [34] F.C. Alcaraz, M.J. Lazo, The exact solution of the asymmetric exclusion problem with particles of arbitrary size: matrix product ansatz, *Braz. J. Phys.* 33 (2003) 533–549.
- [35] O. Golinelli, K. Mallick, Spectral gap of the totally asymmetric exclusion process at arbitrary filling, *J. Phys. A: Math. Gen.* 38 (2005) 1419–1425; K. Mallick, Some exact results for the exclusion process, *J. Stat. Mech.* (2011) P01024.
- [36] P. Devillard, Interface motion in a two-dimensional Ising model with a field, *J. Stat. Phys.* 62 (1991) 443–451; P. Devillard, H. Spohn, Kinetic shape of Ising clusters, *Europhys. Lett.* 17 (1992) 113–118.
- [37] S. Gupta, M. Barma, U. Basu, P.K. Mohanty, Driven k-mers: correlations in space and time, *Phys. Rev. E* 84 (2011) 041102.
- [38] S.F. Edwards, D.R. Wilkinson, The surface statistics of a granular aggregate, *Proc. R. Soc. Lond. Ser. A* 381 (1982) 17–31.
- [39] A.A. Ferreira, F.C. Alcaraz, Anomalous tag diffusion in the asymmetric exclusion model with particles of arbitrary sizes, *Phys. Rev. E* 65 (2002) 052102.
- [40] P.L. Krapivsky, Limiting shapes of Ising droplets, Ising fingers, and Ising solitons, *Phys. Rev. E* 85 (2012) 011152.