



ELSEVIER

Linear Algebra and its Applications 356 (2002) 253–274

**LINEAR ALGEBRA
AND ITS
APPLICATIONS**

www.elsevier.com/locate/laa

Forest matrices around the Laplacian matrix

Pavel Chebotarev*, Rafiq Agaev

*Trapeznikov Institute of Control Sciences of the Russian Academy of Sciences, 65 Profsoyuznaya str.,
Moscow 117997, Russia*

Received 4 November 2001; accepted 16 May 2002

Submitted by W. Haemers

Abstract

We study the matrices Q_k of in-forests of a weighted digraph Γ and their connections with the Laplacian matrix L of Γ . The (i, j) entry of Q_k is the total weight of spanning converging forests (*in-forests*) with k arcs such that i belongs to a tree rooted at j . The forest matrices, Q_k , can be calculated recursively and expressed by polynomials in the Laplacian matrix; they provide representations for the generalized inverses, the powers, and some eigenvectors of L . The normalized in-forest matrices are row stochastic; the normalized matrix of maximum in-forests is the eigenprojection of the Laplacian matrix, which provides an immediate proof of the Markov chain tree theorem. A source of these results is the fact that matrices Q_k are the matrix coefficients in the polynomial expansion of $\text{adj}(\lambda I + L)$. Thereby they are precisely Faddeev's matrices for $-L$. © 2002 Elsevier Science Inc. All rights reserved.

AMS classification: 05C50; 15A48

Keywords: Weighted digraph; Laplacian matrix; Spanning forest; Matrix-tree theorem; Matrix-forest theorem; Leverrier–Faddeev method; Markov chain tree theorem; Eigenprojection; Generalized inverse

1. Introduction

According to the matrix-tree theorem, the (i, j) cofactor of the Laplacian matrix of a weighted digraph equals the total weight of spanning converging trees rooted at vertex i of the digraph.

Fiedler and Sedláček [25] proved that the principal minor of the Laplacian matrix resulting by the removal of the rows and columns indexed by a set \mathcal{J} is equal to the total weight of in-forests with $|\mathcal{J}|$ trees rooted at the vertices of \mathcal{J} .

* Corresponding author.

E-mail addresses: chv@lpi.ru, pchv@rambler.ru (P. Chebotarev), arpo@ipu.rssi.ru (R. Agaev).

These results are generalized by the *all minors matrix tree theorem* [10,17] (see also [53]) which expresses arbitrary minors of the Laplacian matrix in terms of in-forests of the digraph.

We study the matrices, Q_k , of a digraph's in-forests: the (i, j) entry of Q_k is the total weight of in-forests with k arcs where i belongs to a tree converging to j . In this paper, we show that the forest matrices can be recursively calculated and represented by simple polynomials in the Laplacian matrix L ; in turn, the powers of L are linear combinations of Q_k 's. Further, we demonstrate that the forest matrices are useful to interpret a number of expressions that involve the Laplacian matrix, including those of the group and Moore-Penrose inverses, and some eigenvectors. Of special interest is the normalized matrix \tilde{J} of *maximum in-forests* of a digraph previously used [42,43] to represent the long run transition probabilities of Markov chains. We prove that \tilde{J} is the eigenprojection of the Laplacian matrix corresponding to the eigenvalue 0 and study some properties of \tilde{J} .

A seminal result that enables one to give short algebraic proofs to these representations is the fact that matrices Q_k coincide with the matrix coefficients in the polynomial form of $\text{adj}(\lambda I + L)$:

$$\text{adj}(\lambda I + L) = \sum_{k=0}^{n-1} Q_{n-k-1} \lambda^k,$$

where $\text{adj} A$ is the transposed matrix of cofactors of A . This expansion is a corollary to the *parametric matrix-forest theorem* [1,16] which expresses the entries of $(I + \tau L)^{-1}$, $\tau \in \mathbb{R}$ in terms of in-forests.

All results of this paper are applicable to unweighted digraphs (by taking all weights equal to one) and undirected graphs (by considering symmetric digraphs).

The paper is organized as follows. After the notation section, we briefly survey the major known results on the minors of the Laplacian (Kirchhoff) matrix of a weighted digraph (Section 3), give a new proof to the matrix-forest theorem for digraphs (Section 4), present a recursive method for calculating the forest matrices (Section 5), establish polynomial representations of the forest matrices (Section 6), study the normalized matrix \tilde{J} of maximum in-forests (Section 7), consider the linear transformations corresponding to L and \tilde{J} and show that \tilde{J} is the eigenprojection of L , which yields the *Markov chain tree theorem* (Section 8), and finally, express the generalized inverses of L in terms of the forest matrices (Section 9).

2. Notation

2.1. Graph definitions

For graph terminology, we mainly follow [30]. Suppose that Γ is a weighted digraph without loops, $V(\Gamma) = \{1, \dots, n\}$, $n > 1$, is its set of vertices and $E(\Gamma)$ its set

of arcs. The weights of all arcs are strictly positive. Let $W = (w_{ij})$ be the matrix of arc weights of Γ . Its (i, j) entry, w_{ij} , equals zero iff there is no arc from vertex i to vertex j in Γ . If Γ' is a subgraph of Γ , then the weight of Γ' , $w(\Gamma')$, is the product of the weights of all its arcs; if Γ' does not contain arcs, then $w(\Gamma') = 1$. The weight of a nonempty set of digraphs \mathcal{G} is defined as follows:

$$w(\mathcal{G}) = \sum_{H \in \mathcal{G}} w(H); \quad w(\emptyset) = 0. \tag{1}$$

A *spanning* subgraph of Γ is a subgraph of Γ with vertex set $V(\Gamma)$. The *out-degree* of vertex v is the number of arcs that come from v . A *converging tree* is a weakly connected (i.e., its corresponding undirected graph is connected) digraph in which one vertex, called the *root*, has outdegree zero and the remaining vertices have outdegree 1.

A converging tree is said to *converge* to its root. Spanning converging trees are sometimes called *in-arborescences*. A *converging forest* is a digraph all of whose weak components (i.e., maximal weakly connected subgraphs) are converging trees. The roots of these trees are the roots of the converging forest.

Definition 1. An *in-forest* is a spanning converging forest.

Definition 2. An in-forest F of a digraph Γ is called a *maximum in-forest* of Γ if Γ has no in-forest with a greater number of arcs than in F .

Out-forests which *diverge* from their roots and *maximum out-forests* are defined in the same manner. In this paper, we deal with in-forests, but a parallel theory can be developed for out-forests.

The notion of maximum in-forest of a digraph generalizes the concept of spanning converging tree (sometimes called in-arborescence). If spanning converging trees of a digraph exist, they coincide with maximum in-forests; otherwise maximum in-forests inherit some of their properties. These properties were studied in [1].

It is easily seen that every maximum in-forest of Γ has the minimum possible number of converging trees; we call this number the *in-forest dimension* of Γ and denoted it by d . The number of arcs in any maximum in-forest is obviously $n - d$; in general, the number of disjoint trees in a spanning forest with k arcs is $n - k$.

By $\mathcal{F}^{\rightarrow*}(\Gamma) = \mathcal{F}^{\rightarrow*}$ and $\mathcal{F}_k^{\rightarrow*}(\Gamma) = \mathcal{F}_k^{\rightarrow*}$ we denote the set of all in-forests of Γ and the set of all in-forests of Γ with k arcs, respectively; $\mathcal{F}_k^{i \rightarrow *j}$ will designate the set of all in-forests with k arcs where i belongs to a tree converging to j ; $\mathcal{F}^{i \rightarrow *j} = \bigcup_{k=0}^{n-d} \mathcal{F}_k^{i \rightarrow *j}$ is the set of such in-forests with any number of arcs. The notation $\mathcal{F}_{(k)}^{\rightarrow*}$ will be used for the set of in-forests that consist of k trees. The $\rightarrow*$ sign relates to in-forests; the corresponding notation for out-forests is $\mathcal{F}^{*\rightarrow}$, etc.

Let

$$\sigma_k = w(\mathcal{F}_k^{\rightarrow*}), \quad k = 0, 1, \dots, \tag{2}$$

$$\sigma = w(\overline{\mathcal{F}} \rightarrow^*) = \sum_{k=0}^{n-d} \sigma_k. \quad (3)$$

By (2) and (1), $\sigma_k = 0$ whenever $k > n - d$, and $\sigma_0 = 1$.

We will also consider the parametric value

$$\sigma(\tau) = \sum_{k=0}^{n-d} \sigma_k \tau^k, \quad (4)$$

which is the total weight of in-forests in Γ provided that all arc weights are multiplied by τ .

Let

$$s_k = \sum_{j=0}^k \sigma_j, \quad k = 0, \dots, n - d \quad (5)$$

be the total weight of in-forests of Γ with at most k arcs. Then, by definition, $s_{n-d} = \sigma$.

Finally,

$$s_k(\tau) = \sum_{j=0}^k \sigma_j \tau^j, \quad k = 0, \dots, n - d, \quad (6)$$

whence $s_{n-d}(\tau) = \sigma(\tau)$.

2.2. Matrix definitions

For any $n \times n$ matrix A , let $A(\overline{\mathcal{I}} | \overline{\mathcal{J}})$, where $\mathcal{I}, \mathcal{J} \subseteq \{1, \dots, n\}$, be the submatrix of A obtained by the removal of the rows indexed by \mathcal{I} and the columns indexed by \mathcal{J} . For a complex matrix A , A^* is the conjugate transpose (Hermitian adjoint) and A^T the transpose of A .

The *Laplacian* (or *row Laplacian*) matrix of a weighted digraph Γ is the $n \times n$ matrix $L = L(\Gamma) = (\ell_{ij})$ with entries $\ell_{ij} = -w_{ij}$ when $j \neq i$ and $\ell_{ii} = -\sum_{k \neq i} \ell_{ik}$, $i, j = 1, \dots, n$. The *column Laplacian* matrix $L' = L'(\Gamma) = (\ell'_{ij})$ differs from L by the diagonal only: $\ell'_{ij} = -w_{ij}$ when $j \neq i$ and $\ell'_{ii} = -\sum_{k \neq i} \ell'_{ki}$, $i, j = 1, \dots, n$. The *Kirchhoff* (or *row Kirchhoff*) matrix [65] is $K = L'^T$; the *column Kirchhoff* matrix is $K' = L^T$. These four singular matrices are generalizations of the Laplacian (Kirchhoff) matrix of an undirected graph. In what follows, we deal with the Laplacian matrix $L(\Gamma)$ and reformulate for it some results originally obtained for the other matrices.

Throughout let Γ be a fixed digraph. Consider the *matrices*

$$Q_k = (q_{ij}^k), \quad k = 0, 1, \dots,$$

of in-forests of Γ with k arcs: the entries of Q_k are

$$q_{ij}^k = w(\mathcal{F}_k^{i \rightarrow *j}). \tag{7}$$

By (7) and (1), $Q_k = 0$ whenever $k > n - d$, and $Q_0 = I$. The matrix of all in-forests is

$$Q = (q_{ij}) = \sum_{k=0}^{n-d} Q_k \tag{8}$$

with entries $q_{ij} = w(\mathcal{F}^{i \rightarrow *j})$.

We will also consider the *normalized matrices of in-forests*:

$$J_k = \sigma_k^{-1} Q_k, \quad k = 0, \dots, n - d, \tag{9}$$

$$J = \sigma^{-1} Q, \tag{10}$$

and the parametric matrices

$$Q(\tau) = \sum_{k=0}^{n-d} Q_k \tau^k, \tag{11}$$

$$J(\tau) = \sigma^{-1}(\tau) Q(\tau), \quad \tau \geq 0, \tag{12}$$

where σ_k, σ , and $\sigma(\tau)$ are defined by (2)–(4).

The *normalized matrix of maximum in-forests* J_{n-d} will be also denoted by \tilde{J} :

$$\tilde{J} = J_{n-d}.$$

In the case of undirected graphs, the entries of \tilde{J} are the same within every connected component. In the directed case, this matrix possesses nontrivial properties determined by the properties of maximum in-forests, cf. [1].

Proposition 1. *The matrices $J_k, k = 0, \dots, n - d, J$, and $J(\tau)$ are row stochastic.*

Proof. Every row sum of $Q_k, k = 0, \dots, n - d$, is σ_k . Indeed, for every $i = 1, \dots, n$, we have

$$\sum_{j=1}^n q_{ij}^k = \sum_{j=1}^n w(\mathcal{F}_k^{i \rightarrow *j}) \stackrel{(*)}{=} w\left(\bigcup_{j=1}^n \mathcal{F}_k^{i \rightarrow *j}\right) = w(\mathcal{F}_k^{\rightarrow *}) = \sigma_k.$$

In the (*) passage, we used the fact that $\mathcal{F}_k^{i \rightarrow *j_1} \cap \mathcal{F}_k^{i \rightarrow *j_2} = \emptyset$ whenever $j_1 \neq j_2$. Thus, the nonnegative matrices $J_k = \sigma_k^{-1} Q_k$ are row stochastic. Now the stochasticity of J and $J(\tau)$ follows from their definitions. \square

The aim of this paper is to interpret, in terms of the forest matrices, a number of expressions that involve the Laplacian matrix as well as to provide polynomial expressions for the forest matrices themselves.

3. Preliminaries

This section briefly surveys some known results on the minors of a digraph's Laplacian matrix.

The oldest result of this kind is the *matrix-tree theorem* by Tutte [64,65], although some authors (e.g., [10]; cf. [52]) trace it back to Sylvester [63] and its proof to Borchardt [9].

Theorem 1. For every $i, j \in V(\Gamma)$, $\ell^{ij} = w(\mathcal{T}^{\rightarrow *i})$ holds, where ℓ^{ij} is the cofactor of the (i, j) entry of L and $\mathcal{T}^{\rightarrow *i}$ is the set of all spanning trees converging to i in Γ .

As stated in [34], “This small formula opens a world of opportunities”.

Tutte [65] formulated this theorem for the diagonal cofactors of the Kirchhoff matrix. A version that involves all cofactors of the Laplacian and the column Laplacian matrices can be found in [30]. We do not describe multiple analogues of the matrix-tree theorem here.

By definition, L has the form $L = D - W$, where W is the nonnegative matrix of arc weights and D is the diagonal matrix ensuring the zero row sums of L . Therefore, by Geršgorin's theorem, the real part of each nonzero eigenvalue of L is positive. Thus, L is a singular M-matrix (see, e.g., [7, Theorem 4.6 in Chapter 6]). One of the consequences is that all the principal minors of L are nonnegative. Fiedler and Sedláček [25] obtained an interpretation of all principal minors of the Laplacian matrix in terms of spanning forests.

Theorem 2. For any $\mathcal{J} \subseteq \{1, \dots, n\}$, $\det L(\overline{\mathcal{J}} | \overline{\mathcal{J}}) = w(\mathcal{F}^{\rightarrow * \mathcal{J}})$ holds, where $\mathcal{F}^{\rightarrow * \mathcal{J}}$ is the set of in-forests for which \mathcal{J} is the set of roots.

Later this theorem was formulated and proved in [11]. Its special case with undirected graphs and $|\mathcal{J}| = 2$ was discovered and employed earlier in the theory of electrical networks (see, e.g., [55]). Fiedler and Sedláček stated their result for the column Laplacian matrix and out-forests. Generally, to get interpretations for the minors of the column Laplacian matrix $L'(I)$, it suffices to observe that for the digraph obtained from Γ by the reversal of all arcs, the Laplacian matrix coincides with $K(\Gamma) = L^T(\Gamma)$ and the in-forests are in a weight preserving correspondence with the out-forests of Γ .

Let

$$\varphi(\lambda) = \det(\lambda I + L) = \sum_{k=0}^n c_{n-k} \lambda^k \tag{13}$$

be the characteristic polynomial of $-L$ and let σ_k be as defined in (2).

Proposition 2. In (13), $c_k = \sigma_k$, $k = 0, \dots, n$.

In view of Theorem 2, this proposition follows from the fact that c_k is equal to the sum of the $k \times k$ principal minors of L . In the case of undirected unweighted multigraphs, Proposition 2 is due to Kelmans [36,37], who was probably the first [35] to study the Laplacian characteristic polynomial (see also discussion in [52, p. 42] and [21, Sections 1.2, 1.5], and [8, Theorem 7.5]); some extensions are given in [4, the last statement on p. 236] and [20, Theorem 2]. An alternative representation for the coefficients of the Laplacian characteristic polynomial can be found in [26].

Since $\sigma_k = 0$ if and only if $k > n - d$ ($k = 0, 1, \dots$), Proposition 2 implies

Corollary 1. The multiplicity of 0 as the eigenvalue of L is d .

Another immediate consequence of Proposition 2 is

Corollary 2

$$\sigma_k = \sum_{\mathcal{J}: |\mathcal{J}|=k} \prod_{j \in \mathcal{J}} \lambda_j, \quad k = 0, \dots, n,$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of L and \mathcal{J} are the subsets of $\{1, \dots, n\}$.

Chen [17, p. 313, Problems 4.14 and 4.16] proposed an extension of the matrix-tree theorem to additional minors of the Laplacian matrix and Chaiken [10] gave a similar graph interpretation to all minors of L' . Moon [53] obtained a more general expansion which applies to all minors of arbitrary matrices; Chaiken's theorem and a number of W.K. Chen's expansions follow from his result as special cases. Minoux [51] generalized Chaiken's theorem to semirings and Bapat et al. [5] to mixed graphs (where each arc is either directed or undirected). Other useful graph interpretations of minors and determinants are given in [45].

We do not quote these results here, but we employ Chaiken's formulation [10] of the *all minors matrix tree theorem* in the proof of the *matrix-forest theorem* in the following section.

4. The matrix-forest theorem

The following theorem [13,15] provides expressions for the forest matrices Q and J (see (8) and (10)) in terms of the cofactors and the determinant of $I + L$, where I is the identity matrix.

Theorem 3 (the matrix-forest theorem). $Q = \text{adj}(I + L)$ and $\sigma = \det(I + L)$. Thus, $J = (I + L)^{-1}$.

For the properties of $(I + L)^{-1}$, see [15,16,46,47].

It was mentioned in [15] that a quick way to prove the matrix-forest theorem is to employ the all minors matrix tree theorem, more specifically, to apply the first formula (without number) on page 328 in [10]. Below we give a complete inference of Theorem 3 from the all minors matrix tree theorem. Note that a self-contained proof of the matrix-forest theorem for unweighted multigraphs can be found in [62]. Another inference based on some results of [25,36,37,45] was given in [13] for the case of weighted multidigraphs and multigraphs. Undirected and unweighted analogies of Theorem 3 have been presented in [46,47] (with the proof based on Chaiken’s theorem) and [14].

In the following proof of Theorem 3, we employ a standard trick which enables one to reduce many novel statements about forests to known statements about trees or forests. Versions of this trick have been used in many papers, e.g., [4,10,12,17,28,33,34,38,46,47,56].¹ We formalize it by

Definition 3. Let Γ be a weighted digraph. The digraph $\widehat{\Gamma}$ with vertex set $V(\widehat{\Gamma}) = V(\Gamma) \cup \{0\}$, arc set $E(\widehat{\Gamma}) = E(\Gamma) \cup \{(j, 0) : j \in V(\Gamma)\}$, the weights of arcs in $E(\widehat{\Gamma}) \cap E(\Gamma)$ the same as for Γ , and $w((j, 0)) = 1, j \in V(\Gamma)$, will be called the *ground extension* of Γ .²

Observation 1. Let $\widehat{\Gamma}$ be the ground extension of Γ . Let $U = I + L(\Gamma), \widehat{L} = L(\widehat{\Gamma})$. Then for any $\mathcal{I}, \mathcal{J} \subseteq V(\Gamma), U(\mathcal{I} | \mathcal{J}) = \widehat{L}(\mathcal{I} \cup \{0\} | \mathcal{J} \cup \{0\})$ holds.

By virtue of Observation 1, if one has expressions for all minors of the Laplacian matrices L (say, those provided by the all minors matrix tree theorem), then expressions for all minors of matrices $I + L$ are got gratis. The following lemma establishes a correspondence between the forests in Γ and some forests in $\widehat{\Gamma}$. The lemma is formulated here in a form useful for expressing all minors of $I + L$.

Lemma 1. Consider $\mathcal{I} = \{i_1, \dots, i_k\} \subseteq V(\Gamma), \mathcal{J} = \{j_1, \dots, j_k\} \subseteq V(\Gamma), 0 \leq k \leq n$, and the set of in-forests $\mathcal{F} \rightarrow^* \cap (\bigcap_{u=1}^k \mathcal{F}^{i_u \rightarrow^* j_u})$ in Γ . Then there exists a weight preserving one-to-one correspondence between this set and the set $\widehat{\mathcal{F}} \rightarrow^*_{\mathcal{J}}^*$ of in-forests $F \in \widehat{\mathcal{F}}^{0 \rightarrow^* 0} \cap (\bigcap_{u=1}^k \widehat{\mathcal{F}}^{i_u \rightarrow^* j_u})$ in $\widehat{\Gamma}$ such that the F ’s consist of exactly $k + 1$ trees.

¹ Note that one more expedient is to identify the roots of all trees in a forest, which converts the forest into a tree [13,21,25,36,37,54].

² In [34] $\widehat{\Gamma}$ is called the *cone* of Γ .

Proof. Let $F \in \mathcal{F}^{\rightarrow*} \cap (\bigcap_{u=1}^k \mathcal{F}^{i_u \rightarrow *j_u})$. To define the corresponding forest in $\widehat{\mathcal{F}}_{\mathcal{J}}^{\rightarrow*}$, consider the replica F' of F in $\widehat{\Gamma}$ and attach the arcs $(r, 0)$ to it, where the r 's are the roots of F' that are not in \mathcal{J} . The resulting in-forest consists of exactly $k + 1$ trees and belongs to $\widehat{\mathcal{F}}_{\mathcal{J}}^{\rightarrow*}$. Conversely, for any $\widehat{F} \in \widehat{\mathcal{F}}_{\mathcal{J}}^{\rightarrow*}$, consider its restriction to $V(\Gamma)$ as the corresponding forest of Γ . Obviously, this correspondence is one-to-one and the corresponding forests share the weight. \square

Proof of Theorem 3. Consider the ground extension $\widehat{\Gamma}$ of Γ . By Observation 1, if $U = I + L(\Gamma)$, U^{ij} is the (i, j) entry of $\text{adj } U$, and $\widehat{L} = L(\widehat{\Gamma})$, then

$$U^{ij} = (-1)^{i+j} \det U(\overline{\{j\}} \mid \overline{\{i\}}) = (-1)^{i+j} \det \widehat{L}(\overline{\{0, j\}} \mid \overline{\{0, i\}}). \tag{14}$$

Let $\widehat{\mathcal{F}}_{(2)}^{0 \rightarrow *0, i \rightarrow *j}$ be the set of in-forests $F \in \widehat{\mathcal{F}}^{0 \rightarrow *0} \cap \widehat{\mathcal{F}}^{i \rightarrow *j}$ that consist of two trees. Denoting by $\text{inv}\{0 \rightarrow 0, i \rightarrow j\}$ the number of violations of monotonicity in the two-element correspondence $\{0 \rightarrow 0, i \rightarrow j\}$, which is obviously zero, and using the all minors matrix tree theorem [10,53], we get

$$\begin{aligned} \det \widehat{L}(\overline{\{0, j\}} \mid \overline{\{0, i\}}) &= (-1)^{|\{k \in V(\Gamma) : k < j\}| + |\{k \in V(\Gamma) : k < i\}|} \\ &\times \sum_{F \in \widehat{\mathcal{F}}_{(2)}^{0 \rightarrow *0, i \rightarrow *j}} (-1)^{\text{inv}\{0 \rightarrow 0, i \rightarrow j\}} w(F) \\ &= (-1)^{j+i-2} w(\widehat{\mathcal{F}}_{(2)}^{0 \rightarrow *0, i \rightarrow *j}). \end{aligned} \tag{15}$$

In the first passage, we used the fact that $\widehat{\mathcal{F}}^{i \rightarrow *0} = \emptyset$.

Lemma 1 implies $w(\widehat{\mathcal{F}}_{(2)}^{0 \rightarrow *0, i \rightarrow *j}) = w(\mathcal{F}^{i \rightarrow *j})$, so, from (14) and (15), we get

$$U^{ij} = (-1)^{2i+2j-2} w(\mathcal{F}^{i \rightarrow *j}) = w(\mathcal{F}^{i \rightarrow *j}) = q_{ij}.$$

By Observation 1, Theorem 1, and Lemma 1, $\det U = \det \widehat{L}(\overline{\{0\}} \mid \overline{\{0\}}) = w(\widehat{\mathcal{F}}_{(1)}^{0 \rightarrow *0}) = w(\mathcal{F}^{\rightarrow*}) = \sigma$ (cf. [33, Eq. (37)] and [34, 7.2 and 7.3]). This completes the proof. \square

Remark 1. Obviously, the positivity of arc weights is needed for the last statement of Theorem 3 only; the first two statements are preserved for digraphs with arbitrary arc weights.

Remark 2. Note that the cofactors and the determinant of $I + L$, in the case of an unweighted undirected graph G , have been expressed in [28] in terms of spanning trees and 2-forests in the ground extension of G (for the case of weighted graphs, cf. [39, Theorem 2.3]). Ref. [28] also discusses the idea of using graph invariants related

to $(I + L)^{-1}$ in the study of the graph isomorphism problem. We surmise that the forest matrices Q_k also have some potential in this respect.

It is easily seen that $I + \tau L$ with $\tau \geq 0$ are nonsingular M-matrices, so their inverses are nonnegative. In the next section, the following parametric matrix-forest theorem [1,16] will be helpful:

Theorem 3'. For any $\tau \in \mathbb{R}$, $Q(\tau) = \text{adj}(I + \tau L)$ and $\sigma(\tau) = \det(I + \tau L)$. Thus, for any $\tau \geq 0$, $J(\tau) = (I + \tau L)^{-1}$.

To prove this theorem, it suffices to apply Theorem 3 to the weighted digraph $\Gamma'(\tau)$ that differs from Γ in the weights of arcs only: for all $i, j = 1, \dots, n$, $w'_{ij}(\tau) = \tau w_{ij}$. By Remark 1, the nonnegativity of τ is needed for the last statement of Theorem 3' only.

5. A method for calculating Q_1, \dots, Q_{n-d}

We first show that Q_1, \dots, Q_{n-d} are the matrix coefficients in the polynomial expansion of $\text{adj}(\lambda I + L)$.

Proposition 3

$$\text{adj}(\lambda I + L) = \sum_{k=0}^{n-d} Q_k \lambda^{n-k-1}.$$

Proof. If $\lambda = 0$, then the right-hand side is zero whenever $d > 1$ and it reduces to Q_{n-1} when $d = 1$ (we put $\lambda^0 \equiv 1$). In any case, this is equal to $\text{adj}(\lambda I + L)$ by Theorem 1. For any $\lambda \neq 0$, let $\tau = \lambda^{-1}$. Using Theorem 3' we get

$$\begin{aligned} \text{adj}(\lambda I + L) &= \text{adj} \lambda(I + \tau L) = \lambda^{n-1} Q(\tau) \\ &= \lambda^{n-1} \sum_{k=0}^{n-d} Q_k \tau^k = \sum_{k=0}^{n-d} Q_k \lambda^{n-k-1}. \quad \square \end{aligned} \tag{16}$$

Proposition 3 underlies an easy algorithm for calculating Q_1, \dots, Q_{n-d} and $\sigma_1, \dots, \sigma_{n-d}$.

Proposition 4. For any $k = 0, 1, \dots,$

$$Q_{k+1} = (-L)Q_k + \sigma_{k+1}I, \tag{17}$$

$$\sigma_{k+1} = \frac{\text{tr}(LQ_k)}{k+1}. \tag{18}$$

Proof. Since, by Proposition 3, Q_0, \dots, Q_n are the matrix coefficients in the polynomial form of $\text{adj}(\lambda I + L)$, where $\lambda I + L$ is the characteristic matrix of $-L$ and, by Proposition 2, $\sigma_0, \dots, \sigma_n$ are the coefficients of the characteristic polynomial of $-L$, the equations [27, Section 3 of Chapter 4] $Q_{k+1} = \sigma_{k+1}I - LQ_k$, $k = 0, 1, \dots$, take place.

To prove (18), it suffices to take the traces on the left and on the right of (17) and use the fact that

$$\text{tr } Q_k = (n - k)\sigma_k, \quad k = 0, 1, \dots,$$

which holds since every in-forest with k arcs has $n - k$ roots. \square

Note that, by virtue of Propositions 2 and 3, the recurrent application of (18) and (17) starting with $Q_0 = I$ coincides with the Leverrier-Faddeev algorithm [23,27] applied to calculate the characteristic polynomial of $-L$.

Consider now a few corollaries to Proposition 4. First, in what follows we will need a recurrence formula for the row stochastic matrices J_k . It is:

$$J_{k+1} = \frac{\sigma_k}{\sigma_{k+1}}(-L)J_k + I, \quad k = 0, \dots, n - d - 1. \tag{19}$$

Second, the matrices LQ_k prevailing in Proposition 4 have a noteworthy graph interpretation. Let Γ_k be the *digraph of in-forests with k arcs* of Γ , i.e., the digraph on vertex set $V(\Gamma_k) = V(\Gamma)$ whose matrix of arc weights results from Q_k by putting zeros on the main diagonal. In other words,

$$(i, j) \in E(\Gamma_k) \quad \text{whenever } j \neq i \text{ and } q_{ij}^k > 0;$$

q_{ij}^k is the weight of such arc. Evidently, $\Gamma_1 = \Gamma$.

Proposition 5. LQ_k is the Laplacian matrix of Γ_{k+1} , $k = 0, 1, \dots$.

Proof. By Proposition 4, $LQ_k = \sigma_{k+1}I - Q_{k+1}$, so the off-diagonal entries of LQ_k coincide with those of $L(\Gamma_{k+1})$. To complete the proof, note that every row sum of LQ_k is zero, since every row sum of both $\sigma_{k+1}I$ and Q_{k+1} is σ_{k+1} . \square

Finally, Proposition 4 provides a recurrence formula for the Laplacian matrices $L_k := L(\Gamma_k)$:

$$L_{k+1} = L \left(-L_k + \frac{\text{tr } L_k}{k} I \right), \quad k = 1, 2, \dots$$

We are going to discuss the application of digraphs Γ_k to the analysis of Γ elsewhere.

6. Forest matrices as polynomials in the Laplacian matrix

It follows from Proposition 4 that the forest matrices $Q_k, Q,$ and $Q(\tau)$ are polynomials in L . As a corollary, the powers of L are linear combinations of Q_0, \dots, Q_{n-d} . First, it is straightforward to prove:

Proposition 6

$$Q_k = \sum_{i=0}^k \sigma_{k-i}(-L)^i, \quad k = 0, 1, \dots$$

Remark 3. These expressions are closely related to the characteristic polynomial of $-L$ (13) which, by Proposition 2, can be represented as $\varphi(\lambda) = (\dots((\sigma_0\lambda + \sigma_1)\lambda + \sigma_2)\lambda + \dots + \sigma_{n-1})\lambda + \sigma_n$. To find $\varphi(\lambda)$, one can successively calculate $\varphi_0(\lambda) = \sigma_0, \varphi_1(\lambda) = \sigma_0\lambda + \sigma_1, \varphi_2(\lambda) = (\sigma_0\lambda + \sigma_1)\lambda + \sigma_2, \dots, \varphi_n(\lambda) = \varphi(\lambda)$. It is easily seen now that $Q_k = \varphi_k(-L), k = 0, \dots, n$.

Corollary 3. *The matrices $Q_k, k = 0, 1, \dots,$ commute with all matrices with which L commutes, in particular, with $L, Q(\tau),$ and each other.*

By Theorems 3 and 3', $Q = \text{adj}(I + L)$ and $Q(\tau) = \text{adj}(I + \tau L)$. Proposition 6, (8), and (11) provide a polynomial form of Q and $Q(\tau)$.

Proposition 7

$$Q = \sum_{k=0}^{n-d} s_{n-d-k}(-L)^k = \text{adj}(I + L),$$

$$Q(\tau) = \sum_{k=0}^{n-d} s_{n-d-k}(\tau) (-\tau L)^k = \text{adj}(I + \tau L), \tag{20}$$

where s_i and $s_i(\tau)$ are defined in (5) and (6).

By (16), $\text{adj}(\lambda I + L) = \lambda^{n-1} Q(\tau)$, where $\lambda \neq 0$ and $\tau = 1/\lambda$. Combining this with (20) and (6), we obtain

Corollary 4

$$\text{adj}(\lambda I + L) = \sum_{k=0}^{n-d} s'_{n-d-k}(\lambda) (-L/\lambda)^k,$$

where $s'_i(\lambda) = \sum_{j=0}^i \sigma_j \lambda^{n-j-1}, i = 0, \dots, n - d,$ and $\lambda \neq 0$.

Corollary 4 and Proposition 3 can be considered as dual representations of $\text{adj}(\lambda I + L)$.

It follows from Proposition 6 that the powers of L are linear combinations of Q_0, \dots, Q_{n-d} , but the coefficients are more complicated than before.

Proposition 8. For $m = 0, 1, \dots$,

$$(-L)^m = \sum_{k=0}^m \alpha_k Q_{m-k}$$

holds, where $\alpha_0 = 1$,

$$\alpha_k = \sum_{(p_1, \dots, p_k): \sum i p_i = k} (-1)^{\sum p_i} \frac{(\sum p_i)!}{\prod (p_i!)} \prod \sigma_i^{p_i}, \quad k = 1, \dots, m, \quad (21)$$

p_i are nonnegative integers, and all sums and products in (21), except for the first sum, range from $i = 1$ to k .

A nice property of these linear combinations is that the coefficients α_k do not depend on m (similarly to Proposition 6). For instance,

$$\begin{aligned} L &= -(Q_1 - \sigma_1 I), \\ L^2 &= Q_2 - \sigma_1 Q_1 - (\sigma_2 - \sigma_1^2) I, \\ L^3 &= -(Q_3 - \sigma_1 Q_2 - (\sigma_2 - \sigma_1^2) Q_1 - (\sigma_3 - 2\sigma_2 \sigma_1 + \sigma_1^3) I), \\ L^4 &= Q_4 - \sigma_1 Q_3 - (\sigma_2 - \sigma_1^2) Q_2 - (\sigma_3 - 2\sigma_2 \sigma_1 + \sigma_1^3) Q_1 \\ &\quad - (\sigma_4 - 2\sigma_3 \sigma_1 - \sigma_2^2 + 3\sigma_2 \sigma_1^2 - \sigma_1^4) I. \end{aligned}$$

Proof. We first prove, by induction on m , the identity

$$(-L)^m = \sum_{k=0}^m \alpha'_k Q_{m-k} \quad (22)$$

with $\alpha'_0 = 1$ and

$$\alpha'_k = \sum_{(\beta(1), \dots, \beta(n_\beta)): \sum \beta(i) = k} \prod (-\sigma_{\beta(i)}), \quad k = 1, \dots, m, \quad (23)$$

where $\beta(i)$ are positive integers, n_β is the variable number of entries in $(\beta(1), \dots, \beta(n_\beta))$, and the unmarked sum and product range from $i = 1$ to n_β .

For the basis of induction, observe that $(-L)^0 = I = \alpha'_0 Q_0$. Let (22) and (23) be valid for $(-L)^0, \dots, (-L)^{m-1}$. By Proposition 6,

$$(-L)^m = \alpha'_0 Q_m - \sum_{i=0}^{m-1} \sigma_{m-i} (-L)^i. \tag{24}$$

Substituting (22) in the right-hand side of (24) and interchanging the two sums we obtain:

$$(-L)^m = \alpha'_0 Q_m + \sum_{k=1}^m \alpha_k^{(m)} Q_{m-k},$$

where

$$\alpha_k^{(m)} = \sum_{i=1}^k (-\sigma_i) \alpha'_{k-i}, \quad k = 1, \dots, m.$$

It is easily seen that $\alpha_k^{(m)} = \alpha'_k$, $k = 1, \dots, m$, thereby the induction step has succeeded.

Next, for an arbitrary positive integer k , consider any vector $(\beta(1), \dots, \beta(n_\beta))$ with positive integer entries such that $\sum_{i=1}^{n_\beta} \beta(i) = k$ (see (23)). Let $p_j = |\{i : \beta(i) = j\}|$, $j = 1, \dots, k$. Classifying the set of vectors $(\beta(1), \dots, \beta(n_\beta))$ such that $\sum \beta(i) = k$ by the equality of the corresponding vectors (p_1, \dots, p_k) , we see that every such a class contains $(\sum p_i)! / \prod (p_i!)$ members. This implies that $\alpha_k = \alpha'_k$, $k = 0, 1, \dots$ (cf. (21) and (23)) and thus, completes the proof. \square

7. The matrix of maximum in-forests

In this section, we study some properties of the normalized matrix $\tilde{J} = J_{n-d}$ of maximum in-forests. Let $\mu_\lambda(A)$ stand for the multiplicity of λ as the eigenvalue of a square matrix A .

Proposition 9

- (i) $L\tilde{J} = \tilde{J}L = LQ_{n-d} = Q_{n-d}L = 0$;
- (ii) $\tilde{J}J_k = J_k\tilde{J} = \tilde{J}$, $k = 0, \dots, n-d$;
- (iii) \tilde{J} is a projection: $\tilde{J}^2 = \tilde{J}$;
- (iv) $\text{rank } \tilde{J} = \mu_1(\tilde{J}) = \text{tr } \tilde{J} = d$; $\mu_0(\tilde{J}) = n-d$.

Proof. (i) Putting $k = n-d$ in (17) and using the facts that $Q_{n-d+1} = 0$ and $\sigma_{n-d+1} = 0$, we get $LQ_{n-d} = 0$. The other identities follow from Corollary 3 and (9).

(ii) Multiplying (19) by \tilde{J} and using item (i) and Corollary 3, we get the required statement, whose special case is (iii).

(iv) Each maximum in-forest of Γ has d roots, hence $\text{tr } Q_{n-d} = d \sigma_{n-d}$ and $\text{tr } \tilde{J} = \text{tr}(\sigma_{n-d}^{-1} Q_{n-d}) = d$. Since \tilde{J} is idempotent, $\text{rank } \tilde{J} = \mu_1(\tilde{J}) = \text{tr } \tilde{J}$, so $\mu_0(\tilde{J}) = n - d$. \square

The following connection between the spectra of L and $L + \alpha \tilde{J}$, $\alpha \in \mathbb{C}$, will be used in the sequel.

Proposition 10

- (i) The spectrum of $L + \alpha \tilde{J}$ consists of all nonzero eigenvalues of L with their multiplicities and α with $\mu_\alpha(L + \alpha \tilde{J}) = d$.
- (ii) $L + \alpha \tilde{J}$ is nonsingular whenever $\alpha \neq 0$.

Proof. (i) Let $p(\lambda) = \sigma_{n-d}^{-1} \sum_{i=0}^{n-d} \sigma_{n-d-i} (-\lambda)^i$. By Proposition 6, $\tilde{J} = \sigma_{n-d}^{-1} \times Q_{n-d} = p(L)$, so $L + \alpha \tilde{J} = L + \alpha p(L)$. Therefore, by [27, Theorem 3 in Chapt. 4], all eigenvalues of $L + \alpha \tilde{J}$ are $\lambda'_i = \lambda_i + \alpha p(\lambda_i)$, where $\lambda_i, i = 1, \dots, n$, are all eigenvalues of L with their multiplicities. By (i) of Proposition 9, $L \tilde{J} = 0 = L p(L)$, whence $\lambda p(\lambda)$ is an annihilating polynomial for L . Therefore, for each λ_i , a nonzero eigenvalue of L , we have $p(\lambda_i) = 0$, hence $\lambda'_i = \lambda_i$. Otherwise, if $\lambda_i = 0$, then $\lambda'_i = \alpha$, since $p(0) = 1$ by definition of $p(\lambda)$. Finally, by Corollary 1, $\mu_0(L) = d$, thus $\mu_\alpha(L + \alpha \tilde{J}) = d$. This implies (ii).

Proposition 11. $\tilde{J} = \lim_{\tau \rightarrow \infty} J(\tau) = \lim_{\tau \rightarrow \infty} (I + \tau L)^{-1}$.

Proof. Using Theorem 3' and the definition (12) of $J(\tau)$, we have

$$\begin{aligned} \lim_{\tau \rightarrow \infty} (I + \tau L)^{-1} &= \lim_{\tau \rightarrow \infty} J(\tau) = \lim_{\tau \rightarrow \infty} \left(\sum_{k=1}^{n-d} \sigma_k \tau^k \right)^{-1} \sum_{k=1}^{n-d} Q_k \tau^k \\ &= \lim_{\tau \rightarrow \infty} \left(\sum_{k=1}^{n-d} \sigma_k \tau^{k-n-d} \right)^{-1} \sum_{k=1}^{n-d} Q_k \tau^{k-n-d} \\ &= \sigma_{n-d}^{-1} Q_{n-d} = \tilde{J}. \quad \square \end{aligned}$$

8. L and \tilde{J} as “complementary” linear transformations

For a complex matrix A , let $\mathcal{R}(A)$ and $\mathcal{N}(A)$ denote its range and null space, respectively. Recall that the *index* of a square matrix A , $\text{ind } A$, is the smallest non-negative integer k for which $\text{rank}(A^{k+1}) = \text{rank}(A^k)$. The *eigenprojection*³ at 0 of

³ The eigenprojections are also called *principal idempotents* [32,69].

A [57] or, for short, the *eigenprojection of A* [58] is the idempotent matrix B such that $\mathcal{R}(B) = \mathcal{N}(A^\nu)$ and $\mathcal{N}(B) = \mathcal{R}(A^\nu)$, where $\nu = \text{ind } A$. In other words, B is the projection on $\mathcal{N}(A^\nu)$ along $\mathcal{R}(A^\nu)$. The eigenprojection is unique, because an idempotent matrix is uniquely determined by its range and null space (see, e.g., [6, p. 50]).⁴

Since $L\tilde{J} = 0$ (Proposition 9), we have $\mathcal{R}(L^*) \cap \mathcal{R}(\tilde{J}) = \{\mathbf{0}\}$, where $L^* = L^T$. Similarly, $\tilde{J}L = 0$ implies $\mathcal{R}(\tilde{J}^*) \cap \mathcal{R}(L) = \{\mathbf{0}\}$. Consequently, by [44, Theorem 11], L and \tilde{J}^* are rank additive, i.e., $\text{rank}(L + \tilde{J}^*) = \text{rank } L + \text{rank } \tilde{J}^*$. Corollary 1 implies that $\text{rank } L \geq n - d$, whereas, by Proposition 9, $\text{rank } \tilde{J}^* = d$. Since $\text{rank}(L + \tilde{J}^*) \leq n$, we have $\text{rank } L = n - d$ and $\text{rank}(L + \tilde{J}^*) = n$. Now $L\tilde{J} = \tilde{J}L = 0$ implies $\mathcal{N}(L) = \mathcal{R}(\tilde{J})$ and $\mathcal{N}(\tilde{J}) = \mathcal{R}(L)$. Furthermore, by Proposition 10, $\text{rank}(L + \tilde{J}) = n$, hence L and \tilde{J} are rank additive. It follows now from [44, Theorem 11] that $\mathcal{R}(L) \cap \mathcal{R}(\tilde{J}) = \{\mathbf{0}\}$. Since $\mathcal{R}(\tilde{J}) = \mathcal{N}(L)$, we get $\mathcal{R}(L) \cap \mathcal{N}(L) = \{\mathbf{0}\}$, which, by [6, p. 165], implies $\text{ind } L = 1$. The latter fact together with $\mathcal{R}(\tilde{J}) = \mathcal{N}(L)$, $\mathcal{N}(\tilde{J}) = \mathcal{R}(L)$, and $\tilde{J}^2 = \tilde{J}$ imply that \tilde{J} is the eigenprojection of L (alternatively, this follows from Proposition 11 and [48, Theorem 3.1]). We proved

Proposition 12

- (i) $L + \tilde{J}^*$ is nonsingular.
- (ii) $\text{rank } L = n - \text{rank } \tilde{J} = n - d$.
- (iii) $\mathcal{N}(L) = \mathcal{R}(\tilde{J})$ and $\mathcal{R}(L) = \mathcal{N}(\tilde{J})$.
- (iv) $\mathcal{R}(L) \cap \mathcal{R}(\tilde{J}) = \{\mathbf{0}\}$.
- (v) $\text{ind } L = 1$.
- (vi) \tilde{J} is the eigenprojection of L .

It is known [57, p. 194], [59, Theorem 7.a.3] that for every finite homogeneous Markov chain with a transition matrix P , the long run transition matrix $P^\infty = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{t=0}^{k-1} P^t$ is the eigenprojection of P at 1, which is the eigenprojection of $I - P$.⁵ On the other hand, $I - P$ is exactly the Laplacian matrix L of the weighted digraph without loops whose arc weights are equal to the corresponding transition probabilities. Therefore \tilde{J} , the eigenprojection of L , coincides with P^∞ . The fact

⁴ Note that for every $A \in \mathbb{C}^{n \times n}$ s.t. $\text{ind } A = \nu$ and every idempotent matrix B , each of the following conditions is equivalent to B being the eigenprojection of A :

- (i) $\mathcal{R}(B) = \mathcal{N}(A^\nu)$ and $\mathcal{R}(B^*) = \mathcal{N}((A^*)^\nu)$ [57];
 - (ii) $A^\nu B = BA^\nu = 0$ and $\text{rank } A^\nu + \text{rank } B = n$ [66,68];
 - (iii) $AB = BA$ and $A + \alpha B$ is nonsingular for all $\alpha \neq 0$ [41] (cf. (ii) of Proposition 10);
 - (iv) $AB = BA$, $A + \alpha B$ is nonsingular for some $\alpha \neq 0$, and AB is nilpotent [41];
 - (v) $AB = BA$, AB is nilpotent, and $AU = I - B = VA$ for some $U, V \in \mathbb{C}^{n \times n}$ [31];
 - (vi) B commutes with all matrices commuting with A , AB is nilpotent, and $B \neq 0$ if A is singular [40];
- Moreover, the eigenprojection of A is $I - AA^D$, where A^D is the Drazin inverse of A (see Section 9).

⁵ This also follows from Meyer’s Theorem 2.2 in [49]. Indeed, by this theorem, $P^\infty = I - (I - P)(I - P)^\#$, where $(I - P)^\#$ is the group inverse of $I - P$, and the right-hand side is the eigenprojection of $I - P$, as mentioned in the next section.

that P^∞ coincides with the normalized matrix of maximum in-forests of the digraph corresponding to a Markov chain is the so called *Markov chain tree theorem* [42,43]. Thus, item (vi) of Proposition 12 provides an immediate proof of this theorem.

By virtue of Proposition 9, every nonzero column of \tilde{J} (or Q_{n-d}) is an eigenvector of L that corresponds to the zero eigenvalue. Moreover, it follows from $\mathcal{N}(L) = \mathcal{R}(\tilde{J})$ (Proposition 12) that the nonzero columns of \tilde{J} span the null space of L . Since, by (16), $Q(\tau)$ is proportional to $\text{adj}(\lambda I - (-L))$ at $\lambda = \tau^{-1}$, $Q(\tau)$ can be used to generate some eigenvectors of L that correspond to its nonzero eigenvalues. For completeness, we give a proof of this fact.

Proposition 13. *Let $\lambda_i \neq 0$ be an eigenvalue of L . Then every nonzero column of $Q(-\lambda_i^{-1})$ is an eigenvector of L that corresponds to λ_i .*

Proof. Let $X = \lambda_i I - L$. Then $\det X = 0$. Using Theorem 3' and the fact that for every square matrix Y , $Y \text{adj} Y = (\det Y)I$ holds, we get

$$(\lambda_i I - L) Q(-\lambda_i^{-1}) = X \text{adj}(I - \lambda_i^{-1}L) = \lambda_i^{1-n} X \text{adj} X = \lambda_i^{1-n} (\det X)I = 0.$$

This implies the desired statement. \square

9. Forest matrices and generalized inverses of L

The *Moore-Penrose generalized inverse* A^+ of a rectangular complex matrix A is the unique matrix X such that

$$AXA = A, \quad XAX = X, \quad (AX)^* = AX, \quad (XA)^* = XA.$$

For an arbitrary square matrix A , its *Drazin inverse*, A^D , is the unique matrix X satisfying the equations

$$A^{\nu+1}X = A^\nu, \quad XAX = X, \quad AX = XA,$$

where $\nu = \text{ind } A$. If $\nu = 0$, then $A^D = A^{-1}$; if $\nu \leq 1$, then A^D is referred to as the *group inverse*, $A^\#$, i.e., the unique matrix X such that

$$AXA = A, \quad XAX = X, \quad AX = XA.$$

As applied to the Laplacian matrices of graphs, the generalized inverses were considered in connection with the analysis of electrical networks (providing “resistance distance”), Markov chains, and some preference aggregation problems (more specifically, estimation from paired comparisons), in constructing geometrical representations of graphs (with applications to chemistry, social networks, etc.), in control, cluster analysis, and parallel computing. There is a huge literature on generalized inverses within the last years. For multiple representations of the Drazin inverse, see, e.g., [18,19,67].

In this section, we present a few relations between the $L^\#$ and the forest matrices and one representation for L^+ . In the case of symmetric L , where $L^\# = L^+$, some of these expressions are given in [16].⁶

For an arbitrary square matrix A , AA^D is the unique projection on $\mathcal{R}(A^v)$ along $\mathcal{N}(A^v)$ [6, p. 173]. Then $I - AA^D$ is the projection on $\mathcal{N}(A^v)$ along $\mathcal{R}(A^v)$. Therefore, $I - AA^D$ is the eigenprojection of A [57,58]. Combining this with items (v) and (iv) of Proposition 12, we obtain

Proposition 14. $\tilde{J} = I - LL^\#$.

The fact that \tilde{J} is the eigenprojection of L helps interpret, in terms of in-forests, the expressions of generalized inverses of L that involve the eigenprojection of L .

Proposition 15. (i) For any $\alpha \neq 0$, $L^\# = (L + \alpha \tilde{J})^{-1} - \alpha^{-1} \tilde{J}$, whence $L^\# = \lim_{|\alpha| \rightarrow \infty} (L + \alpha \tilde{J})^{-1}$.

(ii) For any $\alpha \neq 0$, $L^\# = (L + \alpha \tilde{J})^{-1} (I - \tilde{J})$.

(iii) $L^\# = \frac{\sigma_{n-d-1}}{\sigma_{n-d}} (J_{n-d-1} - \tilde{J})$.

(iv) $L^\# = \lim_{\tau \rightarrow \infty} \tau (J(\tau) - \tilde{J})$.

Remarks on Proposition 15. (i), (iii), and (iv) were presented in [2]. (i) results by substituting \tilde{J} for the eigenprojection in the expression of group inverse employed in [50, p. 150] (for its proof see [60, Theorem 4.2]; related expressions appeared in [49, Theorem 5.5] and [58, last line on p. 646], where ‘+’ must be replaced by ‘-’). (ii) is obtained by the same substitution in the representation of Drazin inverse given in [40] (the case with $\alpha = 1$ appeared in [58]) or by multiplying (i) by $LL^\# = I - \tilde{J}$. In view of Propositions 2 and 3, (iii) follows from the expression of Drazin inverse discovered independently by Hartwig [32, Eq. (13)] and Gower [29, Theorem 1].

The matrices $L + \alpha \tilde{J}$ are the “complementary perturbations” [50] of L . Matrices of this kind are important for the analysis of M-matrices and singular systems of equations. In particular, a matrix A with eigenprojection B and nonpositive off-diagonal entries is an M-matrix if and only if for some $c > 0$, $(A + \alpha B)^{-1}$ is nonnegative when $\alpha \in (0, c)$ [50]. If A is an M-matrix, then $(A + \alpha B)^{-1}$, $\alpha \in (0, c)$, make up a class of *nonnegative nonsingular commuting weak inverses* for A [50]. $(L + \alpha \tilde{J})^{-1}$ can be represented as a linear combination of forest matrices using (i) and (iii) of Proposition 15:

$$(L + \alpha \tilde{J})^{-1} = \frac{\sigma_{n-d-1}}{\sigma_{n-d}} (J_{n-d-1} + \beta \tilde{J}),$$

⁶ For symmetric L , interesting representations for $L^\# = L^+$ were proposed in [24], [39, Theorem 2.2], and, in case of weighted trees, in [39] and [3, Theorem 3]. In [22, Theorem 3] a combinatorial interpretation of the Campbell-Youla inverse (the symmetric generalized inverse with the zero diagonal) of L is given.

where

$$\beta = \frac{\sigma_{n-d}}{\alpha\sigma_{n-d-1}} - 1.$$

This throws some light on the nonnegativity of $(L + \alpha\tilde{J})^{-1}$: if

$$\alpha \in \left(0, \frac{\sigma_{n-d}}{\sigma_{n-d-1}}\right),$$

then $(L + \alpha\tilde{J})^{-1}$ is a positive combination of J_{n-d-1} and \tilde{J} . Based on this, we termed $(L + \alpha\tilde{J})^{-1}$ the *matrices of dense in-forests* of Γ . These and the inverse “uniform diagonal perturbations” $(L + \alpha I)^{-1}$ can serve to measure proximity between digraph vertices [2]. Note in this connection that by [60, Corollary 4.4], $(L + \alpha\tilde{J})_{ij}^{-1} > 0$ for all $\alpha > 0$ sufficiently small if and only if vertex j is accessible from i in Γ , and the same is true for $(L + \alpha I)_{ij}^{-1}$. By Theorem 3', $(L + \alpha I)^{-1}$ is proportional to $J(\tau)$ with $\tau = 1/\alpha$.

We conclude with one expression for the Moore–Penrose inverse of L .

Consider the matrix $Z := L + \tilde{J}^*$ which is nonsingular by Proposition 12. Using the identity $L\tilde{J} = 0$ (Proposition 9), we get $(Z^*)^{-1}Z^{-1} = (ZZ^*)^{-1} = (\tilde{J}^*\tilde{J} + LL^*)^{-1}$.

Proposition 16 [2]. $L^+ = L^*(ZZ^*)^{-1} = L^*(\tilde{J}^*\tilde{J} + LL^*)^{-1}$.

One method to prove this is to check the conditions in the definition of Moore–Penrose inverse by direct computation using Proposition 9 and the facts that $(ZZ^*)^{-1}$ commutes with LL^* and $\tilde{J}^*\tilde{J}$ and that $LL^*(ZZ^*)^{-1}$ and $\tilde{J}^*\tilde{J}(ZZ^*)^{-1}$ are symmetric [2]. Alternatively, Proposition 16 can be proved by employing the Penrose formula $A^+ = A^*(AA^*)^+$, the fact that $(AA^*)^+ = (AA^*)^\#$ (since AA^* is Hermitian) and an expression of $(AA^*)^\#$ such as those given in (i) and (ii) of Proposition 15.

10. A concluding remark

It is instructive to compare the “Laplacian graph mathematics” we touched upon in this paper with the corresponding results on the adjacency characteristic matrix, see, e.g., [21, Sections 1.4, 1.9.1, 1.9.5 and others] and the articles by Kasteleyn and Ponstein cited therein, [61], and so on. This comparison suggests that the Laplacian mathematics is based on trees in the same sense as the “adjacency graph mathematics” is based on routes and circuits. We mean that a number of expressions related with the adjacency characteristic matrix can be interpreted in terms of routes and circuits, whereas the counterparts of these expressions related with the Laplacian characteristic matrix involve spanning forests for their interpretation.

Acknowledgements

This work was supported in part by the Russian Foundation for Basic Research under grants 01–01–10732-z and 02–01–00614-a. The authors thank a referee for a careful reading of the paper and many thoughtful suggestions.

References

- [1] R.P. Agaev, P.Y. Chebotarev, The matrix of maximum out forests and its applications, *Automation and Remote Control* 61 (2000) 1424–1450.
- [2] R.P. Agaev, P.Y. Chebotarev, Spanning forests of a digraph and their applications, *Automation and Remote Control* 62 (2001) 443–466.
- [3] R.B. Bapat, Moore-Penrose inverse of the incidence matrix of a tree, *Linear and Multilinear Algebra* 42 (1997) 159–167.
- [4] R.B. Bapat, G. Constantine, An enumerating function for spanning forests with color restrictions, *Linear Algebra Appl.* 173 (1992) 231–237.
- [5] R.B. Bapat, J.W. Grossman, D.M. Kulkarni, Generalized matrix tree theorem for mixed graphs, *Linear and Multilinear Algebra* 46 (1999) 299–312.
- [6] A. Ben-Israel, T.N.E. Greville, *Generalized Inverses: Theory and Applications*, Wiley, New York, 1974.
- [7] A. Berman, R. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, Academic Press, New York, 1979.
- [8] N. Biggs, *Algebraic Graph Theory*, Cambridge University Press, Cambridge, 1974.
- [9] C.W. Borchardt, über eine der Interpolation entsprechende Darstellung der Eliminations-Resultante, *Journal für die reine und angewandte Mathematik* 57 (1860) 111–121.
- [10] S. Chaiken, A combinatorial proof of the all minors matrix tree theorem, *SIAM Journal on Algebraic and Discrete Methods* 3 (1982) 319–329.
- [11] S. Chaiken, D.J. Kleitman, Matrix tree theorems, *J. Combin. Theory, Series A* 24 (1978) 377–381.
- [12] P. Chebotarev, E. Shamis, The forest metrics for graph vertices, *Discrete Mathematics*, 2001 (accepted).
- [13] P.Y. Chebotarev, E.V. Shamis, Matrix-forest theorems, 1995 (unpublished).
- [14] P.Y. Chebotarev, E.V. Shamis, On the proximity measure for graph vertices provided by the inverse Laplacian characteristic matrix, 5th Conference of the International Linear Algebra Society, Georgia State University, Atlanta, 1995, pp. 30–31.
- [15] P.Y. Chebotarev, E.V. Shamis, The matrix-forest theorem and measuring relations in small social groups, *Automation and Remote Control* 58 (1997) 1505–1514.
- [16] P.Y. Chebotarev, E.V. Shamis, On proximity measures for graph vertices, *Automation and Remote Control* 59 (1998) 1443–1459.
- [17] W.K. Chen, *Applied Graph Theory, Graphs and Electrical Networks*, second ed., North-Holland, Amsterdam, 1976.
- [18] Y.-L. Chen, Representation and approximation for the Drazin inverse $A^{(d)}$, *Appl. Math. Comput.* 119 (2001) 147–160.
- [19] Y.-L. Chen, X. Chen, Representation and approximation of the outer inverse $A_{T,S}^{(2)}$ of a matrix A , *Linear Algebra Appl.* 308 (2000) 85–107.
- [20] F.R.K. Chung, R.P. Langlands, A combinatorial Laplacian with vertex weights, *J. Combin. Theory, Series A* 75 (1996) 316–327.
- [21] D.M. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs*, Academic Press, New York, 1980.

- [22] S.W. Drury, G.P.H. Styan, G.E. Subak-Sharpe, On spanning trees and terminal weight numbers, Report from the Department of Mathematics and Statistics 94-4, McGill University, Montreal, Queb., Canada, 1994.
- [23] D.K. Faddeev, V.N. Faddeeva, Numerical Methods in Linear Algebra, Dover, New York, 1959.
- [24] M. Fiedler, Moore-Penrose involutions in the classes of Laplacians and simplices, *Linear and Multilinear Algebra* 39 (1995) 171–178.
- [25] M. Fiedler, J. Sedláček, O W -basích orientovaných grafů, *Časopis Pěst. Mat.* 83 (1958) 214–225.
- [26] R. Forman, Determinants of Laplacians on graphs, *Topology* 32 (1993) 35–46.
- [27] F.R. Gantmacher, *Teoriya Matrits*, second ed., Nauka, Moscow, 1966. (The first edition translated as: *The Theory of Matrices*, Vols. I and II, Chelsea, New York, 1959.)
- [28] V.E. Golender, V.V. Drboglav, A.B. Rosenblit, Graph potentials method and its application for chemical information processing, *J. Chem. Inform. Comput. Sci.* 21 (1981) 196–204.
- [29] J.C. Gower, A modified Leverrier-Faddeev algorithm for matrices with multiple eigenvalues, *Linear Algebra Appl.* 31 (1980) 61–70.
- [30] F. Harary, *Graph Theory*, Addison-Wesley, Reading, MA, 1969.
- [31] R.E. Harte, Spectral projections, *Irish Mathematical Society Newsletter* 11 (1984) 10–15.
- [32] R.E. Hartwig, More on the Souriau-Frame algorithm and the Drazin inverse, *SIAM Journal on Applied Mathematics* 31 (1976) 42–46.
- [33] B.D. Jones, B.G. Pittel, J.S. Verducci, Tree and forest weights and their application to nonuniform random graphs, *The Annals of Applied Probability* 9 (1999) 197–215.
- [34] A. Kelmans, I. Pak, A. Postnikov, Tree and forest volumes of graphs, *Rutcor Research Report 47-99*, Rutgers Center for Operations Research, Rutgers University, Piscataway NJ, 1999.
- [35] A.K. Kel'mans, The number of trees in a graph I, II, *Automation and Remote Control* 26 (1965) 2118–2129, 27 (1966) 233–241.
- [36] A.K. Kelmans, On the properties of the characteristic polynomial of a graph, in: A.I. Berg (Ed.), *Kibernetiku – na sluzhbu kommunizmu (Cybernetics Should Serve Communism)*, vol. 4, Energiya, Moscow–Leningrad, 1967, pp. 27–41 (in Russian).
- [37] A.K. Kelmans, V.M. Chelnokov, A certain polynomial of a graph and graphs with an extremal number of trees, *J. Combin. Theory, Series B* 16 (1974) 197–214.
- [38] S.J. Kirkland, M. Neumann, On group inverses of M -matrices with uniform diagonal entries, *Linear Algebra Appl.* 296 (1999) 153–170.
- [39] S.J. Kirkland, M. Neumann, B.L. Shader, Distances in weighted trees and group inverse of Laplacian matrices, *SIAM J. Matrix Anal. Appl.* 18 (1997) 827–841.
- [40] J.J. Koliha, Block diagonalization, *Mathematica Bohemica* 126 (2001) 237–246.
- [41] J.J. Koliha, I. Straškraba, Power bounded and exponentially bounded matrices, *Applications of Mathematics* 44 (1999) 289–308.
- [42] T. Leighton, R.L. Rivest, The Markov chain tree theorem, *Computer Science Technical Report MIT/LCS/TM-249*, Laboratory of Computer Science, MIT, Cambridge, MA, 1983.
- [43] T. Leighton, R.L. Rivest, Estimating a probability using finite memory, *IEEE Transactions on Information Theory* IT-32 (1986) 733–742.
- [44] G. Marsaglia, G.P.H. Styan, Equalities and inequalities for ranks of matrices, *Linear and Multilinear Algebra* 2 (1974) 269–292.
- [45] J.S. Maybee, D.D. Olesky, P. van den Drissche, G. Wiener, Matrices, digraphs, and determinants, *SIAM J. Matrix Anal. Appl.* 10 (1989) 500–519.
- [46] R. Merris, Doubly stochastic graph matrices, *Univ. Beograd. Publ. Electrotechn. Fak., Ser. Mat.* 8 (1997) 64–71.
- [47] R. Merris, Doubly stochastic graph matrices II, *Linear and Multilinear Algebra* 45 (1998) 275–285.
- [48] C.D. Meyer Jr., Limits and the index of a square matrix, *SIAM J. Appl. Math.* 26 (1974) 469–478.
- [49] C.D. Meyer Jr., The role of the group generalized inverse in the theory of finite Markov chains, *SIAM Review* 17 (1975) 443–464.

- [50] C.D. Meyer Jr., M.W. Stadelmaier, Singular M-matrices and inverse positivity, *Linear Algebra Appl.* 22 (1978) 139–156.
- [51] M. Minoux, A generalization of the all minors matrix tree theorem to semirings, *Discrete Mathematics* 199 (1999) 139–150.
- [52] J.W. Moon, Counting Labelled Trees, Canadian Mathematical Congress, Montreal, 1970.
- [53] J.W. Moon, Some determinant expansions and the matrix-tree theorem, *Discrete Mathematics* 124 (1994) 163–171.
- [54] W. Myrvold, Counting k -component forests of a graph, *Networks* 22 (1992) 647–652.
- [55] W.S. Percival, The solution of passive electrical networks by means of mathematical trees, *Proceedings of the Institute of Electrical Engineers* 100 (1953) 143–150.
- [56] J.G. Propp, D.B. Wilson, How to get a perfectly random sample from a generic Markov chain and generate a random spanning tree of a directed graph, *J. Algorithms* 27 (1998) 170–217.
- [57] U.G. Rothblum, Computation of the eigenprojection of a nonnegative matrix at its spectral radius, *Mathematical Programming Study* 6 (1976) 188–201.
- [58] U.G. Rothblum, A representation of the Drazin inverse and characterizations of the index, *SIAM J. Appl. Math.* 31 (1976) 646–648.
- [59] U.G. Rothblum, Expansions of sums of matrix powers, *SIAM Review* 23 (1981) 143–164.
- [60] U.G. Rothblum, Resolvent expansions of matrices and applications, *Linear Algebra Appl.* 38 (1981) 33–49.
- [61] A.J. Schwenk, The adjoint of the characteristic matrix of a graph, *J. Combin., Inform. Syst. Sci.* 16 (1991) 87–92.
- [62] E.V. Shamis, Graph-theoretic interpretation of the generalized row sum method, *Mathematical Social Sciences* 27 (1994) 321–333.
- [63] J.J. Sylvester, On the change of systems of independent variables, *Quart. J. Pure Appl. Math.* 1 (1857) 42–56, reprinted in: *Collected Math. Papers*, Cambridge 2 (1908) 65–85.
- [64] W.T. Tutte, The dissection of equilateral triangles into equilateral triangles, *Proceedings of the Cambridge Philosophical Society* 44 (1948) 463–482.
- [65] W.T. Tutte, *Graph Theory*, Addison-Wesley, Reading, MA, 1984.
- [66] Y. Wei, A characterization and representation of the Drazin inverse, *SIAM J. Matrix Anal. Appl.* 17 (1996) 744–747.
- [67] Y. Wei, H. Wu, The representation and approximation for Drazin inverse, *J. Comp. Appl. Math.* 126 (2000) 417–432.
- [68] L. Zhang, A characterization of the Drazin inverse, *Linear Algebra Appl.* 335 (2001) 183–188.
- [69] J.H.M. Wedderburn, *Lectures on Matrices*, Colloq. Publ., vol. 17, AMS, Providence, RI, 1934.