

lead to a better understanding of the molecular mechanisms responsible for apoptotic cell clearance. If the clearance process is impaired, apoptotic cells may progress to secondary necrosis, resulting in release of harmful cellular contents and tissue inflammation. (IUPUI Membrane Biosciences Signature Center grant.)

2409-Pos

Lipid-Induced Up-Regulation of Acyl-CoA Synthetase 5 Promotes Apoptosis in Human Hepatocytes

Andrea Reinartz^{1,2}, Christopher A. Haynes², Ruth Knuechel¹, Alfred H. Merrill, Jr.², Nikolaus Gassler¹.

¹RWTH Aachen University Hospital, Aachen, Germany, ²Georgia Institute of Technology, Atlanta, GA, USA.

Long chain acyl-CoA synthetases (ACSL) activate fatty acids for utilization by numerous metabolic pathways. Of the five mammalian ACSL isozymes known, ACSL5 is the only one located on mitochondria and thought to be involved in apoptosis. Fatty acids up-regulate ACSL5 and increase apoptosis susceptibility in human hepatocytes, thus, we hypothesize that ACSL5 is a promoting factor in hepatocellular lipoapoptosis. To investigate this mechanism, we have used immunochromatographic and RNA interference as well as liquid chromatography, tandem mass spectrometry (LC-MS/MS). Fatty acid uptake led to up-regulation of ACSL5 expression and enzymatic activity in primary hepatocytes, HepG2 cells and steatotic liver. Over-expression of ACSL5 decreased HepG2 cell viability and increased susceptibility to TRAIL and TNF α , whereas knock down of ACSL5 reduced apoptosis susceptibility in fatty-acid treated HepG2 cells. Apoptosis sensitization was accompanied by enhanced caspase-3/7 activity, but was not associated with up-regulation of DR4, DR5 or TNF-R1. By applying lipidomic techniques, we determined the effect of ACSL5 on the cellular amounts and subspecies of fatty acyl-CoAs as well as on sphingolipids, the downstream metabolites that are known to be important regulators of cell death and survival. High ACSL5 activity in HepG2 cells increased synthesis of long-chain acyl-CoAs by 50%, and enhanced ceramide and sphingomyelin levels by 2 to 3 fold. These results indicate that steatosis-induced up-regulation of ACSL5 increased apoptosis susceptibility in human hepatocytes and that alterations in sphingolipid metabolism might contribute to ACSL5-mediated apoptotic effects. We propose that ACSL5 could play a role in promoting fatty acid-induced lipoapoptosis in hepatocytes as an important mechanism in fatty liver-related disorders.

2410-Pos

Differential Susceptibility of Normal and Transformed Human Leukocytes to Hydrolytic Attack by Secretory Phospholipase A₂

Lynn Anderson, Kelly Damm, Ryan Baker, Joseph Chen, Amy Hamaker, Izadora Izidoro, Eric Moss, Mikayla Orton, Kristin Papworth, Lyndee Sherman, Evan Stevens, Celestine Yeung, Jennifer Nelson, Allan M. Judd, John D. Bell.

Brigham Young University, Provo, UT, USA.

Previous experiments with cultured lymphoma cells demonstrated that secretory phospholipase A₂ (sPLA₂) distinguishes healthy cells from those that are dying by apoptosis or necrosis. This distinction depends on cell membrane properties including the amount of negative charge at the bilayer surface and the strength of interactions among neighboring phospholipids. These results raised two important questions. 1) Does the enzyme's ability to distinguish healthy and dying cells apply to normal human leukocytes? 2) Does sPLA₂ differentiate between normal and tumor cells? These questions were addressed by comparing membrane properties and susceptibility to hydrolysis among cultured transformed leukocytes and freshly-isolated human neutrophils and lymphocytes. Membrane properties were assessed by flow cytometry, merocyanine 540 fluorescence spectra, trimethylammonium diphenylhexatriene fluorescence anisotropy, and two-photon scanning microscopy with lauridan. Similar to the behavior of transformed cells, normal human leukocytes resisted hydrolysis by sPLA₂. Upon addition of a calcium ionophore, ionomycin, the cells became vulnerable to hydrolysis, again analogous to the results observed with tumor cells. However, several important quantitative distinctions were observed. First, the various types of normal leukocytes responded differently to the enzyme; lymphocytes exhibited significantly greater rates of hydrolysis by sPLA₂ compared to granulocytes. Second, hydrolysis was substantially slower in normal cells compared to transformed cells. Third, the time required for ionomycin to induce cells to be attacked by sPLA₂ was greater in normal compared to transformed cells. Likewise, changes in membrane physical properties following ionomycin treatment were more subtle in normal cells than they were in transformed cells. These results suggest the possibility that sPLA₂ could function as a therapeutic ally during cancer chemotherapy to assist with the demise of tumor cells.

2411-Pos

Kinetic Evaluation of Cell Membrane Hydrolysis during Apoptosis by Human Isoforms of Secretory Phospholipase A₂

Jennifer Nelson, Erin Olson, Katalyn Griffith, Michael Streeter, Allan M. Judd, John D. Bell.

Brigham Young University, Provo, UT, USA.

Some isoforms of secretory phospholipase A₂ (sPLA₂) distinguish between healthy and damaged or apoptotic cells. This distinction reflects differences in membrane physical properties. Since various sPLA₂ isoforms respond differently to properties of artificial membranes such as surface charge, they should also behave differently as these properties evolve during a dynamic physiological process such as apoptosis. To test this idea, S49 lymphoma cell death was induced by glucocorticoid (6-48 h), thapsigargin (3-4 h) or calcium ionophore. Rates of membrane hydrolysis catalyzed by various concentrations of snake venom and human groups IIa, V, and X sPLA₂ were compared after each treatment condition. The data were analyzed using a model that evaluates the adsorption of enzyme to the membrane surface and subsequent binding of substrate to the active site. Results were compared temporally to changes in membrane biophysics and composition. Under control conditions, membrane hydrolysis was confined to the few unhealthy cells present in each sample. Increased hydrolysis during apoptosis and necrosis appeared to reflect substrate access to adsorbed enzyme for the snake venom and group X isoforms corresponding to weakened lipid-lipid interactions in the membrane. In contrast, apoptosis promoted initial adsorption of human groups V and IIa concurrent with phosphatidylserine exposure on the membrane surface. However, this observation was inadequate to explain the behavior of the groups V and IIa enzymes toward necrotic cells where hydrolysis was reduced or absent. The response to endoplasmic reticulum stress (thapsigargin) was intermediate between that observed for glucocorticoid and ionomycin. Thus, a combination of changes in cell membrane properties during apoptosis and necrosis capacitates the cell for hydrolysis differently by each isoform.

2412-Pos

VDAC1 Cysteine Residues: Topology and Function in Channel Activity and Apoptosis

Varda Shoshan-Barmatz, Shay Geula, Lior Aram, Nir Arbel.

Department of Life Sciences and the NIBN Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel.

VDAC is proposed to control metabolic cross-talk between mitochondria and the cytosol, as well as apoptotic cell death. It has been suggested that apoptosis is modulated by the oxidation state of VDAC. Since cysteine residues are the major target for oxidation/reduction, we verified whether one or both VDAC1 cysteine residues are involved in VDAC1-mediated transport or apoptosis activities. To assess the function of the VDAC1 cysteines in channel activity and to probe cysteine topology, with respect to facing the pore or the bilayer, we used thiol-modifying agents; membrane permeable N-ethyl-maleimide (NEM); bulky, charged 5-fluorescein-maleimide (5-FM) and cross-linking reagent bis[maleimido]ethane (BMOE). Bilayer-reconstituted VDAC conductance was decreased by 5-FM but not by NEM, while 5-FM had no effect on NEM-labeled VDAC conductance. BMOE formed dimeric VDAC1, suggesting that a VDAC1 cysteine residue is exposed and available for cross-linking. The results suggest that one of the VDAC1 cysteine residues faces the VDAC pore while the second is oriented toward the lipid bilayer. The positions of VDAC1 Cys127 and Cys232 with respect to the membrane and channel pore, were considered in light of proposed VDAC1 topology models. Mutated VDAC1 in which Cys127 and Cys232 were replaced by alanines showed channel activity as of native VDAC1 and when expressed in cells was localized to mitochondria. As with over-expression of native rVDAC1, cysteine-less rVDAC1 induced apoptotic cell death and underwent oligomerization upon apoptosis induction. The results suggest that the two cysteine residues are not required for VDAC1 oligomerization or apoptosis, as induced by H₂O₂, As₂O₃ or selenite, ROS producing agents.

2413-Pos

A Voltage Dependent Na⁺ Channel is Activated during Apoptosis in Xenopus Oocytes

Ulrika H. Englund, Jens Gertow, Fredrik Elinder.

IKE, Linköping, Sweden.

Apoptosis is regulated by a cascade of intracellular biochemical reactions. However, plasmamembrane bound ion channels are also essential for the apoptotic process. In previous studies we and other have found that K, Cl and Na channels of different types are upregulated early in the apoptotic process. Furthermore, block of these channels prevent or delay the apoptosis, suggesting a critical role of the channels in the apoptotic process. In the present investigation we examined whether ion channels are upregulated in oocytes from the