

Cephalometric assessment of the hyoid bone position in Oral Breathing Children

Maria Julia Pereira Coelho Ferraz¹, Darcy Flávio Nouer², José Ricardo Teixeira³, Fausto Bérzin⁴

Keywords: cephalometry, hyoid bone, oral breathing.

Summary

Material and Methods: because of its anatomical and functional relationship with the craniofacial complex, we assessed the cephalometry of the hyoid bone position in relation to the respiratory pattern of these 53 female children, with average age of 10 years; 28 of them are nasal breathers and 25 are oral breathers. Horizontal, vertical and angular cephalometric measures were used in order to determine the hyoid bone location. The Student "t" and the Pearson correlation tests were used in order to compare the groups and the variables. **Results:** We did not see statistically significant differences in mandible and hyoid bone positions and the respiratory pattern. In the hyoid triangle, the 0.40 correlation coefficient was significant between AA-ENP (distance between the Atlas vertebrae and the posterior nasal spine) and C3-H (distance between the third cervical vertebrae and the hyoid bone) showing a positive relation between the bony limits of the upper and lower air spaces. For cranial measures we have suggested a relationship between the hyoid bone position and the mandible morphology. **Conclusion:** The results led us to conclude that the hyoid bone keeps a stable position, probably in order to secure correct ratios in the airways, and it does not depend on the respiratory pattern.

¹ M.S. In Orthodontics. Member of CEBAPE - Biosciences Center Applied to Patients with Special Needs / UNESP. PhD Student in Buccodental Biology - anatomy department - FOP/UNICAMP, DDS.

² PhD. Professor of Orthodontics - Faculdade de Odontologia de Piracicaba/ UNICAMP and Head of the Department of Pediatric Dentistry - Faculdade de Odontologia de Piracicaba/ UNICAMP.

³ MD. Otorhinolaryngologist - AMB. Preceptor at the ENT Medical Residency - Núcleo de ORL de Limeira (NOL).

⁴ PhD. Full Professor at the Department of Morphology - FOP/UNICAMP.

Summary of the MS thesis dissertation carried out at the Department of Pediatric Dentistry - Faculdade de Odontologia de Piracicaba/ UNICAMP.

Mailing Address: Maria Julia Pereira Coelho Ferraz - Alameda João Senra 148 Colinas de São João Limeira SP 13481-299. Tel: (0xx19) 3443-1994/ 9608-2646 - E-mail: mariajulia@respiremehor.com.br

Paper submitted to the ABORL-CCF SGP (Management Publications System) on September 2nd, 2005 and accepted for publication on June 5th, 2006.cod.1028

INTRODUCTION

Based on the complexity of the stomatognathic system, specific knowledge on its anatomy, physiology and craniofacial growth theories are paramount in order to understand its whole functioning in individuals.

According to Meredith¹, an important growth increment happens in the first years of life. At birth, the craniofacial bones of a caucasian American corresponds to 60% of the adult head size, 80% at six months of age, 90% at three years, and 95% at nine years of age. Thus, at 12 years of age, when many orthodontists start treatment, almost all facial growth is completed.

Regarding oral breathing, it is possible to pin it as a cause of malocclusion and related areas. Inadequate breathing patterns cause functional adaptations, promoting facial muscle balance, postural changes such as open lips, posterior head tilt and a lower position for mandible and tongue. Consequent to such unbalances, there may be undesirable changes to the craniofacial morphology. Nonetheless, data behind these statements are limited and obscure²⁻⁷.

The importance of the hyoid bone is related to its single relation, nonetheless, it provides connections to pharynx, mandible and cranial muscles, ligaments and fascia.^{8,9}

Many of the characteristics of the so called Long Face Syndrome (LFS) group and the Short Face Syndrome (SFS) group could be explained based on the clockwise and counter-clock wise rotation of the mandible "in harmony" with the hyoid bone, tongue, pharynx and neck. The vital need to keep the air space pattern at the tongue base may explain this rotation in the LFS¹⁰.

Adenoid tissue or tongue mass may reduce the air space and cause postural adaptations at the level of the oropharynx. A hyoid bone drop in relation to the mandible would represent an attempt to assure a relatively constant air-space diameter in the antero-posterior direction. This neuromuscular recruiting could cause changes in mandibular rest position and neck extension, thus influencing the craniofacial growth pattern¹¹.

Thus, air space pattern and stability would represent major factors responsible for hyoid bone position¹².

Since malocclusion may be caused by inadequate oral habit, such as atypical swallowing and oral breathing - hyoid bone position could serve as an important diagnostic guide⁹.

Numerous authors have studied the hyoid bone morphology and function^{13,14}, and others have investigated hyoid bone position in relation to the cranium and the cervical spine by means of cephalometric techniques^{8,15-30}.

Since the studies regarding hyoid bone position represent an open field in sciences, the present study aimed at investigating the relationship between hyoid bone,

mandible, cranium and cervical spine, by cephalometric means, trying to establish a relationship between hyoid bone position in oral breathing and in nasal breathing patients in order to aid in the diagnosis of alterations in the craniofacial complex, of multidisciplinary character, involving otorhinolaryngologists, orthodontists, functional orthopedists, speech therapists and physical therapists.

MATERIALS AND METHODS

The present investigation only started after it was approved by the Research in Human Beings Ethics Committee (CEP) - FOP-UNICAMP, according to the documentation required by Resolution 196/96 of the National Committee of Research Ethics (CONEP) - National Health Council - Ministry of Health.

For this study, we used side view teleradiographies from 53 Caucasian, female individuals, selected from a pool of 450 duly enrolled children in the Municipal School Network at the city of Limeira, ranging between 9 and 12 years.

We requested authorization from the schools and also the written consent from the parents of the children who participated in this research project.

The children were evaluated by a dentist by means of an anamnesis and standard dental examination from UNICAMP. After that, the children were referred to take a lateral view teleradiography and on the same day underwent a nasofibrosopic exam. After diagnosing the predominant respiratory pattern, issued by the otorhinolaryngologist, the children were divided into control group - predominantly nasal breathers (n = 28), and experimental group - predominantly oral breathers (n = 25). Later on we carried out the cephalometric analysis.

The following criteria were used in order to select the sample from the present group: females, Class I Angle malocclusion and mixed dentition, no orthodontic treatment and/or functional orthopedic treatment of the maxillas, no extensive carious lesions, enough contrast and sharpness for a good visualization and identification of the structures that make up the tegumentary tissue, bone structures, dental elements and the hyoid bone, and no radiographic distortions.

The teleradiographies were taken laterally and in the natural head position (NHP)^{31,32} always by the same technician - person responsible for the Department of Dental Documentation, following the standards established by the School of Dentistry of Piracicaba/UNICAMP.

All the individuals were evaluated by the otorhinolaryngologist in charge of diagnosing the respiratory pattern. He examined the patients' ears nose and throat and did a nasofibrosopic, evaluated the questionnaire answered by the parents, the history filled out by the dentist, as well as the lateral view teleradiography, making up the standard respiratory process and classifying them as predominantly

nasal (clinically normal) or predominantly oral breathers. The following material were employed for nasofibrosco-
py: Samsung 14" monitor; Sony 4 head VCR; Welch Allyn
light source; micro camera for a Toshiba endoscope; the
endoscopes used were from two models, with 0° and
30° visual angle; Pentax flexible nasofibroscope; Sony
videotape; paper towel, gauze, 2% glutaraldehyde; 70%
alcohol and nasal decongestant; 45% lidocaine and 10%
spray xylocaine. The otorhinolaryngologist used the Ianni
Filho³³ and Wang et al.³⁴ protocols in order to present
the nasofibrosopic results and the exams were recorded
in videotape, which became part of the Department of
Pediatric Dentistry's Library at the FOP/UNICAMP.

The hyoid bone was cephalometrically evaluated,
using craniofacial and hyoid measures²¹, adapted to the
present study, and also the measures from the Hyoid
Triangle⁸ (Figures 1, 2,3).

In order to evaluate the reliability of the cepha-
lometric measures, the curves were done twice by one
single researcher in one week interval, keeping the same
environmental conditions and work instruments. We used
the average of the values collected in the two curves. Thirty
days after the curves were made, ten teleradiographies
were randomly selected from individuals enrolled in this
research, aiming at checking the error made between
the two moments by means of calculating the error as
proposed by Dalberg³⁵ and advocated by Houston³⁶. For




Figure 1. Cephalometric measures to determine mandibular position.

1- Ar-Pog	6- BaN.PM
2- PP-Me	7- PTM.PM
3- ENP-PM	8- PO.PM
4- S-PM	9- PPM
5- Goniac angle	




Figure 2. Cephalometric measures to determine the hyoid bone position.

1- d horiz. H	4- PH.PP
2- d vert. H	5- PH.PM
3- PH.BaN	6- PH.PO




Figure 3. Hyoid Triangle linear and angular cephalometric measures.

1- C3-RGn	4- H-H'
2- C3-H	5- Hyoid plane angle
3- H-RGn	6- AA-ENP

data analysis, initially we obtained a descriptive analysis (mean and standard deviation), and later we applied the F test and the t Student test, with 5% significance level. The correlation between variables AA-ENP (antero-posterior distance between the first cervical vertebra and the pos-




Figure 4. Correlation between AA-ENP and C3-H measures.

terior nasal spine) and C3-H (distance between the most anterior-inferior point of the third cervical vertebra and the hyoid bone body), analyzed by the Pearson³⁷ correlation coefficient (1909), mentioned by Stepovich (1965)¹⁹, with a significance level of $\alpha=0.05$.

RESULTS

Calculating the error we noticed that there was no statistical significance among the moments assessed, showing reliability in the curves and measures.

Mandibular cephalometric measures' mean values and standard deviations are presented on Table 1. Average differences, as shown by the results from the "t" tests are presented on the same table, and are not significant ($p > 0.05$).

Average and standard deviation of the measures that characterize the hyoid bone position are presented on Table 2. According to the results from the t tests presented on the same table, the mean cephalometric values from predominantly oral and nasal breathers are not statistically significant ($p > 0.05$).

In the present investigation we also studied cranial values that measure the hyoid bone position according to Bibby & Preston⁸. The data obtained from the 53 patients enrolled in the present investigation and divided in two groups are depicted on Table 3, not presenting statistically significant difference between the groups ($p > 0.05$).

A significant, though weak, correlation ($r=0.40$), between AA-ENP and C3-H was found (Figure 4) representing the upper and lower limits of the air space, respectively.

DISCUSSION

The sample was standardized in relation to gender, since it is necessary to consider sexual differences in skeletal morphology, and the basic size difference happens after puberty, with males growing more and for longer than women³⁸, and most specifically in regards of cervical spine growth^{15,17} and hyoid bone position in relation to the mandible³⁰.

In relation to the age range, mean values were of 10 years for predominantly nasal and oral breathers, we

Table 1. Mean value, Standard deviation and t test of the measures related to mandible position (values in degrees and millimeters).

Measures	Nasal		Oral		p*
	Mean	SD	Mean	SD	
Ar-Pog	101.96	5.69	102.06	5.80	0.95 (NS)
PP-Me	59.59	3.43	59.85	3.90	0.80 (NS)
ENP-PM	42.87	4.45	43.65	3.48	0.49 (NS)
S-PM	44.23	3.05	43.17	2.97	0.21 (NS)
Ang. Gon.	125.16	4.34	123.40	5.04	0.18 (NS)
BaN.PM	53.41	3.51	52.32	5.10	0.37 (NS)
PTM.PM	117.18	3.39	117.80	4.64	0.59 (NS)
PP.PM	27.83	3.50	28.11	4.52	0.81 (NS)
PO.PM	15.08	3.32	14.42	2.74	0.43 (NS)

SD=Standard Deviation

*Student t test

(NS) not significant

Table 2. Mean, Standard deviation and t test of the measures related to hyoid bone position (values in degrees and millimeters).

Measures	Nasal		Oral		p*
	Mean	SD	Mean	SD	
d horiz. H	27.58	2.59	27.83	3.49	0.78 (NS)
d vert. H	52.41	5.45	52.95	6.46	0.74 (NS)
PH.BaN	54.30	7.42	51.44	11.33	0.29 (NS)
PH.PP	28.58	7.29	26.57	10.25	0.61 (NS)
PH.PM	1.03	6.58	-1.44	10.71	0.32 (NS)
PH.PO	16.07	7.32	12.93	10.50	0.22 (NS)

SD=Standard Deviation

*Student t test

(NS) not significant

Table 3. Mean, Standard deviation and t test of the measures related to the Hyoid triangle (values in degrees and millimeters).

Measures	Nasal		Oral		p*
	Mean	SD	Mean	SD	
C3-RGn	64.90	6.68	67.09	6.90	0.55 (NS)
C3-H	31.99	2.99	32.45	2.54	0.55 (NS)
H-RGn	33.70	5.09	35.53	5.76	0.23 (NS)
H-H'	1.53	5.18	2.36	5.12	0.56 (NS)
Ang. PH	22.74	7.94	20.72	12.25	0.49 (NS)
AA-ENP	32.87	3.34	32.86	2.18	0.98 (NS)

SD=Standard Deviation

*Student t test

(NS) not significant

considered the high frequency with which craniofacial alterations justify the search for orthodontic treatment. Meredith¹ commented that at nine years of age, 95% of the craniofacial growth has already happened. This shows the importance of the prevention aspect in the diagnostic process.

Stressing the importance of the multi and interdisciplinary approach among health care professionals involved in the diagnosis and treatment of oral breathers, we chose the nasofibroscope carried out by the ENT, together with the telerradiography taken at a lateral view, dental assessment carried out by the investigator, interview and questionnaires answered by the parents, making up the diagnostic process of the respiratory pattern.

In a general way, the hyoid bone cephalometric measures do not show significant differences among the groups with predominantly nasal and oral breathing. This statement is in agreement with studies from Subtelny & Sakuda¹⁸, Bibby & Preston⁸, Bibby⁹, Kumar et al.²⁵ and Kawashima et al.²⁹, in which the hyoid bone presented a permanent rest position, unrelated with the respiratory pattern. Nonetheless, Adamidis & Spyropoulos²¹ found statistically significant differences in tongue, mandible and hyoid bone position among nasal and oral breathers. According to findings from the author who included oral breathers among Class I malocclusion children when compared to a control group with ideal occlusion, without evidences of nasopharynx obstruction, the respiratory pattern impacted mandible and hyoid bone positions. Authors such as Behlfelt et al.^{22,23}, Shintani et al.²⁶, Finkelstein et al.²⁷ found differences in hyoid bone vertical position.

In regards of the Hyoid Triangle measures, horizontal linear measures C3-H, C3-Rgn and H-Rgn did not show significant differences between oral and nasal breathers, respectively, in accordance with Bibby⁹ and Kawashima et al.²⁹. Nonetheless, qualitatively speaking, when we compared the mean values, we noticed that the C3-Rgn antero-posterior dimension (67.09mm and 64.90mm) was greater for the group of oral breathers when compared to nasal breathers. The antero-posterior dimension between the first cervical vertebra (AA) and the PNS (posterior nasal spine) was constant for both groups, since it is established in initial childhood stages¹⁵. Mean value was of 32.87mm and standard deviation of 3.34 for predominantly nasal breathers; and of 32.86mm and standard deviation of 2.18 for predominantly oral breathers. These results corroborate the ones from Bibby & Preston⁸, Ferraz et al.³⁰, with mean values of 32.91mm, 32.83mm and standard deviation of 3.66 and 2.69, respectively. The correlation coefficient between these two horizontal measures AA-ENP and C3-H was significant and positive ($r=0.40$); however, less so when compared to the results from Bibby & Preston⁹ ($r=0.98$) and Ferraz et al.³⁰ ($r=0.56$) who defined the hyoid bone as the pharynx anterior bony limit at a lower level

than the PNS.

Considering mean values, it has been noticed that the hyoid bone vertical behavior in relation to C3-RGN was positioned more caudally in predominantly oral breathers, with mean value of 2.36mm and standard deviation of 5.12 when compared to nasal breathers, whose mean values was of 1.53mm and standard deviation of 5.18. This lower position, under the qualitative view point, could be interpreted as a postural adaptation at the level of the oropharynx, in an attempt to keep constant the antero-posterior diameter^{11,12,22}.

Finally, the need to keep the correct ratios in the airways is one of the key factors that control hyoid bone position in individuals with a different respiratory pattern. The craniofacial complex tries to achieve a more favorable position in order to carry out its function and, therefore, adapts itself, according to its possibilities, aiming at what is easier in order to breathe properly.

CONCLUSION

Considering the characteristics of the sample used, the methodology employed and the rigorous data analysis, it was possible to conclude that respiratory pattern did not interfere in mandibular position, or in hyoid bone position, which was maintained stable, most likely in order to assure the correct proportions of air space.

In the present investigation, quantitative studies were necessary in order to investigate changes in hyoid bone position, considering possible clinical implications of neuromuscular adaptations of oral breathing vis-à-vis the cervical spine and body posture.

REFERENCES

1. Meredith Pediatrics HV. Growth in head width during the first twelve years of life. *Pediatr* 1953;12(4):411-29.
2. Harvold EP, Vargervik K, Chierici G. Primate experiments on oral sensation and dental malocclusions. *Am J Orthod* 1973;63(5):494-508.
3. Rubin RM. Mode of respiration and facial growth. *Am J Orthod* 1980;78(5):504-10.
4. Bresolin D, Shapiro GG, Shapiro PA, Dassel SWD, Furukawa CT, Pierson WE et al. Facial characteristics of children who breathe through the mouth. *Pediatrics* 1984;73(5):622-5.
5. McNamara JA. A method of cephalometric evaluation. *Am J Orthod* 1984;86(6) p.449-69.
6. Solow B, Siersbaek-Nielsen S, Greve E. Airway adequacy, head posture, and craniofacial morphology. *Am J Orthod* 1984;86(3):214-23.
7. Vargervik K, Miller AJ, Chierici G, Harvold E, Tomer BS. Morphologic response to changes in neuromuscular patterns experimentally induced by altered modes of respiration. *Am J Orthod* 1984;85(2):115-24.
8. Bibby RE, Preston CB. The hyoid triangle. *Am J Orthod* 1981;80(1):92-7.
9. Bibby RE. The hyoid bone position in mouth breathers and tongue-thrusters. *Am J Orthod* 1984;85(5):431-3.
10. Opdebeeck MD, Bell WH, Eisenfeld J, Mischelevich D. Comparative study between the SFS and LFS rotation as a possible morphogenic mechanism. *Am J Orthod* 1978;74(5):509-21.

11. Tourné LP. Growth of the pharynx and its physiologic implications. *Am J Orthod Dentofacial Orthop* 1991;99(2):129-39.
12. Graber LW. Hyoid change following orthopedic treatment of mandibular prognathism. *Angle Orthod* 1978;48(1):33-8.
13. Brodie AG. Anatomy and physiology of head and neck musculature. *Am J Orthod* 1950;36:831-44.
14. Durzo CA, Brodie AG. Growth of the hyoid bone. *Angle Orthod* 1962;32(3):193-204.
15. King EW. A roentgenographic study of pharyngeal growth. *Angle Orthod* 1952;22(1):23-37.
16. Grant LE. A radiographic study of hyoid bone position in Angle's Class I, II, and III malocclusions [Master's thesis]. Kansas: University of Kansas City; 1959. Apud Stepovich ML. A cephalometric positional study of the hyoid bone. *Am J Orthod* 1965;51(12):882-900.
17. Bench RW. Growth of the cervical vertebrae as related to tongue, face, and denture behavior. *Am J Orthod* 1963;49(3):183-214.
18. Subtelny JD, Sakuda M. Open-bite: diagnosis and treatment. *Am J Orthod* 1964;50(5):337-58.
19. Stepovich ML. A cephalometric positional study of the hyoid bone. *Am J Orthod* 1965;51(12):882-90.
20. Ingerval B, Carlsson GE, Helkimo M. Change in location of hyoimandibular positions. *Acta Odontol Scand* 1970;28(3):337-61.
21. Adamidis IP, Spyropoulos MN. The effects of lymphadenoid hypertrophy on the position of the tongue, the mandible and the hyoid bone. *Eur J Orthod* 1983;5(4):287-94.
22. Behlfelt K, Linder-Aronson S, Neander P. Posture of the head, the hyoid bone, and the tongue in children with and without enlarged tonsils. *Eur J Orthod* 1990a;12(4):458-67.
23. Behlfelt K, Linder-Aronson S, McWilliam J, Neander P, Laage-Hellman J. Cranio-facial morphology in children with and without enlarged tonsils. *Eur J Orthod* 1990b;12(3):233-43.
24. Haralabakis NB, Toutountzakakis NM, Yiagtzis SC. The Hyoid bone position in adult individuals with open bite and normal occlusion. *Eur J Orthod* 1993;15(4):265-71.
25. Kumar R, Sidhu SS, Kharbanda OP, Tandon, DA. Hyoid Bone and atlas vertebra in established mouth breathers: a cephalometry study. *J Clin Pediatr Dent* 1995;19(3):191-4.
26. Shintani T, Asakura K, Kataura A. Evaluation of the role of adenotonsillar hypertrophy and facial morphology in children with obstructive sleep apnea. *ORL* 1997;59(5):286-91.
27. Finkelstein Y, Wexler D, Berger G, Nachmany A, Shapiro-Feinberg M, Ophir D. Anatomical basis of sleep-related breathing abnormalities in children with nasal obstruction. *Arch Otolaryngol Head Neck Surg* 2000;126(5):593-600.
28. Yamaoka M, Furusawa K, Uematsu T, Okafuji N, Kayamoto D, Kurihara S. Relationship of the hyoid bone and posterior surface of the tongue in prognathism and micrognathia. *J Oral Rehabil* 2003;30(9):914-20.
29. Kawashima S, Peltomäki T, Sakata H, Mori K, Happonen R-P, Rönning O. Absence of facial type differences among preschool children with sleep-related breathing disorder. *Acta Odontol Scand* 2003;61(2):65-71.
30. Ferraz MJPC, Sousa MA, Pereira Neto JS, Nouer DF, Magnani MBBA. Avaliação cefalométrica o triângulo hióideo em brasileiros da região de Piracicaba. In: *Anais da Reunião da Sociedade Brasileira de Pesquisa Odontológica*, 2003. Águas de São Pedro. São Paulo: SBPQO; 2003. p.161. [Resumo Pb017].
31. Solow B, Tallgren A. Natural head position in standing subjects. *Acta Odontol Scand* 1971;29(5):591-607.
32. Viazis AD. A cephalometric analysis based on natural head position. *J Clin Orthod* 1991;25(3):172-81.
33. Ianni Filho D, Raveli DB, Raveli RB, de Castro Monteiro Loffredo L, Gandini LG. A comparison of nasopharyngeal endoscopy and lateral cephalometric radiography in the diagnosis of nasopharyngeal airway obstruction. *Am J Orthod Dentofacial Orthop* 2001;120(4):348-52.
34. Wang DY, Bernheim N, Kaufman L, Clement P. Assessment of adenoid size in children by fiberoptic examination. *Clin Otolaryngol allied Sci* 1997;22(2):172-77.
35. Dahlberg G. Statistical methods for medical and biological students. New York; 1940. Apud Houston WJ. The analysis of errors in orthodontic measurements. *Am J Orthod* 1983;83(5):382-90.
36. Houston WJ. The analysis of errors in orthodontic measurements. *Am J Orthod* 1983;83(5):382-90.
37. Pearson FG. Topography and morphology of the human hyoid. *J Anat Physiol* 1909;43:279-90. Apud Stepovich ML. A cephalometric positional study of the hyoid bone. *Am J Orthod* 1965;51(12):882-900.
38. Enlow DH, Hans MG. *Noções básicas sobre crescimento facial*. São Paulo: Santos; 1998, 304 pp.