

Available online at www.sciencedirect.com

ScienceDirect

Procedia Computer Science 86 (2016) 132 – 135

Procedia
Computer Science

2016 International Electrical Engineering Congress, iEECON2016, 2-4 March 2016, Chiang Mai, Thailand

Non-thermal plasma for germination enhancement of radish seeds

Khanit Matra*

*Department of electrical engineering, Engineering faculty, Srinakharinwirot University,
63 M.7 Rangsit-Nakhonnayok road, Ongkharak, Nakhonnayok 26120, Thailand*

Abstract

This paper is aiming to present the atmospheric non-thermal plasma for the germination improving of radish seeds. Plasma generation power and plasma irradiation time were studied in order to obtain the proper conditions for the germination enhancement of seeds. The growth rate was determined by measuring the length of root and dry weight of the plants treated by plasma jet. From the experimental results, it could be confirmed that plasma treatment could enhance the growth rate of radish seeds 9-12 percent of the dry weight of radishes, approximately.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the Organizing Committee of iEECON2016

Keywords: non-thermal, plasma jet, germination enhancement, seeds ;

1. Introduction

Nowadays, non-thermal atmospheric gas discharge or plasma has shown great interest among scientists owing to its many advantages such as low temperature, scalable size, low operation cost, flexible operation, and high electron and reactive specie density. Non-thermal plasma has been principally used for surface modification because of its energetic electron and reactive species. Recently, non-thermal plasma has been promoted for applying in biotechnology, medical treatment, food preservation, and agriculture. This is due to the fact that non-thermal plasma does not cause thermal damage to the living organisms, and it is also a green technology^{1,2}.

Regarding the literature reviews of the seed growth enhancement by plasma treatment, the results show worthy outcomes; in addition, Thailand is an agriculture country^{1,2,3,4}. Therefore, our research group has come across idea to develop and improve the agriculture technology by our non-thermal plasma jet model. In this work, the preliminary

* Corresponding author. Tel.: +662-649-5000; fax: +663-732-2602.

E-mail address: khanit@g.swu.ac.th

results of seed germination enhancement by plasma treatment are present. The radish seeds were chosen to study the effect of plasma treatment on seed germination. To obtain the proper condition for the germination enhancement of radish seeds, plasma generation power and plasma irradiation time were optimized. The growth rate of seeds was determined by measuring the length of root and dry weight of the plants treated by plasma jet.

2. Experimental set-up

2.1. Design of plasma jet model

The schematic drawing of the experiment set up for radish seed germination enhancement by plasma treatment is illustrated in Fig. 1(a). The experiments were conducted in a room environment. Argon gas from a gas cylinder tank was controlled by a pressure gauge and fed through the plastic tube to the copper tube cathode. Plastic tube and diameters of 6.4 mm copper tube cathode were hooked together by the plastic cork on the stand and base. A 1.5 mm diameter copper rod was inserted into the center of copper tube cathode and was also hooked by a plastic cork as an anode. The copper rod tip was shorter than the end of copper tube by 1.75 mm, approximately. The 10x15x20 mm³ acrylic box as a plasma reactor was worn at the end of the copper tube cathode. A positive source voltage (V_s) of 0-30 kV was applied to the anode by a dc high voltage power source (Matsusada, AU-30P10) through 5 or 7 mega ohms ballast resistors. The characteristics of discharge voltage and current can be monitored by an oscilloscope (SIGLENT, SDS2304). A high voltage probe (Pintek HVP-28HF) was connected across two electrodes to observe the discharge voltage (V_d). The copper tube cathode was grounded through a 22 ohms monitored resistor to detect the discharge current waveform (I_d).

2.2. Plasma Treatment

In this research, the radish seeds (*Raphanus sativus*); which is one of the industrial crops of Thailand, were chosen to study the effect of plasma treatment on seed germination since the physical characteristics of the radish seed is small shaped and thin skinned. The radish seeds were obtained from the Ongkarak market. When the Argon gas was fed through the tube and the source voltage was supplied to the system sufficiently high enough, the plasma jet was generated between the tips of copper tube and ejected to the acrylic box. From the experimental results, it shows that 4 liters per minute (LPM) of Argon gas was most suitable for plasma generation for our system due to the lowest breakdown voltage and plasma irradiating area. Plasma generation power and plasma irradiation time were optimized to obtain the proper experimental condition for the germination enhancement of radish seeds. The plasma generation power was varied from 0 (the controlled condition), 90, and 140 watts. The 10 radish seeds per one experimental condition were exposed to plasma for 2, 4, and 6 minutes of plasma treating time in each plasma generation power. Plasma was generated at the source voltage (V_s) of 21.2, and 30 kV for 90, and 140 watts of plasma generation power, respectively. The characteristic waveforms of V_d and I_d at 90 watts of plasma generation power are present in Fig. 1(b). It should be mentioned here that in Fig. 1(b), due to the limitation of the high voltage probe, only the waveforms of 90 watts of plasma generation power are present. The discharge waveforms were in self-pulsing discharge mode which was suitable for seed treatment. The peak discharge current pulses were ranging from 0.0072 - 0.186 amperes with a life time of 45 micro second, approximately. The self-pulsing frequency was increased by increasing of the source voltage. The explanation of this situation is in^{5,6}.

After the plasma treatment process, 6 groups of treated radish seeds and 1 controlled group were soaked in the water for 2 hours. Seeds were cultivated in the dark of environmental room condition. The growth rate of seeds was determined by the length of root and dry weight of the radish sprouts. The radish sprouts were dehydrated at 105 degree Celsius for 3 hours by a universal oven (MEMMERT UM 500) and weighed by an analytical balance (Mettler Toledo PL303 Milligram Balance) after measuring the 7 day radish sprout root length.




Fig. 1. (a) the experimental set up for plasma treatment; (b) discharge voltage and current waveforms of plasma generation at 90 watts during the self-pulsing interval.

3. Experimental results and discussion

3.1. Germination enhancement: root length

From the experimental result, it shows that all the radish seeds of every experimental case could be sprouted. The radish sprouts after 1 day of cultivation had an average root length of 1.5 millimeters, approximately. After 2 days of cultivation, the cotyledon could be observed. However, to determine the root length and compare it with other conditions in this state were difficult to notice; in addition, the roots of radish sprouts were very fragile to straighten out for proper measurement. Therefore, the length of radish sprout roots was measured after 7 days of cultivation.

After 7 days of cultivation, it could be observed clearly that the root length and stem length of plasma treatment cases were rather longer than that of the controlled group. However, for plasma treatment cases, the root length and stem length were not significantly different. Fig. 2(a) shows the image of radish cultivation after 140 watts of plasma treatment for 0, 2, 4 and 6 minutes. It could be noticed that the root and stem of the radish sprout were grown straight for the first 5 days after cultivation and, after that they begin to bend. This was because the stem substantially absorbs water and the basement of cultivation place was not suitable for holding the weighty sprouts. The average root length of a controlled case and plasma treatment cases were 4.5 and 7.5 cm, respectively. However, the length of the root and stem of a radish sprout was not strong enough to judge that plasma could improve the germination rate. Therefore, to confirm the growth enhancement effect from plasma treatment, it is necessary to analyze together with another process to determine the total mass of sprout.

3.2. Growth enhancement: weight

The radish sprouts were brought in to dehydrate after measuring the length of root of radish sprout. The radish sprouts were dehydrated at 105 degree Celsius for 3 hours by a MEMMERT UM 500 universal oven and weighed by a Mettler Toledo PL303 analytical balance. The comparative results of the total mass of dehydrated radish sprout from various experimental conditions are illustrated in Fig. 2(b). The total mass was calculated in percentages from the ratio of the weight of sprout between pre and post dehydration process. From the results, it could be observed that 4 cases out of 6 had a greater total mass compared to the controlled case. For the other 2 cases, they had a slightly lower total mass compared to the controlled case. This could be implied that plasma treatment has a positive effect on germination enhancement. The 140 watts plasma generation power case had slightly higher total mass compared to the 90 watts case. It could be clearly noticed that the total mass of 140 watts plasma treatments for every treatment time had a higher total mass compared to the controlled case of 9-12 percent, approximately. It seems that increasing the discharge time could enhance the germination rate as noticed in the 90 watts of plasma generation; however, increasing of discharge time for the higher generation of power could lessen the germination rate. This problem is possibly caused by the fact that an excessively high plasma power could heat and damage some parts of the seeds. For the detailed explanation of this reason, it will be discussed in the further study of this research topic, which this plasma jet model will be applied to. Even though the mechanism on how the plasma affected seed

germination enhancement is still unclear at this moment, one possible reason that plasma can enhance the growth rate is that the skin of the seed irradiated by plasma will be sputtered by plasma. The thin and soft skin allows the seed to absorb water and nutrition efficiently. Moreover, the radical species from plasma would cause a chemical reaction on the skin of seed^{1,2,3,4,7}.




Fig. 2. (a) radish cultivation after a 140 watts plasma treatment for 5 and 7 days, which the treating times are (A) 2, (B) 4, (C) 6, and (D) 0 minutes; (b) percentage of total mass of dehydrated radish sprout from various experimental conditions.

4. Summary

The preliminary result of an atmospheric non-thermal plasma jet for germination enhancement of radish seeds was presented. The plasma reactor was a simple design; connect a small acrylic box at the tip of the plasma jet model. From the experimental results, it could be confirmed that plasma treatment has a positive effect on improving the growth rate of radish seeds. At the experimental condition of 140 watts of power generation, the growth rate of plasma treated seeds were 9-12 percent of the dry radish weight, approximately higher than that of the un-treated seeds. The average root lengths of plasma treatment cases were also longer than that of the controlled case by 3 centimeters, approximately.

Acknowledgements

This work was supported by Srinakharinwirot University. Moreover, the author would like to express thanks to all students belonging to the Atmospheric Plasma Engineering Laboratory (APEL).

References

1. Kitazaki S, Koga K, Shiratani M, Hayashi N. Growth Enhancement of Radish Sprouts Induced by Low Pressure O₂ Radio Frequency Discharge Plasma Irradiation, *Jpn. J. Appl. Phys.*, 2012; **51**: 01AE01.
2. Dobrin D, Magureanu M, Mandache N B, Ionita M.-D. The effect of non-thermal plasma treatment on wheat germination and early growth, *Innov. food sci. & emerg. technol.*, 2015; **29**: 255–260.
3. Dubinov A, Lazarenko E, Selemir V, Effect of glow discharge air plasma on grain crops seed. *IEEE Trans. Plasma Sci.*, 2000; **28**: 180-183.
4. Ling L, Jiafeng J, Jiangang L, Minchong S, Xin H, Hanliang S, Yuanhua D. Effects of cold plasma treatment on seed germination and seedling growth of soybean. *Scientific reports*, 2014; **4**: 5859.
5. Matra K, Mizobuchi Y, Furuta H, Hatta A. Local sputter etching by micro plasma jet in SEM, *Vacuum*, 2013; **87**: 132-135.
6. Matra K, Furuta H, Hatta A, Current-Voltage Characteristics of DC Discharge in Micro Gas Jet Injected into Vacuum Environment, *J Phys Conf Ser.*, 2013; **441**: 012021
7. Jiang J, He X, Li L, Li J, Shao H, Xu Q, Dong Y. Effect of Cold Plasma Treatment on Seed Germination and Growth of Wheat, *Plasma Sci. Technol.*, 2014; **16**: 54-58.