



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: www.elsevier.com/locate/apjtm



Document heading doi: 10.1016/S1995-7645(14)60017-1

Tracheal relaxation of five medicinal plants used in Mexico for the treatment of several diseases

Amanda Sánchez–Recillas¹, Paul Mantecón–Reyes¹, Patricia Castillo–España², Rafael Villalobos–Molina³, Maximiliano Ibarra–Barajas³, Samuel Estrada–Soto^{1*}

¹Facultad de Farmacia, Universidad Autónoma del Estado de Morelos, Avenida Universidad 1001, Col. Chamilpa, Cuernavaca, Morelos 62209, México

²Centro de Investigación en Biotecnología, Universidad Autónoma del Estado de Morelos, Avenida Universidad 1001, Col. Chamilpa, Cuernavaca, Morelos 62209, México

³Unidad de Biomedicina, Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, Tlalnepantla, Estado de México 54090, México

ARTICLE INFO

Article history:

Received 13 November 2013

Received in revised form 20 January 2014

Accepted 15 February 2014

Available online 20 March 2014

Keywords:

Asthma

Mexican medicinal plants

Trachea

Relaxant effect

ABSTRACT

Objective: To assess the relaxant effect of several organic extracts obtained from *Agastache mexicana* (*A. mexicana*), *Cochlospermum vitifolium* (*C. vitifolium*), *Cordia morelosana* (*C. morelosana*), *Lepechinia caulescens* (*L. caulescens*) and *Talauma mexicana* (*T. mexicana*) used in Mexican traditional medicine for the treatment of several diseases. **Methods:** Extracts were obtained by maceration at room temperature using hexane, dichloromethane and methanol for each plant material. The organic extracts were evaluated *ex vivo* to determine their relaxant activity on the contractions induced by carbachol (cholinergic receptor agonist, 1 μ mol/L) in isolated rat tracheal rings. **Results:** A total of 15 extracts were evaluated (three for each species). All test samples showed significant relaxant effect, in a concentration–dependent manner, on the contractions induced by 1 μ mol/L carbachol, with exception of extracts from *C. morelosana*. Active extracts were less potent than theophylline [phosphodiesterase inhibitor, EC₅₀: (28.79 \pm 0.82) μ g/mL] that was used as positive control. Concentration–response curves revealed that the extracts with more significant effects were dichloromethanic extracts of *T. mexicana* [E_{max}: (103.03 \pm 3.32)% and EC₅₀: (159.39 \pm 3.72) μ g/mL] and *C. vitifolium* [E_{max}: (106.58 \pm 2.42)% and EC₅₀: (219.54 \pm 7.61) μ g/mL]. Finally, hexanic and dichloromethanic extracts from *A. mexicana* were fully effective but less potent than *T. mexicana* and *C. vitifolium*. **Conclusions:** Less polar extracts obtained from *A. mexicana*, *T. mexicana* and *C. vitifolium* exhibited greater relaxant effect on tracheal rat rings, which allows us to suggest them as sources for the isolation of bioactive molecules with potential therapeutic value in the treatment of asthma.

1. Introduction

Asthma is a chronic inflammatory disease of the lung characterized by episodic airway obstruction and increased bronchial responsiveness[1]; the structural features include

epithelial shedding, increased airway smooth muscle mass, mucus gland hyperplasia, sub epithelial fibrosis and infiltration of the bronchial wall with inflammatory cells[2]. Bronchial obstruction is the most important symptom because of the potential risk of severe asthma attacks; however, other non–obstructive symptoms are frequent and may also cause severe suffering and poor quality of life[3]. The current treatment does not resolve the disease and only focuses on countering episodes of bronchospasm and controlling the underlying inflammation[4]. Together, the high level tolerance and resistance of asthmatic patients, as a result of frequent drug use, increase the interest and

*Corresponding author: Dr. S. Estrada Soto, Facultad de Farmacia, Avenida Universidad 1001, Col. Chamilpa, Cuernavaca, Morelos 62209, México.

Tel/fax: +52 777 329 7089

E-mail: enoch@uaem.mx

Foundation project: This study was financed by a grant from “Apoyo a la Mejora del Perfil Individual del profesorado de tiempo completo (Fondo para la Consolidación de las Universidades Públicas Estatales y con Apoyo Solidario Ejercicio 2009)”, Faculty of Pharmacy budgets (2010 and 2011), and grants from CONACyT–CIENCIA BÁSICA CB–2011–01 (167044).

need to develop new therapeutic strategies with specific, unexplored targets acting on the mechanisms that generate the asthmatic episode.

Mexico has over 4 000 species of medicinal plants, that have been used empirically by 80% of the indigenous population to treat health problems like respiratory diseases[5]. Current research was carried out in order to screen the pharmacological properties of *Agastache mexicana* (*A. mexicana*), *Cochlospermum vitifolium* (*C. vitifolium*), *Cordia morelosana* (*C. morelosana*), *Lepechinia caulescens* (*L. caulescens*) and *Talauma mexicana* (*T. mexicana*) to find candidates for further bio-guided studies, which allows the discovery of novel molecules with therapeutic properties for treating asthma, in particular, those which involve a contractile process of the smooth muscle cells.

2. Materials and methods

2.1. Chemicals and drugs

Carbamylcholine (carbachol), teophylline, dimethylsulfoxide (DMSO) and ethyl ether were purchased from Sigma–Aldrich Co. (St. Louis, MO, USA). All other reagents were of analytical grade from local sources. Stock solutions of extracts were made using distilled water and DMSO and freshly prepared at the same day of experimentation.

2.2. Plant material

Plant species were selected using an ethnomedical criterion[6] and were collected and identified by Dr. Patricia Castillo España of the Center for Research in Biotechnology, UAEM. A sample of each specimen was deposited in the HUMO–herbarium of the Center for Research and Environmental Studies “Sierra de Huautla”(CEAMISH) of Morelos State University. Details of the collection, traditional uses, and voucher number of each species, among others, are mentioned in Table 1.

2.3. Extraction of plant material

The air-dried plant material was ground into powder. Crude extracts were prepared by successive room temperature–maceration of 30 g of dried plant material in 600 mL of hexane, dichloromethane and methanol for 72 h, respectively. After filtration, extracts were concentrated in vacuo at 45 °C. Finally, a yield was obtained for each extract.

To carry out the experiments, all extracts were dissolved in a mixture of dimethylsulfoxide:water (50:50, v:v).

2.4. Pharmacological evaluations

2.4.1. Rats

Healthy male Wistar rats (250–300 g) were used and maintained under standard laboratory conditions with free access to food and water. All animal procedures were conducted in accordance with our Federal Regulations for Animal Experimentation and Care (SAGARPA, NOM–062–ZOO–1999, Mexico) and approved by the Institutional Animal Care and Use Committee based on US National Institute of Health publication (No. 85–23, revised 1985). All experiments were carried out using six animals per group. Animals used were euthanized by cervical dislocation.

2.4.2. Rat trachea ring test

Previous protocol was used[7]. The trachea was dissected, cleaned out of connective tissue and mucus, and immediately cut into 4–5 mm long rings. Then, tissue segments were mounted in stainless steel hooks, under optimal tension of 2 g, in 10 mL organ baths containing warmed (37 °C) and oxygenated (O₂/CO₂, 95:5, v:v) Krebs solution (118.0 mmol/L NaCl, 4.7 mmol/L KCl, 2.5 mmol/L CaCl₂, 1.2 mmol/L MgSO₄, 1.2 mmol/L KH₂PO₄, 25.0 mmol/L NaHCO₃, 0.026 mmol/L EDTA, and 11.1 mmol/L glucose, pH 7.4). Changes in tension were recorded by Grass–FT03 force transducers (Astromed, West Warwick, RI, USA) connected to a MP100 analyzer (BIOPAC Instruments, Santa Barbara, CA, USA) as previously described[7]. After equilibration, rings were contracted by 1 μmol/L carbachol and washed every 40 min for 2 h. After contraction with carbachol, the test samples (organic extracts, vehicle and positive control) were added to the bath in a volume of 100 μL; then cumulative concentration–response curves were obtained for each ring (3.03–1 000.00 μg/mL). The relaxant effect of extracts and positive control (theophylline, an inhibitor of phosphodiesterase, 1.67–550.00 μmol/L) were determined by comparing the muscular tone of the contraction before and after addition of the test materials. Muscular tone was calculated from the tracings, using Acknowledge software (BIOPAC®).

2.5. Data analysis

The experimental results are expressed as mean of six experiments ± standard error of mean (SEM). Concentration–response curves were plotted and experimental data of

the concentration–response curves were adjusted by the nonlinear, curve–fitting program Microcal™ origin 8.0 (Microcal Software Inc., USA). Significance was evaluated using the ANOVA [8]. Values of $P < 0.05$ imply significance of the pharmacological effects in the experiments.

3. Results

The plant material was subjected to extraction process by maceration, and details of their yields are shown in Table 2. A total of 15 extracts were evaluated (three for each species). All test samples showed significant relaxant effect

in a concentration–dependent manner on the contractions induced by $1 \mu\text{mol/L}$ carbachol, with exception of extracts from *C. morelosana* (Table 2). Active extracts were less potent than theophylline [phosphodiesterase inhibitor, EC_{50} : $(28.79 \pm 0.82) \mu\text{g/mL}$] used as positive control. Concentration–response curves revealed that the extracts with more significant effects were dichloromethanic extracts of *T. mexicana* [E_{max} : $(103.03 \pm 3.32)\%$ and EC_{50} : $(159.39 \pm 3.72) \mu\text{g/mL}$] and *C. vitifolium* [E_{max} : $(106.58 \pm 2.42)\%$; EC_{50} : $(219.54 \pm 7.61) \mu\text{g/mL}$] (Figure 1). Finally, hexanic and dichloromethanic extracts from *A. mexicana* were fully effective but less potent than *T. mexicana* and *C. vitifolium* (Figure 2).

Table 1

Detailed information about plant species screened.

Plant name	Family	Common name	Voucher number	Traditional uses	Place of collection
<i>A. mexicana</i> subsp. xolocotziiana Bye, Lin & Ram.	Lamiaceae	Toronjil	26336	Anxiety, stress, antihypertensive, heart disease, antispasmodic, stomachache, vomit, kidney inflammation, asthma	Felipe Neri, Morelos State
<i>C. vitifolium</i> (Willd)	Bixaceae	Pánicua	14628	Antihypertensive, diabetes, hepatobiliary disorders	Sierra de Huatla, Morelos State
<i>C. morelosana</i> Standley	Boraginaceae	Anacahuite	Pending	Respiratory diseases	Amatlán, Morelos State
<i>L. caulescens</i> (Ortega) Epling	Lamiaceae	Bretónica, Brenilla	20386	Kidney diseases, diarrhea, vomit, diabetes	Amatlán, Morelos State
<i>T. mexicana</i> (DC.) G. Don	Magnoliaceae	Yolloxóchitl	1986–012 A	Anxiety, sleeplessness, heart disease, stress	Jalapa, Veracruz




Figure 1. Concentration–response curves of the relaxant effect of organic extracts on trachea rings pre–contracted with $1 \mu\text{mol/L}$ carbachol. All results are expressed as the mean \pm SEM of six experiments ($P < 0.05$).

Table 2Extraction yield and relaxant effect of plant extracts on rat tracheal rings pre-contracted with carbachol 1 μ mol/L.

Plant name	Extract	Extraction yield (%)	E _{max} (%)	EC ₅₀ (μ g/mL)
Positive control	Theophylline		106.85±0.89	28.79±0.82
<i>A. mexicana</i>	Hexane	1.68	100.16±1.71	219.00±7.40
	Dichloromethane	2.15	97.78±3.34	320.87±5.72
	Methanol	6.38	75.54±6.03	644.44±4.53
<i>C. vitifolium</i>	Hexane	0.45	72.71±6.63	507.14±5.30
	Dichloromethane	0.49	106.60±2.42	152.83±7.61
	Methanol	1.23	61.91±3.03	631.52±3.42
<i>C. morelosana</i>	Hexane	2.93	42.46±5.56	ND
	Dichloromethane	1.89	35.80±1.54	ND
	Methanol	9.59	29.07±1.71	ND
<i>L. caulescens</i>	Hexane	0.81	85.51±1.42	516.69±3.72
	Dichloromethane	0.70	59.81±4.39	801.23±4.53
	Methanol	2.51	51.67±5.03	ND
<i>T. mexicana</i>	Hexane	4.00	97.45±1.04	477.28±5.19
	Dichloromethane	6.50	103.03±3.32	159.39±8.7
	Methanol	7.00	91.77±5.19	430.71±5.04

E_{max} and EC₅₀ indicate efficacy and potency of the extract, respectively. ND: Not determined.




Figure 2. Concentration–response curves of the relaxant effect of the most potent organic extracts on trachea rings pre-contracted with 1 μ mol/L carbachol.

All results are expressed as the mean±SEM of six experiments ($P<0.05$). HE: Hexane extract; DE: Dichloromethane extract; ME: Methanol extract.

4. Discussion

Medicinal plants *A. mexicana*, *C. vitifolium*, *C. morelosana*, *L. caulescens* and *T. mexicana* are used in Mexican traditional medicine for the treatment of several diseases[6]. These plants were selected based on ethnomedical criterion[6] and previous reports related with relaxation of several smooth muscle tissues[9–15], with exception of *C. morelosana* which is used as an anti-asthmatic agent and related respiratory diseases in traditional medicine. Current strategies for anti-asthmatic drug development include anti-inflammatory and/or bronchodilator drugs. Thus, we decided to explore the direct airway relaxant effects of

hexanic, dichloromethanic and methanolic extracts from these selected plants using carbachol–precontracted rat tracheal rings, an emergent model related to the widely used guinea-pig tracheal model[16].

In this framework, the extracts were obtained by maceration at room temperature and, as we expected, the methanolic extracts were those that showed higher yields in the extraction process for all plant species, because the metabolic content of medicinal plants are mainly compounds with medium to high polarity[17]. All extracts evaluated showed significant relaxant effect in the range of concentrations used, with exception of extracts of *C. morelosana*. However, we cannot discard the anti-asthmatic properties of *C. morelosana*, since this species may act through modulating the inflammatory process in asthmatic episodes[18], modulating airway hyperresponsiveness and hypersensitivity development as a result of asthmatics' status, or inhibiting inflammatory mediator release[19]. It is important to mention that *C. morelosana* is a plant species that has been traditionally used for the treatment of respiratory diseases and/or asthma, and it is necessary to accomplish other pharmacological tests to confirm its traditional use in the population, such as the evaluation of the asthmatic effect in mice sensitized with ovalbumin[4]. On the other hand, extracts from *L. caulescens* showed moderated relaxant effect being hexane extract of *L. caulescens* the most active extract from this plant. Last suggest that previous isolated low polarity bioactive compounds from *L. caulescens* may be responsible for the tracheal relaxant effect such as ursolic and oleanolic acids, spathulenol, 9 α ,13 α -epidioxyabiet-8(14)-en-18-oic acid methyl ester, dehydroabietic acid, and 9 β -hydroxydehydroabietyl alcohol which showed relaxant effect on various smooth muscle tissue[9,20]. On the other hand, *C. vitifolium*, *A. mexicana* and *T. mexicana* are used in traditional Mexican medicine for the treatment of cardiovascular diseases. These species were the most

active tracheal relaxant agents, with the dichloromethanic extracts from *C. vitifolium* and *T. mexicana* being the most active samples, following by the dichloromethanic and hexanic extracts from *A. mexicana*. Previous reports showed that *C. vitifolium* and *T. mexicana* have significant vasorelaxant effect on partially endothelium-dependent and concentration-dependent manners on aorta rat rings. Flavonoids and triterpenic derivatives were found to be the responsible bioactive agents, with NO/cGMP system involved as main vasorelaxant mode of action^[12–15]. Moreover, there is an evidence demonstrating that flavonoids have relaxing properties over smooth muscle of the rat and guinea pig trachea^[21,22]. Finally, *T. mexicana* has been used since the 15th century for the treatment of cardiovascular diseases, but this species has few pharmacological studies that support the traditional uses of the plant^[23]. Further phytochemical experiments are necessary to isolate the responsible of the relaxant effect on tracheal rat rings.

In conclusion, we have shown that various extracts of *T. mexicana*, *C. vitifolium* and *A. mexicana* induced a relaxant effect on rat-isolated trachea, which may propose them as potential sources for the isolation of anti-asthmatic agent(s).

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This study was financed by a grant from “Apoyo a la Mejora del Perfil Individual del profesorado de tiempo completo (Fondo para la Consolidación de las Universidades Públicas Estatales y con Apoyo Solidario Ejercicio 2009)”, Faculty of Pharmacy budgets (2010 and 2011) and grants from CONACYT-CIENCIA BÁSICA CB-2011-01 (167044). This article was taken in part from the Master in Pharmacy thesis of A. Sánchez-Recillas.

References

- [1] Qutayba H. Pathogenesis of small airways in asthma. *Respiration* 2012; **84**: 4–11.
- [2] Dunnill MS. The pathology of asthma, with special reference to changes in the bronchial mucosa. *J Clin Pathol* 1960; **13**: 27–33.
- [3] Löwhagen O. Diagnosis of asthma: A new approach. *Allergy* 2012; **67**: 713–717.
- [4] Torres R, Picado C, Mora F. Discovering asthma origin through the mouse. A review to the pathogenesis of mice models of asthma and their similarity with human asthma. *Arch Bronconeumol* 2005; **41**(3): 141–152.
- [5] Waizel S, Waizel J. Algunas plantas utilizadas en México para el tratamiento del asma. *An ORL Méx* 2009; **54**(4): 145–171.
- [6] Castillo-España P, Monroy-Ortiz C. *Plantas Medicinales Utilizadas en el Estado de Morelos*. 2nd ed. Universidad Autónoma del Estado de Morelos; 2007.
- [7] Estrada-Soto S, Sánchez-Recillas A, Navarrete-Vázquez JG, Castillo-España P, Villalobos-Molina R, Ibarra-Barajas M. Relaxant effects of *Artemisia ludoviciana* on isolated rat smooth muscle tissues. *J Ethnopharmacol* 2012; **139**: 513–518.
- [8] Woodson RF. Statistical methods for the analysis of biomedical data. In: *Wiley series in probability and mathematical statistics*. New York: Wiley; 1987, p. 315–316.
- [9] Aguirre-Crespo F, Vergara-Galicia J, Villalobos-Molina, López-Guerrero JJ, Navarrete-Vázquez G, Estrada-Soto S. Ursolic acid mediates the vasorelaxant activity of *Lepechinia caulescens* via NO release in isolated rat thoracic aorta. *Life Sci* 2006; **79**: 1062–1068.
- [10] Estrada-Soto S, Navarrete-Vázquez G, León-Rivera I, Rios MY, Aguilar-Guadarrama B, Castillo-España P, et al. Antihypertensive effect of *Lepechinia caulescens* extract on spontaneously hypertensive rats. *Phytopharmacol* 2012; **2**: 170–178.
- [11] Estrada-Soto S, Rodríguez-Avilez A, Castañeda-Ávila C, Castillo-España P, Hernández de Jesús MLP, Villalobos-Molina R, et al. Spasmolytic action of *Lepechinia caulescens* is through calcium channel blockade and NO release. *J Ethnopharmacol* 2007; **114**: 364–370.
- [12] Sánchez-Salgado JC, Castillo-España P, Ibarra-Barajas M, Villalobos-Molina R, Estrada-Soto S. *Cochlospermum vitifolium* induces vasorelaxant and antihypertensive effects mainly by activation of NO/cGMP signaling pathway. *J Ethnopharmacol* 2010; **130**: 477–484.
- [13] Sánchez-Salgado JC, Ortiz-Andrade R, Aguirre-Crespo F, Vergara-Galicia J, León-Rivera I, Montes S, et al. Hypoglycemic, vasorelaxant and hepatoprotective effects of *Cochlospermum vitifolium* (willd.) Sprengel: a potential agent for the treatment of metabolic syndrome. *J Ethnopharmacol* 2007; **109**: 161–164.
- [14] Hernández-Abreu O, Castillo-España P, León-Rivera I, Ibarra-Barajas M, Villalobos-Molina R, González-Christen J, et al. Antihypertensive and vasorelaxant effects of tilianin isolated from *Agastache mexicana* are mediated by NO/cGMP pathway and potassium channel opening. *Biochem Pharmacol* 2009; **78**: 54–61.
- [15] González-Trujano ME, Ventura-Martínez R, Chávez M, Díaz-Reval I, Pellicer F. Spasmolytic and antinociceptive activities of ursolic acid and acacetin identified in *Agastache mexicana*. *Planta Med* 2012; **78**: 793–796.
- [16] Irié-N’guessan G, Champy P, Kouakou-Siransy G, Koffi A, Kablan BJ, Leblais V. Tracheal relaxation of five Ivorian anti-asthmatic plants: Role of epithelium and K⁺ channels in the effect of the aqueous-alcoholic extract of *Dichrostachys cinerea* root bark. *J Ethnopharmacol* 2011; **138**: 432–438.
- [17] Jones WP, Kinghorn AD. Extraction of plant secondary metabolites. *Methods Mol Biol* 2012; **864**: 341–366.
- [18] Ramos-Barbón D. Asthma from biology to clinic. *Arch Bronconeumol* 2007; **43**(Supl 2): 3–14.
- [19] Ganguly T, Sainis KB. Inhibition of cellular immune responses by *Tylophora indica* in experimental models. *Phytotherapy* 2001; **8**(5): 348–355.
- [20] Pérez-Hernández N, Ponce-Monter H, Medina JA, Joseph-Nathan P. Spasmolytic effect of constituents from *Lepechinia caulescens* on rat uterus. *J Ethnopharmacol* 2008; **115**: 30–35.
- [21] Ko WC, Liu PY, Chen JL, Leu IJ, Shih CM. Relaxant effects of flavonoids in isolated guinea pig trachea and their structure-activity relationships. *Planta Med* 2003; **69**(12): 1086–1090.
- [22] Chiwororo WD, Ojewole JA. Dual effect of quercetin on rat isolated portal vein smooth muscle contractility. *Cardiovasc J Afr* 2010; **21**(3): 132–136.
- [23] Waizel-Bucay J. Uso tradicional e investigación científica de *Talauma mexicana* (D.C.) Don., o flor del corazón. *Rev Mex Cardiol* 2002; **13**(1): 31–38.