

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

J. Math. Anal. Appl. 296 (2004) 309–313

Journal of
 MATHEMATICAL
 ANALYSIS AND
 APPLICATIONS

www.elsevier.com/locate/jmaa

Integral representation of a series which includes the Mathieu **a**-series

Tibor K. Pogány

Department of Sciences, Faculty of Maritime Studies, University of Rijeka, Studentska 2, 51000 Rijeka, Croatia

Received 25 February 2004

Available online 10 June 2004

Submitted by B.C. Berndt

Abstract

Integral expression is deduced for the series

$$\mathfrak{S}(r, \mu, \nu, \mathbf{a}) = \sum_{n=1}^{\infty} \frac{{}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2} + 1; \nu + 1; -\frac{r^2}{a(n)^2}\right)}{a(n)^{\nu-\mu+1}(a(n)^2 + r^2)^{\mu-1/2}},$$

where $r > 0$, $\mu > 1/2$, $\nu + 1 < \mu$ and $\mathbf{a}: 0 < a(1) < a(2) < \dots < a(n) \uparrow \infty$, and ${}_2F_1$ is the Gauß hypergeometric function. The result precizes the integral expression for the generalized Qi type Mathieu **a**-series $S(r, p, \mathbf{a}) = \sum_{n=0}^{\infty} a(n)(a(n)^2 + r^2)^{-p-1}$ given in [J. Inequal. Pure Appl. Math. 4 (2003), (4.5)] generalizing some other results by Cerone and Lenard, Tomovski and Qi as well. Bounding inequalities are given for $\mathfrak{S}(r, \mu, \nu, \mathbf{a})$ using the derived integral expression.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Bessel function of first kind; Dirichlet series; Generalized Mathieu series; Gauß hypergeometric function; Laplace integral; Mathieu **a**-series

1. Introduction

In this article we consider the series

$$\mathfrak{S}(r, \mu, \nu, \mathbf{a}) = \sum_{n=1}^{\infty} \frac{{}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2} + 1; \nu + 1; -\frac{r^2}{a(n)^2}\right)}{a(n)^{\nu-\mu+1}(a(n)^2 + r^2)^{\mu-1/2}}, \quad (1)$$

E-mail address: poganj@brod.pfri.hr.

where ${}_2F_1$ denotes the Gauß hypergeometric function and $(r, \mu - 1/2) \in \mathbb{R}_+^2$, $\nu < \mu - 1$, while the real sequence \mathbf{a} : $0 < a(1) < a(2) < \dots < a(n) \uparrow \infty$. The special case of (1) is the Qi type Mathieu \mathbf{a} -series

$$S(r, p, \mathbf{a}) = \sum_{n=1}^{\infty} \frac{a(n)}{(a(n)^2 + r^2)^{p+1}}, \quad r, p + 1 > 0, \quad (2)$$

treated differently by the author in [5]. Indeed, it is $\mathfrak{S}(r, p + 3/2, p - 1/2, \mathbf{a}) = S(r, p, \mathbf{a})$, since ${}_2F_1(-1/2, 0; p + 1/2; -(r/a(n))^2) \equiv 1$.

In this article we give integral expression for $\mathfrak{S}(r, \mu, \nu, \mathbf{a})$. Our main tools are the closed (integral) form expression for a Dirichlet's series $\sum_{n=1}^{\infty} \exp\{-a(n)x\}$, $x > 0$, and \mathbf{a} is positive, monotonous increasing divergent with $\lim_{n \rightarrow \infty} a(n) = \infty$. The sources for this are the suitable chapters of classical books [3,4].

In the sequel $[u]$ denotes the integral part of u ; $\mathcal{L}_s\{f\}$, $\mathcal{M}_z\{f\}$ stand for Laplace and Mellin transform of f and $J_\nu(x)$ is the ν th order Bessel function of first kind. Finally, $\mathbb{1}_B(t)$ denotes the characteristic function of the set B .

2. Integral representation of $\mathfrak{S}(r, \mu, \nu, \mathbf{a})$

Consider the formula

$$\int_0^{\infty} e^{-\alpha x} x^{\mu-1} J_\nu(\beta x) dx = \frac{\beta^\nu \Gamma(\nu + \mu) {}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2} + 1; \nu + 1; -\frac{\beta^2}{\alpha^2}\right)}{2^\nu \alpha^{\nu-\mu+1} \Gamma(\nu + 1) (\alpha^2 + \beta^2)^{\mu-1/2}}, \quad (3)$$

valid for $\alpha, \beta, \nu + 1, \mu - 1/2 > 0$, such that is listed (in equivalent form) as [2, 6.621 W421(3)], then specify $\alpha \equiv a(n)$, $\beta \equiv r$. Finally, the summation of (3) with respect to $n \in \mathbb{N}$ results in

$$\begin{aligned} & \int_0^{\infty} \left(\sum_{n=1}^{\infty} e^{-a(n)x} \right) x^{\mu-1} J_\nu(rx) dx \\ &= \left(\frac{r}{2} \right)^\nu \frac{\Gamma(\nu + \mu)}{\Gamma(\nu + 1)} \sum_{n=1}^{\infty} \frac{{}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2} + 1; \nu + 1; -\frac{r^2}{a(n)^2}\right)}{a(n)^{\nu-\mu+1} (a(n)^2 + r^2)^{\mu-1/2}}. \end{aligned} \quad (4)$$

Denoting the series in (4) by $\mathfrak{S}(r, \mu, \nu, \mathbf{a})$ we finish the first part of our exposition.

The Dirichlet series in the integrand of (4) possesses Laplace integral form

$$\sum_{n=1}^{\infty} e^{-a(n)x} = x \int_0^{\infty} e^{-xt} A(t) dt, \quad (5)$$

where the so-called *counting function* $A(t)$ is given as

$$A(t) = \sum_{a(n) \leq t} 1 = [a^{-1}(t)], \quad (6)$$

see [3, IV], [4, Part C, I.1, V.5] or [5, §1, 2]. Here a^{-1} denotes the inverse of the function $a(x)$, $x \in \mathbb{R}_+$, whose restriction \mathbf{a} is mentioned in $\mathfrak{S}(r, \mu, \nu, \mathbf{a})$. Therefore, replacing (6) into (5) we deduce the following result.

Theorem 1. *Let $r, \mu > 1/2$, $\nu < \mu - 1$ and let \mathbf{a} be the restriction of the monotonous increasing function $a : \mathbb{R}_+ \mapsto \mathbb{R}_+$, $a(0) = 0$ to \mathbb{N}_0 , such that*

$$0 < a(1) < a(2) < \dots < a(n) \uparrow \infty \quad \text{as } n \rightarrow \infty.$$

Then it holds true

$$\mathfrak{S}(r, \mu, \nu, \mathbf{a}) = \left(\frac{2}{r}\right)^\nu \frac{\Gamma(\nu + 1)}{\Gamma(\nu + \mu)} \int_0^\infty \int_{a(1)}^\infty e^{-xt} x^\mu J_\nu(rx) [a^{-1}(t)] dx dt. \tag{7}$$

Proof. We remark that a has the unique inverse being a^{-1} monotonous. So, by the same reason $[a^{-1}(t)] = 0$ for all $a(0) = 0 \leq a^{-1}(t) < 1$, i.e., for $0 \leq t < a(1)$. \square

Bearing on mind the Laplace and Mellin transform notations, the integral representation (7) becomes

$$\mathfrak{S}(r, \mu, \nu, \mathbf{a}) = \left(\frac{2}{r}\right)^\nu \frac{\Gamma(\nu + 1)}{\Gamma(\nu + \mu)} \mathcal{M}_{\mu+1} \{ \mathcal{L}_x \{ [a^{-1}(t)] \} J_\nu(rx) \}. \tag{8}$$

We will consider the case $\mathfrak{S}(r, p + 3/2, p - 1/2, \mathbf{a})$ separately because it represents the Qi type Mathieu \mathbf{a} -series $S(r, p, \mathbf{a})$. We remark that S is discussed from different points of view in [5–7].

Corollary 2.1. *For all $r > 0$, $p + 1 > 0$ and \mathbf{a} being the same as in Theorem 1, we have*

$$S(r, p, \mathbf{a}) = \frac{2\sqrt{\pi}}{(2r)^{p-1/2} \Gamma(p + 1)} \int_0^\infty \int_{a(1)}^\infty e^{-xt} x^{p+3/2} J_{p-1/2}(rx) [a^{-1}(t)] dx dt. \tag{9}$$

Proof. Having on mind the earlier connection between \mathfrak{S} and S , an application of the duplication identity for the Γ -function finishes the proof of (9). \square

It is not hard to see that (9) is the desired closed integral expression for the relation [1, (4.5)] by Cerone and Lenard. In the same time

$$F(\mathbf{a}) = \sum_{n=1}^\infty e^{-a(n)x} = x \int_{a(1)}^\infty e^{-xt} [a^{-1}(t)] dt, \quad x > 0, \tag{5&6}$$

is an answer to the question of closed form expression for the series $F(\mathbf{a})$, posed as open problem in the same article, for monotonous \mathbf{a} .

Specified values of the parameters r, μ, ν and assumptions upon the behavior of the sequence \mathbf{a} cover certain questions, posed as open problems by Feng Qi, and by Tomovski, compare [6–8].

Theorem 2. Let the situation be the same as in Theorem 1. Then we have the sharp inequality

$$0 \leq \left(\frac{2}{r}\right)^{\nu} \frac{\Gamma(\nu+1)}{\Gamma(\nu+\mu)} M(r, \mu, \mathbf{a}) - \mathfrak{S}(r, \mu, \nu, \mathbf{a}) < \frac{{}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2}+1; \nu+1; -\frac{r^2}{a(1)^2}\right)}{a(1)^{\nu-\mu+1}(a(1)^2+r^2)^{\mu-1/2}}, \quad (10)$$

where

$$M(r, \mu, \mathbf{a}) = \mathcal{M}_{\mu+1} \left\{ \mathcal{L}_x \left\{ a^{-1}(t) \mathbb{I}_{[a(1), \infty)}(t) \right\} J_{\nu}(rx) \right\}.$$

Proof. Applying the well-known (sharp) bilateral inequality

$$a^{-1}(t) - 1 < [a^{-1}(t)] \leq a^{-1}(t), \quad t \in \mathbb{R}_+,$$

to the integrand in (7) we clearly deduce the sharp bilateral bounding inequality which lower bound is equal to

$$\begin{aligned} & \left(\frac{2}{r}\right)^{\nu} \frac{\Gamma(\nu+1)}{\Gamma(\nu+\mu)} \int_0^{\infty} x^{\mu} J_{\nu}(rx) dx \int_0^{\infty} e^{-xt} (a^{-1}(t) - 1) \mathbb{I}_{[a(1), \infty)}(t) dt \\ &= \left(\frac{2}{r}\right)^{\nu} \frac{\Gamma(\nu+1)}{\Gamma(\nu+\mu)} \left(M(r, \mu, \mathbf{a}) - \int_0^{\infty} e^{-a(1)x} x^{\mu-1} J_{\nu}(rx) dx \right). \end{aligned} \quad (11)$$

The integral in (11) we calculate putting $\alpha = a(1)$, $\beta = r$ into (3). So, (11) takes the value

$$\left(\frac{2}{r}\right)^{\nu} \frac{\Gamma(\nu+1)}{\Gamma(\nu+\mu)} M(r, \mu, \mathbf{a}) - \frac{{}_2F_1\left(\frac{\nu-\mu+1}{2}, \frac{\nu-\mu}{2}+1; \nu+1; -\frac{r^2}{a(1)^2}\right)}{a(1)^{\nu-\mu+1}(a(1)^2+r^2)^{\mu-1/2}}. \quad (12)$$

The upper bound we get by the same procedure, and it is equal to the first addend in (12). Now, collecting these estimates easy calculations lead to the asserted bilateral bounding inequality (10). \square

Finally, it could be mentioned that this results enable to give bilateral bounding inequalities for Qi type Mathieu \mathbf{a} -series by using suitable values for parameters μ and ν in (10). However, this, and the caused calculations we leave to the interested reader.

References

- [1] P. Cerone, C.T. Lenard, On integral forms of generalized Mathieu series, J. Inequal. Pure Appl. Math. 4 (2003), available online at http://jipam.vu.edu.au/v4n5/100_01.html.
- [2] I.S. Gradshteyn, I.M. Ryzhik, Table of Integrals, Series and Products, supervised by Yu.V. Geronimus, M. Yu. Tseitlin, fourth ed., Fizmatgiz, Moscow, 1963 (in Russian).
- [3] G.H. Hardy, M. Riesz, The General Theory of Dirichlet's Series, Cambridge Univ. Press, Cambridge, 1915.
- [4] J. Karamata, Theory and Praxis of Stieltjes Integral, Srpska Akademija Nauka, Posebna izdanja CLIV, Knjiga I, Matematički Institut, Beograd, 1949 (in Serbian).

- [5] T.K. Pogány, Integral representation of Mathieu (a, λ) -series, RGMIA Research Report Collection 7 (2004), available online at <http://rgmia.vu.edu.au/v7n1.html>.
- [6] F. Qi, Inequalities for Mathieu series, RGMIA Research Report Collection 4 (2001) 187–193, available online at <http://rgmia.vu.edu.au/v4n2.html>.
- [7] F. Qi, Integral expression and inequalities of Mathieu type series, RGMIA Research Report Collection 6 (2003), available online at <http://rgmia.vu.edu.au/v6n2.html>.
- [8] Ž. Tomovski, New double inequalities for Mathieu type series, RGMIA Research Report Collection 6 (2003), available online at <http://rgmia.vu.edu.au/v6n2.html>.