



King Saud University

Saudi Journal of Biological Sciences

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Mitochondria from rat uterine smooth muscle possess ATP-sensitive potassium channel

Olga B. Vadzyuk^{*}, Serhiy O. Kosterin¹

Department of Muscles Biochemistry, Palladin Institute of Biochemistry of the National Academy of Sciences of Ukraine, Leontovych str. 9, 01601 Kyiv, Ukraine

Received 16 September 2015; revised 10 December 2015; accepted 31 January 2016

KEYWORDS

ATP-sensitive potassium channel;
Diazoxide;
5-hydroxydecanoate;
Myometrium;
Mitochondria;
Mitochondrial swelling;
Mitochondrial membrane potential;
ROS

Abstract The objective of this study was to detect ATP-sensitive K^+ uptake in rat uterine smooth muscle mitochondria and to determine possible effects of its activation on mitochondrial physiology. By means of fluorescent technique with usage of K^+ -sensitive fluorescent probe PBF1 (potassium-binding benzofuran isophthalate) we showed that accumulation of K^+ ions in isolated mitochondria from rat myometrium is sensitive to effectors of K_{ATP} -channel (ATP-sensitive K^+ -channel) – ATP, diazoxide, glibenclamide and 5HD (5-hydroxydecanoate). Our data demonstrates that K^+ uptake in isolated myometrium mitochondria results in a slight decrease in membrane potential, enhancement of generation of ROS (reactive oxygen species) and mitochondrial swelling. Particularly, the addition of ATP into incubation medium led to a decrease in mitochondrial swelling and ROS production, and an increase in membrane potential. These effects were eliminated by diazoxide. If blockers of K_{ATP} -channel were added along with diazoxide, the effects of diazoxide were removed. So, we postulate the existence of K_{ATP} -channels in rat uterus mitochondria and assume that their functioning may regulate physiological conditions of mitochondria, such as matrix volume, ROS generation and polarization of mitochondrial membrane.

© 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The mitochondria are involved in a variety of cellular metabolic functions, ranging from ATP supply to triggering of

apoptosis (Galluzi et al., 2012). Thus, the proper mitochondrial functioning is essential for cellular physiology. K^+ -transporting channels, which were discovered in mitochondrial inner membrane in last decades, exert influence not only on mitochondrial physiology, but also on cell functioning both under normal and stress conditions (Costa et al., 2006; Wojtovich and Brookes, 2008; Garg and Hu, 2007). The ATP-sensitive mitochondrial K^+ -channels (mito K_{ATP}) are the most studied among the types of mitochondrial K^+ -channels. The mito K_{ATP} channels are presumed to regulate different characteristics of mitochondria, such as matrix volume, respiration rate, etc., and mediate cell survival under stress conditions (Costa et al., 2006; Wojtovich and Brookes,

^{*} Corresponding author. Tel./fax: +380 44 2796365.

E-mail address: olga_vadzyuk@hotmail.com (O.B. Vadzyuk).

¹ Tel./fax: +380 44 2796365.

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<http://dx.doi.org/10.1016/j.sjbs.2016.01.045>

1319-562X © 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Vadzyuk, O.B., Kosterin, S.O. Mitochondria from rat uterine smooth muscle possess ATP-sensitive potassium channel. Saudi Journal of Biological Sciences (2016), <http://dx.doi.org/10.1016/j.sjbs.2016.01.045>

2008; Garg and Hu, 2007; Garlid and Halestrap, 2012; Foster et al., 2012; Hu et al., 2009; Flagg et al., 2010). In spite of the absence of accurate data regarding the subunits composition and their arrangement in membrane, the functioning of mitoK_{ATP} has been studied using selective effectors of plasma membrane K_{ATP}-channels. The mitoK_{ATP} channels were described in many tissues by different techniques, both on isolated mitochondria and mitoplasts (Wojtovich and Brookes, 2008; Garg and Hu, 2007; Foster et al., 2012; Hu et al., 2009; Flagg et al., 2010). The studies have also been conducted on selectivity and regulation of mitoK_{ATP} reconstituted into planar lipid membrane (Mironova et al., 2004). MitoK_{ATP} channels from uterus cells have not been described yet, although there are a great number of publications regarding this type of channels from plasma membrane (Brainard et al., 2007; Zhu et al., 2013). The present study was designed to establish the existence of ATP-sensitive K⁺-transporter in isolated rat uterus mitochondria and to determine its main functions. Conventional techniques, including fluorescence spectrophotometry, light scattering and direct measurements of K⁺ uptake with K⁺-sensitive fluorescent probe were used to achieve the goal. Our data allow us to propose the existence of ATP-sensitive K⁺-transporter in rat uterus mitochondria, similar to the one described in liver, brain and heart. We assume that mitoK_{ATP} may play an important role in uterus cells by means of regulating essential mitochondrial processes, such as mitochondrial membrane voltage, regulation of matrix volume and ROS production.

2. Material and methods

2.1. Isolation of mitochondria from rat uterus

All procedures performed on animals were approved by the Institutional Commission on Humane Treatment of Laboratory Animals of the Palladin Institute of Biochemistry of the National Academy of Sciences of Ukraine. The animals were housed in plastic cages and fed standard laboratory diet and water *ad libitum*. The study was carried out on sexually mature white female laboratory rats of no particular breed, average weight 150–200 g.

Animals were anesthetized with diethyl ether or chloroform and decapitated. For the extraction of mitochondria, the tissue of uterus was cleaned up from blood and fat, minced and homogenized on ice in 8 ml of isolation buffer containing 250 mM of sucrose, 1 mM of EDTA, and 10 mM of HEPES (pH 7.2 buffered with TRIS). The homogenate has been centrifuged for 7 min at 1000 g and temperature of 4 °C. The supernatant has been separated and centrifuged for 7 min at 12,000g and temperature of 4 °C. The pellet was resuspended in isolation buffer (containing no EDTA) and kept on ice (4 °C). Protein content was determined by Bradford method.

All fluorimetric measurements were conducted on the PTI Quanta Master 40 spectrofluorimeter (Canada) equipped with thermostated cuvette holder and magnetic stirrer. The standard incubation medium contained 10 mM HEPES (pH 7.2 at 28 °C, buffered with TRIS), 125 mM KCl, 5 mM K₂HPO₄, 1 mM MgCl₂, 5 mM succinate, 5 μM rotenone, and oligomycin 2 μg/ml. Oligomycin, an inhibitor of F₁F₀-ATPase, was used to exclude the influence of this enzyme on polarization of the inner mitochondrial membrane after the addition of ATP.

The K⁺-free medium contained all the above-mentioned components with the exception of KCl, which has been substituted isotonicly by 125 mM NaCl. Final protein concentration in all measurements was 55–60 μg/ml.

Stock solutions of glibenclamide had been prepared in DMSO and added into cuvette immediately prior to measurement in volumes of 1 μl per 1.8 ml of working volume (the final concentration of glibenclamide was 10 μM). 1 μl of DMSO was added to the control sample. Concentrated solution of 5-HD had been prepared in deionized water and added in volume of 5 μl per 1.8 ml of working volume (final concentration of 5-HD was 200 μM). Mitochondrial extraction buffer and working solutions had been prepared on deionized water. Aliquots of all substances had been added to the cuvette prior to measurement before the addition of mitochondria.

2.2. Mitochondrial membrane potential assay

Mitochondrial membrane potential was assayed with voltage-sensitive fluorescent probe TMRM (tetramethylrhodamine acetoxymethyl ester, 50 nM). Fluorescence signal was excited at 540 nm and registered at the wavelength of emission of 580 nm. The measurements started with the addition of aliquot of mitochondrial suspension. Complete depolarization of mitochondria was achieved by the addition of protonophore CCCP (carbonyl cyanide *m*-chlorophenyl hydrazone) to a final concentration of 2.5 μM or 100 μM CaCl₂.

2.3. Measurements of K⁺ influx into mitochondria loaded with PBFI-AM

Mitochondria were loaded with PBFI as described in (Costa et al., 2006). Briefly, pellets after second centrifugation were resuspended in 0.5 ml of buffered medium (pH 7.2, 25 °C) containing sucrose (250 mM) and pyruvate (10 mM). This suspension was incubated with 40 μM PBFI under gentle stirring for 10 min. An aliquot of PBFI solution in DMSO was previously mixed in 2:1 ratio (vol/vol) with the non-ionic surfactant F-127 before adding to the mitochondrial suspension. The mitochondrial suspension was then diluted with TEA⁺ medium containing (in mM): 175 sucrose, 10 HEPES (pH 7.2, 25 °C), 5 succinate, 5 Na₂HPO₄, 1 MgCl₂, 50 tetraethylammonium chloride (TEA-Cl) and incubated for 2 min under gentle stirring at 25 °C. This incubation was designed to substitute matrix K⁺ with TEA⁺. Then the suspension was diluted with 7 ml of ice-cold isolation buffer, centrifuged at 12,000g for 8 min, resuspended in isolation buffer and kept on ice. PBFI fluorescence was measured with excitation ratio technique (excitation wavelength 340/380 nm, emission wavelength 480 nm), in which the signal at 340 nm corresponds to the maximal sensitivity of the probe to K⁺ and the signal at 380 nm corresponds to the isosbestic point of the probe. Ratiometric measurements reduce variations in the measured fluorescence intensity that may arise from competing factors, probe concentration, etc (Costa et al., 2006). Measurements were conducted in the standard incubation medium (see above for details).

2.4. Mitochondrial swelling

Changes in mitochondrial matrix volume were followed using light-scattering technique. Measurements were conducted on

the PTI Quanta Master 40 spectrofluorimeter (Canada). Registrations of side light scattering at 520 nm were done in standard and K^+ -free incubation media with composition described above (see above for details). All effectors were added before the addition of the aliquot of mitochondrial suspension. Registrations of light scattering were started just after addition of the aliquot of mitochondrial suspension.

2.5. Measurement of ROS generation

Measurements had been conducted using ROS-sensitive probe DCF-DA (2,7-dichlorofluorescein-diacetate) (Gomes et al., 2005). A small aliquot of stock solution of DCF-DA was added into the incubation medium immediately prior to the measurement of the final concentration of 4 μ M. Wavelengths of the excitation and emission of fluorescence were 498 nm and 523 nm respectively. Registration of fluorescence was started at the 1 s. after the addition of aliquot of the suspension of mitochondria. Measurements were done in standard and K^+ -free incubation media with the composition described above (see above for details).

2.6. Statistical analysis

Analysis of the results and graph generation has been conducted by Microcal Origin software, version 5.0 (Microcal Software Inc., USA). Data were analyzed using Student's *t*-test. A value of $P < 0.05$ was considered statistically significant.

2.7. Chemicals

Sucrose, HEPES, KCl, CCCP, rotenone, oligomycin, succinate, glibenclamide, 5HD, 2,7-DCF-DA, TMRM were obtained from Sigma – Aldrich (USA); PBFI-AM – Molecular probes (USA), TRIS, EDTA, NaCl CaCl₂ (1 M solution) were obtained from Fluka (Switzerland); other chemicals were obtained from a domestic manufacturer (Ukraine).

3. Results

3.1. Measurements of the mitochondrial membrane potential

Experimental work has been performed on isolated fraction of rat uterus mitochondria prepared as described in the Methods section. Fluorescent probe TMRM is widely used for measurements of the inner mitochondrial membrane potential. Cationic molecule of TMRM easily moves through inner membrane toward its negative potential which can reach -200 mV in well coupled mitochondria (Suski et al., 2012). The mitochondria were coupled and maintained membrane potential as it can be seen in Fig. 1A. Accumulation of TMRM in the mitochondrial matrix after the addition of aliquot of mitochondrial suspension into the standard incubation medium supplied with respiration substrate results in fluorescence quenching with subsequent setting of the steady-state level. Protonophore CCCP (2.5 μ M) or CaCl₂ (100 μ M) caused rapid a dissipation of mitochondrial potential and restored the initial level of fluorescence signal. The same level of fluorescent signal was obtained in samples previously incubated with

2.5 μ M CCCP ((Fig. 1A). These results confirm the well-coupled conditions of the mitochondrial suspension, we used for this study.

As it can be seen in Fig. 1B the ATP-sensitive K^+ transport in uterus mitochondria leads to a decrease in $\Delta\psi$. Addition of ATP (200 μ M) caused a small increase in mitochondrial potential in the standard incubation medium if compared to control conditions. This effect was eliminated by the addition of diazoxide (50 μ M) along with ATP. Neither ATP nor diazoxide influenced the membrane potential in K^+ -free medium (Fig. 1B). Thus K^+ transport in mitochondria through the ATP-sensitive pathway results in slight depolarization of the inner mitochondrial membrane.

3.2. Measurement of K^+ influx into PBFI-loaded mitochondria

K^+ -sensitive fluorescent indicator PBFI is effectively used for direct measurement of K^+ transporting processes in living objects (Costa et al., 2006). Costa et al. demonstrated that PBFI can be used to detect K^+ uptake directly in the mitochondrial matrix in isolated rat heart mitochondria (Costa et al., 2006). Using the methodology described in this paper (Costa et al., 2006) we studied the K^+ uptake in isolated rat uterus mitochondria. Mitochondria loaded with PBFI accumulate K ions in the standard incubation medium as shown in Fig. 2. We had ensured that the intensity of fluorescence emission of PBFI loaded to mitochondria was directly proportional to the concentration of K ions absorbed from the incubation media with different concentration of KCl (data not shown). The K^+ uptake into mitochondrial matrix was sensitive to ATP (200 μ M), diazoxide (50 μ M) and glibenclamide (1 μ M). Addition of ATP (200 μ M) slowed down the increase in PBFI fluorescence signal in isolated rat uterus mitochondria in the standard incubation medium. The blocking effect of ATP on K^+ uptake was prevented by diazoxide, while glibenclamide eliminated the effect of mitoK_{ATP} activator (Fig. 2).

3.3. Light scattering studies of isolated mitochondria

The mitochondrial swelling could be detected by means of registration of light scattering signal on 520 nm. This technique is widely used for the studying of mitoK_{ATP} activation in isolated mitochondria (Costa et al., 2006; Garlid and Halestrap, 2012). Light scattering values are inversely dependent on the volume of mitochondria. This measurement is based on the fact that mitochondria isolated in K^+ -free medium are depleted of K ions, easily accumulate them and swell when added to K^+ -rich medium, resulting in a loss of light scattering. Fig. 3A contains typical light scattering traces from rat uterus mitochondria respiring in K^+ medium. In the absence of ATP, the matrix swelled to a higher steady-state volume. Addition of 200 μ M ATP resulted in a slower swelling rate and consequently in a lower steady-state volume, which means that ATP-sensitive K^+ uptake is blocked under these conditions. This effect of ATP was reversed by diazoxide (50 μ M). Another K_{ATP}-channel blocker glibenclamide (10 μ M) added along with diazoxide and ATP lowered the rate of swelling and steady-state volume (Fig. 3A and BI). But ATP and other effectors caused such effects only when K ions were present in the incubation medium. There was no dependency of light




Figure 1 Changes of membrane potential analyzed using TMRM 50 nM in isolated rat uterus mitochondria. (A) Polarization of mitochondrial inner membrane in the standard incubation medium supplied with succinate. Addition of either CCCP 2.5 mcM (1) or CaCl_2 100 mcM (2) (signed by arrow) caused full depolarization in coupled mitochondria; 3-CCCP 2.5 mcM was added at 0 s of measurements, prior to the addition of mitochondrial suspension; (B) effectors of $\text{mitoK}_{\text{ATP}}$ change polarization of mitochondrial membrane. (1) Polarization of mitochondrial membrane in the standard incubation medium. ATP 200 mcM alone (2) or alongside with diazoxide 50 mcM (3), were added to incubation medium before the addition of aliquot of mitochondrial suspension. In the end of each trace CCCP 2.5 mcM was added to ensure full depolarization (signed by arrow). Additions of aliquots of mitochondrial suspension signed by arrow (mito). Results of typical experiment are shown. Each curve is representative of at least three independent experiments.




Figure 2 K^+ uptake in isolated rat uterus mitochondria loaded with K^+ -sensitive fluorescent probe PBFI. Representative traces of PBFI fluorescence in the standard incubation medium. Effectors of $\text{mitoK}_{\text{ATP}}$ were present at the beginning of each trace, where indicated, in following concentrations: ATP 200 mcM, diazoxide (diaz) 50 mcM, glibenclamide (glyb) 1 mcM.

scattering of mitochondria on ATP, diazoxide or glibenclamide in K^+ -free medium (Fig. 3BII).

3.4. ROS measurements in isolated mitochondria

Under basic conditions in the standard incubation medium supplied with a substrate of respiration the isolated mitochondria produce moderate level of ROS as normal by-products of electron-transporting chain (Fig. 4A). Application of ATP (200 μM) attenuated ROS production. The rate of ROS

production was restored when diazoxide was added along with ATP. The stimulating effect of diazoxide on the generation of ROS was fully eliminated by 5-HD (200 μM) (Fig. 4A, BI). The same results were obtained with 1 μM glibenclamide (data not shown). All the effectors mentioned above had no influence on the rate of ROS production in the absence of K^+ ions (Fig. 4BII).

4. Discussion and conclusions

Although functioning and regulation of $\text{mitoK}_{\text{ATP}}$ -channel from other tissues is studied widely, the problem of existence of this structure in mitochondria from uterus has not been discussed yet. So, the major objective of this study was to reveal the ATP-sensitive K^+ transporting pathway in mitochondria from rat uterus. To accomplish it we studied the K^+ -dependent mode of action of the effectors of K_{ATP} -channels on mitochondrial physiology and directly measured K^+ uptake into the matrix of mitochondria from rat uterus.

The existence and physiological role of K_{ATP} -channel in plasma membrane of myometrium cells is well defined (Brainard et al., 2007; Zhu et al., 2013). This type of potassium channels is one of the most abundant in myometrium plasma membrane. Its main function, as in other types of smooth muscles, is to regulate excitability and provide for relaxation of uterus smooth muscle, which is crucial for its proper functioning (Brainard et al., 2007; Zhu et al., 2013). It was shown that both the known subunits of the K_{ATP} -channel are expressed in plasma membrane of human and rat myometrium (Brainard et al., 2007). Although the structure and mechanisms of regulation of uterus plasma membrane K_{ATP} -channel are actively investigated and discussed, there are very few data about this




Figure 3 Measurements of mitochondrial swelling in standard K^+ -rich medium (A, BI) or in K^+ -free medium with NaCl (equimolar substitution) (II). (A) Typical traces of light scattering changes in the standard incubation medium. (B) Averaged swelling rates relative to controls (with no additions, columns 1 and 2 respectively). Effectors of $\text{mitoK}_{\text{ATP}}$ were added as follows: (1)-none; (2)-ATP; (3)-ATP and diazoxide; (4)-ATP, diazoxide and glibenclamide. Concentrations of substances were: ATP 200 μM , diazoxide 50 μM , and glibenclamide 10 μM . Additions of aliquots of mitochondrial suspension signed by arrow (mito). At the 0.05 level, the means are significantly different. @-column 2I is significantly different from II, #-column 3I is significantly different from 2I, &-column 4I is significantly different from 3I.




Figure 4 Mitochondrial reactive oxygen species production increases with K^+ uptake. (A) Relative fluorescence of 2,7-dichlorofluorescein-diacetate expressed in arbitrary units (a.u.) is plotted against time (results of typical experiments are shown). ATP (200 μM) was present in all traces except for the trace marked "none". Diazoxide (diaz) and 5HD were added where mentioned in concentration 50 and 200 μM respectively. (B) Rates of ROS production by isolated rat uterus mitochondria are plotted as percentage of control rate obtained in control conditions in the standard incubation medium with no additions (II). I-The standard incubation medium, II- K^+ free medium, with KCl substituted with equimolar NaCl. (1)-No additions, (2)-ATP, (3)-ATP and diazoxide, (4)-ATP, diazoxide and 5HD, concentrations of effectors are mentioned above. Additions of aliquots of mitochondrial suspension signed by arrow (mito). At the 0.05 level, the means are significantly different. @-column 2I is significantly different from II, #-column 3I is significantly different from 2I, &-column 4I is significantly different from 3I.

channel in the inner membrane from uterus mitochondria (Vadzyuk and Kosterin, 2015), though there are plenty of data regarding $\text{mitoK}_{\text{ATP}}$ in mitochondria from other tissues, such as heart, liver, brain, and vessels (Garlid and Halestrap, 2012; Brainard et al., 2007; Zhu et al., 2013).

Study of $\text{mitoK}_{\text{ATP}}$ in living cell is complicated by the fact that both $\text{mitoK}_{\text{ATP}}$ and plasma membrane K_{ATP} -channel are regulated by the same effectors with the exception of 5-HD (Garlid and Halestrap, 2012). Also there are some contradictory data about the specificity of diazoxide toward

the K_{ATP} -channels (Garlid and Halestrap, 2012). We conducted the current study on the isolated mitochondrial fraction, so that data we obtained cannot be related to plasma membrane K_{ATP} -channels. In order to assume the potassium specificity of the substances we used, we repeated the same experiments in K^+ -free medium and made sure that all effects of ATP, diazoxide, glibenclamide and 5-HD on studied physiological processes in mitochondria appeared only if potassium ions were present. This is a strong indication that their influence on mitochondrial physiology is mediated by K^+ -dependent processes. Also, we directly demonstrated that K^+ uptake into the matrix of isolated mitochondria from rat uterus loaded with PBFI is sensitive to diazoxide and blockers of K_{ATP} -channel in the way that it is activated by diazoxide after blocking by ATP, and blocked if another blocker glibenclamide is added along with ATP and diazoxide (Fig. 2).

Thus, we studied K^+ -influx pathway in rat uterus mitochondria by means of four different assays, mitochondrial swelling, measurements of mitochondrial membrane voltage and ROS production, and direct measurement of K^+ uptake with K^+ -sensitive fluorescent probe. Taking into account all of the above, we may assume that all effects of activator and inhibitors of K_{ATP} -channel on studied physiological processes in mitochondria are mediated by the transporting of K ions through the ATP-sensitive pathway in rat uterus mitochondria, namely $mitoK_{ATP}$, similar to those from other tissues (Costa et al., 2006; Garg and Hu, 2007).

Our data allow us to assume that consequences of the activation of ATP-sensitive K^+ uptake in rat uterus mitochondria are (1) mild mitochondrial depolarization, (2) mitochondrial matrix swelling and (3) slight enhancement of ROS production. It should be said that our data regarding the influences of activators and blockers on functioning of mitochondria from myometrium are in concordance with those described in mitochondria isolated from other tissues (Costa et al., 2006; Garg and Hu, 2007).

The inner membrane potential is an important indicator of the mitochondrial energetic state (Suski et al., 2012). As it is a driving force for ATP synthesis and Ca^{2+} uptake, its value determines the proper functioning of the mitochondria and the cell as a whole. Thus, one of our aims was to determine whether the ATP-sensitive K^+ transport in uterus mitochondria could influence the mitochondrial inner membrane potential. It should be said that small depolarizing effect of $mitoK_{ATP}$ activation on mitochondrial inner membrane was reported on isolated vascular and renal mitochondria (Katakam et al., 2014; Cancherini et al., 2003). The same results were obtained for uterus mitochondria (Fig. 1B). This could be explained in that way, that K^+ uptake results in an increase in mitochondrial matrix volume, which in turn activates K^+/H^+ exchanger, promotes H^+ extrusion out of matrix, and provides a mild uncoupling effect (Costa et al., 2006). This uncoupling stimulates electron transporting chain. The result of it could be enhancement of ATP synthesis, as well as amplification of ROS production (Suski et al., 2012). It should be noticed that mitochondrial uncoupling could have protective effect by itself (Minners et al., 2000; Hausenloy et al., 2004). Minners et al. showed that mitochondrial uncoupling agents confer protection under stress conditions (Minners et al., 2000).

Another beneficial consequence of partial mitochondrial depolarization as a result of K^+ transporting into mitochondria

is attenuation of Ca^{2+} uptake. As Ca^{2+} influx into mitochondria is driven primarily by $\Delta\psi$, a small change in $\Delta\psi$ can induce a large change in matrix Ca^{2+} concentration (Ishida et al., 2001). Indeed, Holmuhamedov et al. showed that the pharmacological activation of cardiac $mitoK_{ATP}$ by diazoxide decreased the accumulation of Ca ions into the mitochondrial matrix (Holmuhamedov et al., 1999). It is known that excessive Ca^{2+} accumulation in mitochondrial matrix may trigger the induction of the mitochondrial permeability transition pore opening and cell death, and the prevention of Ca^{2+} overload may protect mitochondria from the induction of apoptosis (Holmuhamedov et al., 1999). Therefore, we assume that the mild uncoupling effect of $mitoK_{ATP}$ activation could have beneficial consequences on physiology of uterus mitochondria under normal as well as under stress conditions, and effectively prevent Ca^{2+} overload.

There is a general agreement that K^+ uptake into mitochondria, accompanied by the movement of anions, evokes osmotic entry of water. This leads to an increase in matrix volume with further activation of K^+/H^+ -exchange and settlement of new steady-state matrix volume without the disruption of the outer mitochondrial membrane. The primary physiological consequence of the mitochondrial matrix swelling is the enhancement of respiration and contraction of intermembrane space, resulting in an increased membrane permeability to ADP and, as a result, the facilitation of ATP synthesis (Costa et al., 2006). Thus, mitochondrial matrix swelling after activation of $mitoK_{ATP}$ could occur in response to an increased cellular need in ATP. These effects could be crucial for mitochondrial functioning under ischemic conditions, as well as under conditions of normoxia. As we showed earlier (Vadzyuk and Kosterin, 2015) the activation of $mitoK_{ATP}$ has a protective effect on rat uterus cell under oxidative stress conditions. The explanation of this phenomenon may consist, as well, in mild uncoupling and mitochondrial swelling as the consequences of the activation of $mitoK_{ATP}$ in rat uterus cells.

ROS are natural by-products of electron transporting chain, and intensity of their generation directly depends on the mitochondrial respiration rate (Suski et al., 2012). ROS are known signaling molecules, which trigger different pathways resulting in stress adaptation or cell death (Brown and Griendling, 2015). Thus, their production under conditions of normoxia should be tightly controlled. We assume that the activation of ATP-sensitive K^+ uptake in mitochondria could be one of mode of regulation of the intensity of ROS production in uterus mitochondria. As data of Fig. 4 show, when $mitoK_{ATP}$ channel is closed by ATP or 5-HD, ROS production is substantially lower than in conditions where ATP-sensitive K^+ uptake is allowed. It is known that ROS regulates homeostasis of Ca^{2+} in smooth muscle and mediates muscle relaxation (Brown and Griendling, 2015; Zhang et al., 2013), so we assume that the activation of $mitoK_{ATP}$ channel in rat uterus cells may contribute to the relaxation of myometrium by means of amplification of ROS generation. On the other hand, ROS induced by activation of $mitoK_{ATP}$ could promote cell adaptation to environmental stress (Brown and Griendling, 2015). It was showed that ROS scavengers blocked protection induced by diazoxide under conditions of ischemia (Semirabet et al., 2008). This clearly indicates that a relationship exists between $mitoK_{ATP}$, ROS and cell protection. Therefore the impact of $mitoK_{ATP}$ activation on ROS production may have a protective effect under stress conditions and

regulate contraction and relaxation under normal conditions. We have shown earlier that diazoxide mildly increased ROS production under normal conditions in rat uterus cells, but decreased severe peroxidation under chemically induced oxidative stress conditions if cells had been incubated with diazoxide prior to the induction of stress. This effect may be explained also by ROS-mediated cytoprotection and may involve other intermediates, such as protein kinases.

Our results allow us to assume that ATP-sensitive K^+ -channel in rat uterine smooth muscle mitochondria exists and exerts versatile effects on mitochondrial physiology – regulates matrix swelling, ROS production and change inner membrane potential, which may be a response of mitochondria on cell demands under different metabolic challenges, and may have an important role in physiology of the smooth muscle as a whole.

Acknowledgements

Authors wish to thank Dr. Dmytro Petukhov from Palladin Institute of Biochemistry for his kind help in editing the manuscript. This work was supported by grant No. 0114U003213 from the fundamental research program of the National Academy of Sciences of Ukraine.

References

- Brainard, A., Korovkina, V.P., England, S.K., 2007. Potassium channels and uterine function. *Semin. Cell Dev. Biol.* 18, 332–339.
- Brown, D.I., Griendling, K.K., 2015. Regulation of signal transduction by reactive oxygen species in the cardiovascular system. *Circ. Res.* 116, 531–549.
- Cancherini, D.V., Trabuco, L.G., Reboucas, N.A., Kowaltowski, A.J., 2003. ATP-sensitive K^+ channels in renal mitochondria. *Am. J. Physiol. Renal Physiol.* 285, F1291–F1296.
- Costa, A.D.T., Casey, L., Andrukhiv, A., West, I.C., Jaburek, M., Garlid, K.D., 2006. The direct physiological effects of $\text{mitoK}_{\text{ATP}}$ opening on heart mitochondria. *Am. J. Physiol. Heart Circ. Physiol.* 290, H406–H415.
- Flagg, T.P., Enkvetchakul, D., Koster, J.C., Nichols, C.G., 2010. Muscle KATP channels: recent insight to energy sensing and myoprotection. *Physiol. Rev.* 90, 799–829.
- Foster, D.B., Ho, A.S., Rucker, J., Garlid, A.O., Chen, L., Sidor, A., Garlid, K.D., O'Rourke, B., 2012. Mitochondrial ROMK channel is a molecular component of $\text{mitoK}_{\text{ATP}}$. *Circ. Res.* 111, 446–454.
- Galluzi, L., Kepp, O., Trojel-Hansen, C., Kroemer, G., 2012. Mitochondrial control of cellular life, stress, and death. *Circ. Res.* 111, 1198–1207.
- Garg, V., Hu, K., 2007. Protein kinase C isoform-dependent modulation of ATP-sensitive K^+ channels in mitochondrial inner membrane. *Am. J. Physiol. Circ. Heart Physiol.* 293, H322–H332.
- Garlid, K.D., Halestrap, A.P., 2012. The mitochondrial K_{ATP} channel – fact or fiction? *J. Mol. Cell. Cardiol.* 52, 578–583.
- Gomes, A., Fernandes, E., Lima, J.L.F.C., 2005. Fluorescence probes used for detection of reactive oxygen species. *J. Biochem. Biophys. Methods* 65, 45–80.
- Hausenloy, D.J., Yellon, D.M., Mani-Babu, S., Duchon, M.R., 2004. Preconditioning protects by inhibiting the mitochondrial permeability transition. *Am. J. Physiol. Heart Circ. Physiol.* 287, H841–H842.
- Holmuhamedov, E.L., Wang, L., Terzic, A., 1999. ATP-sensitive K^+ channel openers prevent Ca^{2+} overload in rat cardiac mitochondria. *J. Physiol.* 519, 347–360.
- Hu, L.F., Lu, M., Wong, P.T.H., Bian, J.S., 2009. Hydrogen sulfide inhibits rotenone-induced apoptosis via preservation of mitochondrial function. *Mol. Pharmacol.* 75, 27–34.
- Ishida, H., Hirota, Y., Genka, C., Nakazawa, H., Nakaya, H., Sato, T., 2001. Opening of mitochondrial KATP channels attenuates the ouabain-induced calcium overload in mitochondria. *Circ. Res.* 89, 856–858.
- Katakam, P.V.G., Gordon, A.O., Sure, V.N.L.R., Rutkai, I., Busija, A., 2014. Diversity of mitochondria-dependent dilator mechanisms in vascular smooth muscle of cerebral arteries from normal and insulin-resistant rats. *Am. J. Physiol. Circ. Heart Physiol.* 307, H493–H503.
- Minners, J., Bos, E.J., Yellon, D.M., Schwalb, H., Opie, L.H., Sack, M.N., 2000. Dinitrophenol, cyclosporine A, and trimetazidine modulate preconditioning in the isolated rat heart: support for a mitochondrial role in cardioprotection. *Cardiovasc. Res.* 47, 68–73.
- Mironova, G.D., Negoda, A., Marinov, B.S., Paucek, P., Costa, A.D. T., Grigoriev, S.M., Skarga, Y.Y., Garlid, K.D., 2004. Functional distinction between the mitochondrial ATP-dependent K^+ channel ($\text{mitoK}_{\text{ATP}}$) and its inward rectifier subunit (mitoKIR). *J. Biol. Chem.* 279, 32562–32568.
- Semirabet, M., Robin, E., Aristi, I., Adamczyk, S., Tavernier, B., Vallet, B., 2008. Preconditioning by an in situ administration of hydrogen peroxide: involvement of reactive oxygen species and mitochondrial ATP-dependent potassium channel in a cerebral ischemia-reperfusion model. *Brain Res.* 1240, 177–184.
- Suski, J.M., Lebiecinska, M., Bonora, M., Adamczyk, S., Tavernier, B., Vallet, B., Bordet, R., Lebuffe, G., 2012. Relation between mitochondria membrane potential and ROS formation. In: Palmeira, C.M., Moreno, A.J. (Eds.), *Mitochondrial Bioenergetics: Methods and Protocols, Methods in Molecular Biology*, 810. Springer Science + Business Media LLC, New York, pp. 183–205.
- Vadzyuk, O., Kosterin, S., 2015. Diazoxide attenuates ROS generation and exerts cytoprotection under conditions of ROS overproduction in rat uterus cells. *Turk. J. Biol.* 39, 447–454.
- Wojtovich, A.P., Brookes, P.S., 2008. The endogenous mitochondrial complex II inhibitor malonate regulates mitochondrial ATP-sensitive potassium channels: implication for ischemic preconditioning. *Biochim. Biophys. Acta* 1777, 882–889.
- Zhang, H.A., Gomes, A.M., Wang, X., Yan, Y., Zheng, M., Cheng, H., 2013. ROS regulation of microdomain Ca^{2+} signaling at the dyads. *Cardiovasc. Res.* 98, 248–258.
- Zhu, R., Xiao, D.L., Zhang, L., 2013. Potassium channels and uterine vascular adaptation to pregnancy and chronic hypoxia. *Curr. Vasc. Pharmacol.* 11, 737–747.