



ELSEVIER

Available at

www.ElsevierMathematics.com

POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 125 (2003) 131–144

<http://www.elsevier.com/locate/jat>JOURNAL OF
Approximation
Theory

Strong asymptotic behavior for extremal polynomials with respect to varying measures on the unit circle[☆]

M. Bello Hernández* and J. Mínguez Cenicerros

Dpto. de Matemáticas y Computación, Universidad de La Rioja, Edificio J. L. Vives, Luis de Ulloa, s/n, 26004 Logroño, Spain

Received 21 December 2002; accepted in revised form 15 September 2003

Communicated by Guillermo López Lagomasino

Abstract

We give a Szegő-type theorem for L^p -extremal polynomials with respect to varying measures on $|z| = 1$. Also, we present a density theorem and a generalization of the main result to closed rectifiable Jordan curves and to $|z| = 1$ with the possible addition of a finite number of mass points.

© 2003 Elsevier Inc. All rights reserved.

1. Introduction

Orthonormal polynomials with respect to varying measures were introduced about 25 years ago by Gonchar and López Lagomasino [5] in connection with a systematic study of the convergence properties of interpolating rational functions with free poles to Markov functions. In a more general context, such approximants are called multipoint Padé approximants. In [5], an analogue of the classical Markov theorem in the theory of continued fractions was proved. Meanwhile, a number of research papers have been devoted to the subject of orthonormal polynomials with

[☆]The work has been supported by Dirección General de Investigación (Ministerio de Ciencia y Tecnología) of Spain under Grant BFM2000-0206-C04-03.

*Corresponding author.

E-mail addresses: mbello@dmc.unirioja.es (M. Bello Hernández), judit.minguez@dmc.unirioja.es (J. Mínguez Cenicerros).

respect to varying measures; surveys can be found in [12,17], and Chapter 6 of [15]. The emphasis in the present paper is on strong asymptotic, which in the unweighted situation is known as Szegő asymptotics for L^p -extremal polynomials [4]. Strong asymptotic results for orthonormal polynomials with respect to varying measures have been proved in various degree of generality in [2,3,11,14,17].

The classical theory of strong asymptotics has its most simple and perhaps also its most natural form for orthonormal polynomials with respect to a measure supported on the unit circle $\mathbb{T} = \{|z| = 1\}$ [16, Chapters XI, XII], or [13, Chapter 3]. Similar results for orthonormal polynomials on a real interval are then usually deduced from the results on \mathbb{T} in a second step. In [10] López–Lagomasino presents orthogonal polynomials with respect to varying measures in a such a way that unifies the theory for the cases of measures with bounded and unbounded support. He also proves asymptotic results for orthogonal polynomials respect to (a fixed) measure with unbounded support using orthogonal polynomials respect to varying measures on \mathbb{T} .

Also, these polynomials are very important in Potential Theory. The solution of a number of problems in approximation theory can be reduced to finding the equilibrium distribution of a charge on a “conductor” in the presence of certain “external fields”. Problems of this type arise, for example, in the theory of convergence of Padé approximants. In [6], Gonchar and Rakhmanov proved a general theorem which characterizes the limit distribution of the zeros of orthogonal polynomials with respect to varying measures. Also, Gonchar and Rakhmanov proved in [7] a general result concerning the exact rate of best rational approximation for a large class of analytic functions. This result was stated in terms of equilibrium distributions in the presence of external fields and the proof is based on the construction of convenient multipoint Padé approximants whose convergence properties in turn reduces to the study of the limit distribution of zeros of sequences of polynomials which satisfy complex orthogonal relations with respect to varying measures.

In this paper, we give a Szegő-type theorem for L^p -extremal polynomials with respect to varying measures on $|z| = 1$. This result will be stated in this section and proved in Section 3. Section 2 is devoted to some auxiliary statements. In Section 4, we present a theorem on density of rational functions and finally, in Section 5, we give some generalizations of the main result. We begin introducing some notations.

Let μ be a finite positive Borel measure on $[0, 2\pi)$ whose support contains an infinite set of points. In the sequel, we consider $\{W_n\}$, $n \in \mathbb{N}$, a sequence of polynomials such that, for each $n \in \mathbb{N}$, W_n has degree n ($\deg W_n = n$), all its zeros $\{w_{n,i} : 1 \leq i \leq n\}$ lie in $\mathbb{D} = \{z : |z| < 1\}$, and they satisfy

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n (1 - |w_{n,i}|) = +\infty. \tag{1}$$

We want to study the asymptotic behavior of polynomials that solve the extremal problem

$$\tau_{n,p} = \inf_{Q_n(z)=z^n+\dots} \left\| \frac{Q_n}{W_n} \right\|_p = \inf_{Q_n \in \Pi_n, Q_n(0)=1} \left\| \frac{Q_n}{W_n^*} \right\|_p, \tag{2}$$

where Π_n is the set of polynomials of degree at most n , $W_n^*(z) = z^n \overline{W_n(\frac{1}{z})}$, and

$$\|f\|_p = \left\{ \frac{1}{2\pi} \int |f|^p d\mu \right\}^{1/p}.$$

Note that the zeros of $W_n^*(z)$ are $\{\frac{1}{w_{n,i}}\}_{i=1,\dots,n} \subset \mathbb{E} = \{|z| > 1\}$. From now on, $P_{n,p}$ denotes a polynomial such that

$$\left\| \frac{P_{n,p}}{W_n} \right\|_p = \tau_{n,p}.$$

Let μ' be the Radon–Nikodym derivate of μ with respect to the Lebesgue measure. Assume that $\log \mu' \in L^1$; let $D_p(\mu, z)$ denote the corresponding Szegő function; that is,

$$D_p(\mu, z) = \exp \left\{ \frac{1}{2p\pi} \int_0^{2\pi} \frac{\zeta + z}{\zeta - z} \log \mu'(\theta) d\theta \right\}, \quad \zeta = e^{i\theta}, z \in \mathbb{D}.$$

Set

$$K_p(\mu, z) = \begin{cases} \frac{D_p(\mu, 0)}{D_p(\mu, z)} & \text{if } z \in S_a \cup \mathbb{D}, \\ 0 & \text{if } z \in S_s, \end{cases} \tag{3}$$

where S_a and S_s give a disjoint decomposition of the unit circle such that μ' and μ_s live on these sets, respectively. Hereafter, μ_s denotes the singular part of μ with respect to the Lebesgue measure. H^p , $0 < p < 1$, is defined as the class of all functions f analytic in \mathbb{D} such that

$$\sup_{0 < r < 1} \left(\int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

It is well known that H^p can be identified with the closure in L^p of the set of polynomials in $e^{i\theta}$.

The function $D_p(\mu, z)$ satisfies the following properties:

- (1) $D_p(\mu, z)$ is analytic in \mathbb{D} ; more precisely, $D_p(\mu, z) \in H^p$,
- (2) $D_p(\mu, z) \neq 0$ in \mathbb{D} , and $D_p(\mu, 0) > 0$,
- (3) $|D_p(\mu, e^{i\theta})|^p = \mu'(\theta)$ almost everywhere (a.e.) in $[0, 2\pi]$.

$H^p(\mu)$ is defined as the $L^p(\mu)$ closure of the polynomials in $e^{i\theta}$. $L_s^p(\mu) = \{f \in L^p(\mu) : f = 0, \mu'_a \text{ a.e.}\}$ and $L_a^p(\mu) = \{f \in L^p(\mu) : f = 0, \mu_s \text{ a.e.}\}$. Similarly, we define $H_s^p(\mu)$ and $H_a^p(\mu)$.

Moreover, if $f \in H^p(\mu)$, then there exist unique functions \tilde{f}, f_s such that

$$f = K_p \tilde{f} + f_s, \quad \tilde{f} \in H^p, \text{ and } f_s \in L_s^p(\mu). \tag{4}$$

A proof of this result can be seen in [1].

The main theorem of this paper which will be proved in Section 3 is the following:

Theorem 1. For $0 < p < \infty$, the following statements are equivalent:

- (i) μ satisfies the Szegő condition; that is, $\log \mu' \in L^1$.
- (ii) The following limit exists and is positive

$$\lim_{n \rightarrow \infty} \tau_{n,p} > 0.$$

- (iii) There exists a function $S \in H^p(\mu)$ with $\|S\|_p \neq 0$, such that

$$\lim_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} - S \right\|_p = 0.$$

- (iv) There exists a function T analytic in \mathbb{D} such that

$$\lim_{n \rightarrow \infty} \frac{\frac{P_{n,p}^*(z)}{W_n^*(z)}}{\left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p} = T(z)$$

holds uniformly on each compact subset of \mathbb{D} .

Moreover, if (i) holds, then

$$\lim_{n \rightarrow \infty} \tau_{n,p} = D_p(\mu, 0),$$

and the functions in (iii) and (iv) are $S(z) = K_p(\mu, z)$ and $T(z) = \frac{1}{D_p(\mu, z)}$.

2. Auxiliary results

Before we can prove the theorems in the following sections, we need to establish several auxiliary results.

Let K be a compact set and $\{\alpha_{n,1}, \dots, \alpha_{n,n}\} \subset \mathbb{C} \setminus K$ be a given set of points. Let F_n be the set of functions of the form

$$\pi_n(z) = \frac{b_{n,0}z^n + b_{n,1}z^{n-1} + \dots + b_{n,n}}{(z - \alpha_{n,1})(z - \alpha_{n,2}) \cdots (z - \alpha_{n,n})}. \tag{5}$$

Let f be a continuous function on K . Denote by $r_n(f)$ the best approximation to $f(z)$ on K in the class F_n in the sense of Tchebycheff; that is,

$$\|f - r_n(f)\| = \min\{\|f - \pi_n\| : \pi_n \in F_n\}$$

with $\|\cdot\|$ the supremum norm on K .

Theorem 2. (See Walsh [18, pp. 246–247]). *Let the points $\alpha_{n,k}$ satisfy $|\alpha_{n,k}| > 1$. A necessary and sufficient condition such that*

$$\lim_{n \rightarrow \infty} r_n(f)(z) = f(z), \text{ uniformly in } |z| \leq 1, \tag{6}$$

for every such function f analytic in $\{|z| \leq 1\}$ is that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \left(1 - \frac{1}{|\alpha_{n,k}|} \right) = +\infty. \tag{7}$$

A result due to Keldysh is also useful in the proofs that follow. This theorem appears in [9]. An extension of Keldysh’s theorem is the following.

Theorem 3. (See Bello Hernández et al. [1]).¹ *Let $\{z_i\}_{i=1, \dots, \Lambda}$ be a set of points in \mathbb{D} , where Λ can be finite or infinite. Let β be a finite positive Borel measure on $[0, 2\pi)$ satisfying the Szegő condition and $\{f_n\} \subset H^p(\beta)$ (from (4), $f_n = K_p(\beta, \cdot) \tilde{f}_n + f_{n,s}$), $0 < p < \infty$, such that*

- (i) $\lim_{n \rightarrow \infty} \tilde{f}_n(0) = 1$;
- (ii) $\lim_{n \rightarrow \infty} \tilde{f}_n(z_i) = 0, i = 1, 2, \dots$;
- (iii) $\sum_{i=1}^{\Lambda} (1 - |z_i|) < +\infty$;
- (iv) $\lim_{n \rightarrow \infty} \left(\int |f_n(e^{i\theta})|^p d\beta(\theta) \right)^{1/p} = \frac{D_p(\beta, 0)}{\prod_{i=1}^{\Lambda} |z_i|}$.

Then

(a) $\lim_{n \rightarrow \infty} \tilde{f}_n(z) = \prod_{i=1}^{\Lambda} \frac{z - z_i}{\bar{z}_i z - 1} \frac{\bar{z}_i}{|z_i|^2}$ holds uniformly on each compact subset of \mathbb{D} ,

and

(b) $\lim_{n \rightarrow \infty} \int \left| f_n - K_p(\beta, z) \prod_{i=1}^{\Lambda} \frac{z - z_i}{\bar{z}_i z - 1} \frac{\bar{z}_i}{|z_i|^2} \right|^p d\beta = 0$.

If Λ is an empty set then the right-hand side of (a) is equal to 1; that is, $\prod_{i=1}^{\Lambda} \frac{z - z_i}{\bar{z}_i z - 1} \frac{\bar{z}_i}{|z_i|^2} \equiv 1$.

3. Proof of Theorem 1

Before proving Theorem 1, we show an intermediate result.

¹We wish to point out a missprint in condition (iv) of the mentioned reference. It is necessary to substitute $\prod_{i=1}^{\Lambda} |z_i|^p$ by $\prod_{i=1}^{\Lambda} |z_i|$.

Theorem 4. For $0 < p < \infty$

$$\lim_{n \rightarrow \infty} \left\| \frac{P_{n,p}}{W_n} \right\|_p = D_p(\mu, 0),$$

where 0 replaces $D_p(\mu, 0)$ if $\log \mu'(\theta)$ is not integrable.

Proof. Let $\Lambda \subset \mathbb{N}$ be an indexed sequence such that

$$\lim_{n \in \Lambda} \left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p = \limsup_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p. \tag{8}$$

From a result due to Szegő (see [16, p. 297]), we know that if $T_{n,2}$ are the extremal polynomials such that

$$\|T_{n,2}\|_2 = \min\{\|Q_n\|_2 : Q_n \text{ monic of degree } n\},$$

then

$$\lim_n \|T_{n,2}\|_2^2 = \lim_n \|T_{n,2}^*\|_2^2 = D_2(\mu, 0)^2, \tag{9}$$

with $D_2(\mu, 0) = 0$ if $\log \mu'(\theta)$ is not integrable. Since the zeros of $T_{n,2}^*$ lie in \mathbb{E} , $\varphi_n(z) = (T_{n,2}^*(z))^{2/p}$ is analytic in $\{|z| \leq 1\}$, so as (1) holds, from Theorem 2 there exists a sequence $\{R_{m_n}^*\}_{m_n \in \Lambda' \subset \Lambda}$ such that

$$\lim_{n \in \Lambda} \sup_{|z| \leq 1} \left| \frac{R_{m_n}^*(z)}{W_{m_n}^*(z)} - \varphi_n(z) \right| = 0.$$

In particular, since $\varphi_n(0) = (T_{n,2}^*(0))^{p/2} = 1 = W_{m_n}^*(0)$, we have $\lim_{n \in \Lambda} R_{m_n}^*(0) = 1$. Hence

$$\lim_{n \in \Lambda} \left\| \frac{R_{m_n}^*}{W_{m_n}^*} \right\|_p^p = \lim_{n \in \Lambda} \|\varphi_n\|_p^p = \lim_{n \in \Lambda} \|T_{n,2}^*\|_2^2 = D_2(\mu, 0)^2 = D_p(\mu, 0)^p$$

and

$$\limsup_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p \leq D_p(\mu, 0). \tag{10}$$

On the other hand, using Jensen’s inequality

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{P_{n,p}^*(e^{i\theta})}{W_n^*(e^{i\theta})} \right|^p d\mu(\theta) &\geq \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{P_{n,p}^*(e^{i\theta})}{W_n^*(e^{i\theta})} \right|^p \mu'(\theta) d\theta \\ &\geq \exp \left\{ \frac{1}{2\pi} \int_0^{2\pi} \log \left| \frac{P_{n,p}^*(e^{i\theta})}{W_n^*(e^{i\theta})} \right|^p d\theta \right\} \\ &\quad \times \exp \left\{ \frac{1}{2\pi} \int_0^{2\pi} \log \mu'(\theta) d\theta \right\} \\ &\geq \left| \frac{P_{n,p}^*(0)}{W_n^*(0)} \right|^p D_p(\mu, 0)^p = D_p(\mu, 0)^p. \end{aligned}$$

Therefore,

$$\liminf_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p \geq D_p(\mu, 0). \tag{11}$$

With (10) and (11) the theorem is proved. \square

Remark 1. We also proved that

$$\lim_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} \right\|_p = \lim_{n \rightarrow \infty} \left(\frac{1}{2\pi} \int \left| \frac{P_{n,p}^*(z)}{W_n^*(z)} \right|^p \mu'(\theta) d\theta \right)^{1/p} = D_p(\mu, 0).$$

Proof of Theorem 1.

Proof. (i) \Leftrightarrow (ii): It follows from Theorem 4.

(i) \Rightarrow (iii): We consider the function

$$h_n(z) = \frac{P_{n,p}^*(z) D_p(\mu, z)}{W_n^*(z) D_p(\mu, 0)}, \tag{12}$$

that belongs to H^p . Since $h_n(0) = 1$ and $|D_p(\mu, e^{i\theta})|^p = \mu'(\theta)$, from Theorem 4, we have

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} |h_n(e^{i\theta})|^p d\theta \right\} = 1. \tag{13}$$

Applying Theorem 3 (here $\Lambda = \emptyset$ and β is the Lebesgue measure) it follows that

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} |h_n(e^{i\theta}) - 1|^p d\theta \right\} = 0.$$

Hence

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{P_{n,p}^*(e^{i\theta}) D_p(\mu, e^{i\theta})}{W_n^*(e^{i\theta}) D_p(\mu, 0)} - 1 \right|^p d\theta \right\} = 0$$

and then

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{P_{n,p}^*(e^{i\theta})}{W_n^*(e^{i\theta})} - \frac{D_p(\mu, 0)}{D_p(\mu, e^{i\theta})} \right|^p \mu'(\theta) d\theta \right\} = 0.$$

Therefore, using (13) and again Theorem 4, we obtain (iii) where $S(z) = K_p(\mu, z)$.

(iii) \Rightarrow (i): It follows from the relation

$$\liminf_{n \rightarrow \infty} \left\| \frac{P_{n,p}^*}{W_n^*} - \psi \right\|_p = 0, \tag{14}$$

where $\psi(\zeta)$ is such that $\|\psi\|_p \neq 0$.

Indeed, according to (14) we have that there exists a subsequence $\{n_v\}$ such that

$$\lim_{v \rightarrow \infty} \left\| \frac{P_{n_v,p}^*}{W_{n_v}^*} - \psi \right\|_p = 0.$$

If (i) does not hold, from Theorem 4

$$\lim_{v \rightarrow \infty} \left\| \frac{P_{n_v,p}^*}{W_{n_v}^*} \right\|_p = 0,$$

and we obtain $\|\psi\|_p = 0$, which is a contradiction.

(iii) \Rightarrow (iv): The sequence of functions $\{h_n\}$ as in (12) satisfies the hypothesis of Theorem 3, hence $\lim_{n \rightarrow \infty} h_n(z) = 1$ holds uniformly on each compact subset of \mathbb{D} . Now, since (i) is equivalent to (iii), using again Theorem 4, we obtain

$$\lim_{n \rightarrow \infty} \frac{\frac{P_{n,p}^*(z)}{W_n^*(z)}}{\left\| \frac{P_{n,p}^*(z)}{W_n^*(z)} \right\|_p} = \frac{1}{D_p(\mu, z)}.$$

(iv) \Rightarrow (i). From (iv) and Theorem 4, we have

$$\lim_{n \rightarrow \infty} \frac{\frac{P_{n,p}^*(0)}{W_n^*(0)}}{\left\| \frac{P_{n,p}^*(z)}{W_n^*(z)} \right\|_p} = T(0) = \frac{1}{D_p(\mu, 0)} < \infty,$$

but this is true if and only if (i) holds. \square

4. Density Theorem

In this section we give a density theorem that can be seen as an “application” of the main result.

We introduce the notation: $R_{n,k} = \left\{ \frac{h}{W_n^*} : h \in \Pi_{n-k} \right\}$.

Theorem 5. Assume that μ is an absolutely continuous measure μ_a and satisfies Szegő's condition, then the following statements are equivalent:

(i) For each $j \in \mathbb{Z}_+$

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{P_{n,n-j,p}^*(e^{i\theta})}{W_n^*(e^{i\theta})} \right|^p d\mu_a(\theta) = D_p(\mu_a, 0)^p,$$

where $P_{n,n-j,p}$ is the monic extremal polynomial; that is,

$$\left\| \frac{P_{n,n-j,p}}{W_n} \right\|_p = \min \left\{ \left\| \frac{Q_{n-j}}{W_n} \right\|_p : Q_{n-j} \in \Pi_{n-j}, \text{ monic} \right\}.$$

(ii) For each $k \in \mathbb{Z}_+$, $R_{n,k}$ is dense in $H^p(\mu_a)$.

Proof. (ii) \Rightarrow (i): Here we use the same technique as in the proof of Theorem 4. It is well known that (ii) \Leftrightarrow (i) is true when $p = 2$ and $W_n(z) = z^n$ (see, for example, [16, p. 297]). Let $T_{n-k,2}$ denote the monic extremal polynomials in this case, then given $k \geq 0$ there exist polynomials R_{m_n-k} of degree $m_n - k$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \left| (T_{n-k}^*(z))^{2/p} - \frac{R_{m_n-k}(z)}{W_{m_n}^*(z)} \right|^p \mu_a'(\theta) d\theta = 0.$$

Notice that the functions $\{(T_{n-k}^*)^{2/p}\}_n$ are analytic in an open set containing $\{|z| \leq 1\}$ because T_{n-k}^* has no zeros in $\{|z| \leq 1\}$. Then

(a) $\lim_{n \rightarrow \infty} R_{m_n-k}(0) = 1;$

(b) $\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{R_{m_n-k}(z)}{W_{m_n}^*(z)} \right|^p \mu_a'(\theta) d\theta = D_p(\mu_a, 0)^p.$

Given $\Lambda \subset \mathbb{N}$ an indexed sequence, from (ii) we observe that the sequence $\{m_n\}$ can be chosen in Λ . Therefore, (i) follows from (a) and (b).

(i) \Rightarrow (ii): Set $i, j \in \mathbb{Z}_+$, using (i) and Theorem 3, we have

$$\frac{P_{n,n-(i+j),p}^*(z)}{W_n^*(z)} \rightarrow K_p(\mu_a, z)$$

in $L^p(\mu_a)$. Then

$$\frac{z^i P_{n,n-(i+j),p}^*(z)}{W_n^*(z)} \rightarrow z^i K_p(\mu_a, z),$$

in $H^p(\mu_a)$. Since $H^p(\mu_a) = H^p \cdot K_p(\mu_a, \cdot)$ and H^p is the closure of the polynomials in L^p , $R_{n,j}$ satisfies (ii). \square

5. Generalizations

In this section we give two generalizations of Szegő’s theorem for extremal rational functions. First, for a closed rectifiable Jordan curve C , and second, for sets of the form $\mathbb{F} = \mathbb{T} \cup \{z_1, z_2, \dots, z_N\}$, with $z_i \in \mathbb{D}$. Szegő’s theorem on \mathbb{F} respect to a fixed measure was proved by Kaliaguine [8].

With the same techniques used in Theorem 1, we can prove an analogous theorem for closed rectifiable Jordan curves. We begin introducing the necessary notation. Let $\alpha \in \mathbb{C}$ and let C be a closed rectifiable Jordan curve with length l in the z -plane in whose interior lies α . Let $\sigma(s)$ be a finite positive measure on $[0, l]$. As usual, by $L^p(C, \sigma)$ we denote the space of complex measurable functions on C , such that

$$\|f\|_{p,\sigma} = \left\{ \frac{1}{2\pi} \int_C |f(\zeta)|^p d\sigma(s) \right\}^{1/p} < \infty, \quad \zeta = \zeta(s), \quad s \in [0, l],$$

with $\zeta = \zeta(s)$ a parametrization of C . Let B denote the interior of C and consider the conformal transformation

$$x = \varphi(z) = \alpha + z + b_2 z^2 + \dots, |z| < 1, \quad \alpha \in B$$

which maps \mathbb{D} onto B , such that $\alpha = \varphi(0)$ and $\varphi'(0) > 0$. From Caratheodory’s theorem φ can be extended continuously to $\{|z| \leq 1\}$ so that φ is a bijection from $\{|z| = 1\}$ to C whose inverse we denote by $z = \gamma(x)$. Then, the measure σ induces an image measure μ on $|z| = 1$ by $\mu(E) = \sigma(\zeta^{-1}(\gamma^{-1}(E))) = \sigma((\gamma \circ \zeta)^{-1}(E))$, thus

$$\sigma'(s) |d\zeta| = \sigma'(s) |\varphi'(e^{i\theta})| d\theta = \mu'(\theta) d\theta.$$

Let J be the unbounded component of the complement of C and $z = \phi(x)$ the conformal transformation which maps J onto \mathbb{E} so that the points at infinity correspond to each other and $\phi'(\infty) > 0$. Let C_R denote generically the curve $|\phi(x)| = R > 1$ in J .

We define $d\sigma_n = \frac{d\sigma}{|Y_n|^p}$, where $\{Y_n\}$, $n \in \mathbb{N}$ is a sequence of polynomials such that for each n , Y_n has exactly degree n , and all its zeros $(\alpha_{n,i})_{i=1, \dots, n \in \mathbb{N}}$ have no limit points interior to C_A , $A > 1$ and $Y_n(\alpha) = 1$.

We want to study the asymptotic behavior of the polynomials $\tilde{P}_{n,p}$, that solve the following extremal problem:

$$\rho_{n,p} = \inf_{Q_n(\alpha)=1} \|Q_n\|_{p,\sigma_n} = \inf_{Q_n(\alpha)=1} \left\{ \frac{1}{2\pi} \int_C \left| \frac{Q_n(\zeta)}{Y_n(\zeta)} \right|^p d\sigma(s) \right\}^{1/p}. \tag{15}$$

First we quote a result analogous to Theorem 2.

Theorem 6 (See Walsh [18], pp. 252–253). *Let C be a closed rectifiable Jordan curve and let the points $(\alpha_{n,i})_{i=1, \dots, n \in \mathbb{N}}$ be given with no limit points interior to C_A . If f is an analytic function on and within C_T , there exists a sequence r_n of functions of the form (5) such that*

$$\lim_{n \rightarrow \infty} r_n(x) = f(x),$$

uniformly for x on each closed subset of interior of C_R , where $R = \frac{A^2 T + T + 2A}{2AT + A^2 + 1}$.

Using this result we can obtain a theorem analogous to Theorem 1:

Theorem 7. For $0 < p < \infty$, the following statements are equivalent:

- (i) σ satisfies the Szegő condition.
- (ii) The following limit exists and is positive:

$$\lim_{n \rightarrow \infty} \rho_{n,p} > 0.$$

- (iii) There exists $S \in L^p(C, \sigma)$ with $\|S\|_{p,\sigma} \neq 0$, such that

$$\lim_{n \rightarrow \infty} \left\| \frac{\tilde{P}_{n,p}(x)}{Y_n(x)} - S(x) \right\|_{p,\sigma} = 0.$$

- (iv) There exists a function T analytic in B such that

$$\lim_{n \rightarrow \infty} \frac{\frac{\tilde{P}_{n,p}(x)}{Y_n(x)}}{\left\| \frac{\tilde{P}_{n,p}}{Y_n} \right\|_{p,\sigma}} = T(x)$$

holds uniformly on each compact subset of B .

Moreover, if (i) holds, $\lim_{n \rightarrow \infty} \rho_{n,p} = \Delta_p(\sigma, \alpha)$, $S(x) = J_p(\sigma, x)$, and $T(x) = \frac{1}{\Delta_p(\sigma, x)}$, where $\Delta_p(\sigma, x) = D_p(\mu, \gamma(x))$ and $J_p(\sigma, x) = K_p(\mu, \gamma(x))$.

This theorem can also be given for sets of the form $\mathbb{F} = \mathbb{T} \cup \{z_1, z_2, \dots, z_N\}$ with $z_k \in \mathbb{D}$. Let β_n be a varying measure such that $\beta_n = \mu_n + \eta$, where η is a discrete measure with mass $A_k > 0$ at the point z_k , $k = 1, \dots, N$, and $d\mu_n = \frac{d\mu}{|W_n|^p}$ the varying measure with μ and W_n as in Section 1.

Here, we study the asymptotic behavior of the polynomials $T_{n,p}^*(z, \beta_n)$, that solve the extremal problem

$$\lambda_{n,p} = \min_{P_n(0)=1} \left\{ \frac{1}{2\pi} \int_{|z|=1} \left| \frac{P_n(z)}{W_n^*(z)} \right|^p d\mu(\theta) + \sum_{k=1}^N |P_n(z_k)|^p A_k \right\}^{1/p}. \tag{16}$$

Theorem 8. For $0 < p < \infty$ the following statements are equivalent:

- (i) μ satisfies the Szegő condition.
- (ii) The following limit exists and is positive:

$$\lim_{n \rightarrow \infty} \lambda_{n,p} > 0.$$

(iii) There exists a function $S \in H^p(\mu)$ with $\|S\|_{p,\mu} \neq 0$ such that

$$\lim_{n \rightarrow \infty} \left\| \frac{T_{n,p}^*(z, \beta_n)}{W_n^*(z)} - S(z) \right\|_{p,\mu} = 0.$$

(iv) There exists a function T analytic in \mathbb{D} such that

$$\lim_{n \rightarrow \infty} \frac{\frac{T_{n,p}^*(z, \beta_n)}{W_n^*(z)}}{\left\| \frac{T_{n,p}^*(\cdot, \beta_n)}{W_n^*} \right\|_{p,\mu}} = T(z)$$

holds uniformly on each compact subset of \mathbb{D} .

Moreover, if (i) holds

$$\lim_{n \rightarrow \infty} \lambda_{n,p} = D_p(\mu, 0)$$

and the functions in (iii) and (iv) are $S(z) = \frac{D_p(\mu, 0)}{D_p(\mu, z)}$ and $T(z) = \frac{1}{D_p(\mu, z)}$, respectively.

Proof. We will only prove (i) \Leftrightarrow (ii) because the rest of the proof is similar to that given in Theorem 1.

Since $A_k > 0$,

$$\lambda_{n,p} \geq \inf_{Q_n(0)=1} \left\{ \frac{1}{2\pi} \int_{|z|=1} \left| \frac{Q_n(z)}{W_n^*(z)} \right|^p d\mu(\theta) \right\}^{1/p},$$

and from Theorem 1

$$\liminf_{n \rightarrow \infty} \lambda_{n,p} \geq D_p(\mu, 0). \tag{17}$$

Now, let V_N be the polynomial whose zeros are z_1, z_2, \dots, z_N and let $T_{n-N,p}^*$, with $T_{n-N,p}^*(0) = 1$, be the extremal polynomial for the measure $\left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu_n$; that is,

$$\begin{aligned} \lambda_{n,p}^p &\leq \inf_{Q_{n-N}(0)=1} \int_{|z|=1} \left| \frac{Q_{n-N}(z)}{W_n^*(z)} \right|^p \left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu \\ &= \int_{|z|=1} \left| \frac{T_{n-N,p}^*(z)}{W_n^*(z)} \right|^p \left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu. \end{aligned}$$

Using the Theorem 4, we have

$$\lim_{n \rightarrow \infty} \int_{|z|=1} \left| \frac{T_{n-N,p}^*(z)}{W_n^*(z)} \right|^p \left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu = D_p \left(\left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu, 0 \right)^p.$$

From the properties of the Szegő function, we obtain that

$$D_p \left(\left| \frac{V_N(z)}{V_N(0)} \right|^p d\mu, 0 \right) = D_p(\mu, 0)$$

and hence

$$\limsup_{n \rightarrow \infty} \lambda_{n,p} \leq D_p(\mu, 0). \tag{18}$$

The result follows from (17) and (18). \square

References

- [1] M. Bello Hernández, F. Marcellán, J. Mínguez Ceniceros, Pseudo-uniform convexity in H^p and some extremal problems on Sobolev spaces, *Complex Var. Theory Appl.* 48 (5) (2003) 429–440.
- [2] M. Bello Hernández, E. Mina Díaz, Strong asymptotic behavior and weak convergence of polynomials orthogonal on an arc of the unit circle, *J. Approx. Theory* 111 (2) (2001) 233–255.
- [3] B. de la Calle, G. López Lagomasino, Strong asymptotics of orthogonal polynomials with varying measures and Hermite–Padé approximants, *J. Comput. Math.* 99 (1–2) (1998) 91–103.
- [4] Ya.L. Geronimus, On some extremal problems in the space L^p_σ , *Mat. Sbornik NS* 31 (73) (1952) 3–26 (In Russian).
- [5] A.A. Gonchar, G. López Lagomasino, Markov’s theorem for multipoint Padé approximants, *Mat. Sbornik NS* 105(147)(4) (1978), 512–524 (Russian).
- [6] A.A. Gonchar, E.A. Rakhmanov, The equilibrium measure and distribution of zeros of extremal polynomials, *Mat. Sbornik NS* 125(167) 1(9) (1984) 117–127 (Russian).
- [7] A.A. Gonchar, E.A. Rakhmanov, Equilibrium distributions and the rate of rational approximation of analytic functions, *Mat. Sbornik NS* 134(176)(3) (1987), 306–352 (Russian).
- [8] V.A. Kaliaguine, On asymptotics of L_p extremal polynomials on a complex curve ($0 < p < \infty$), *J. Approx. Theory* 74 (2) (1993) 226–236.
- [9] M.V. Keldysh, *Selected Papers*, Academic Press, Moscow, 1985 (Russian).
- [10] G. López Lagomasino, Comparative asymptotics for polynomials that are orthogonal on the real axis, *Mat. Sbornik NS* 137(179)(4) (1988) 500–525 (Russian).
- [11] G. López Lagomasino, Szegő’s theorem for polynomials orthogonal with respect to varying measures, *Orthogonal Polynomials and their Applications* (Segovia, 1986), *Lecture Notes in Math.*, Vol. 1329, Springer, Berlin, 1988, pp. 225–260.
- [12] G. López Lagomasino, E.A. Rakhmanov, Rational approximation, orthogonal polynomials and equilibrium distributions, *Orthogonal Polynomials and their Applications* (Segovia, 1986), *Lecture Notes in Math.*, Vol. 1329, Springer, Berlin, 1988, pp. 125–157.
- [13] E.M. Nikishin, V.N. Sorokin, *Rational Approximations and Orthogonality*, *Translations of Math. Monographs*, Vol. 92, Amer. Math. Soc., Providence, RI, 1991.
- [14] H. Stahl, Strong asymptotics for orthonormal polynomials with varying weights, *Acta Sci. Math. (Szeged)* 66 (2000) 147–192.
- [15] H. Stahl, V. Totik, *General Orthogonal Polynomials*, in: *Encyclopedia of Mathematics*, Vol. 43, Cambridge University Press, Cambridge, 1992.
- [16] G. Szegő, *Orthogonal Polynomials*, Vol. 23, 4th Edition, American Math. Society Colloquium Publications, Amer. Math. Soc., Providence, RI, 1975.

- [17] V. Totik, *Weighted Approximation with Varying Weight*, Lecture Notes in Math. Vol. 1569, Springer, 1994.
- [18] J.L. Walsh, *Interpolation and Approximation by Rational Functions in the Complex Domain*, Vol. XX, 3rd Edition, Amer. Math. Soc. Colloquium Publications, Amer. Math. Society, Providence 6, RI, 1960.