

Linear Inequalities for Flags in Graded Partially Ordered Sets¹

Louis J. Billera²

Department of Mathematics, Cornell University, Ithaca, New York

E-mail: billera@math.cornell.edu

and

Gábor Hetyei³

Mathematics Department, University of Kansas, Lawrence, Kansas

E-mail: hetyei@math.ukans.edu

Communicated by the Managing Editors

Received May 15, 1999

The closure of the convex cone generated by all flag f -vectors of graded partially ordered sets is shown to be polyhedral. In particular, we give the facet inequalities to the polar cone of all nonnegative chain-enumeration functionals on this class of partially ordered sets. These are in one-to-one correspondence with antichains of intervals on the set of ranks and thus are counted by Catalan numbers. Furthermore, we prove that the convolution operation introduced by Kalai assigns extreme rays to pairs of extreme rays in most cases. We describe the strongest possible inequalities for graded partially ordered sets of rank at most 5. © 2000 Academic Press

Key Words: partially ordered set; chain; flag; flag f -vector.

INTRODUCTION

An initial step in obtaining a characterization of f -vectors of some class of objects is to determine the linear equations and inequalities that they must satisfy. The former give a description of the linear span of all such f -vectors, while the latter describe the closure of the convex cone they

¹ Research done while both authors were at the Mathematical Sciences Research Institute in Berkeley, CA, during the academic year 1996–1997. Research at MSRI is supported in part by NSF Grant DMS-9022140.

² Supported in part by NSF Grant DMS-9500581.

³ On leave from the Mathematical Research Institute of the Hungarian Academy of Sciences. Partially supported by Hungarian National Foundation for Scientific Research Grant No. F 023436.

generate. In most cases where this has been done successfully, the description of the linear equations proved to be the more difficult part. Once this was done, and an appropriate basis found for the linear span of all f -vectors (for example, the f -vector in the case of simplicial complexes, the h -vector in the case of Cohen-Macaulay simplicial complexes, or the g -vector in the case of simplicial convex polytopes), the desired cone turned out to be an orthant, that is, only nonnegativity of the basic invariants could be asserted. (See, for example, [7, Theorems II.2.1, II.3.3, and III.1.1].)

The situation of flag f -vectors seems to be quite different. While it is true that flag h -vectors of balanced Cohen-Macaulay complexes (even Cohen-Macaulay graded posets—see [7, Section III.4]) span an orthant, the more basic case of flag f -vectors of all graded posets behaves quite differently. Here, it is the first of these that is simple: there are no equations. However, the cone generated in this case turns out to be quite a bit more complicated. In this paper, we give a description of this cone by giving its minimal generating set. It is already a nontrivial statement that this generating set is finite. Equivalently, we give a finite list of linear inequalities that describe the cone polar to that generated by all flag f -vectors of graded posets. For posets of rank $n + 1$, these inequalities are in one-to-one correspondence with antichains of intervals in the linearly ordered set $\{1, 2, \dots, n\}$, and so are counted by a Catalan number. Thus while the space of flag f -vectors has dimension 2^n , the cone they generate will have on the order of $(2n)^n$ generators.

The proof of the fact that our list of linear inequalities describes a set of generators for the cone of flag f -vectors relies on two ingredients: an explicit construction of sequences of graded posets $P(n, \mathcal{I}, N)$ of rank $n + 1$ yielding the extreme rays of the closure as limits, and an explicit partitioning of the maximal chains of every graded poset P which allows to show the sufficiency of our list of conditions. The chain partitioning used may be generalized to a construction showing that every graded poset satisfies a generalized condition of lexicographic shellability, allowing an explicit description of the order complex of every graded poset. This will be the subject of a subsequent paper [1].

While the description of the extreme rays of the closure of the convex cone generated by the flag f -vectors of graded posets of a given rank is fairly tractable, finding even the number of the facets of the same cone seems to be highly difficult. These represent the strongest possible inequalities holding for flag f -vectors of graded posets of a given rank, and correspond to the extreme rays of the polar cone. In Section 5 we show operations which yield higher-rank extremes from lower rank ones. There are lifting operations, embedding the cones of inequalities into each other as faces, and we give an exact description of those situations where the convolution operation introduced by Kalai in [6] assigns an extreme inequality to a

pair of extreme inequalities. These results allow a short description of the extreme inequalities up to rank 5.

1. PRELIMINARIES

Here we enumerate the basic definitions and results used in this paper.

1.1. Graded Partially Ordered Sets

DEFINITION 1.1. A *graded partially ordered set* (or *graded poset*) P is a finite partially ordered set with a unique minimum element $\hat{0}$, a unique maximum element $\hat{1}$, and a *rank function* $\text{rank}: P \rightarrow \mathbb{N}$ where

- (i) $\text{rank}(\hat{0}) = 0$, and
- (ii) $\text{rank}(y) - \text{rank}(x) = 1$ whenever $y \in P$ covers $x \in P$.

We call $\text{rank}(\hat{1}) \geq 1$ the rank of the poset P . Given a graded poset P of rank $n + 1$, and a subset S of $\{1, 2, \dots, n\}$ we define the *S-rank selected subposet of P* to be the poset

$$P_S := \{x \in P : \text{rank}(x) \in S\} \cup \{\hat{0}, \hat{1}\}.$$

We denote by $f_S(P)$ the number of maximal chains of P_S . Equivalently, $f_S(P)$ is the number of chains $x_1 < \dots < x_{|S|}$ in P such that $\{\text{rank}(x_1), \dots, \text{rank}(x_{|S|})\} = S$. The function

$$f: 2^{\{1, 2, \dots, n\}} \rightarrow \mathbb{N}$$

$$S \mapsto f_S(P)$$

is called the *flag f-vector* of P . Whenever it does not cause confusion we will write $f_{s_1 \dots s_k}$ rather than $f_{\{s_1, \dots, s_k\}}$; in particular, $f_{\{m\}}$ will always be denoted f_m .

A simple example is perhaps helpful here. Consider graded posets of rank 2. In addition to elements $\hat{0}$ and $\hat{1}$ of rank 0 and 2, respectively, such a poset will have only elements of rank 1. For reasons that will become clear later, we wish to consider the closure of the convex cone generated by

$$\{(f_{\emptyset}(P), f_1(P)) \mid \text{rank}(P) = 2\} = \{(1, n) \mid n \geq 1\}.$$

The polar of this cone is generated by the functionals $h_{\emptyset} := f_{\emptyset}$ and $h_1 := f_1 - f_{\emptyset}$.

1.2. The Ring of Chain Operators

We adopt the following terminology and results from [2]. The *chain operators* f_S^{n+1} assign $f_S(P)$ to every graded poset P of rank $n+1$, and zero to all other graded posets. These operators are linearly independent and hence they generate a vector space A_{n+1} of dimension 2^n over \mathbb{R} . We set $A_0 = \mathbb{R}$. The vector space $A := \bigoplus_{n \geq 0} A_n$ may be made into a graded non-commutative ring by introducing the *convolution operation* (first considered by Kalai in [6])

$$f_S^m * f_T^n := f_{S \cup \{m\} \cup (T+m)}^{m+n}$$

for $m, n \geq 1$, and by making the generator 1 of A_0 to be the unit of A . The interest of this convolution operation is the following. (cf. [2, Proposition 1.3])

PROPOSITION 1.2. *The convolution $F * G$ of two nonzero linear combinations of chain operators is nonnegative on all graded posets if and only if both F and G are simultaneously nonnegative or nonpositive on all graded posets.*

According to [2, Proposition 1.1] the chain operators are linearly independent. Moreover, in [2, Section 2] we find the following:

PROPOSITION 1.3. *The ring A is a free graded associative algebra over the set of variables $\{f_\varnothing^n : n \geq 1\}$.*

It follows from [3, Theorem 3] that the semigroup of homogeneous polynomials of a free graded associative algebra has unique factorization. Hence we have the following

PROPOSITION 1.4. *Up to nonzero linear factors, every form $F \in A_n$ may be uniquely written as a product $F = F^{n_1} * \dots * F^{n_k}$ of homogeneous forms $F^{n_i} \in A_{n_i}$ where $n = n_1 + \dots + n_k$ and the F^{n_i} 's cannot be written as a product of two homogeneous forms of lower degree.*

1.3. Blockers of Families of Sets

DEFINITION 1.5. Let \mathcal{S} be an arbitrary family of subsets of a finite set X . A subset $T \subseteq X$ is a *blocker* of \mathcal{S} , if for every $S \in \mathcal{S}$ we have $S \cap T \neq \emptyset$.

We denote the set of blockers of \mathcal{S} in X by $\mathbf{B}_X(\mathcal{S})$. In particular, for $\mathcal{S} = \emptyset$ every subset of X is a blocker and so we have $\mathbf{B}_X(\emptyset) = \{T : T \subseteq X\}$.

LEMMA 1.6. *Let \mathcal{S}_1 and \mathcal{S}_2 be families of subsets of the same finite set X . Then $\mathbf{B}_X(\mathcal{S}_1) \subseteq \mathbf{B}_X(\mathcal{S}_2)$ if and only if every $S_2 \in \mathcal{S}_2$ contains some $S_1 \in \mathcal{S}_1$.*

Proof. If every set from \mathcal{S}_2 contains a set from \mathcal{S}_1 then every blocker of \mathcal{S}_1 also blocks \mathcal{S}_2 . Assume that there is an element S_2 of \mathcal{S}_2 not containing any element of \mathcal{S}_1 . Then the set $X \setminus S_2$ blocks \mathcal{S}_1 , but it is disjoint from S_2 and so does not block \mathcal{S}_2 . Hence we have $\mathbf{B}_X(\mathcal{S}_1) \not\subseteq \mathbf{B}_X(\mathcal{S}_2)$. ■

COROLLARY 1.7. *Let \mathcal{S}_1 and \mathcal{S}_2 be families of subsets of the same finite set X . Then $\mathbf{B}_X(\mathcal{S}_1) = \mathbf{B}_X(\mathcal{S}_2)$ if and only if the minimal sets with respect to inclusion are the same in \mathcal{S}_1 and \mathcal{S}_2 .*

The minimal sets in a family of sets form an *antichain* (or Sperner family, or clutter), i.e., a family of sets such that no set contains another one. On the other hand, a family of sets \mathcal{S} on a set X is a *dual ideal* if for every $S \in \mathcal{S}$ all $S' \subseteq X$ containing S belong to \mathcal{S} . For an arbitrary family of sets \mathcal{S} on X the *dual ideal generated by \mathcal{S}* is the family

$$\mathcal{S}^+ := \{T \subseteq X : \exists S \in \mathcal{S} (S \subseteq T)\}.$$

PROPOSITION 1.8. *For any family of sets \mathcal{S} on a set X , we have $\mathbf{B}_X(\mathcal{S}) = \mathbf{B}_X(\mathcal{S}^+)$.*

Proof. Since \mathcal{S} is a subfamily of \mathcal{S}^+ , we have $\mathbf{B}_X(\mathcal{S}) \supseteq \mathbf{B}_X(\mathcal{S}^+)$. The other inclusion follows from Lemma 1.6 using the fact that every set from \mathcal{S}^+ contains some set from \mathcal{S} .

It is shown in [4] that whenever \mathcal{S} is an antichain, then the family of minimal elements of $\mathbf{B}_X(\mathbf{B}_X(\mathcal{S}))$ is \mathcal{S} . This statement and repeated application of Proposition 1.8 yields

$$\mathbf{B}_X(\mathbf{B}_X(\mathcal{S})) = \mathcal{S}^+$$

for every family of sets \mathcal{S} . From this and Proposition 1.8 we have the following.

PROPOSITION 1.9. *Let \mathcal{S}_1 and \mathcal{S}_2 be families of sets on the same set X . Then*

$$\mathbf{B}_X(\mathcal{S}_1) = \mathbf{B}_X(\mathcal{S}_2) \quad \text{if and only if} \quad \mathcal{S}_1^+ = \mathcal{S}_2^+.$$

For more on blockers, we refer the reader to [4], or to Section 8.1 of [5].

2. THE MAIN THEOREM

This section contains our main result, which characterizes all linear inequalities holding for the flag f -vectors of all graded posets of rank $n + 1$. Every such linear inequality may be written as

$$F(P) := \sum_{S \subseteq \{1, \dots, n\}} a_S \cdot f_S^{n+1}(P) \geq 0,$$

where the coefficients a_S are real numbers. Moreover, $\sum_{S \subseteq \{1, \dots, n\}} a_S \cdot f_S^{n+1}(P)$ is zero if $\text{rank}(P) \neq n + 1$. Hence we are interested in determining the subset

$$K_{n+1} := \left\{ F = \sum_{S \subseteq \{1, \dots, n\}} a_S \cdot f_S^{n+1} \in A_{n+1} : \forall P (F(P) \geq 0) \right\}$$

of A_{n+1} . For convenience, we will let $K_0 = A_0$.

In this section we show that K_{n+1} is a polyhedral cone, that is, the intersection of finitely many half spaces. We give these half spaces in terms of *interval systems* on the linearly ordered set $\{1, 2, \dots, n\} \subset \mathbb{N}$.

A subset of a partially ordered set P is an *interval*, if it is empty, or of the form

$$[p, q] := \{x \in P : p \leq x \leq q\} \quad \text{for some } p, q \in P.$$

In particular, an *interval* in \mathbb{N} is a finite (possibly empty) set of consecutive natural numbers. An *interval system on a partially ordered set P* is a family \mathcal{I} of nonempty intervals. We consider the empty family also as an interval system. The following theorem gives the list of inequalities that determine K_{n+1} .

THEOREM 2.1. *An expression $\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1}$ is nonnegative on all graded posets of rank $n + 1$ if and only if we have*

$$\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0 \quad \text{for every interval system } \mathcal{I} \text{ on } [1, n]. \quad (1)$$

First we will show the necessity of condition (1) by constructing for every interval system \mathcal{I} on $\{1, 2, \dots, n\}$ a family of posets $\{P(n, \mathcal{I}, N) : N \in \mathbb{N}\}$ of rank $n + 1$ such that we have

$$\lim_{N \rightarrow \infty} \frac{1}{f_{[1, n]}(P(n, \mathcal{I}, N))} \cdot \sum_{S \subseteq [1, n]} a_S \cdot f_S(P(n, \mathcal{I}, N)) = \sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S.$$

Then we will prove the sufficiency by using an appropriate partitioning of the set of maximal chains for every graded poset P .

DEFINITION 2.2. Let n and N be positive integers, and let $\mathcal{I} = \{I_1, I_2, \dots, I_k\}$ be an interval system on $[1, n]$. We define the elements of the poset $P(n, \mathcal{I}, N)$ to be all arrays $(i; p_1, \dots, p_k)$ such that

- (i) $i \in [0, n + 1]$, and
- (ii) for every $j \in [1, k]$ we have

$$p_j \in \begin{cases} [1, N] & \text{whenever } i \in I_j, \\ \{*\} & \text{otherwise} \end{cases}$$

Here $*$ is a special symbol, different from all integers.

We set $(i; p_1, p_2, \dots, p_k) \leq (i'; p'_1, p'_2, \dots, p'_k)$ if

- (1) $i \leq i'$, and
- (2) for every $j \in [1, k]$ we have either $p_j = p'_j$ or $*$ $\in \{p_j, p'_j\}$.

Observe that $P(n, \mathcal{I}, N)$ has a unique minimum element $\hat{0} = (0; *, \dots, *)$ and a unique maximum element $\hat{1} = (n + 1; *, \dots, *)$.

EXAMPLE 2.3. Let $n = 3$, $\mathcal{I} = \{[1, 2], [2, 3]\}$, and $N = 2$. Then $P(3, \{[1, 2], [2, 3]\}, 2)$ is the poset shown on Fig. 1.

EXAMPLE 2.4. If $\mathcal{I} = \emptyset$ then $P(n, \emptyset, N)$ is the same chain $\hat{0} = 0 < 1 < 2 \dots < n < n + 1 = \hat{1}$ for every positive integer N .




FIG. 1. $P(3, \{[1, 2], [2, 3]\}, 2)$.

PROPOSITION 2.5. *Let n and N be positive integers and $\mathcal{I} := \{I_1, \dots, I_k\}$ a nonempty interval system on $[1, n]$. Then $P(n, \mathcal{I}, N)$ is a graded poset of rank $n + 1$ and we have*

$$f_S(P(n, \mathcal{I}, N)) = N^{|\{j \in [1, k] : S \cap I_j \neq \emptyset\}|} \quad \text{for all } S \subseteq [1, n].$$

Proof. First we show that P is graded with the rank function given by

$$\text{rank}((i; p_1, \dots, p_k)) = i.$$

Obviously, for every $(i; p_1, \dots, p_k) < (j; q_1, \dots, q_k)$ we have $i < j$. We only need to show that whenever $i + 1 < j$ also holds then there is an element $(i + 1; p'_1, \dots, p'_k) \in P(n, \mathcal{I}, N)$ strictly between $(i; p_1, \dots, p_k)$ and $(j; q_1, \dots, q_k)$. Let us set

$$p'_i = \begin{cases} * & \text{if } i + 1 \notin I_l, \\ p_l & \text{if } i + 1 \in I_l \text{ and } i \in I_l, \\ q_l & \text{if } i + 1 \in I_l \text{ and } j \in I_l, \\ \text{an arbitrary element of } \{1, 2, \dots, N\} & \text{if } i + 1 \in I_l \text{ and } i, j \notin I_l. \end{cases}$$

Observe that whenever we have $\{i, j\} \subseteq I_l$ we also have $i + 1 \in I_l$ and $p_l = q_l$, hence there is no contradiction in this definition of p'_i . It is easy to verify that $(i + 1; p'_1, \dots, p'_k) \in P(n, \mathcal{I}, N)$ is strictly between $(i; p_1, \dots, p_k)$ and $(j; q_1, \dots, q_k)$.

Next we compute $f_S(P(n, \mathcal{I}, N))$, which is, by definition, the number of chains $x_1 < \dots < x_{|S|}$ satisfying $\{\text{rank}(x_1), \dots, \text{rank}(x_{|S|})\} = S$. If for some $j \in [1, k]$ we have $S \cap I_j = \emptyset$ then every element $x_t = (i, p_1, \dots, p_k)$ of such a chain must satisfy $p_j = *$. If $S \cap I_j \neq \emptyset$ then for the elements $x_t = (i, p_1, \dots, p_k)$ satisfying $\text{rank}(x_t) \in S \cap I_j$ we must have $p_j \in \{1, 2, \dots, N\}$, and by the definition of the partial order on $P(n, \mathcal{I}, N)$ the value of p_j must be the same for all such x_t 's. All other x_t 's must satisfy $p_j = *$. Conversely, let us fix a vector (q_1, \dots, q_k) such that q_j is an arbitrary element of $\{1, 2, \dots, N\}$ whenever $S \cap I_j \neq \emptyset$ and $q_j = *$ otherwise. Then the set $\{(s; p(s)_1, \dots, p(s)_k) : s \in S\}$ defined by

$$p(s)_j = \begin{cases} q_j & \text{if } s \in I_j, \\ * & \text{otherwise} \end{cases}$$

is a chain in $P(n, \mathcal{I}, N)$ satisfying $\{\text{rank}((s; p(s)_1, \dots, p(s)_k)) : s \in S\} = S$. This shows that $f_S(P(n, \mathcal{I}, N))$ equals to the number of possible choices of (q_1, \dots, q_k) , i.e., $N^{|\{j : S \cap I_j \neq \emptyset\}|}$. ■

COROLLARY 2.6. *Let n be a positive integer, and \mathcal{I} an interval system on $[1, n]$. Then we have*

$$\lim_{N \rightarrow \infty} \frac{1}{f_{[1, n]}(P(n, \mathcal{I}, N))} \cdot \sum_{S \subseteq [1, n]} a_S \cdot f_S(P(n, \mathcal{I}, N)) = \sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S.$$

In fact, for a non-empty interval system $\mathcal{I} = \{I_1, \dots, I_k\}$ we have

$$\begin{aligned} & \frac{1}{f_{[1, n]}(P(n, \mathcal{I}, N))} \cdot \sum_{S \subseteq [1, n]} a_S \cdot f_S(P(n, \mathcal{I}, N)) \\ &= \frac{1}{N^k} \cdot \sum_{S \subseteq [1, n]} a_S \cdot N^{|\{j: S \cap I_j \neq \emptyset\}|}. \end{aligned}$$

In this expression a_S is multiplied by 1 if and only if S intersects every interval of the system \mathcal{I} , otherwise it is multiplied by a negative power of N . Finally, when $\mathcal{I} = \emptyset$ then, as it was observed in Example 2.4, every poset $P(n, \emptyset, N)$ is a chain of rank $n + 1$, and we have

$$\frac{1}{f_{[1, n]}(P(n, \emptyset, N))} \cdot \sum_{S \subseteq [1, 2, \dots, n]} a_S \cdot f_S(P(n, \emptyset, N)) = \sum_{S \subseteq [1, n]} a_S$$

immediately. We can conclude that condition (1) is necessary:

COROLLARY 2.7. *Suppose $\sum_{S \subseteq [1, n]} a_S \cdot f_S(P) \geq 0$ for every graded poset of rank $n + 1$. Then for every interval system \mathcal{I} on $[1, n]$,*

$$\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0.$$

Hence we are left to show the sufficiency of (1).

PROPOSITION 2.8. *Assume that the set of coefficients $\{a_S: S \subseteq [1, n]\}$ satisfies condition (1). Then we have*

$$\sum_{S \subseteq [1, n]} a_S \cdot f_S(P) \geq 0$$

for every graded poset of rank $n + 1$.

Proof. Let P be a graded poset of rank $n + 1$. For every $i \in [1, n]$ let us fix an arbitrary numbering of the elements of rank i . Given an interval $[p, q]$ of P , let $\phi([p, q])$ denote the first atom in $[p, q]$. (Note that all atoms of $[p, q]$ have the same rank, namely $\text{rank}(p) + 1$.) We will need the following two elementary observations:

1. If y covers x then $\phi([x, y]) = y$.
2. If $p \in [x, y] \subseteq [x, z]$ and $p = \phi([x, z])$ then $p = \phi([x, y])$.

For every $S \subseteq [1, n]$ we define an operation $M_S: [1, n] \rightarrow [1, n]$ by

$$M_S(i) := \min\{j \in [i, n+1] : j \in S \cup \{n+1\}\}.$$

In other words, M_S assigns to i the smallest element of S which is not less than i , if such an element exists. Otherwise, it assigns $n+1$ to i .

Consider the set of maximal chains

$$F_S := \{\hat{0} = p_0 < p_1 < \cdots < p_n < p_{n+1} = \hat{1} : \\ \forall i \in [1, n](p_i = \phi([p_{i-1}, p_{M_S(i)}]))\}.$$

We claim that F_S contains exactly $f_S(P)$ elements. For every $i \in S$ we have $M_S(i) = i$ and so, by our second elementary observation $p_i = \phi([p_{i-1}, p_{M_S(i)}]) = \phi([p_{i-1}, p_i])$ is trivially satisfied. Hence it is sufficient to show that every chain $\{p_s : s \in S\}$ satisfying $\{\text{rank}(p_s) : s \in S\} = S$ may be uniquely extended to a maximal chain $\{p_0, p_1, \dots, p_n\} \in F_S$. The only possible choice for p_0 is $\hat{0}$. Assume by induction that we have found a unique possible value for p_0, \dots, p_m . Let i be the smallest rank above m such that $i \notin S$. Then the only possible value of p_i is $\phi([p_{i-1}, p_{M_S(i)}])$, and choosing this value we obtain that $\{p_1, \dots, p_i\} \cup \{p_s : s \in S\}$ is a chain. This recursive algorithm shows that we have at most one extension of $\{p_s : s \in S\}$ to a maximal chain in F_S . On the other hand, at the end of the algorithm we obtain a maximal chain belonging to F_S , since all defining conditions are satisfied. In particular, we obtain that $F_{[1, n]}$ contains all maximal chains, while F_\emptyset contains the unique maximal chain for which every p_i ($i = 1, 2, \dots, n$) is the first element among all elements of rank i covering p_{i-1} .

Let us fix now a maximal chain $C := \{\hat{0} = p_0 < p_1 < \cdots < p_n < p_{n+1} = \hat{1}\}$, and determine all those sets of ranks $S \subseteq [1, n]$ for which C belongs to F_S . In view of our two observations, for every $i \in [1, n]$ there is a largest $j \in [i, n+1]$ such that $p_i = \phi([p_{i-1}, p_j])$. Let us denote this largest j by $\psi(C, i)$. Obviously, $p_i = \phi([p_{i-1}, p_{M_S(i)}])$ is satisfied if and only if we have $M_S(i) \leq \psi(C, i)$, or equivalently

$$(S \cup \{n+1\}) \cap [i, \psi(C, i)] \neq \emptyset.$$

Introducing

$$\mathcal{I}_C := \{[i, \psi(C, i)] : i \in [1, n], \psi(C, i) \neq n+1\}$$

we may say that $C \in F_S$ if and only if S blocks the system \mathcal{I}_C .

For every $S \subseteq [1, n]$, let us put a weight a_S on every $C \in F_S$. On the one hand, the sum of all weights put on the chains is $\sum_{S \subseteq [1, n]} a_S \cdot f_S(P)$. On the other hand, the total weight associated to an individual chain C is $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I}_C)} a_S$. Hence we obtain

$$\sum_{S \subseteq [1, n]} a_S \cdot f_S(P) = \sum_{C \in F_{[1, n]}} \sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I}_C)} a_S,$$

and so $\sum_{S \subseteq [1, n]} a_S \cdot f_S(P)$ is a sum of nonnegative terms, if (1) is satisfied. \blacksquare

3. THE FACETS OF THE CONE K_{n+1}

In this section we show that the inequalities in (1) give facets of the cone K_{n+1} . We also show that the number of facets is a Catalan number.

PROPOSITION 3.1. *Every condition of the form $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0$ defines a facet of the cone K_{n+1} .⁴*

Proof. Assume by way of contradiction that $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0$ is not a facet. Then, by Farkas' lemma, there exists nonnegative numbers c_1, \dots, c_k and interval systems $\mathcal{I}_1, \dots, \mathcal{I}_k$ such that we have

$$\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S = \sum_{i=1}^k c_i \cdot \sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I}_i)} a_S,$$

and the conditions $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I}_i)} a_S \geq 0$ are different from $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0$. Since the set $[1, n]$ blocks every interval system, the coefficient of $a_{[1, n]}$ is 1 on the left hand side and $\sum_{i=1}^k c_i$ on the right hand side. Hence we must have

$$\sum_{i=1}^k c_i = 1.$$

The coefficient of every other a_S is zero or one on the left hand side, and a convex combination of zeros and ones on the right hand side. Hence we must have

$$\mathbf{B}_{[1, n]}(\mathcal{I}) = \mathbf{B}_{[1, n]}(\mathcal{I}_1) = \mathbf{B}_{[1, n]}(\mathcal{I}_2) = \dots = \mathbf{B}_{[1, n]}(\mathcal{I}_k),$$

contradicting our assumption. \blacksquare

⁴ It is because of this Proposition that we decided to exclude $\mathcal{I} = \{\emptyset\}$ from the definition of interval systems. When \mathcal{I} contains the empty set, then no subset of $[1, n]$ blocks \mathcal{I} and $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0$ reduces to $0 \geq 0$, a true but trivial inequality.

More directly, Proposition 3.1 follows from the fact that a subset of vertices of the 2^n -dimensional cube are vertices of their convex hull.

The number of facets is thus equal to the number of families of sets of the form $\mathbf{B}_{[1,n]}(\mathcal{I})$ where \mathcal{I} is an arbitrary interval system on $[1,n]$. In order to count the number of these families, observe that by Proposition 1.9 we may replace \mathcal{I} by \mathcal{I}^+ . Hence the number of facets of the cone K_{n+1} is equal to the number of dual ideals of subsets of $[1,n]$ generated by intervals. There is a bijection between such dual ideals and Ferrers shapes contained in the shape of the partition $(n, n-1, \dots, 1)$, defined as follows. Given an $n \times n$ square, write the interval $[i, j] \subseteq [1, n]$ into the box in the j th row and i th column. The boxes into which we have written an interval form the Ferrers shape of the partition $(n, n-1, \dots, 1)$. Clearly an interval system \mathcal{I} on $[1, n]$ is the family of all intervals in a dual ideal, if and only if for every box representing an interval $I \in \mathcal{I}$ all boxes above and to the left are marked with an interval from \mathcal{I} . Equivalently, the boxes representing \mathcal{I} form a Ferrers shape. Figure 2 shows the Ferrers shape representation for the dual ideal $\{[1, 2], [2, 3], [4, 4]\}^+$ on $[1, 4]$.

The number of Ferrers shapes contained in the shape of the partition $(n, n-1, \dots, 1)$ is equal to the number of those lattice paths from the lower left corner of the box marked with $[1, 1]$ to the upper right corner of the box marked with $[n, n]$ which use only unit steps up and to the right and which never leave the shape of the partition $(n, n-1, \dots, 1)$.

COROLLARY 3.2. *The number of facets of the cone K_{n+1} is the Catalan number $(1/(n+2))\binom{2(n+1)}{n+1}$.*

Let us also note the following consequence of Corollary 1.7 which gives a description of the facets of K_{n+1} which is more useful in practice.




FIG. 2. Representation of the dual ideal $\{[1, 2], [2, 3], [4, 4]\}^+$ on $[1, 4]$.

TABLE 1

The value of $\sum_{S \in \mathbf{B}_{[1,3]}(\mathcal{I})} a_S$ for $\sum_{S \subseteq [1,3]} a_S \cdot f_S = f_{13} - f_1 + f_2 - f_3$

\mathcal{I}	$\sum_{S \in \mathbf{B}_{[1,3]}(\mathcal{I})} a_S$	\mathcal{I}	$\sum_{S \in \mathbf{B}_{[1,3]}(\mathcal{I})} a_S$
\emptyset	0	$\{[1], [2, 3]\}^+$	1
$\{[1, 3]\}^+$	0	$\{[2]\}^+$	1
$\{[1, 2]\}^+$	1	$\{[1, 2], [3]\}^+$	1
$\{[2, 3]\}^+$	1	$\{[1], [2]\}^+$	0
$\{[1]\}^+$	0	$\{[1], [3]\}^+$	1
$\{[1, 2], [2, 3]\}^+$	2	$\{[2], [3]\}^+$	0
$\{[3]\}^+$	0	$\{[1], [2], [3]\}^+$	0

COROLLARY 3.3. *Every inequality in (1) may be written as*

$$\sum_{S \in \mathbf{B}_{[1,n]}(\mathcal{I})} a_S \geq 0$$

for some antichain of intervals \mathcal{I} on $[1, n]$.

EXAMPLE 3.4. This inequality was found by Billera and Liu (see [2]). For every graded poset P of rank 4 we have

$$f_{13}(P) - f_1(P) + f_2(P) - f_3(P) \geq 0.$$

In fact, we may apply Theorem 2.1 for $a_{13} = 1$, $a_1 = -1$, $a_2 = 1$, $a_3 = -1$, and $a_S = 0$ for every other subset S of $\{1, 2, 3\}$. There are 14 dual ideals generated by intervals on the set $[1, 3]$, which are given by the antichain of their minimal intervals in Table 1. Evaluating $\sum_{S \in \mathbf{B}_{[1,3]}(\mathcal{I})} a_S$ for these 14 interval systems we see that Condition (1) is satisfied.

4. PROJECTIONS AND CONVOLUTIONS

In this section we present linear projections $\pi_m^{n+1}: A_{n+1} \rightarrow A_n$ which allow us to describe the cone K_{n+1} in terms of the cone K_n . We show that these projections are nicely compatible with the convolution operation defined in [2].

DEFINITION 4.1. Let $n > 0$ and $m \in [0, n]$ be integers. For an arbitrary form $\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ we define its m th projection into A_n by

$$\begin{aligned} & \pi_m^{n+1} \left(\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \right) \\ & := \sum_{S \subseteq [1, n-1]} (\chi((S \cup \{0\}) \cap [m, n-1]) \neq \emptyset) \cdot a_S + a_{S \cup \{n\}} \cdot f_S^n. \end{aligned}$$

We extend this definition to $n=0$ and to negative m 's by setting $\pi_0^1(f_\emptyset^1) := 1$ and $\pi_m^{n+1} := \pi_0^{n+1}$ whenever $m < 0$.

Equivalently, the effect of the projections π_m^{n+1} on the operators f_S^{n+1} may be given by

$$\pi_m^{n+1}(f_S^{n+1}) = \begin{cases} f_{S \setminus \{n\}}^n & \text{if } n \in S, \\ \chi((S \cup \{0\}) \cap [m, n-1]) \neq \emptyset \cdot f_S^n & \text{if } n \notin S. \end{cases} \quad (2)$$

In particular, for $m=0$, $(S \cup \{0\}) \cap [0, n-1] \neq \emptyset$ holds for every $S \subseteq [1, n-1]$, and we have

$$\pi_0^{n+1} \left(\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \right) = \sum_{S \subseteq [1, n-1]} (a_S + a_{S \cup \{n\}}) \cdot f_S^n.$$

and

$$\pi_0^{n+1}(f_S^{n+1}) = f_{S \setminus \{n\}}^n. \quad (3)$$

At the other extreme, for $m=n$ the interval $[m, n-1]$ is the empty set, and we have

$$\pi_n^{n+1} \left(\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \right) = \sum_{S \subseteq [1, n-1]} a_{S \cup \{n\}} \cdot f_S^n.$$

Given an arbitrary interval system \mathcal{I} on $[1, n-1]$ and $m \in [1, n]$, a set $T \subseteq [1, n]$ blocks $\mathcal{I} \cup \{[m, n]\}$ if and only if $S := T \setminus \{n\} \subseteq [1, n-1]$ blocks \mathcal{I} and either $n \in T$, i.e., $T = S \cup \{n\}$, or $n \notin T$ and $S = T$ blocks $[m, n-1]$. Moreover, $(S \cup \{0\}) \cap [m, n-1] \neq \emptyset$ is equivalent to $S \cap [m, n-1] \neq \emptyset$. Hence we have

$$\begin{aligned} & \sum_{T \in \mathbf{B}_{[1, n]}(\mathcal{I} \cup \{[m, n]\})} a_T \\ & = \sum_{S \in \mathbf{B}_{[1, n-1]}(\mathcal{I})} (\chi((S \cup \{0\}) \cap [m, n-1]) \neq \emptyset) \cdot a_S + a_{S \cup \{n\}}. \end{aligned} \quad (4)$$

Similarly, for $m = 0$, a set $T \subseteq [1, n]$ blocks \mathcal{J} if and only if $S := T \setminus \{n\} \subseteq [1, n-1]$ blocks \mathcal{J} . Moreover, as noted earlier, $\chi((S \cup \{0\}) \cap [0, n-1] \neq \emptyset) = 1$ for every $S \subseteq [1, n-1]$. These observations yield

$$\sum_{T \in \mathbf{B}_{[1, n]}(\mathcal{J})} a_T = \sum_{S \in \mathbf{B}_{[1, n-1]}(\mathcal{J})} (\chi((S \cup \{0\}) \cap [0, n-1] \neq \emptyset) \cdot a_S + a_{S \cup \{n\}}). \tag{5}$$

Using Eqs. (4) and (5) we may show the following.

THEOREM 4.2. *A form $\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ belongs to K_{n+1} if and only if for every $m \in [0, n]$ the projection $\pi_m^{n+1}(\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1})$ belongs to K_n .*

Proof. The necessity is evident in view of of the Eqs. (4) and (5). To prove sufficiency observe that by Corollary 3.3 it suffices to verify the nonnegativity conditions

$$\sum_{T \in \mathbf{B}_{[1, n]}(\mathcal{J})} a_T \geq 0 \tag{6}$$

for every antichain of intervals \mathcal{J} on $[1, n]$. Such an antichain is either also an interval system on $[1, n-1]$ or the union of the antichain $\mathcal{J} := \{I \in \mathcal{J} : I \subseteq [1, n-1]\}$ and of the singleton $\{[m, n]\}$ where $[m, n]$ is the unique interval in \mathcal{J} containing n . In either case, we may use (5) or (4), respectively, to show that (6) holds. ■

We now introduce some linear operators acting on A which will be useful in giving a simple expression for π_m^{n+1} for $m > 0$.

DEFINITION 4.3. Let n be a positive integer and $k \in [0, n-1]$. We define $\rho_k^{n+1} : A_{n+1} \rightarrow A_n$ by setting

$$\rho_k^{n+1}(f_S^{n+1}) := \chi(S \subseteq [1, k]) \cdot f_{S \cap [1, k]}^n$$

for every $S \subseteq [1, n]$. We extend this definition to $n = 0$ and to negative k 's by setting $\rho_0^1(f_\emptyset^1) := 0$ and $\rho_k^{n+1} := 0$ whenever $k < 0$.

In particular, for $n > 0$ we have

$$\rho_0^{n+1}(f_S^{n+1}) = \begin{cases} f_\emptyset^n & \text{if } S = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

It is an easy consequence of (2) that we have

$$\pi_m^{n+1} = \pi_0^{n+1} - \rho_{m-1}^{n+1} \quad \text{for } n \in \mathbb{N} \quad \text{and} \quad m \in [0, n]. \tag{7}$$

Let us consider now the effect of the projection operations π_m^{n+1} on a convolution of two chain operators.

PROPOSITION 4.4. *We have*

$$\pi_0^{m+n}(f_S^m * f_T^n) = f_S^m * \pi_0^n(f_T^n)$$

for all positive m, n and sets $S \subseteq [1, m-1]$, $T \subseteq [1, n-1]$.

Proof. Assume first that we have $n \geq 2$. A simple substitution into the definitions and (3) yields

$$\begin{aligned} \pi_0^{m+n}(f_S^m * f_T^n) &= \pi_0^{m+n}(f_{S \cup \{m\} \cup (T+m)}^{m+n}) = f_{S \cup \{m\} \cup (T+m) \setminus \{m+n-1\}}^{m+n-1} \\ &= f_{S \cup \{m\} \cup (T \setminus \{n-1\}) + m}^{m+n-1} = f_S^m * \pi_0^n(f_T^n). \end{aligned}$$

For $n = 1$ we have

$$\begin{aligned} \pi_0^{m+1}(f_S^m * f_\emptyset^1) &= \pi_0^{m+1}(f_{S \cup \{m\}}^{m+1}) = f_{S \cup \{m\} \setminus \{m\}}^m \\ &= f_S^m = f_S^m * 1 = f_S^m * \pi_0^1(f_\emptyset^1) \end{aligned}$$

by $\pi_0^1(f_\emptyset^1) = 1$. ■

PROPOSITION 4.5. *Let m and n be positive integers, $S \subseteq [1, m-1]$, $T \subseteq [1, n-1]$. Then for every $k \in [0, m+n-2]$ we have*

$$\rho_k^{m+n}(f_S^m * f_T^n) = f_S^m * \rho_{k-m}^n(f_T^n).$$

Proof. If k is negative then both sides are identically zero. Assume $0 \leq k \leq m-1$. Then we have

$$\begin{aligned} \rho_k^{m+n}(f_S^m * f_T^n) &= \rho_k^{m+n}(f_{S \cup \{m\} \cup (T+m)}^{m+n}) \\ &= \chi(S \cup \{m\} \cup (T+m) \subseteq [1, k]) \\ &\quad \cdot f_{(S \cup \{m\} \cup (T+m)) \cap [1, k]}^{m+n-1} = 0 \end{aligned}$$

since $m \notin [1, k]$. On the other hand, ρ_{k-m}^n is identically zero by definition, and so the equality of the both sides holds trivially.

Assume finally that $m \leq k \leq m+n-2$ holds. Then we must have $n \geq 2$, S is a subset of $[1, k]$ and $S \cup \{m\} \cup (T+m) \subseteq [1, k]$ is equivalent to $T \subseteq [1, k-m]$. Therefore we have

$$\begin{aligned}
 \rho_k^{m+n}(f_S^m * f_T^n) &= \rho_k^{m+n}(f_{S \cup \{m\} \cup (T+m)}^{m+n}) \\
 &= \chi(S \cup \{m\} \cup (T+m) \subseteq [1, k]) \cdot f_{(S \cup \{m\} \cup (T+m)) \cap [1, k]}^{m+n-1} \\
 &= \chi(T \subseteq [1, k-m]) \cdot f_{S \cup \{m\} \cup (T \cap [1, k-m] + m)}^{m+n-1} \\
 &= \chi(T \subseteq [1, k-m]) \cdot f_S^m * f_{T \cap [1, k-m]}^{n-1} \\
 &= f_S^m * \rho_{k-m}^n(f_T^n). \quad \blacksquare
 \end{aligned}$$

As a corollary of Propositions 4.4 and 4.5, and of Eq. (7) we obtain

COROLLARY 4.6. *The equality*

$$\pi_k^{m+n}(f_S^m * f_T^n) = f_S^m * \pi_{k-m}^n(f_T^n)$$

holds for $m, n > 0$, $k \in [0, m+n-1]$, $S \subseteq [1, m-1]$, and $T \subseteq [1, n-1]$.

In Section 5 we will need to consider the maximum element of the *support* of a form $\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1}$.

DEFINITION 4.7. The *support* of a form $F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ is the set

$$\text{supp}(F) := \bigcup_{\substack{S \subseteq [1, n] \\ a_S \neq 0}} S.$$

We call the maximum element of $\text{supp}(F)$ the *largest letter occurring in F*.

LEMMA 4.8. *The projections π_m^{n+1} do not increase the largest letter occurring in a form.*

Proof. Let us denote the largest occurring letter in $F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ by l . If $l = n$ then there is nothing to prove, so we may assume $l < n$. Evidently $k \in [1, n]$ is greater than l if and only if for every $S \subseteq [1, n]$ containing k we have $a_S = 0$. Hence it is sufficient to show the following: if for every $S \subseteq [1, n]$ containing some $k > l$ we have $a_S = 0$ then for every $S \subseteq [1, n-1]$ containing some $k > l$ we have

$$\chi((S \cup \{0\}) \cap [m, n-1] \neq \emptyset) \cdot a_S + a_{S \cup \{n\}} = 0.$$

The second term in this sum is zero since $S \cup \{n\}$ contains the letter n which larger than l by assumption. The first term is zero too, since S contains some letter $k > l$. \blacksquare

5. EXTREME RAYS OF K_{n+1}

In this section we study the extreme rays of the cones K_{n+1} where $n \in \mathbb{N}$. First we show that f_{\emptyset}^{n+1} is an extreme ray of the cone K_{n+1} for every $n \geq 0$. Then we show for $k = 1, 2, \dots, n$ that the inclusions

$$\begin{aligned} \sigma_k^{n+1}: A_{n+1} &\rightarrow A_{n+2} \\ f_S^{n+1} &\mapsto f_{\sigma_k(S)}^{n+2} \end{aligned} \tag{8}$$

induced by the *shift operators*

$$\begin{aligned} \sigma_k: \mathbb{P} &\rightarrow \mathbb{P} \\ i &\mapsto \begin{cases} i & \text{if } i \leq k \\ i+1 & \text{if } i \geq k+1 \end{cases} \end{aligned} \tag{9}$$

embed K_{n+1} into A_{n+2} as a face of K_{n+2} . Finally we describe completely those situations when the convolution $F * G$ of an extreme ray F of the cone K_m and an extreme ray G of the cone K_n is an extreme ray of the cone K_{m+n} . The main result of this section is the following theorem.

THEOREM 5.1. *Let $F \in K_m$ and $G \in K_n$ be extreme rays in their respective cones. Then the convolution $F * G \in K_{m+n}$ is an extreme ray, unless $F = F' * f_{\emptyset}^k$ and $G = f_{\emptyset}^l * G'$ for some $k \leq m$, $l \leq n$, $F' \in K_{m-k}$, and $G' \in K_{n-l}$.*

To interpret this result, note that Proposition 1.2 is equivalent to the following.

PROPOSITION 5.2. *Let $F \in A_m$ and $G \in A_n$. The convolution $F * G \in A_{m+n}$ belongs to K_{m+n} if and only if exactly one of the following holds:*

- (i) $F \in K_m$ and $G \in K_n$, or
- (ii) $-F \in K_m$ and $-G \in K_n$.

Thus the convolution of two extreme rays is surely in the cone of valid inequalities. If, say, G is a positive linear combination of G' and G'' from K_n then $F * G$ will be the positive linear combination of $F * G'$ and $F * G''$ from K_{m+n} . (Recall that according to Proposition 1.3 the ring A is a free associative algebra, and so it has no zero divisors.) Thus only the convolution of extreme rays may yield an extreme ray. It may happen that the convolution of extremes is not extreme: for every $m, n > 0$ the operators f_{\emptyset}^m and f_{\emptyset}^n are extreme rays in K_m and K_n respectively, yet

$$f_{\emptyset}^m * f_{\emptyset}^n = f_m^{m+n} = (f_m^{m+n} - f_{\emptyset}^{m+n}) + f_{\emptyset}^{m+n}$$

where both $f_m^{m+n} - f_{\emptyset}^{m+n}$ and f_{\emptyset}^{m+n} belong to K_{m+n} . Theorem 5.1 affirms that, essentially, only such anomalies may occur.

According to Proposition 1.4 the semigroup of homogeneous polynomials of A has unique factorization. In view of Proposition 5.2, an expression $F \in A_n$ belongs to K_n if and only if every factor $F_i \in A_{n_i}$ in its complete homogeneous factorization $F = F_1 * \dots * F_k$ may be chosen to belong to K_{n_i} . (Since uniqueness holds up to a choice of nonzero constant factors, we may change the signs of the factors of two F_i 's at a time.) When F is an extreme ray of K_n then every F_i must be an extreme ray in its cone, and no two consecutive factors can be of the form $F_i = f_{\emptyset}^{n_i}$, $F_{i+1} = f_{\emptyset}^{n_{i+1}}$. Theorem 5.1 implies that the converse is true as well: every convolution of extreme rays such that no two consecutive factors are of the form $F_i = f_{\emptyset}^{n_i}$, $F_{i+1} = f_{\emptyset}^{n_{i+1}}$, is an extreme ray.

One way of showing $F \in K_{n+1}$ is an extreme ray is to give $2^n - 1$ linearly independent facets of K_{n+1} containing F . (Note that since K_{n+1} contains the positive orthant of A_{n+1} , it is full dimensional.) Hence it is useful to introduce the following operators.

DEFINITION 5.3. Let P be an arbitrary graded poset of rank $n + 1$, and \mathcal{I} an arbitrary interval system on $[1, n]$. The operators $\varepsilon_P^{n+1}, \varepsilon_{\mathcal{I}}^{n+1} \in A_{n+1}^*$ are given by

$$\begin{aligned} \varepsilon_P^{n+1}(f_S^{n+1}) &:= f_S^{n+1}(P), \quad \text{and} \\ \varepsilon_{\mathcal{I}}^{n+1}(f_S^{n+1}) &:= \begin{cases} 1 & \text{if } \forall I \in \mathcal{I} \quad (S \cap I \neq \emptyset), \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

As a consequence of this definition we have for $F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ that

$$\varepsilon_{\mathcal{I}}^{n+1}(F) = \sum_{S \in \mathbf{B}[1, n](\mathcal{I})} a_S,$$

hence the hyperplanes determining the facets of K_{n+1} are the kernels of the $\varepsilon_{\mathcal{I}}^{n+1}$'s. Let us also note that Corollary 2.6 may be rewritten as

$$\varepsilon_{\mathcal{I}}^{n+1} = \lim_{N \rightarrow \infty} \frac{1}{f_{[1, n]}(P(n, \mathcal{I}, N))} \cdot \varepsilon_{P(n, \mathcal{I}, N)}^{n+1}.$$

It is easy to show that the operators $\varepsilon_{\mathcal{I}}^{n+1}$, where \mathcal{I} runs over all interval systems on $[1, n]$, contain a basis of the vector space A_{n+1}^* .

DEFINITION 5.4. For every $S \subseteq [1, n]$ let \mathcal{I}_S denote the interval system $\{\{s\} : s \in S\}$. As a shorthand for $\varepsilon_{\mathcal{I}_S}^{n+1}$ we will use ε_S^{n+1} .

LEMMA 5.5. For every $S, T \subseteq [1, n]$ we have

$$\varepsilon_S^{n+1}(f_T^{n+1}) = \begin{cases} 1 & \text{if } S \subseteq T, \\ 0 & \text{otherwise.} \end{cases}$$

The proof is straightforward.

COROLLARY 5.6. The set $\{\varepsilon_S^{n+1} : S \subseteq [1, n]\}$ is a basis of A_{n+1}^* .

PROPOSITION 5.7. The chain operator f_{\emptyset}^{n+1} is an extreme ray of the cone K_{n+1} for every $n \geq 0$.

Proof. Since $f_{\emptyset}^{n+1}(P) > 0$ holds for every partially ordered set of rank $n+1$, we have $f_{\emptyset}^{n+1} \in K_{n+1}$. We only need to show that f_{\emptyset}^{n+1} lies on at least $2^n - 1$ linearly independent facets of K_{n+1} . This is true, since by Lemma 5.5 we have $\varepsilon_S^{n+1}(f_{\emptyset}^{n+1}) = 0$, whenever S is not the empty set. ■

The facets of the form $\text{Ker}(\varepsilon_S^{n+1})$ are also useful in proving the following proposition.

PROPOSITION 5.8. Let $n > 0$ be an integer and $k \in [1, n]$. Then the set $\sigma_k^{n+1}(K_{n+1})$ is a face of K_{n+2} .

Proof. Evidently we have

$$\text{Im}(\sigma_k^{n+1}) = \bigcap_{\substack{S \subseteq [1, n+1] \\ k \in S}} \text{Ker}(\varepsilon_S^{n+1}),$$

since both vector spaces are spanned by those chain operators f_S^{n+2} for which $k \notin S$. Thus we only need to show that $\sigma_k^{n+1}(K_{n+1}) \subseteq K_{n+2}$. The cone $\sigma_k^{n+1}(K_{n+1})$ is then contained in the 2^n -dimensional face

$$K_{n+2} \cap \left(\bigcap_{\substack{S \subseteq [1, n+1] \\ k \in S}} \text{Ker}(\varepsilon_S^{n+1}) \right)$$

of K_{n+2} , and, having the same dimension, it is also equal to it. Equivalently, we have to prove that

$$\varepsilon_{\mathcal{J}}^{n+2}(\sigma_k^{n+1}(F)) \geq 0$$

holds for every interval system \mathcal{J} on $[1, n+1]$.

Assume first that \mathcal{J} contains the interval $\{k\}$. Then we have

$$\varepsilon_{\mathcal{J}}^{n+2}(\sigma_k^{n+1}(F)) = 0$$

for every $F \in A_{n+1}$. If \mathcal{I} does not contain $\{k\}$ then consider the interval system

$$\mathcal{I}' := \{(I \cap [1, k-1]) \cup (I \cap [k, n+1] - 1) : I \in \mathcal{I}\}$$

on $[1, n]$. It is easy to verify that we have

$$\varepsilon_{\mathcal{I}'}^{n+2}(\sigma_k^{n+1}(F)) = \varepsilon_{\mathcal{I}}^{n+1}(F)$$

for every $F \in A_{n+1}$, and so $F \in K_{n+1}$ implies $\varepsilon_{\mathcal{I}'}^{n+2}(\sigma_k^{n+1}(F)) \geq 0$. ■

COROLLARY 5.9. *Given $n \geq 1$ and $k \in [1, n]$, the form $F \in A_{n+1}$ belongs to K_{n+1} if and only if $\sigma_k^{n+1}(F)$ belongs to K_{n+2} . Moreover, $F \in K_{n+1}$ is an extreme ray if and only if $\sigma_k^{n+1}(F) \in K_{n+2}$ is an extreme ray.*

Corollary 5.9 implies that every extreme ray $F \in K_{n+2}$ with $\text{supp}(F) \neq [1, n+1]$ is obtained by lifting an extreme ray of K_{n+1} using an embedding σ_k^{n+1} . Iterated use of Corollary 5.9 yields the following.

THEOREM 5.10. *Let $F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \in A_{n+1}$ be a form with $\text{supp}(F) = \{i_1, i_2, \dots, i_k\}$ and let $\gamma: [1, k] \rightarrow \text{supp}(F)$ be the bijection $j \mapsto i_j$. Then F belongs to K_{n+1} if and only if*

$$F' := \sum_{S \subseteq \text{supp}(F)} a_{\gamma^{-1}(S)} \cdot f_{\gamma^{-1}(S)}^{k+1}$$

belongs to K_{k+1} . Moreover $F \in K_{n+1}$ is an extreme ray if and only if $F' \in K_{k+1}$ is an extreme ray.

It follows from Theorem 5.1 and our calculation of the rank 2 case in Subsection 1.1 that the functionals $h_i := f_i - f_\emptyset$, $1 \leq i \leq n$, will all be extreme in rank $n+1$.

The crucial step in the proof of Theorem 5.1 is the following.

LEMMA 5.11. *Let $F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} \neq f_\emptyset^{n+1}$ be an extreme ray of K_{n+1} and let m be the largest letter occurring in F . Assume that F cannot be written as $G * f_\emptyset^{n+1-m}$ for some $G \in K_m$. Then there exists integers k, l satisfying $0 \leq k < l \leq m$ and an interval system \mathcal{I} on $[1, m-1]$ such that*

$$\sum_{S \cup \{0\} \in \mathbf{B}[0, n](\mathcal{I} \cup \{[k, m]\})} a_S = 0 \quad \text{and} \quad \sum_{S \in \mathbf{B}[1, n](\mathcal{I} \cup \{[l, m]\})} a_S \neq 0 \quad \text{hold.}$$

Proof. Let us show first that, without loss of generality, we may restrict ourselves to the case $m = n$. Since m is the largest occurring letter, we have

$$F = \sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} = \sum_{S \subseteq [1, m]} a_S \cdot f_S^{n+1}.$$

The form F is of the form $G * f_{\emptyset}^{n+1-m}$ for some $G \in K_m$ if and only if the form

$$\tilde{F} := \sum_{S \subseteq [1, m]} a_S \cdot f_S^{m+1}$$

is of the form $G * f_{\emptyset}^1$ for the same $G \in K_m$. Given an arbitrary interval system \mathcal{J} on $[1, m]$ we have

$$\varepsilon_{\mathcal{J}}^{n+1}(F) = \varepsilon_{\mathcal{J}}^{m+1}(\tilde{F}).$$

Therefore, if the lemma holds for the case $m = n$ then we may use the result on \tilde{F} to prove the same result for F .

Hence we may assume that $m = n$ holds. The statement is equivalent to saying that there exists a facet $\text{Ker}(\varepsilon_{\mathcal{J}})$ of K_n and integers $0 \leq k < l \leq n$ such that $\pi_k^{n+1}(F)$ belongs to this facet but $\pi_l^{n+1}(F)$ does not.

In the contrary event every facet of K_n containing some $\pi_k^{n+1}(F)$ also contains all forms $\pi_l^{n+1}(F)$ for every $l > k$. Let $G \in A_n$ be an extreme ray of the intersection of all facets containing $\pi_n^{n+1}(F)$. (Note that n being the largest occurring letter, $\pi_n^{n+1}(F) = \sum_{S \subseteq [1, n-1]} a_{S \cup \{n\}} \cdot f_S^n$ is not the zero form.) By our assumptions, this form is also an extreme ray of the intersection of all facets containing $\pi_i^{n+1}(F)$ for $i = 0, 1, \dots, n-1$. Hence we may choose a small positive number $q \in \mathbb{R}$ such that

$$\pi_i^{n+1}(F) - q \cdot G \in K_n \quad \text{holds for } i = 0, 1, \dots, n.$$

By Corollary 4.6 we have $\pi_i^{n+1}(G * f_{\emptyset}^1) = G$ for $i = 1, 2, \dots, n$ and so we obtain

$$\pi_i^{n+1}(F - q \cdot G * f_{\emptyset}^1) \in K_n$$

for $i = 0, 1, \dots, n$. By Theorem 4.2 we obtain that $F - q \cdot G * f_{\emptyset}^1$ belongs to K_{n+1} . Since F is an extreme ray, and by Proposition 1.2 we have $G * f_{\emptyset}^1 \in K_{n+1}$, F must be equal to a nonzero constant multiple of $G * f_{\emptyset}^1$, contrary to our assumptions.

COROLLARY 5.12. *Let $F \in K_{n+1}$ be an extreme satisfying the conditions of Lemma 5.11. Then there exists integers k, l satisfying $0 \leq k < l \leq m$ and an interval system \mathcal{I} on $[1, m-1]$ such that*

$$\sum_{S \cup \{0\} \in \mathbf{B}[0, n](\mathcal{I} \cup \{[k, n]\})} a_S = 0 \quad \text{and} \quad \sum_{S \in \mathbf{B}[1, n](\mathcal{I} \cup \{[l, n]\})} a_S \neq 0 \quad \text{hold.}$$

In fact, replacing the intervals $[k, m]$ and $[l, m]$ with $[k, n]$ and $[l, n]$ respectively does not change the sums involved, since whenever S contains a letter larger than m , a_S is zero.

We now proceed to the proof of Theorem 5.1. Since applying the chain operators to the dual of every poset yields an anti-isomorphism of the graded ring A which sends products of the form $F * f^l_{\emptyset}$ into products of the form $f^l_{\emptyset} * \hat{F}$, it is sufficient to show the following “half” of the original statement.

PROPOSITION 5.13. *Assume $F \in K_m$ and $G \in K_n$ are extreme rays such that F is not a convolution of the form $F' * f^l_{\emptyset}$. Then $F * G$ is an extreme ray of K_{m+n} .*

Proof. By Proposition 5.2 we know that $F * G$ belongs to K_{m+n} . In order to show that it is an extreme ray, it is sufficient to find $2^{m+n-1} - 1$ interval systems $\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_{2^{m+n-1}-1}$ such that the operators $\varepsilon_{\mathcal{I}_1}^{m+n}, \varepsilon_{\mathcal{I}_2}^{m+n}, \dots, \varepsilon_{\mathcal{I}_{2^{m+n-1}-1}}^{m+n}$ are linearly independent, and they all vanish on $F * G$.

Assume we have

$$F = \sum_{S \subseteq [1, m-1]} a_S^m \cdot f_S^m \quad \text{and} \quad G = \sum_{S \subseteq [1, n-1]} a_S^n \cdot f_S^n.$$

Then

$$F * G = \sum_{S \subseteq [1, m+n-1]} a_S^{m+n} \cdot f_S^{m+n}$$

is given by the following formula:

$$a_S^{m+n} = \begin{cases} a_{S \cap [1, m-1]}^m \cdot a_{(S \cap [m+1, m+n-1]) - m}^n & \text{if } m \in S, \\ 0 & \text{if } m \notin S. \end{cases}$$

Since for every nonzero a_S^{m+n} we must have $m \in S$, for an arbitrary interval system \mathcal{I} on $[1, m+n-1]$ we have

$$\varepsilon_{\mathcal{I}}^{m+n}(F * G) = \varepsilon_{\{I \in \mathcal{I} : m \notin I\}}^{m+n}(F * G),$$

i.e., we may remove every interval containing m from \mathcal{I} without changing the effect of $\varepsilon_{\mathcal{I}}^{m+n}$ on $F * G$. The remaining intervals are either contained in $[1, m-1]$ or in $[m+1, m+n-1]$. Introducing

$$\begin{aligned}\mathcal{I}' &:= \{I \in \mathcal{I} : I \subseteq [1, m-1]\} & \text{and} \\ \mathcal{I}'' &:= \{I - m : I \in \mathcal{I}, I \subseteq [m+1, m+n-1]\}\end{aligned}$$

we obtain

$$\varepsilon_{\mathcal{I}}^{m+n}(F * G) = \varepsilon_{\mathcal{I}'}^m(F) \cdot \varepsilon_{\mathcal{I}''}^n(G). \quad (10)$$

Since $F \in K_m$ is an extreme ray, there exist interval systems $\mathcal{I}'_1, \mathcal{I}'_2, \dots, \mathcal{I}'_{2^{m-1}}$ on $[1, m-1]$ such that the operators $\varepsilon_{\mathcal{I}'_1}^m, \varepsilon_{\mathcal{I}'_2}^m, \dots, \varepsilon_{\mathcal{I}'_{2^{m-1}}}^m$ are linearly independent, $\varepsilon_{\mathcal{I}'_j}^m(F) = 0$ for $j \leq 2^{m-1} - 1$, and $\varepsilon_{\mathcal{I}'_{2^{m-1}}}^m(F)$ is strictly positive. Similarly, the fact of $G \in K_n$ being an extreme ray implies that there exist interval systems $\mathcal{I}''_1, \mathcal{I}''_2, \dots, \mathcal{I}''_{2^{n-1}}$ on $[1, n-1]$ such that the operators $\varepsilon_{\mathcal{I}''_1}^n, \varepsilon_{\mathcal{I}''_2}^n, \dots, \varepsilon_{\mathcal{I}''_{2^{n-1}}}^n$ are linearly independent, $\varepsilon_{\mathcal{I}''_j}^n(F) = 0$ holds for $j \leq 2^{n-1} - 1$, and $\varepsilon_{\mathcal{I}''_{2^{n-1}}}^n(F)$ is strictly positive. Moreover, since F is not of the form $F' * f_{\mathcal{I}'}^{l, 2^{m-1}}$, by Corollary 5.12 we may assume that $\mathcal{I}'_{2^{m-1}}$ is of the form $\mathcal{I}' \cup \{[l, m-1]\}$ where \mathcal{I}' is an interval system on $[1, m-1]$, $l \in [1, m-1]$ and either we have $\varepsilon_{\mathcal{I}'}^m(F) = 0$, or there exist a $k \in [1, l-1]$ such that $\varepsilon_{\mathcal{I}' \cup \{[k, m-1]\}}^m(F) = 0$ holds.

Consider now the following interval systems:

- (i) All interval systems of the form $\mathcal{I}'_i \cup (\mathcal{I}''_j + m)$, where at least one of $i \neq 2^{m-1}$ and $j \neq 2^{n-1}$ holds.
- (ii) All interval systems of the form $\mathcal{I}'_i \cup \{m\} \cup (\mathcal{I}''_j + m)$, where at least one of $i \neq 2^{m-1}$ and $j \neq 2^{n-1}$ holds.
- (iii) The interval system $\mathcal{I}' \cup \{[l, m]\} \cup (\mathcal{I}''_{2^{n-1}} + m)$ if we have $\varepsilon_{\mathcal{I}'}^m(F) = 0$, or the interval system $\mathcal{I}' \cup \{[k, m-1]\} \cup \{[l, m]\} \cup (\mathcal{I}''_{2^{n-1}} + m)$ if $\varepsilon_{\mathcal{I}' \cup \{[k, m-1]\}}^m(F) = 0$ holds.

The above list contains $2^{m+n-1} - 1$ interval systems: $2^{m+n-2} - 1$ of them are of type (i), $2^{m+n-2} - 1$ of them are of type (ii), and there is exactly one system listed at item (iii). It is easy to see using (10) that they all vanish on $F * G$. We are left to show that the operators they define are linearly independent.

First we show that the operator defined by the last system is not in the span of the operators defined by all others. For this purpose consider the form $H := \sum_{S \subseteq [1, m+n-1]} \tilde{a}_S \cdot f_S^{m+n}$ given by

$$\tilde{a}_S = \begin{cases} 0 & \text{if } m \in S, \\ a_{S \cap [1, m-1]}^m \cdot a_{(S \cap [m+1, m+n-1]) - m}^n & \text{if } m \notin S. \end{cases}$$

It is easy to see that we have

$$\varepsilon_{\mathcal{J}'_i \cup (\mathcal{J}''_j + m)}^{m+n}(H) = \varepsilon_{\mathcal{J}'_i}^m(F) \cdot \varepsilon_{\mathcal{J}''_j}^n(G)$$

and so the operators defined by the systems of type (i) vanish on H . The operators defined by the systems of type (ii) vanish on H too, since we have $\tilde{a}_S = 0$ whenever $m \in S$. By the same reason we also have

$$\begin{aligned} \varepsilon_{\mathcal{J}' \cup \{[k, m-1]\} \cup \{[l, m]\} \cup (\mathcal{J}''_{2^{n-1}+m})}^{m+n}(H) \\ = \varepsilon_{\mathcal{J}' \cup \{[k, m-1]\} \cup \{[l, m-1]\} \cup (\mathcal{J}''_{2^{n-1}+m})}^{m+n}(H) \end{aligned}$$

Since k is less than l , the dual ideal of intervals generated by $\mathcal{J}' \cup \{[k, m-1]\} \cup \{[l, m-1]\} \cup (\mathcal{J}''_{2^{n-1}+m})$ is the same as the dual ideal generated by $\mathcal{J}' \cup \{[l, m-1]\} \cup (\mathcal{J}''_{2^{n-1}+m}) = \mathcal{J}'_{2^{m-1}} \cup (\mathcal{J}''_{2^{n-1}+m})$. Hence we have

$$\varepsilon_{\mathcal{J}' \cup \{[k, m-1]\} \cup \{[l, m]\} \cup (\mathcal{J}''_{2^{n-1}+m})}^{m+n}(H) = \varepsilon_{\mathcal{J}'_{2^{m-1}}}^m(F) \cdot \varepsilon_{\mathcal{J}''_{2^{n-1}}}^m(G) \neq 0.$$

Next we show that the intersection of the subspace generated by the operators associated to the interval systems listed in (i) with the subspace generated by the operators associated to the interval systems listed in (ii) is zero. For this purpose observe that A_n may be written as a direct sum of two 2^{m+n-2} -dimensional vector spaces, $A_n = A'_n \oplus A''_n$, where

$$A'_n := \langle f_{S \cup \{m\}}^{m+n} - f_S^{m+n} : S \subseteq [1, m+n-1], m \notin S \rangle$$

and

$$A''_n := \langle f_{S \cup \{m\}}^{m+n} : S \subseteq [1, m+n-1], m \notin S \rangle.$$

Evidently, linear combinations of operators associated to interval systems of type (i) vanish on A'_n while linear combinations of operators associated to interval systems of type (ii) have the same values on the respective generators of A'_n and A''_n . Assume now that $\varepsilon \in A_n^*$ is simultaneously a linear combination of operators associated to interval systems of type (i) and of operators associated to interval systems of type (ii). Then ε vanishes on A'_n and hence on A''_n , so $\varepsilon = 0$.

Finally, let $F_1, \dots, F_{2^{m-1}} \in A_m$ and $G_1, \dots, G_{2^{n-1}} \in A_n$ be dual bases to $\{\varepsilon_{\mathcal{J}'_i}^m\}$ and $\{\varepsilon_{\mathcal{J}''_j}^n\}$, respectively, that is, $\varepsilon_{\mathcal{J}'_i}^m(F_j) = \delta_{ij}$ and $\varepsilon_{\mathcal{J}''_k}^n(G_l) = \delta_{kl}$. Then by (5.3) we have

$$\varepsilon_{\mathcal{J}'_i \cup (\mathcal{J}''_k + m)}(F_j * G_l) = \varepsilon_{\mathcal{J}'_i \cup \{m\} \cup (\mathcal{J}''_k + m)}(F_j * G_l) = \delta_{ij} \delta_{kl},$$

showing the interval systems of types (i) and (ii) to be linearly independent. ■

We can describe all extremes of the cone K_n for $n \leq 5$. Equivalently, these represent the strongest linear inequalities holding for the flag f -vectors of graded posets of these ranks.

$n = 1$: The only extreme ray of K_1 is $h^1_{\emptyset} = f^1_{\emptyset}$.

$n = 2$: As seen in Subsection 1.1, the extreme rays for K_2 are $h^2_{\emptyset} = f^2_{\emptyset}$ and $h^2_1 = f^2_1 - f^2_{\emptyset}$.

$n = 3$: By repeated use of Theorems 5.1 and 5.10, we may generate five extremes for K_3 : h^3_{\emptyset} , h^3_1 , h^3_2 , $h^1_{\emptyset} * h^2_1 = f^3_{12} - f^3_1$ and $h^2_1 * h^1_{\emptyset} = f^3_{12} - f^3_2$. Direct computation shows these are all the extremes in this case.

$n = 4$: Direct calculation shows that there are 13 extremes for K_4 . All except one can be obtained by repeated use of Theorems 5.1 and 5.10. The remaining one is $f^4_{13} - f^4_1 + f^4_2 - f^4_3$, which represents the inequality given in Example 3.4.

$n = 5$: Again, direct calculation reveals that K_5 has 41 extremes. All but seven of these arise from lifting and convolution as above. The remainder are

1. $f^5_{134} - f^5_{14} + f^5_{24} - f^5_{34} - f^5_2 + f^5_3$
2. $f^5_{124} - f^5_{12} + f^5_{13} - f^5_{14} + f^5_2 - f^5_3$.
3. $f^5_{1234} - f^5_{123} - f^5_{234} + f^5_{13} - f^5_{14} + f^5_{23} + f^5_{24} - f^5_2$.
4. $f^5_{1234} - f^5_{123} - f^5_{234} + f^5_{13} - f^5_{14} + f^5_{23} + f^5_{24} - f^5_3$.
5. $f^5_{124} + f^5_{234} - f^5_{12} + f^5_{13} - f^5_{14} - f^5_{23} - f^5_{24} + f^5_2$.
6. $f^5_{123} + f^5_{134} - f^5_{34} + f^5_{24} - f^5_{14} - f^5_{23} - f^5_{13} + f^5_3$.
7. $f^5_{134} + f^5_{124} - f^5_{13} - f^5_{14} + f^5_{23} - f^5_{24}$.

For rank 6 direct calculation yields 796 extreme rays. Only 131 of them come from earlier extremes via lifting and convolution; the remaining 665 are new. An interesting problem would be to find a reasonable characterization of the extreme rays of K_n .

6. THE FLAG ℓ -VECTOR

In this section we present a transformed version of the flag f -vector which allows a sparse encoding of K_{n+1} .

DEFINITION 6.1. The chain operators ℓ_S^{n+1} (where $S \subseteq [1, n]$) are defined by the formula

$$\ell_S^{n+1} := (-1)^{n-|S|} \sum_{T \supseteq [1, n] \setminus S} (-1)^{|T|} \cdot f_T^{n+1}.$$

We call the vector $(\ell_S^{n+1}(P): S \subseteq [1, n])$ the *flag ℓ -vector* of the graded poset P .

Straightforward substitution in the definition shows

$$f_S^{n+1} = \sum_{T \subseteq [1, n] \setminus S} \ell_T^{n+1}. \tag{11}$$

The interest of the flag ℓ -vector is that the facet inequalities of K_{n+1} take a very simple form in the flag ℓ basis.

LEMMA 6.2. *If $\sum_{S \subseteq [1, n]} a_S \cdot f_S^{n+1} = \sum_{S \subseteq [1, n]} c_S \cdot l_S^{n+1}$ then*

$$a_S = (-1)^{|S|} \sum_{T \subseteq [1, n] \setminus S} (-1)^{n-|T|} \cdot c_T \quad \text{and} \quad c_S = \sum_{T \subseteq [1, n] \setminus S} a_T.$$

Proof. The equations follow from

$$\begin{aligned} \sum_{T \subseteq [1, n]} c_T \cdot l_T^{n+1} &= \sum_{T \subseteq [1, n]} c_T \cdot (-1)^{n-|T|} \sum_{S \subseteq [1, n] \setminus T} (-1)^{|S|} \cdot f_S^{n+1} \\ &= \sum_{S \subseteq [1, n]} f_S^{n+1} \cdot (-1)^{|S|} \sum_{T \subseteq [1, n] \setminus S} (-1)^{n-|T|} \cdot c_T, \end{aligned}$$

and

$$\begin{aligned} \sum_{T \subseteq [1, n]} a_T \cdot f_T^{n+1} &= \sum_{T \subseteq [1, n]} a_T \cdot \sum_{S \subseteq [1, n] \setminus T} f_S^{n+1} \\ &= \sum_{S \subseteq [1, n]} f_S^{n+1} \cdot \sum_{T \subseteq [1, n] \setminus S} a_T. \quad \blacksquare \end{aligned}$$

PROPOSITION 6.3. *Let $\mathcal{I} = \{I_1, \dots, I_k\}$ be an antichain of intervals. The inequality $\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S \geq 0$ is equivalent to*

$$\sum_{L \subseteq [1, k]} (-1)^{|L|} c_{\cup_{j \in L} I_j} \geq 0.$$

Proof. By the sieve formula we have

$$\sum_{S \in \mathbf{B}_{[1, n]}(\mathcal{I})} a_S = \sum_{L \subseteq [1, k]} (-1)^{|L|} \cdot \sum_{\substack{S \subseteq [1, n] \\ S \cap (\cup_{j \in L} I_j) = \emptyset}} a_S,$$

and we may apply Lemma 6.2. \blacksquare

For a “typical” antichain of intervals, k will be considerably less than n and so the vectors representing facet inequalities of K_{n+1} include many more zero entries in the flag ℓ basis than in the flag f basis. In fact, at most

2^k terms will be nonzero in the flag ℓ basis, while in the flag f basis any subset of $[1, n]$ containing a fixed k -element blocking set is a blocker, thus the number of nonzero entries is at least $k \cdot 2^{n-k}$ which is much larger than k whenever $k \leq n/2$.

We expect that the flag ℓ -vector will be very useful in expressing linear inequalities holding for the flag numbers of special subclasses of graded partially ordered sets as well.

Let us note finally that the formula connecting the flag ℓ -vector with the flag h vector is an involution. Straightforward calculation shows

$$\ell_S^{n+1} = (-1)^{|S|} \sum_{T \supseteq S} h_T^{n+1}, \quad \text{and} \quad (12)$$

$$h_S^{n+1} = (-1)^{|S|} \sum_{T \supseteq S} \ell_T^{n+1}. \quad (13)$$

ACKNOWLEDGMENT

We are grateful to Margaret Bayer, Vesselin Gasharov and Swapneel Mahajan for pointing out misprints in previous versions of this paper.

REFERENCES

1. L. J. Billera and G. Hetyei, Decompositions of partially ordered sets, preprint, 1998.
2. L. J. Billera and N. Liu, Noncommutative enumeration in graded posets, *J. Algebraic Combin.*, to appear.
3. P. M. Cohn, On subsemigroups of free semigroups, *Proc. Amer. Math. Soc.* **13** (1962), 347–351.
4. J. D. Edmonds and D. R. Fulkerson, Bottleneck extrema, *J. Combin. Theory* **8** (1970), 299–306.
5. M. Grötschel, L. Lovász, and A. Schrijver, “Geometric Algorithms and Combinatorial Optimization,” Springer-Verlag, Berlin, 1988.
6. G. Kalai, A new basis for polytopes, *J. Combin. Theory Ser. A* **49** (1988), 191–208.
7. R. P. Stanley, “Combinatorics and Commutative Algebra,” 2nd ed., Birkhäuser, Boston, 1996.