


Cell Metabolism

Cdk5rap1-Mediated 2-Methylthio Modification of Mitochondrial tRNAs Governs Protein Translation and Contributes to Myopathy in Mice and Humans

Graphical Abstract



Authors

Fan-Yan Wei, Bo Zhou, ..., Yuichi Oike, Kazuhito Tomizawa

Correspondence

tomikt@kumamoto-u.ac.jp

In Brief

Wei et al. report that Cdk5rap1 is responsible for 2-methylthio (ms²) modifications of mammalian mt-tRNAs. The modification is critical for efficient mitochondrial translation in mice and dysregulated in MELAS patients.

Highlights

- Cdk5rap1 catalyzes 2-methylthio (ms²) modification of four mitochondrial tRNAs
- The ms² modifications optimize mitochondrial translation and OXPHOS activity
- Deficiency of ms² modification accelerates myopathy and cardiac dysfunction in mice
- The ms² modification levels are reduced in patients with mitochondrial disease



Cdk5rap1-Mediated 2-Methylthio Modification of Mitochondrial tRNAs Governs Protein Translation and Contributes to Myopathy in Mice and Humans

Fan-Yan Wei,^{1,7} Bo Zhou,^{1,7} Takeo Suzuki,³ Keishi Miyata,² Yoshihiro Ujihara,⁴ Haruki Horiguchi,² Nozomu Takahashi,¹ Peiyu Xie,¹ Hiroyuki Michiue,⁵ Atsushi Fujimura,⁵ Taku Kaitsuka,¹ Hideki Matsui,⁵ Yasutoshi Koga,⁶ Satoshi Mohri,⁴ Tsutomu Suzuki,³ Yuichi Oike,² and Kazuhito Tomizawa^{1,*}

¹Department of Molecular Physiology

²Department of Molecular Genetics

Faculty of Life Sciences, Kumamoto University, Kumamoto 860-8556, Japan

³Department of Chemistry and Biotechnology, School of Engineering, The University of Tokyo, Tokyo 113-8656, Japan

⁴First Department of Physiology, Kawasaki Medical School, Okayama 701-0192, Japan

⁵Department of Physiology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan

⁶Department of Pediatrics and Child Health, Kurume University Graduate School of Medicine, Fukuoka 830-0011, Japan

⁷Co-first author

*Correspondence: tomikt@kumamoto-u.ac.jp

<http://dx.doi.org/10.1016/j.cmet.2015.01.019>

SUMMARY

Transfer RNAs (tRNAs) contain a wide variety of post-transcriptional modifications that are important for accurate decoding. Mammalian mitochondrial tRNAs (mt-tRNAs) are modified by nuclear-encoded tRNA-modifying enzymes; however, the physiological roles of these modifications remain largely unknown. In this study, we report that Cdk5 regulatory subunit-associated protein 1 (Cdk5rap1) is responsible for 2-methylthio (ms^2) modifications of mammalian mt-tRNAs for Ser(UCN), Phe, Tyr, and Trp codons. Deficiency in ms^2 modification markedly impaired mitochondrial protein synthesis, which resulted in respiratory defects in Cdk5rap1 knockout (KO) mice. The KO mice were highly susceptible to stress-induced mitochondrial remodeling and exhibited accelerated myopathy and cardiac dysfunction under stressed conditions. Furthermore, we demonstrate that the ms^2 modifications of mt-tRNAs were sensitive to oxidative stress and were reduced in patients with mitochondrial disease. These findings highlight the fundamental role of ms^2 modifications of mt-tRNAs in mitochondrial protein synthesis and their pathological consequences in mitochondrial disease.

INTRODUCTION

Transfer RNA (tRNA) is a key molecule in the translational apparatus to decode genetic information into proteins. A unique feature of tRNAs is the presence of a variety of chemical modifications of their nucleotides (Machnicka et al., 2013). These modifications are critical for efficient and accurate decoding (Agris, 2004; Suzuki, 2005). To date, more than 100 modified nucleotides

have been identified in tRNAs from the three domains of life, indicative of the universal importance of tRNA modifications (Machnicka et al., 2013).

Given the critical roles of tRNA modifications in cells, it is not surprising that tRNA modification deficiencies have been associated with human diseases (Torres et al., 2014). Genetic variations in the gene encoding Cdk5 regulatory subunit associated protein 1-like-1 (CDKAL1), which inserts a 2-methylthio (ms^2) group into the N^6 -threonylcarbamoyl adenosine (t^6A) of cytosolic tRNA^{Lys}(UUU), have been associated with the development of type 2 diabetes (Steinthorsdottir et al., 2007; Arragain et al., 2010). A deficiency in the ms^2 modification of tRNA^{Lys}(UUU) resulted in aberrant proinsulin synthesis, which ultimately led to impaired glucose metabolism and insulin secretion in Cdkal1 knockout (KO) mice and in human subjects carrying T2D-associated alleles of *CDKAL1* (Wei et al., 2011; Xie et al., 2013).

In mitochondrial tRNAs (mt-tRNAs), 15 species of modified nucleotides at 118 positions have been identified in bovine (Suzuki and Suzuki, 2014). Some of these modifications have been associated with the development of mitochondrial diseases, such as Mitochondrial myopathy, encephalopathy, lactic acidosis, stroke-like episodes (MELAS), and myoclonus epilepsy with ragged-red fibers (MERRF) (Suzuki et al., 2011). mt-tRNA^{Leu} and mt-tRNA^{Lys} contain 5-taurinomethyl (τm^5) and 5-taurinomethyl-2-thio (τm^5s^2) modifications, respectively, at U34 (Yasukawa et al., 2001; Suzuki et al., 2002), and both of these modifications are critical for decoding their cognate codons (Kirino et al., 2004; Yasukawa et al., 2001). The absence of these modifications has been observed in MELAS patients carrying the A3243G mutation in mt-tRNA^{Leu} and in MERRF patients carrying the A8344G mutation in mt-tRNA^{Lys} (Yasukawa et al., 2001). These results suggest a critical role for mt-tRNA modifications in the pathogenesis of human diseases. Nevertheless, knowledge regarding the physiological roles of tRNA modifications is incomplete, and a complete investigation of the individual types of mt-tRNA modifications is required to fully understand the physiological function and molecular pathology of tRNA modifications in human diseases.

In mammalian mt-tRNAs, 2-methylthio- N^6 -isopentenyl adenosine (ms^2i^6A) is a unique modification that is conserved in all three domains of life (Machnicka et al., 2013). In bacteria, the ms^2 modification of ms^2i^6A contributes to accurate decoding by improving tRNA binding to codons (Urbonavicius et al., 2001; Jenner et al., 2010). However, the physiological importance of ms^2 modifications in mammals is unknown. We have previously shown that Cdk5rap1 might be responsible for the ms^2 modification of ms^2i^6A because of its homology to Cdkal1, which catalyzes ms^2t^6A modification (Arragain et al., 2010). Recently, Cdk5rap1 was proposed to catalyze ms^2 group insertion in both cytosolic RNAs and mt-tRNAs; however, the exact substrate tRNA of Cdk5rap1 in mammalian cells remains unclear (Reiter et al., 2012).

Given the exclusive mitochondrial localization of ms^2i^6A in mammals and the implication of this localization in the decoding process, we hypothesize that Cdk5rap1 might specifically catalyze the ms^2 modification of mt-tRNAs and contribute to mitochondrial function in vivo. In this study, we validated this hypothesis through a thorough investigation of the physiological function of the ms^2 modification in Cdk5rap1 KO mice. Furthermore, we investigated the pathological implication and the molecular mechanism of the ms^2 modification in MELAS patients.

RESULTS

Cdk5rap1 Catalyzes the Conversion of i^6A to ms^2i^6A in Mitochondrial tRNAs

Based on the high homology between mammalian Cdk5rap1 and bacterial MiaB, we and another group previously showed that Cdk5rap1 might be a mammalian methylthiotransferase that catalyzes the conversion of N^6 -isopentenyl adenosine (i^6A) to 2-methylthio- N^6 -isopentenyl adenosine (ms^2i^6A) (Figure 1A and see Figure S1A available online; Arragain et al., 2010; Reiter et al., 2012). To investigate this hypothesis, we transformed Cdk5rap1 in MiaB-deficient bacteria (Δ MiaB), which do not contain ms^2 modifications. As expected, the transformation of Cdk5rap1 restored ms^2 modifications (Figure 1B). The conserved cysteine residues in the UPF domain and the radical SAM domain of MiaB are critical for ms^2 modification through their interaction with two [4Fe-4S] clusters (Forouhar et al., 2013). Similar to MiaB, the mutation of cysteine residues in the UFP domain or the radical SAM domain of Cdk5rap1 completely abolished the ms^2 modification (Figure 1B).

To prove that Cdk5rap1 is a mitochondrial methylthiotransferase and to identify its exact substrates, we generated Cdk5rap1 KO mice (Figures S1B–S1D). As expected, the ms^2 modification was completely abolished in KO mice (Figures S1E and S1F). Cdk5rap1 colocalized with Mitotracker in HeLa cells (Figure S2A). Cdk5rap1 with a deletion of mitochondrial localization signal at the N terminus exhibited cytosolic localization but retained its enzyme activity (Figures S2A and S2B). However, no ms^2 modification was detected in Cdk5rap1-deficient mouse embryonic fibroblast (MEF) cells expressing the cytosolic form of Cdk5rap1 (Figures S2C and S2D). These results indicate that Cdk5rap1 localizes on mitochondria and specifically modifies mt-tRNAs. To identify the exact substrate of Cdk5rap1, individual mt-tRNA was isolated from WT and KO mice and subjected to mass spectrometric analysis. The ms^2 modification

was completely absent at A37 in mt-tRNA^{Phe}, mt-tRNA^{Trp}, mt-tRNA^{Tyr}, and mt-tRNA^{Ser(UCN)} isolated from KO mice (Figures 1C–1F). The absence of an ms^2 modification did not affect the nearby τm^5 modification at U34 in mt-tRNA^{Trp} (Figure S1F). These results clearly demonstrate that Cdk5rap1 is a two [4Fe-4S] cluster-containing mitochondrial methylthiotransferase that specifically converts i^6A to ms^2i^6A at A37 of mt-tRNA^{Phe}, mt-tRNA^{Trp}, mt-tRNA^{Tyr}, and mt-tRNA^{Ser(UCN)} in mammalian cells.

The ms^2 Modification Controls Codon-Specific Decoding Fidelity in a Translation Rate-Dependent Manner

A previous study demonstrated that the ms^2 group of 2-methylthio- N^6 -hydroxyisopentenyl adenosine (ms^2io^6A) is critical for the accurate decoding of Tyr and Phe codons (Urbonavicius et al., 2001). To thoroughly investigate the role of ms^2 modifications of tRNAs during the decoding of their cognate codons, we utilized a luciferase-based reporter and transformed the plasmid into a WT strain or the Δ MiaB strain to detect frameshifting in the presence or absence of the ms^2 modification. The firefly luciferase gene was properly translated only when frameshifting occurred at the cognate codons read by tRNA^{Phe}, tRNA^{Trp}, tRNA^{Tyr}, or tRNA^{Ser(UCN)} (Figure S2E). A deficiency in ms^2 modification induced frameshifting at Phe(TTT) and Tyr(TAT) codons (Figure 1G). Induction of protein translation by isopropyl-beta-D-thiogalactopyranoside (IPTG) exaggerated the overall frameshifting rate in both the WT and Δ MiaB strains. In addition to the Phe(TTT) and Tyr(TAT) codons, there was a significant increase in ms^2 -dependent frameshifting at the Tyr(TAC), Ser(TCT), Ser(TCC), and Ser(TCG) codons, for which frameshifting was not observed without IPTG induction (Figures S1G–S1I). Importantly, these results show that ms^2 -dependent frameshifting specifically occurred during the translation of wobble codons, such as Phe(TTT), Tyr(TAT), Ser(TCT), Ser(TCC), and Ser(TCG), with the exception of the Tyr(TAC) codon, but not the cognate codons, such as Phe(TTC), Ser(TCA), and Trp(TGG) (shown in red in Figures S1F–S1I). Furthermore, the ms^2 -deficiency-evoked frameshifting was fully reversed by transformation with active Cdk5rap1, but not dominant-negative Cdk5rap1 (Figure 1G). These results demonstrate that ms^2 modification is critical for the accurate decoding of wobble codons corresponding to tRNA^{Phe}, tRNA^{Tyr}, or tRNA^{Ser(UCN)} in a translation rate-dependent manner.

Deficiency of the Mitochondrial ms^2 Modification Attenuates Mitochondrial Translation

To examine mitochondrial protein synthesis, WT or KO MEF cells were labeled with ³⁵S-methionine for 1 hr and subjected to pulse chase. The mitochondrial protein synthesis was substantially decreased in KO MEF cells (Figure 2A). Furthermore, MEF cells were radioactively labeled and subjected to blue native PAGE to examine the formation of respiratory complexes. The incorporation of mitochondrial proteins into complexes I, III, and IV was substantially decreased in KO MEF, whereas complex V was unaffected (Figures 2B and 2C). These results suggest that the deficiency in ms^2 modification greatly attenuated mitochondrial protein synthesis, resulting in impaired complex assembly.

The maintenance of mitochondrial OXPHOS subunits is critical for the electron transport chain, which maintains the resting




Figure 1. Cdk5rap1 Is The Mammalian mt-tRNA Methylthiotransferase

(A) Structures of N⁶-isopentenyladenosine (i^6A) and 2-methylthio-N⁶-isopentenyladenosine (ms^2i^6A) and the corresponding m/z values are shown. The ms^2 group (S-CH₃) is shown in red.

(B) GST-Cdk5rap1 or GST-Cdk5rap1 with Cys-to-Ala mutations (3CA) was transformed into the MiaB-deficient strain ($\Delta MiaB$). The arrows and arrowhead indicate the peaks corresponding to i^6A and ms^2i^6A , respectively.

(C–F) Examination of the ms^2i^6A modification in mt-tRNA^{Phe} (C), mt-tRNA^{Trp} (D), mt-tRNA^{Tyr} (E), and mt-tRNA^{Ser(UCN)} (F) isolated from WT and KO mice by mass spectrometry. The mass chromatograms show the peaks corresponding to fragments containing i^6A or ms^2i^6A .

(G) Frameshift assay. WT and $\Delta MiaB$ bacteria were transformed with Cdk5rap1 or inactive Cdk5rap1-3CA). The relative ratio of firefly luciferase activity to renilla luciferase activity (F/R) represents the decoding error. n = 4. Data are mean \pm SEM. **p < 0.01, ***p < 0.0001.

mitochondrial membrane potential and drives respiration. We thus investigated the mitochondrial membrane potential in WT and KO MEF cells. There was a marked increase in cell populations with very low membrane potential in KO MEF cells (Figure 2D). Consequently, the oxygen consumption rate in KO cells was significantly lower than that in WT MEF cells (Figure 2E). Furthermore, the KO cells quickly lost their mitochondrial membrane potential after treatment with very low doses of rotenone and FCCP, which had little effect on WT cells (Figures 2F and

2G). These results thus demonstrate that ms^2 modifications of mt-tRNAs are critical for maintaining efficient mitochondrial translation and respiratory chain.

Deficiency of Mitochondrial ms^2 Modification Impairs OXPHOS in Skeletal Muscle and Heart Tissue without Affecting Basal Metabolism

To investigate the physiological role of mitochondrial ms^2 modification, we examined the phenotype of KO mice in vivo. Despite




Figure 2. Deficiency in the ms^2 Modifications Impaired Mitochondrial Protein Synthesis and Mitochondrial Functions

(A) WT and KO MEF cells were labeled with ^{35}S -Met/Cys and chased for 0 or 24 hr in the presence of emetine for measurement of mitochondrial protein synthesis (left panels). MEF cells were labeled with ^{35}S -Met/Cys for 1 hr, and total protein synthesis was measured (right panels). CBB staining of gel was used as loading control.

(B and C) The autoradiogram of blue native PAGE shows a decrease in the incorporation of mitochondrial proteins in complexes I, III, and IV in KO MEF cells (B). The relative intensity of complex I, III, and IV versus complex V was quantified (C).

(D) The histogram and inserted graph show that KO MEF cells contain a number of mitochondria with low membrane potentials; $n = 110$ each.

(E) KO cells showed a significant decrease in the oxygen consumption rate (OCR); $n = 10$ each.

(E–G) Cells were stained with TMRM for measuring membrane potential. The relative membrane potential in the presence of rotenone (F) or FCCP (G) was analyzed; $n = 66$ for WT and $n = 30$ for KO (F); $n = 43$ for WT and $n = 32$ for KO (G). Data are mean \pm SEM. * $p < 0.05$.




Figure 3. The Deficiency in ms^2 Modification Impaired Mitochondrial Function In Vivo

(A) Steady-state levels of complex I (CI), complex II (CII), complex III (CIII), complex IV (CIV), and complex V (CV) in mitochondria isolated from skeletal muscle and heart tissues were examined by BN-PAGE.

(B) Steady-state levels of representative proteins of CI-CIV were examined by western blotting. CBB staining was used as loading control.

(C) The activities of CS, CI-CIV in skeletal muscle (left panel), and heart (right panel) of WT and KO mice were examined. $n = 4-5$.

(D and E) Respiratory coupling was decreased in *Cdk5rap1*-deficient mitochondria isolated from skeletal muscle and heart tissue when examined using XF24 Flux Analyzer; $n = 4-5$.

(F) The steady-state levels of ATP in the skeletal muscle and heart tissue of WT and KO mice were examined; $n = 4$ each. Data are mean \pm SEM. * $p < 0.05$.

the mitochondrial dysfunction in KO cells, the *Cdk5rap1* KO mice developed normally without obvious morphological changes in major tissues (Figures S3A and S3B). The energy expenditure and glucose metabolism in KO mice were compatible with those of WT mice (Figures S3C–S3E). Furthermore, there was no difference in neurological behaviors between KO and WT mice (Figures S3F–S3I).

Of all tissues, skeletal muscle and heart tissue are the most susceptible tissues to mitochondrial dysfunction (DiMauro and Schon, 2003). We therefore closely examined these two tissues in KO mice. Substantial decreases in the steady-state levels of complex I and IV were observed in both skeletal muscle and heart tissues of KO mice compared with WT mice, with complex I being markedly affected (Figure 3A). Accordingly, the steady-state levels of complex I and IV proteins, such as NDUFB8 and

MTCO1, respectively, were markedly decreased in skeletal muscle and heart tissues of KO mice compared with WT mice (Figure 3B). As a result, complex I activity was significantly impaired in KO mice (58.6% of WT for skeletal muscle and 51.5% of WT for heart tissue) (Figure 3C). There was a mild but significant decrease in complex III and complex IV activity in KO mice (muscle: complex III, 88.8% of WT; complex IV, 80.5% of WT; heart: complex III, 80.7% of WT, complex IV, 79.8% of WT) (Figure 3C). The decrease in mitochondrial activity in the skeletal muscle of KO mice was also confirmed by cytochrome oxidase (COX) staining (Figure S3J). Accordingly, the oxygen consumption elicited by ADP and FCCP in *Cdk5rap1*-deficient mitochondria was significantly lower than that in WT mitochondria, indicating that electron transport and respiratory coupling were impaired in the skeletal muscle and heart tissue of KO mice (respiratory

control ratio of the hearts of WT and KO: 4.9 and 3.2, skeletal muscles of WT and KO: 9.4 and 6.5, respectively; Figures 3D and 3E). Consequently, the steady-state ATP level in skeletal muscle and heart tissue of KO mice was lower than that in WT mice (Figure 3F).

Impairment of mitochondrial function usually exaggerates mitochondrial remodeling as a compensation mechanism. There was an increase in the mitochondrial mass in the skeletal muscle of KO mice as examined by electron microscopy and Gomori Trichrome staining (Figures 4A, 4B, and S3J). Strikingly, the mitochondria were abnormally enlarged in heart tissue of KO mice (Figures 4C and 4D). A progressive disruption of cristae was occasionally observed in the mitochondria of the cardiac muscle of KO mice (arrow in Figure 4C). Furthermore, there was a significant increase in citrate synthase activity (Figure 3C) as well as relative mtDNA content (Figure 4E) in the skeletal muscle of KO mice.

Reactive oxygen species (ROS) are byproducts of mitochondrial electron transport and mainly generated from complexes I and III (Murphy, 2009). A deficiency of complexes I and III accelerates the leakage of ROS from electron transport chain and contributes to the development of mitochondrial diseases. Given the marked decrease in complex I protein level in KO mice, we investigated ROS production in KO mice (Figure 4F). The ROS level was slightly but significantly higher in KO MEF cells than that in WT MEF cells. This finding was corroborated by a moderate increase in protein carbonylation (Figure 4G) as well as oxidative stress-related gene expression in both skeletal muscle and heart tissues of KO mice (Figure 4H).

To further investigate the impact of deficiency of ms^2 modification on physiological function, we examined muscular and cardiac function in vivo. However, the treadmill performance of KO mice was comparable with that of WT mice (Figure S3K). The echocardiography examination indicated that no apparent cardiac defects were present in the KO mice (Figure S3L). Taken together, these results demonstrate that the deficiency of ms^2 modifications in mt-tRNAs impairs mitochondrial protein synthesis, which leads to a reduction of respiratory activity and increase in ROS in skeletal muscle and heart tissue. However, considering the overall phenotypes, mice seem to tolerate an up to 50% reduction of complex I activity due to the loss of ms^2 modifications under sedentary conditions.

Loss of ms^2 Modifications Accelerates OXPHOS Defects under Stressed Conditions

The mild phenotype of KO mice prompted us to challenge the mice with a ketogenic diet (KD; very high fat and ultra-low carbohydrate). Ketone bodies from KD bypass glycolysis and generate energy mostly through fatty acid oxidation in mitochondria (Lafel, 1999). Adaptation to this metabolic pressure is accompanied by mitochondrial rearrangement (Grimsrud et al., 2012). Therefore, it is conceivable that the accurate regulation of mitochondrial protein synthesis by ms^2 modification is particularly important for mitochondrial remodeling under stressed conditions.

As expected, KD treatment accelerated OXPHOS defects in the skeletal muscle and heart tissue of KO mice (Figures 4A and 4B). Complex I activity was significantly impaired in KD-fed KO mice (48.6% of the KD-fed WT for skeletal muscle and 47.7% of the KD-fed WT for heart tissue) (Figures 5A and 5B).


In addition, accelerated decreases in complex III and IV activities were observed in the KD-fed KO mice (muscle: complex III, 82.9% of the KD-fed WT; complex IV, 62.9% of the KD-fed WT; heart: complex III, 75% of the KD-fed WT; complex IV, 57.8% of the KD-fed WT).

The OXPHOS defect after KD treatment exaggerated the mitochondrial remodeling pathway in both WT and KO mice. There was a ~ 3 -fold and ~ 1.5 -fold increase in mtDNA content in the skeletal muscle and heart tissue of both WT and KO mice fed a KD, respectively (Figure 5C). However, there was no difference in the mtDNA content in skeletal muscle and heart tissue between WT and KO mice (Figure 5C). Accordingly, subsequent electron microscopic examination revealed a marked increase in mitochondrial mass (Figures 5D and 5F). In the skeletal muscles of KO mice fed a KD, mitochondrial proliferation was observed in both intermyofibrillar and subsarcolemmal mitochondria, with the latter drastically increased (Figure 5D). Importantly, KD-fed KO mice exhibited a considerable population of mitochondria with disrupted cristae in the skeletal muscle tissue (arrowheads in Figure 5D). The enlargement of mitochondria and the disruption of cristae were even more prominent in the heart tissue of KO mice fed a KD (arrows in Figure 5D). These results demonstrate that Cdk5rap1-dependent ms^2 modification is crucial for the maintenance of OXPHOS activity and mitochondrial morphology under stress.

The acceleration of the OXPHOS defect in KO mice may be due to the indirect lipotoxicity from the very high-fat diet. However, the body weight and serum metabolic profiles of KO mice fed a KD were the same as those of WT mice fed a KD (Figures S4A–S4C). There was no difference in the locomotor activity or energy expenditure between the WT and KO mice fed a KD (Figures S4D and S4E). Interestingly, the glucose level in the KD-fed KO mice was somewhat lower than that in the KD-fed WT mice (Figure S4F). Taken together, these results indicate that the progressive OXPHOS defects and mitochondrial degeneration in KD-fed KO mice directly resulted from a deficiency in Cdk5rap1-dependent ms^2 modification during mitochondrial remodeling.

Loss of ms^2 Modification Accelerates Muscular and Cardiac Dysfunction under Stress

The KD-induced OXPHOS defect markedly accelerated the dysfunction of skeletal muscle and heart tissue in the KO mice. In a treadmill test, KO mice fed a KD showed a significant increase in the number of falls and became exhausted as early as 30 min into the test (Figure 5G). The KO mice also showed moderate cardiac hypertrophy, as indicated by an increase in heart volume, heart weight, and left ventricle posterior wall thickness (Figures S5A–S5C). The percentage of fractional shortening (FS%) in KO mice fed a KD was significantly lower than that in WT mice fed a KD (WT, 43% versus KO, 34%; Figure 5H). In addition to a KD-induced stress model, we utilized a transverse aortic constriction (TAC) model, which is a standard model for inducing cardiac dysfunction by pressure overload. Because the TAC model is also accompanied by global mitochondrial remodeling (Dai et al., 2012), we expected that a deficiency in ms^2 modification would further accelerate cardiac dysfunction. Indeed, chronic TAC resulted in a progressive cardiac hypertrophy, as indicated by an increase in heart weight and left ventricle posterior wall thickness in KO mice (WT,



7.5 mg/g body weight; KO, 11.3 mg/g body weight; [Figures S5A and S5D–S5F](#)). In WT mice, the FS% dropped from 51.2% to 34.5% 1 week after TAC but was then maintained until 10 weeks (5 weeks, 32.4; 10 weeks, 31.1; [Figure 5I](#)). In contrast, the FS% in KO mice continuously decreased after TAC and eventually decreased to as low as 21.7% ([Figure 5I](#)). Further examination in isolated cardiomyocytes revealed that the cardiac dysfunction observed in the TAC model of KO mice was associated with a decrease in calcium influx and contraction rate ([Figures 5J and S5G](#)). These results demonstrate that a deficiency of the ms^2 modifications in mt-tRNAs can cause a catastrophic defect in muscle and heart tissue under stressed conditions.

Deficiencies in ms^2 Modification Compromise the Quality of Mitochondria

Next, we investigated the molecular mechanism underlying the stress-induced acceleration of OXPHOS defects and cardiomyopathy in KO mice. Adaptation to mitochondrial stress requires coordinated protein synthesis ([Dai et al., 2012](#)). Because ms^2 modification controls decoding fidelity in a translation-rate-dependent manner, it is conceivable that a deficiency in ms^2 modification under stressed conditions might markedly compromise mitochondrial quality as well as integrity, which would result in severe OXPHOS defects and ultimately lead to myopathy and cardiac dysfunction. Indeed, a moderate increase in the complex I level was observed in WT mice treated with KD or TAC surgery ([Figures 6A and 6B](#)). In contrast, the steady-state levels of complexes I and IV levels were somewhat decreased in KD-fed and TAC KO mice when compared with NC-fed KO mice. The mitochondrial stresses increased protein carbonylation in both WT and KO mice. As a result, the protein carbonylation level in stressed heart tissues of KO mice was moderately higher than that in the stressed WT mice ([Figure S6A](#)). Impaired mitochondrial proteostasis exaggerates mitochondrial unfolded protein response (mt-UPR) ([Durieux et al., 2011; Houtkooper et al., 2013](#)). Accordingly, proteins involved in mtUPR, such as Yme111, Afg3l2, and Lonp1, were upregulated in mitochondria isolated from the hearts of KO mice treated with KD and TAC compared with WT mice ([Figure 6C](#)). Furthermore, a marked increase in polyubiquitinated proteins was observed in mitochondria isolated from the hearts of KO mice under stressed conditions ([Figure 6D](#)). Interestingly, the levels of polyubiquitination were proportional to the levels of cardiac function (FS%) in stressed KO mice (TAC > KD > NC; [Figure 6D](#); also see [Figures 5H and 5I](#)).

Mitophagy is the hallmark of the existence of compromised mitochondria. Parkin, an E3 ubiquitin ligase, primes mitophagy

by translocation to mitochondria with low membrane potentials and ubiquitination of mitochondrial proteins ([Kubli and Gustafsson, 2012](#)). Because cells with ms^2 modification deficiencies had a low basal mitochondrial membrane potential and were susceptible to stress-induced depolarization ([Figures 2F and 2G](#)), we hypothesized that mitochondrial stress might exaggerate the recruitment of Parkin to mitochondria and the acceleration of mitophagy in Cdk5rap1 KO cells. In KO cells treated with FCCP, most of the Parkin translocated to the mitochondria as soon as 2 hr after treatment, whereas similar translocation was not observed in WT cells treated with FCCP ([Figure 6E](#); also see the separated imaged in [Figure S6B](#)). A number of large mitochondrial aggregates were surrounded by the autophagosomal membrane protein LC3 in KO cells treated with FCCP, which is indicative of acceleration of mitophagy ([Figure 6F](#); also see the separated imaged in [Figure S6C](#)). Furthermore, we observed a number of degenerated mitochondria, with some mitochondria being degraded in autophagic vacuoles in KD-fed and TAC KO mice, by electron microscopic examination ([Figure 6G](#)). These results demonstrate that stress-induced mitochondrial remodeling impaired mitochondrial protein synthesis and accelerated the decomposition of respiratory complexes, which triggered the mtUPR and mitophagy in KO mice. Thus, the accumulation of compromised mitochondria ultimately contributes to the development of myopathy and cardiac dysfunction.

Association of ms^2 Modification with Mitochondrial Disease

Because the pathological phenotypes of KO mice resembled those of mitochondrial disease, we speculated that ms^2 modification might be involved in mitochondrial disease. We investigated the ms^2 modification level in peripheral blood cells collected from MELAS patients who carry the A3243G mutation ([Figure S7A](#)). Because of the limited number of clinical RNA samples, we adapted the quantitative PCR-based method ([Xie et al., 2013](#)), which was originally developed to examine the ms^2 level of ms^2t^6A in cytosolic tRNA^{Lys(UUU)}, to sensitively examine the ms^2 modification level of ms^2t^6A in mt-tRNAs ([Figures S7B–S7E](#)). Strikingly, the heteroplasmy level of mutant mt-DNA was significantly correlated with the ms^2 modification levels of four mt-tRNAs, but not with the cytosolic tRNA^{Lys(UUU)} ([Figures 7A–7D and S7F](#)). Interestingly, the mutant mtDNA level was not correlated with the expression level of *CDK5RAP1*, suggesting that the decrease in ms^2 modifications was not due to a deficiency in *CDK5RAP1* ([Figures S5D–S5G](#)). Because the A3243G mutation is located in the mtDNA region corresponding to mt-tRNA^{Leu(UUR)}, the decrease in the ms^2 levels of tRNA^{Trp}, tRNA^{Phe}, tRNA^{Tyr}, and tRNA^{Ser(UCN)} was likely not caused by the A3243G

Figure 4. Aberrant Mitochondrial Morphology and ROS Metabolism in KO Mice


(A–D) Mitochondria in skeletal muscle and heart tissue were examined by electron microscopy. KO mice exhibit disrupted mitochondrial morphology (A and C) and increased mitochondrial mass (B and D). Bars in (A) and the left panels of (C), 10 μ m. Bars in the right panels of (C), 0.5 μ m. WT, n = 100; KO, n = 112 in (B) and n = 126; KO, n = 132 in (D).

(E) The relative contents of mtDNA in muscle and heart tissue were examined; n = 6–9.

(F) ROS levels were analyzed by measuring the fluorescent intensity of CM-H2DCFDA in WT and MEF cells (left panel). The intensity was quantified (right panel); n = 3.

(G) Protein carbonylation levels were increased in the mitochondria of skeletal muscle and heart tissue in KO mice. NC, negative control.

(H) Heatmap showing the differentially regulated genes involved in oxidative stress response in skeletal muscle and heart tissue of WT and KO mice; n = 4. Data are mean \pm SEM. * $p < 0.5$, **** $p < 0.0001$.



(legend on next page)

mutation but, rather, was due to secondary effects. Cells bearing A3243G mutations in mtDNA exhibit a marked reduction of mitochondrial protein synthesis and an increase in the oxidative stress level (Crimi et al., 2005; Ishikawa et al., 2005). Because Cdk5rap1 contains highly oxidation-sensitive [4Fe-4S] clusters (Arragain et al., 2010), we speculated that the excess oxidative stress originated from mutant mitochondria might result in a collateral inhibition of Cdk5rap1 activity. Indeed, cells treated with sublethal doses of H₂O₂ showed a rapid decrease in ms² modification, which was completely reversed by adding 10 mM pyruvate, which serves as an antioxidant (Figures 7E–7G). In addition, treatment of cells with an NO donor such as SNAP and NOC18 significantly reduced the ms² modification level, which was reversed by the addition of the NO scavenger PTIO (Figure 7H). Taken together, these results suggest that oxidative stress-induced decreases in ms² modifications might compromise the quality of the mitochondria and contribute to the progression of mitochondrial disease.

DISCUSSION

Regulation of Mitochondrial Protein Synthesis by Cdk5rap1-Mediated ms² Modification

In the present study, we revealed the important physiological functions of ancient mitochondrial ms² modifications in mice and human. Using Cdk5rap1 KO mice, we provide direct evidence that Cdk5rap1 catalyzes the ms² modifications of mt-tRNA^{Phe}, mt-tRNA^{Trp}, mt-tRNA^{Tyr}, and mt-tRNA^{Ser(UCN)} in mammalian cells. The ms² group at the A37 of tRNA can directly participate in crossstrand stacking with the first nucleotide of the codon of the mRNA to maintain the reading frame (Jenner et al., 2010). Indeed, a deficiency in the ms² modification of ms²i⁶A impaired reading frame maintenance in bacteria and caused defective mitochondrial protein synthesis in Cdk5rap1 KO mice. Interestingly, nuclear-encoded mitochondrial protein, such as NDUF8 in complex I, was also decreased in KO mice. The indirect decrease of NDUF8 is most likely due to the poorly assembled respiratory complexes in KO mice. A previous study has shown that a deficiency in a single subunit in complex I could compromise complex formation and cause the proteolysis of other subunits (Karamanlidis et al., 2013). Our results thus demonstrate that the ms² modification of mt-tRNAs is indispensable for mitochondrial protein synthesis and the proper assembly of respiratory complexes.

Whereas only one transcript of *Cdk5rap1* has been found in mice, multiple splicing variants of human *CDK5RAP1* are listed in the database (Figure S1A). One transcript of human *CDK5RAP1* encodes a short form of CDK5RAP1 without a mitochondrial localization signal (Q95SZ6-2 in Figure S1A), which raises the possibility that CDK5RAP1 might regulate cellular function by modifying cytosol RNAs (Reiter et al., 2012). However, there was no detectable ms²i⁶A in total RNA isolated from KO MEF cells expressing the cytosolic form of Cdk5rap1 with the enzyme activity preserved. These results clearly suggest that Cdk5rap1 does not modify nuclear DNA-derived RNAs in murine cells. Furthermore, the defective mitochondrial protein synthesis observed in Cdk5rap1 KO mice may be directly caused by the loss of ms² modifications in mt-tRNAs.

Deficiency of ms² Modification and Its Physiological Outcome

This study revealed unique phenotypic outcomes of Cdk5rap1 KO mice in response to distinct environmental conditions. Under sedentary conditions, the skeletal and cardiac functions of the Cdk5rap1 KO mice were compatible with those of the WT mice, despite the marked decrease in respiratory activities and regardless of the increase of oxidative stress. The mitochondrial dysfunction in KO mice might be compensated by the remodeling of mitonuclear protein balance, which serves as a protective mechanism by inducing mtUPR (Houtkooper et al., 2013). Indeed, in contrast to the decrease of mitochondrial protein synthesis, several cytosolic proteins appear to be upregulated in KO cells (right panels in Figure 2A). This mitonuclear protein imbalance might contribute to the upregulation of basal mtUPR in muscle and heart tissues of KO mice (Figure 6C). Furthermore, a collective increase of ROS metabolism genes, including ROS scavenger genes such as *ApoE*, *DHCR24*, and *SRXN1*, might ameliorate oxidative stress and protect the muscular and cardiac functions in KO mice. Similar to our results, previous studies have found no adverse phenotypes under basal conditions in transgenic mice with respiratory defects (Karamanlidis et al., 2013; Wenz et al., 2009). Our results thus support the current perspective that mitochondrial dysfunction, depending on its degree, may not immediately produce a pathological phenotype under sedentary conditions.

In contrast, under stressed conditions, Cdk5rap1 KO mice exhibited apparent skeletal muscle and heart dysfunctions. The

Figure 5. Mitochondrial Stresses Accelerated Myopathy and Cardiac Dysfunction in ms²-Deficient Mice

(A and B) WT and KO mice at 8 weeks old were fed for KD for 10 weeks. The relative activities of CS, CI-CIV in skeletal muscle (A), and heart (B) were examined; n = 5–7 each.

(C) The relative mtDNA contents in skeletal muscle and heart tissue of KD-fed WT and KO mice were examined; n = 6–7. The dashed lines represent the relative mtDNA content in NC-fed WT mice.


(D–F) Electron microscopy examination of skeletal muscle and heart tissue show disrupted mitochondrial architecture (D) and a marked increase in mitochondrial mass (E and F) in KD-fed KO mice. Arrowheads and arrows indicate mitochondria with abnormal cristae in skeletal muscle and heart tissues of KD-fed KO mice, respectively; bars, 10 μm; WT, n = 152; KO, n = 144 in (E) and n = 161; KO, n = 130 in (F).

(G) A treadmill test performed at the end of 10 weeks of KD feeding showed that the KD induced a higher number of falls in the KO mice during acute exercise compared with the WT mice; n = 5 each.

(H) The fractional shortening (FS) rate in KO mice fed with KD for 10 weeks was significantly lower than that in KO-fed WT mice; n = 10–11.

(I) WT and KO mice at 8 weeks old were subject to TAC surgery. The KO mice showed a significant decrease in FS after TAC surgery.

(J) Cardiomyocytes were isolated from WT and KO mice 10 weeks after TAC. Calcium imaging revealed a decrease in the peak calcium influx in KO cardiomyocytes. The inserted graph shows the normalized peak amplitude of calcium influx in cardiomyocytes from WT and KO mice; n = 13 for WT and n = 6 for KO. Data are the mean ± SEM. *p < 0.05. **p < 0.01, ***p < 0.001, ****p < 0.0001.



(legend on next page)

defective mitochondrial protein synthesis caused a marked decrease in protein levels and activities of complexes I and IV in KO mice under stressed conditions. The progressive disruption of respiratory complexes, which exaggerates mtUPR and mitophagy, thus largely compromised mitochondria quality and led to myopathy in KO mice. Recent studies have shown that Parkin-mediated mitophagy is critical for the removal of damaged mitochondria and thus protects cardiac function under stressed conditions (Hoshino et al., 2013; Chen and Dorn, 2013). However, given the observation of a number of degenerated mitochondria in KO mice (Figures 4D and 5G), the extent of mitochondria damage in KO mice was likely beyond the maintenance capacity of mitophagy, which ultimately led to catastrophic mitochondrial dysfunction and myopathy. In addition, the acceleration of complex I defect was associated with a modest increase of oxidative stress, which could trigger mtUPR and cause cytotoxicity in stressed KO mice (Runkel et al., 2013). However, compared with the progressive impairment of mitochondria quality, the degree of increase of ROS after mitochondrial stress was rather small in stressed KO mice. Our results thus suggest that ROS might also contribute to the pathogenesis, but to a limited extent. The accumulation of malfunctioning mitochondria is likely the primary cause of the progression of myopathy in KO mice.

Regulation of ms^2 Modification by Oxidative Stress and Its Association with Human Disease

An important finding of this study is that the ms^2 modification levels were reduced in MELAS patients carrying the A3243G mutation in mt-tRNA^{Leu}. This result is surprising because mt-tRNA^{Leu} does not contain an ms^2 modification. Previous studies have shown that the deficiency of taurine modification in mt-tRNA^{Leu} carrying the A3243G mutation is the primary cause of MELAS (Kirino et al., 2005; Yasukawa et al., 2001). Given the significant association of the heteroplasmy level with the ms^2 modification level in MELAS patients, our results suggest that the myopathy in MELAS is caused not only by a decoding error at the Leu codon but also by decoding errors occurring at multiple codons, including Leu, Phe, Tyr, Trp, and Ser codons.

The reason A3243G in mt-tRNA^{Leu} is associated with decreased modifications in other mt-tRNAs remains unclear. Although further studies are required to reveal the molecular mechanism, our results suggest that oxidative stress may be one of the reasons for this finding. Cdk5rap1 requires two [4Fe-4S] clusters for ms^2 group insertion (Forouhar et al., 2013); therefore, it is conceivable that ROS, such as H₂O₂ or

ONOO⁻, may oxidize these [4Fe-4S] clusters and inactivate Cdk5rap1. In support of our hypothesis, ROS-treated cells exhibited a rapid decrease in ms^2 modification that was effectively reversed by antioxidants. Thus, ROS generated by the mutation in mt-tRNA^{Leu} might impair Cdk5RAP1-mediated ms^2 modification, which might further amplify mitochondrial dysfunction and ultimately accelerate myopathy in MELAS patients. In addition to mitochondrial disease, ms^2 modifications might be involved in a wide variety of human diseases in which ROS have been previously implicated, such as cardiac dysfunction and cancer (Schieber and Chandel, 2014).

In conclusion, this study reveals a unique quality control system in mitochondria by which the ms^2 modification of mt-tRNAs dynamically regulates mitochondrial protein synthesis and contributes to the development of myopathy in vivo. Our findings have important physiological implications for the basic mechanism of mitochondrial protein synthesis and provide insights into the pathological mechanism of mitochondrial disease.

EXPERIMENTAL PROCEDURES

Please see the [Supplemental Experimental Procedures](#) for additional details.

Animals

Cdk5rap1 KO mice were generated by crossing transgenic mice with exon 5 and 6 of *Cdk5rap1* floxed with LoxP sequence, with transgenic mice expressing Cre recombinase under the control of the CAG promoter. Mice were backcrossed to C57BL6/J mice for at least seven generations to eliminate Cre transgene and control genetic background. Littermates of WT and KO mice (8–12 weeks old) were used for experiments unless otherwise specified. Animals were housed at 25°C with 12 hr light and 12 hr dark cycles. A KD was purchased from Research Diets (D12369B). All animal procedures were approved by the Animal Ethics Committee of Kumamoto University (Approval ID, C25-163). Detailed information on genotyping can be found in the [Supplemental Experimental Procedures](#).

Luciferase Assay

E. coli colonies were transformed with plasmids encoding dual luciferase for detecting decoding error, *GST-Cdk5rap1*, or dominant-negative *GST-Cdk5rap1*. Colonies were cultured at 37°C, and isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to the cultures at a final concentration of 1 mM. After 1 hr of incubation, the cultures were harvested for the luciferase assay using the Dual-Luciferase Reporter Assay System (Promega). Detailed procedures for detecting decoding error can be found in the [Supplemental Experimental Procedures](#).

Cell Culture and Transfection

Mammalian cells were grown in DMEM high-glucose medium (GIBCO) supplemented with 10% fetal bovine serum (FBS, HyClone) at 37°C and 5% CO₂.

Figure 6. The Deficiency in ms^2 Modification Compromises Mitochondrial Protein Quality under Stressed Conditions

- (A) Steady-state levels of CI, CIII, CIV, and CV in heart tissue in WT and KO mice treated with NC, KD, and TAC surgery were examined by BN-PAGE.
- (B) The steady-state levels of CI protein NDUFB8, CIV protein MTCOI, and CIII protein UQCRC2 in heart tissue in WT and KO mice treated with NC, KD, and TAC surgery were examined by BN-PAGE followed by western blotting. UQCRC2 was used as a loading control.
- (C) The protein levels of Yme111, Afg 3l2, and Lonp1 were examined in heart tissues from WT and KO mice treated with NC, KD, and TAC surgery. Membranes stained with CBB were used as a loading control.
- (D) Enhanced polyubiquitination was observed in the mitochondria in the hearts of KO mice under each stress.
- (E) WT and KO cells transfected with Parkin-YFP were treated with 10 μM FCCP for 2 and 4 hr; bar, 10 μm.
- (F) WT and KO cells were treated with 10 μM FCCP for 2 and 4 hr. The cells were then stained with an anti-LC3 antibody and Mitotracker. The inserted box shows mitochondria surrounded by the LC3 protein and is magnified in the bottom panels; bar, 10 μm.
- (G) Electron microscopy of mitochondria in heart tissue from WT and KO mice treated with KD or TAC surgery. Arrows indicate the autophagic vacuoles; bar, 10 μm.




Figure 7. Association of ms² Modifications with MELAS

(A–D) Negative correlation of the ms² modification level of mt-tRNA^{Trp} (A), mt-tRNA^{Phe} (B), mt-tRNA^{Ser(UCN)} (C), or mt-tRNA^{Tyr} (D) with the heteroplasmy level in MELAS patients; n = 18 each.

(E) Treatment with H₂O₂ reduced the level of ms² modification of mt-tRNAs in HeLa cells. Cells were treated with 10 μ M or 50 μ M H₂O₂ for 24 hr, and the ms² modification levels were examined by qPCR; n = 4 each.

(F) Time-dependent decreases in ms² modification after H₂O₂ treatment in HeLa cells. Cells were treated with 50 μ M H₂O₂ for 6, 12, and 24 hr; n = 4 each.

(G) HeLa cells were treated with 50 μ M H₂O₂ for 24 hr in the presence or absence of 10 mM pyruvate. The decrease in the ms² modification level was prevented by pyruvate; n = 4 each.

(H) HeLa cells were treated with NO donors, 100 μ M SNAP or 100 μ M NOC18 for 24 hr in the presence or absence of PTIO. The ms² modification level of mt-tRNA^{Trp} was examined by qPCR; n = 4 each. Data are the mean \pm SEM. *p < 0.05. **p < 0.01, ****p < 0.0001.

map (Cheadle et al., 2003). The sequences of primers used can be found in the [Supplemental Experimental Procedures](#).

Analysis of tRNA Modification

Total RNAs were isolated from bacteria and tissues using Trizol reagent (Invitrogen). RNA was digested with Nuclease P1 (Sigma) and subjected to mass spectrometry (Agilent 6460). For detecting tRNA modification using the qPCR-based method, we adapted a protocol described previously (Xie et al., 2013). Detailed procedures for the mass spectrometry and qPCR method can be found in the [Supplemental Experimental Procedures](#). To measure the tRNA modification level in blood samples, blood samples were collected from MELAS patients using standard procedures approved by Kurume University (IRB#9715).

ATP Measurement

Small pieces of skeletal muscle and heart tissue were immediately dissected after sacrificing mice

and snap frozen in liquid nitrogen until measurement. ATP was measured using the ATP Bioluminescence Assay Kit following the manufacturer's protocol (TA100, WAKO). The luminescence was measured using a Centro XS³ LB960 (Berthold) and normalized to total protein concentration.

Cardiac Function Examination

Echocardiographs were examined in M-mode while the mice were under anesthesia using the Vevo2100 system (Fujifilm VisualSonics, Inc.) according to the manufacturer's instructions.

Statistical Analysis

Statistical analyses were performed using Prism 6 Software (GraphPad Software). An unpaired Student t test was used to test the differences between two groups. Analysis of variance (one-way ANOVA or two-way ANOVA) was used to test the difference among multiple groups followed by a post hoc examination of the p value between two groups. A two-tailed p value of 0.05 was considered statistically significant.

Transfection of the plasmid DNA was performed with Lipofectamine 2000 (Invitrogen).

Oxygen Consumption

The oxygen consumption rate in MEF cells and intact mitochondria was measured using an XF24 Analyzer (Seahorse Bioscience). The oxygen consumption rate was normalized to the total protein concentration for measurement in cells. Detailed procedures for the respiratory assay can be found in the [Supplemental Experimental Procedures](#).

Gene Expression Assay

RNA was extracted from tissues using Trizol (Invitrogen) following the manufacturer's instructions. Quantitative PCR (qPCR) was performed using SYBR Premix Ex Taq (TAKARA). For examination of the expression levels of oxidative response genes, the results were normalized to the geometric mean of multiple reference genes (Hprt1, RPL13A, B2M, GAPDH, ACT). Then, a Z-transformation was applied to the results to calculate the Z score and construct a heat-

SUPPLEMENTAL INFORMATION

Supplemental Information includes seven figures and Supplemental Experimental Procedures and can be found with this article at <http://dx.doi.org/10.1016/j.cmet.2015.01.019>.

AUTHOR CONTRIBUTIONS

F.-Y.W. and K.T. designed the experiments and wrote the manuscript. F.-Y.W. and B.Z. performed the experiments. Takeo Suzuki and Tsutomu Suzuki performed the mass spectrometry experiments. H.H., K.M., and Y.O. performed cardiac examinations. Y.U. and S.M. performed TAC surgery and cardiac examinations. H. Michiue, A.F., and H. Matsui performed electron microscopy. Y.K. provided the blood samples. N.T., P.X., and T.K. performed the qPCR-based examination of tRNA modifications.

ACKNOWLEDGMENTS

We thank Nobuko Maeda for the technical assistance. This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, by the Japan Society for the Promotion of Science (JSPS) through its "Funding Program for Next Generation World-Leading Researchers," and by the Takeda Science Foundation.

Received: September 5, 2014

Revised: November 24, 2014

Accepted: January 26, 2015

Published: March 3, 2015

REFERENCES

- Agris, P.F. (2004). Decoding the genome: a modified view. *Nucleic Acids Res.* *32*, 223–238.
- Arragain, S., Handelman, S.K., Forouhar, F., Wei, F.Y., Tomizawa, K., Hunt, J.F., Douki, T., Fontecave, M., Mulliez, E., and Atta, M. (2010). Identification of eukaryotic and prokaryotic methylthiotransferase for biosynthesis of 2-methylthio-*N*⁶-threonylcarbamoyladenine in tRNA. *J. Biol. Chem.* *285*, 28425–28433.
- Cheadle, C., Vawter, M.P., Freed, W.J., and Becker, K.G. (2003). Analysis of microarray data using Z score transformation. *J. Mol. Diagn.* *5*, 73–81.
- Chen, Y., and Dom, G.W., 2nd. (2013). PINK1-phosphorylated mitofusin 2 is a Parkin receptor for culling damaged mitochondria. *Science* *340*, 471–475.
- Crimi, M., Bordoni, A., Menozzi, G., Riva, L., Fortunato, F., Galbiati, S., Del Bo, R., Pozzoli, U., Bresolin, N., and Comi, G.P. (2005). Skeletal muscle gene expression profiling in mitochondrial disorders. *FASEB J.* *19*, 866–868.
- Dai, D.F., Hsieh, E.J., Liu, Y., Chen, T., Beyer, R.P., Chin, M.T., MacCoss, M.J., and Rabinovitch, P.S. (2012). Mitochondrial proteome remodeling in pressure overload-induced heart failure: the role of mitochondrial oxidative stress. *Cardiovasc. Res.* *93*, 79–88.
- DiMauro, S., and Schon, E.A. (2003). Mitochondrial respiratory-chain diseases. *N. Engl. J. Med.* *348*, 2656–2668.
- Durieux, J., Wolff, S., and Dillin, A. (2011). The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell* *144*, 79–91.
- Forouhar, F., Arragain, S., Atta, M., Gambarelli, S., Mouesca, J.M., Hussain, M., Xiao, R., Kieffer-Jaquinod, S., Seetharaman, J., Acton, T.B., et al. (2013). Two Fe-S clusters catalyze sulfur insertion by radical-SAM methylthiotransferases. *Nat. Chem. Biol.* *9*, 333–338.
- Grimsrud, P.A., Carson, J.J., Hebert, A.S., Hubler, S.L., Niemi, N.M., Bailey, D.J., Jochem, A., Stapleton, D.S., Keller, M.P., Westphal, M.S., et al. (2012). A quantitative map of the liver mitochondrial phosphoproteome reveals post-translational control of ketogenesis. *Cell Metab.* *16*, 672–683.
- Hoshino, A., Mita, Y., Okawa, Y., Ariyoshi, M., Iwai-Kanai, E., Ueyama, T., Ikeda, K., Ogata, T., and Matoba, S. (2013). Cytosolic p53 inhibits Parkin-mediated mitophagy and promotes mitochondrial dysfunction in the mouse heart. *Nat. Commun.* *4*, 2308.
- Houtkooper, R.H., Mouchiroud, L., Ryu, D., Moullan, N., Katsyuba, E., Knott, G., Williams, R.W., and Auwerx, J. (2013). Mitonuclear protein imbalance as a conserved longevity mechanism. *Nature* *497*, 451–457.
- Ishikawa, K., Kimura, S., Kobayashi, A., Sato, T., Matsumoto, H., Ujii, Y., Nakazato, K., Mitsugi, M., and Maruyama, Y. (2005). Increased reactive oxygen species and anti-oxidative response in mitochondrial cardiomyopathy. *Circ. J.* *69*, 617–620.
- Jenner, L.B., Demeshkina, N., Yusupova, G., and Yusupov, M. (2010). Structural aspects of messenger RNA reading frame maintenance by the ribosome. *Nat. Struct. Mol. Biol.* *17*, 555–560.
- Karamanlidis, G., Lee, C.F., Garcia-Menendez, L., Kolwicz, S.C., Jr., Suthammarak, W., Gong, G., Sedensky, M.M., Morgan, P.G., Wang, W., and Tian, R. (2013). Mitochondrial complex I deficiency increases protein acetylation and accelerates heart failure. *Cell Metab.* *18*, 239–250.
- Kirino, Y., Yasukawa, T., Ohta, S., Akira, S., Ishihara, K., Watanabe, K., and Suzuki, T. (2004). Codon-specific translational defect caused by a wobble modification deficiency in mutant tRNA from a human mitochondrial disease. *Proc. Natl. Acad. Sci. USA* *101*, 15070–15075.
- Kirino, Y., Goto, Y., Campos, Y., Arenas, J., and Suzuki, T. (2005). Specific correlation between the wobble modification deficiency in mutant tRNAs and the clinical features of a human mitochondrial disease. *Proc. Natl. Acad. Sci. USA* *102*, 7127–7132.
- Kubli, D.A., and Gustafsson, Å.B. (2012). Mitochondria and mitophagy: the yin and yang of cell death control. *Circ. Res.* *111*, 1208–1221.
- Laffel, L. (1999). Ketone bodies: a review of physiology, pathophysiology and application of monitoring to diabetes. *Diabetes Metab. Res. Rev.* *15*, 412–426.
- Machnicka, M.A., Milanowska, K., Osman Oglou, O., Purta, E., Kurkowska, M., Olchowik, A., Januszewski, W., Kalinowski, S., Dunin-Horkawicz, S., Rother, K.M., et al. (2013). MODOMICS: a database of RNA modification pathways—2013 update. *Nucleic Acids Res.* *41* (Database issue), D262–D267.
- Murphy, M.P. (2009). How mitochondria produce reactive oxygen species. *Biochem. J.* *417*, 1–13.
- Reiter, V., Matschkal, D.M., Wagner, M., Globisch, D., Kneutinger, A.C., Müller, M., and Carell, T. (2012). The CDK5 repressor CDK5RAP1 is a methylthiotransferase acting on nuclear and mitochondrial RNA. *Nucleic Acids Res.* *40*, 6235–6240.
- Runkel, E.D., Liu, S., Baumeister, R., and Schulze, E. (2013). Surveillance-activated defenses block the ROS-induced mitochondrial unfolded protein response. *PLoS Genet.* *9*, e1003346.
- Schieber, M., and Chandel, N.S. (2014). ROS function in redox signaling and oxidative stress. *Curr. Biol.* *24*, R453–R462.
- Steinthorsdottir, V., Thorleifsson, G., Reynisdottir, I., Benediktsson, R., Jonsdottir, T., Walters, G.B., Styrkarsdottir, U., Gretarsdottir, S., Emilsson, V., Ghosh, S., et al. (2007). A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat. Genet.* *39*, 770–775.
- Suzuki, T. (2005). Biosynthesis and function of tRNA wobble modifications. *Topics Curr. Genet.* *12*, 24–69.
- Suzuki, T., and Suzuki, T. (2014). A complete landscape of post-transcriptional modifications in mammalian mitochondrial tRNAs. *Nucleic Acids Res.* *42*, 7346–7357.
- Suzuki, T., Suzuki, T., Wada, T., Saigo, K., and Watanabe, K. (2002). Taurine as a constituent of mitochondrial tRNAs: new insights into the functions of taurine and human mitochondrial diseases. *EMBO J.* *21*, 6581–6589.
- Suzuki, T., Nagao, A., and Suzuki, T. (2011). Human mitochondrial tRNAs: biogenesis, function, structural aspects, and diseases. *Annu. Rev. Genet.* *45*, 299–329.
- Torres, A.G., Battle, E., and Ribas de Pouplana, L. (2014). Role of tRNA modifications in human diseases. *Trends Mol. Med.* *20*, 306–314.
- Urbanavicius, J., Qian, Q., Durand, J.M., Hagervall, T.G., and Björk, G.R. (2001). Improvement of reading frame maintenance is a common function for several tRNA modifications. *EMBO J.* *20*, 4863–4873.

- Wei, F.Y., Suzuki, T., Watanabe, S., Kimura, S., Kaitsuka, T., Fujimura, A., Matsui, H., Atta, M., Michiue, H., Fontecave, M., et al. (2011). Deficit of tRNA(Lys) modification by Cdkal1 causes the development of type 2 diabetes in mice. *J. Clin. Invest.* *121*, 3598–3608.
- Wenz, T., Luca, C., Torraco, A., and Moraes, C.T. (2009). mTERF2 regulates oxidative phosphorylation by modulating mtDNA transcription. *Cell Metab.* *9*, 499–511.
- Xie, P., Wei, F.Y., Hirata, S., Kaitsuka, T., Suzuki, T., Suzuki, T., and Tomizawa, K. (2013). Quantitative PCR measurement of tRNA 2-methylthio modification for assessing type 2 diabetes risk. *Clin. Chem.* *59*, 1604–1612.
- Yasukawa, T., Suzuki, T., Ishii, N., Ohta, S., and Watanabe, K. (2001). Wobble modification defect in tRNA disturbs codon-anticodon interaction in a mitochondrial disease. *EMBO J.* *20*, 4794–4802.