



## ORIGINAL ARTICLE

# Effects of temperature, water activity and incubation time on fungal growth and aflatoxin B1 production by toxinogenic *Aspergillus flavus* isolates on sorghum seeds



Amani Lahouar<sup>a,b</sup>, Sonia Marin<sup>b</sup>, Ana Crespo-Sempere<sup>b</sup>, Salem Saïd<sup>a</sup>,  
Vicente Sanchis<sup>b,\*</sup>

<sup>a</sup> Laboratory of Biochemistry, Unity of Mycotoxicology, Faculty of Medicine of Sousse, University of Sousse, Sousse, Tunisia

<sup>b</sup> Food Technology Department, University of Lleida, XaRTA-UTPV, Agrotecnico center, Lleida, Spain

Received 23 May 2015; accepted 23 October 2015

Available online 23 February 2016

## KEYWORDS

Water activity;  
Incubation time;  
Temperature;  
*Aspergillus flavus*;  
Aflatoxin B1;  
Fungal growth

**Abstract** Sorghum, which is consumed in Tunisia as human food, suffers from severe colonization by several toxigenic fungi and contamination by mycotoxins. The Tunisian climate is characterized by high temperature and humidity that stimulates mold proliferation and mycotoxin accumulation in foodstuffs. This study investigated the effects of temperature (15, 25 and 37 °C), water activity ( $a_w$ , between 0.85 and 0.99) and incubation time (7, 14, 21 and 28 d) on fungal growth and aflatoxin B1 (AFB1) production by three *Aspergillus flavus* isolates (8, 10 and 14) inoculated on sorghum grains. The Baranyi model was applied to identify the limits of growth and mycotoxin production. Maximum diameter growth rates were observed at 0.99  $a_w$  at 37 °C for two of the isolates. The minimum  $a_w$  needed for mycelial growth was 0.91 at 25 and 37 °C. At 15 °C, only isolate 8 grew at 0.99  $a_w$ . Aflatoxin B1 accumulation could be avoided by storing sorghum at low water activity levels ( $\leq 0.91 a_w$ ). Aflatoxin production was not observed at 15 °C. This is the first work on the effects of water activity and temperature on *A. flavus* growth and AFB1 production by *A. flavus* isolates on sorghum grains.

© 2015 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [vsanchis@tecal.udl.cat](mailto:vsanchis@tecal.udl.cat) (V. Sanchis).

**PALABRAS CLAVE**

Actividad de agua;  
Tiempo de  
incubación;  
Temperatura;  
*Aspergillus flavus*;  
Aflatoxina B1;  
Crecimiento fúngico

### Efectos de la temperatura, la actividad de agua y el tiempo de incubación en el crecimiento fúngico y la producción de aflatoxina B1 por aislados toxicogénicos de *Aspergillus flavus* en sorgo

**Resumen** El sorgo, que se consume en Túnez como alimento humano, puede sufrir la colonización severa de varios hongos toxicogénicos, con la consiguiente bioacumulación de micotoxinas. Además, el clima de Túnez, caracterizado por las altas temperaturas y humedad, estimula el crecimiento fúngico y la acumulación de micotoxinas en los productos alimenticios. Este estudio investigó los efectos de la temperatura (15, 25 y 37 °C), la actividad de agua ( $a_w$ ) (entre 0,85 y 0,99) y el tiempo de incubación (7, 14, 21 y 28 días) sobre el crecimiento y la producción de aflatoxina B1 (AFB1) de 3 aislados de *Aspergillus flavus* (designados como 8, 10 y 14) que se inocularon sobre granos de sorgo. El modelo Baranyi se aplicó para identificar los límites del crecimiento y la producción de micotoxinas. Las tasas máximas de crecimiento para 2 de los aislados se observaron en la combinación 0,99  $a_w$  y 37 °C. La  $a_w$  mínima necesaria para el crecimiento del micelio fue de 0,91 a 25 °C y 37 °C. A 15 °C, solo el aislado 8 creció a 0,99  $a_w$ , pero fue incapaz de producir la aflatoxina B1. Es posible evitar la acumulación de aflatoxina B1 en el sorgo almacenándolo a baja actividad de agua ( $\leq 0,91 a_w$ ). Este es el primer trabajo que ha estudiado el efecto de la actividad del agua y la temperatura sobre el crecimiento de aislados de *A. flavus* y su producción de aflatoxina B1 en granos de sorgo.

© 2015 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Aflatoxins are a group of toxic chemical compounds that are characterized by their immunotoxic, mutagenic and carcinogenic effects<sup>17</sup>. Aflatoxins are classified as group I carcinogens by the International Agency for Research on Cancer<sup>38</sup>. Among them, aflatoxin B1 (AFB1) is the most frequent and most potent toxin. *A. flavus* and to a lesser extent *Aspergillus parasiticus* are among the most widely studied fungal species, as a result of their ability to produce aflatoxins and their potential to persist pre- and post-harvest as a pathogen and saprophyte in the food supply.

The growth of molds and the accumulation of mycotoxins in food and feedstuff are influenced by multiple variables, such as water activity ( $a_w$ ), temperature, pH, atmosphere composition, substrate, interaction among species, and time<sup>3,37</sup>. Generally, relative humidity and temperature are considered to be the most critical factors during drying and storage<sup>14</sup>. Various physical and chemical methods have been recommended to reduce mycotoxins, but only a few have been accepted for practical use. Thus, the prevention of fungal growth and mycotoxin production represent key steps in risk management. Many kinetic models have been developed and used to model the growth of toxigenic molds in various food substrates. The effect of biotic and abiotic factors on growth and aflatoxin production have been widely studied using mathematical modeling<sup>4,6,14,16,18,20,28,39</sup>; however, no studies exist concerning their growth on sorghum seeds.

Sorghum (*Sorghum bicolor* (L.) Moench) is a drought-resistant crop and an important food resource in terms of nutritional and economic values, especially in semi-arid environments<sup>7</sup>. Sorghum grains can be colonized by several fungal genera during the panicle and grain development stages<sup>21,22</sup>. Several species of *Aspergillus*, *Alternaria*, *Fusarium*, *Cladosporium*, *Curvularia* and *Penicillium* are

among the most prevalent grain mold pathogens in sorghum<sup>34</sup>. Moreover, a number of fungal isolates from this cereal primarily belonging to the genera *Fusarium* (*F. verticillioides*, *F. graminearum*, *F. equiseti*), *Alternaria* (*A. alternata*), *Aspergillus* (*A. flavus*), and *Penicillium* (*P. funiculosum*) have been reported to produce mycotoxins<sup>27,30,36</sup>.

The Tunisian climate is characterized by high temperature and humidity levels that seem to stimulate toxigenic mold growth and mycotoxin production. The Tunisian population consumes large amounts of cereals and cereal-based products, such as wheat, barley and sorghum. The occurrence of aflatoxins in commercialized sorghum in Tunisia was reported for the first time in 1977<sup>2</sup>, and several studies confirmed aflatoxin contamination in Tunisian sorghum<sup>1,8,9,10,19</sup>. According to Oueslati *et al.*<sup>10</sup>, AFB1-contaminated cereals represent a real problem for Tunisian consumers. Therefore, it is important to develop prevention and control strategies to minimize aflatoxin contamination in sorghum.

The objectives of the present work were to determine: (i) the effects of the interactions of water activity, temperature and incubation time on fungal growth and aflatoxin B1 production by Tunisian *Aspergillus flavus* isolates cultured on sorghum seeds and (ii) to identify the  $a_w$  and temperature limiting conditions for aflatoxin B1 production. The Baranyi model<sup>2</sup> was applied to predict the growth of aflatoxigenic isolates of *A. flavus* on sorghum seeds as a function of water activity and temperature. Elucidating the ecology and physiology of *A. flavus* will provide tools to manage AFB1 in sorghum.

## Materials and methods

### Fungal isolates

Three isolates of *A. flavus* (8, 10 and 14) were used in this study. The three isolates were obtained from sorghum sam-

ples collected from the retail market of the Sahel region in Tunisia in 2011 and were kept in the Food Technology department collection of the University of Lleida. All isolates were previously found to produce AFB<sub>1</sub> when cultivated in Czapeck Yeast Extract Agar (CYA) and sorghum seeds. The identification of the *A. flavus* isolates was done microscopically<sup>29</sup>. Molecular characterization of the isolates was also performed by PCR amplification using the primer pair AfAflT-F and AfAflT-R<sup>35</sup>. Two DNA extracts of *A. flavus* [CECT (Spanish Type Culture Collection) 2695 and *A. parasiticus* CECT 2681] were used as references.

### Experimental design and statistical analysis

A full factorial design of three temperatures (15, 25 and 37 °C) × six water activities ( $a_w$ ) (0.85, 0.88, 0.91, 0.94, 0.97 and 0.99) × three isolates (8, 10 and 14) was used. Colony diameters and AFB<sub>1</sub> production after 7, 14, 21 and 28 d were the dependent variables. Three replicates were included for each treatment. The effects of temperature and  $a_w$  on mycelial growth and AFB<sub>1</sub> production after 7, 14, 21 and 28 d were statistically analyzed by analysis of variance (ANOVA) using Statgraphics Plus 5.1 (Manugistics, Inc., MD, USA).

### Preparation of sorghum

The study was performed *in vitro* on sorghum grains. To determine the growth and aflatoxin patterns on sorghum grains, different amounts of water (calculated from a moisture absorption curve for the sorghum batch used) were added to subsamples of 100 g of grain in flasks. The sorghum was allowed to equilibrate at 4 °C for 48 h with periodic shaking. Once sealed, the flasks were autoclaved for 20 min at 121 °C, and then single layers of grain were carefully placed in 9 cm sterile plastic Petri dishes on a flow bench. The final  $a_w$  levels were 0.85, 0.88, 0.91, 0.94, 0.97 and 0.99. All  $a_w$  values were verified with an AquaLab 3 (Decagon Devices, Inc., WA, USA) with an accuracy of ±0.003 before, during and at the end of the experiment.

### Inocula preparation and incubation

*A. flavus* isolates were sub-cultured on potato dextrose agar (PDA) plates and incubated at 25 °C for 7 d to enable significant sporulation. After incubation, a sterile inoculation loop was used to remove the conidia from the PDA plates. The conidia were suspended in an aqueous solution of 0.05% (w/v) Tween 80. After homogenizing, the spore concentrations were determined using the Thoma cell counting chamber. The suspensions were diluted to adjust the final concentration to vary between 1 and 5 × 10<sup>5</sup> spores/ml. Triplicate plates with single layers of sorghum for each water activity and temperature combination were inoculated centrally with a needlepoint load and incubated for 28 d. Previous experiments showed that the number of spores inoculated through this technique was 10–100 spores on each inoculation point. The incubation temperatures for all three isolates were 15, 25 and 37 °C, and the  $a_w$  values were 0.85, 0.88, 0.91, 0.94, 0.97 and 0.99. Plates with the same temperature and  $a_w$  level were stacked in plastic

chambers together with 250 ml of glycerol/water solution of the same  $a_w$  to maintain the relative humidity equilibrium in the chambers as constant as possible.

### Growth measurement and growth rate calculation

The diameter of each growing colony was determined by measurement of two right-angled rays of the colony. Measurements were recorded daily during the 28 d of growth. The growth rate ( $\mu$ ) (mm/d) and growth time ( $\lambda$ ) (d) were evaluated for each isolate at each different combination of  $a_w$  and temperature by plotting the colony diameter extension (mm) against time (d) and fitting the Baranyi<sup>2</sup> model using Statgraphics Plus. This model represents the three growth phases (lag, linear, and asymptotic phase). The proposed primary model has the general form of:

$$y = y_0 + \mu_{\max} A - \ln \left( 1 + \frac{[\exp(\mu_{\max} [A]^{-1})]}{\exp(y_{\max} - y_0)} \right)$$

$$A = t + \left( \frac{1}{\mu_{\max}} \right) \ln [\exp(-\mu_{\max} t) + \exp(-\mu_{\max} \lambda) - \exp(-\mu_{\max} t - \mu_{\max} \lambda)]$$

Where  $y$  is the colony diameter (or radius),  $y_0$  is the initial colony diameter (or radius, usually zero),  $y_{\max}$  is the maximum colony diameter (or radius) attained (asymptotic value),  $\mu_{\max}$  is the maximum specific growth rate (defined as the slope of the growth curve at the point of inflexion),  $\lambda$  is the lag period (*i.e.* the intersection of the line defining the maximum specific growth rate with the x-axis), and  $t$  is the time.

### Aflatoxin B<sub>1</sub> production and quantification

AFB<sub>1</sub> production was determined after 7 and 14 d of incubation under the optimal growth conditions [at 25 °C (0.97 and 0.99  $a_w$ ) and at 37 °C (0.94, 0.97, and 0.99  $a_w$ )] and after 21 and 28 d of incubation under suitable but not optimal conditions [at 37 °C (0.91  $a_w$ ), at 25 °C (0.91 and 0.94  $a_w$ ) and at 15 °C (0.99  $a_w$ )] (Table 1). AFB<sub>1</sub> was extracted from the sorghum samples (20 g) with 100 ml of 4% acetonitrile aqueous solution containing potassium chloride (9:1). The mixture was adjusted to a pH of 1.5 with undiluted hydrochloric acid, shaken for 20 min and filtered through a Whatman paper. The filtrate was defatted with 100 ml of hexane. The solution was shaken for 10 min. After separation, the upper phase (hexane) was discarded. This step was repeated with 50 ml of hexane. The lower phase was extracted with 50 ml of chloroform and 50 ml of deionized water, and the solution was shaken for 10 min. After separation, the lower phase (chloroform) was collected. The upper phase was re-extracted twice with 25 ml of chloroform and 25 ml of deionized water. The chloroform extracts were evaporated to near dryness using a rotary evaporator at 40 °C. AFB<sub>1</sub> was resuspended in 2 ml of methanol and evaporated to dryness under nitrogen. Finally, the sample was analyzed with HPLC.

**Table 1** Minimal conditions for aflatoxin B1 (AFB1) production by three Tunisian *Aspergillus flavus* isolates on sorghum seeds

Isolate	Temperature (°C)	Water activity ( $a_w$ )	Incubation time (d)	AFB1 concentration (ng/g) <sup>a</sup>
8	37	0.94	7	0.23 ± 0.34
	25	0.94	21	0.11 ± 0.02
10	37	0.97	7	1.33 ± 0.28
	25	0.97	14	0.22 ± 0.08
14	37	0.97	7	1.64 ± 0.28
	25	0.97	14	0.17 ± 0.06

<sup>a</sup> Detection limit = 0.075 ng/g.

The determination of AFB1 was performed using a Waters (Milford, MA, USA) chromatograph equipped with a reverse phase C<sub>18</sub> silica gel column (Water Spherisorb 3 μm ODS2 4.6 × 150 mm, Milford, MA, USA), followed by fluorescence detection ( $\lambda_{exc}$  362 nm;  $\lambda_{em}$  425 nm) (Waters 2475 Fluorescence Detector, Waters, Milford, MA, USA). The mobile phase consisted of water–acetonitrile–methanol (700:170:170) with a flow rate of 1 ml/min. The detection limit (LOD) of the method was 0.075 ng/g. The injection volume was 10 μl, and the retention time was 8 min. Post-column derivatization was achieved using a photochemical reactor for enhanced detection (PHRED) (LCTech UVE, Dorfen, Germany).

## Results and discussion

Previous works noted that data obtained regarding mycotoxin production on culture media could not be directly extrapolated to the natural substrates<sup>40</sup>. Thus, the present study was performed directly on sorghum grains.

### Effects of water activity and temperature on mycelial growth


Figure 1 shows an example of the observed colony diameters for *A. flavus* isolates 8, 10 and 14 at two water activities (0.90 and 0.97) and their respective growth adjusted to the Baranyi model. Under optimum growth conditions, the colony diameters increased linearly until the colonies reached the edge of the Petri dishes. Under limiting growth conditions, a clear lag time was observed prior to growth (Fig. 2). The Baranyi model was used to determine two growth parameters: the lag phases (d) and the growth rates (mm/d). The percentage of variance explained by the model ( $R^2$  coefficient) obtained for the different treatments (combinations) varied between 85.85% and 99.83%, with 31 combinations showing no growth and only two combinations exhibiting 54 < 90%. The low MSEs (0.017–0.673) and the high  $R^2$  values (85.85–99.83%) indicated that the evaluated model could sufficiently determine the growth parameters.

Analysis of variance for the growth rate showed that the single factors  $a_w$  and temperature and the interactions isolate × temperature,  $a_w$  × temperature and isolate ×  $a_w$  × temperature had significant influences on the growth rate ( $p < 0.05$ ). On average, the three isolates behaved similarly, as noted by Mohale et al.<sup>24</sup> with atoxigenic and toxigenic strains of *A. flavus*. The combined effects of  $a_w$  and temperature on the growth rates (mm/d) of the three isolates tested are shown in Fig. 2. The maximum

growth rates achieved were 16.29, 14.91 and 14.22 mm/d by isolates 14, 8 and 10, respectively, at 37 °C and 0.99  $a_w$ . In this study, the optimum temperature for growth was 37 °C for all  $a_w$  levels tested (Fig. 2). Additionally, 25 °C was a very favorable temperature and yielded high growth rates. Similar results for *A. flavus* isolates were found by Horn<sup>23</sup>. The extent of seed colonization by *Aspergillus* section *Flavi* grew with the increase of water activity and temperature to the optimal conditions of  $a_w = 0.96$  and 37 °C. Samapundo et al.<sup>33</sup> showed that *A. flavus* had optimum growth on corn at 30 °C. Schindler et al.<sup>28</sup> reported that growth of *A. flavus* was most favored in wort agar between 29 and 35 °C. Temperatures between 25 and 37 °C were not tested in our study; however, the mycelial extension of Tunisian *A. flavus* isolates was significantly higher at 37 °C than at 25 °C ( $p < 0.05$ ). Figure 2 showed that very short lag phases were encountered at high temperatures (25 and 37 °C) and  $a_w$  (0.97 and 0.99  $a_w$ ). The shorter lag phase period was observed for isolates 10 and 8 at 25 °C and for isolate 14 at 37 °C. Samapundo et al.<sup>33</sup> showed that an increase in temperature from 30 to 37 °C resulted in a decrease in the colony growth rate and an increase in the lag phase.

No growth was detected at 15 °C at any of the  $a_w$  conditions tested for isolates 10 and 14; however, isolate 8 grew at 0.99  $a_w$ . The minimum growth rate observed was 0.9 mm/d, which was calculated for isolate 14 at 37 °C and 0.91  $a_w$ . No growth was detected for isolates 10 and 14 at 25 °C and 0.91  $a_w$ . However, at 37 °C all isolates grew at 0.91  $a_w$ . No growth at 0.85 and 0.88  $a_w$  was detected after 28 d at all tested temperatures. However, Pitt and Miscamble<sup>11</sup> reported that the minimum water activity for growth of *A. flavus* was between 0.80 and 0.83. The discrepancy in these results may be due to differences in the media used and the availability of nutrients. Some components of the pericarp layer of the grain, such as tannins and high levels of phenol-based pigments, and some antifungal proteins can affect fungal growth<sup>11</sup>. Similar studies have also indicated that the availability of nutrients may affect the chances of growth at marginal  $a_w$  values<sup>13</sup>.

Maximum growth rates were observed for the three isolates at 25 and 37 °C at 0.97 and 0.99  $a_w$ . Additionally, the *A. flavus* isolates assayed grew faster at 0.99  $a_w$  than at 0.97  $a_w$  at 25 and 37 °C, with the exception of isolate 14, where the optimum water activity for growth was 0.97  $a_w$  at 25 °C. Similar results were reported by Marin et al.<sup>18</sup> for *A. flavus* grown on maize extract agar, where the optimal  $a_w$  for growth was 0.994. However, the optimum water activity for growth in some studies was somewhat different from the conditions demonstrated by our data. For instance,



**Figure 1** Effect of water activity (0.99 and 0.97) and temperature 37 °C (◆) and 25 °C (■) on colony diameter (cm) of three *A. flavus* isolates (8, 10 and 14) on single layers of sterile sorghum grains. Values of three replicated are plotted and adjusted with Baranyi's model.

Horn<sup>23</sup> found that the optimum water activity for growth was 0.96  $a_w$ . These differences can be attributed to intraspecific and regional variability among isolates.

### Effects of water activity and temperature on aflatoxin B1 production


AFB1 was the only metabolite considered in the data analysis because AFB2 was rarely detected and only in trace amounts whereas aflatoxins G1 and G2 were never detected. AFB1 has been reported to be the major AFB in natural substrates such as sorghum<sup>1,9,10</sup>. A similar situation was reported by Giorni et al<sup>41</sup>.

Analysis of variance of the effects of  $a_w$  and temperature on AFB1 production by the three *A. flavus* isolates revealed that the single factors  $a_w$  and temperature and the interactions temperature  $\times$  isolates,  $a_w \times$  incubation time and temperature  $\times$  incubation time had a significant effect on

AFB1 production after 7 and 14 d of growth on sorghum seeds ( $p < 0.05$ ). After 21 and 28 d, only the single factors  $a_w$  and temperature and the interaction between water activity and isolates significantly affected AFB1 accumulation ( $p < 0.05$ ).

The maximum accumulation of AFB1 (2.26 ppb) was observed for isolate 8 at 37 °C and 0.99  $a_w$  after 7 d of incubation (Fig. 3). Generally, at 37 °C and 25 °C AFB1 accumulation grew with increasing water activities (0.97 and 0.99  $a_w$ ). No significant differences were found between 0.99 and 0.97  $a_w$  ( $p < 0.05$ ) at 37 °C after 7 d of incubation. At 37 °C, significantly higher amounts of AFB1 ( $p < 0.05$ ) were produced by the three *A. flavus* isolates at 0.97 and 0.99  $a_w$ . However, Montani et al.<sup>32</sup> found that the most favorable  $a_w$  for aflatoxin production was 0.90  $a_w$  at all incubation times studied. In this study, the only isolate that produced AFB1 at 0.94 was isolate 8, which produced 0.23 ng/g at 37 °C after 7 d of incubation and 0.105 ng/g at 25 °C after 21 d. To determine the minimum  $a_w$  that allowed aflatoxin production, we tested  $a_w$  values of 0.85, 0.88 and 0.91  $a_w$ . AFB1 was not detected at






**Figure 2** Effect of water activity and temperature on growth rate (mm/day) and lag phases (day) of three *A. flavus* isolates (8, 10 and 14) on single layers of sterile sorghum grains at different temperatures 37 (■), 25 (■) and 15 °C (■). If all letters next to a bar are different from the letters in another bar, there are significant differences among the treatments, if they have a common letter, they are not significantly different ( $p < 0.05$ ).

25 and 37 °C, whereas growth occurred at 0.91  $a_w$  and 37 °C. AFB1 was detected under a narrower range of water activity than the range that allowed growth (0.94 and 0.99  $a_w$ ). Consequently, it may be possible to avoid aflatoxin accumulation from these strains by storing the commodities at "safe" water activity levels ( $\leq 0.91 a_w$ ). At 15 °C, AFB1 was not detected (Table 1). Hunter<sup>31</sup> proposed 0.84  $a_w$  as the minimal value for aflatoxin production in corn, and Astoreca *et al.*,<sup>26</sup> proposed 0.83 as the minimum value in corn extract medium.

Several researchers reported a different behavior of the *A. flavus* isolates regarding temperature, with optimum values varying from 25 to 30 °C regardless of the media<sup>5,12,25</sup>. For instance, Schindler *et al.*<sup>28</sup>, reported that maximal production of aflatoxins on wort media occurred at 24 °C, whereas the optimal temperature for AFB1 production was 35 °C.

The incubation period that allowed higher AFB1 accumulation depended on the temperature, water activity and isolate tested. At the optimal temperature (37 °C) and 0.99  $a_w$ , all isolates required 7 d to reach maximum

AFB1 production; no AFB1 production was detected after 14 d under the same conditions. At water activities above 0.94  $a_w$ , AFB1 accumulated at 37 °C was lower after 14 d of incubation on sorghum grains than after 7 d ( $p < 0.0001$ ). The results showed that the optimal  $a_w$  for AFB1 accumulation at 25 °C was 0.99  $a_w$ . At 25 °C, the amounts of AFB1 detected at 0.97 and 0.99  $a_w$  after 14 d of incubation increased compared to the amounts produced after 7 d of incubation, with the exception of isolate 8 at 0.99  $a_w$ , whose AFB1 accumulation was higher after 7 d of incubation ( $p < 0.001$ ). AFB1 decrease following the peak of maximum production was observed under the optimal conditions (37 °C and 0.97–0.99  $a_w$ ). The decrease in AFB1 concentration may be explained by its degradation caused by the mold itself. The detection of AFB1 after 21 d and 28 d of incubation showed that only isolate 8 produced AFB1 at 0.94  $a_w$  and 37 °C and 25 °C after 7 and 21 d, respectively. *A. flavus* species comprise a high number of isolates with certain genetic variability, which explains why the expression of genes involved in the aflatoxin metabolic pathway is dependent on the environmental conditions.



**Figure 3** Comparison aflatoxin B1 (AFB1) production by 3 *Aspergillus flavus* isolates after 7 days of incubation on sorghum seeds at different water activities (0.94, 0.97 and 0.99) (A, B, C, D). If all letters next to a bar are different from the letters in another bar, there are significant differences among the treatments, if they have a common letter, they are not significantly different ( $p < 0.05$ ).

In our study, maximum AFB1 production could not be linked to a specific time point after inoculation, but overall 7 d of incubation was generally sufficient for *A. flavus* to achieve important AFB1 accumulation under the optimal conditions for AFB1 production (37 °C and 0.97–0.99  $a_w$ ). Nevertheless, the optimum incubation period and temperature required for maximum AFB1 production varied, probably due to the set of conditions tested, the nature and composition of the substrate and intraspecific and regional variability among isolates.

In this study, maximal growth was correlated with maximal aflatoxin production (0.99  $a_w$  and 37 °C). However, other studies have shown that good growth may occur without any production of aflatoxins. For example, Rabie and Smalley<sup>15</sup> demonstrated that the maximal production of aflatoxins occurred at 24 °C, but the maximal growth of *A. flavus* isolates occurred at 29 and 35 °C.

This is the first study concerning the effect of water activity and temperature on *A. flavus* growth and AFB1 production on sorghum grains.

The optimal conditions for mycelial growth and AFB1 production were 0.99  $a_w$  and 37 °C. There was neither *A. flavus* growth nor AFB1 production at 0.85 and 0.88  $a_w$  at all temperatures investigated. Consequently, aflatoxin accumulation could be avoided by storing sorghum at low water activity levels ( $\leq 0.91 a_w$ ). Our results showed that *A. flavus* isolated from Tunisian sorghum was able to grow in a wide range of water activities (0.91–0.99  $a_w$ ) and temperatures (15–37 °C), however, the production of AFB1 occurred at a narrower range of water activities (0.94–0.99  $a_w$ ) and temperatures (25–37 °C). Within the range of  $a_w$  evaluated in this study, 0.94 was the limiting level for AFB1 production. The investigation of trends for growth and AFB1 production at different temperatures and water activities in sorghum seeds can provide important information concerning the contamination of sorghum by AFB1. Our study contributes to the understanding of the ecology and physiology of Tunisian *A. flavus* isolates.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this investigation.

**Confidentiality of data.** The authors declare that no patient data appears in this article.

**Right to privacy and informed consent.** The authors declare that no patient data appears in this article.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

We thank M. Prim for her encouragement and technical assistance. The authors are grateful to the European Union (MYCORED KBBE-2007-2-5-05 project) and Tunisian Government for financial support.

## References

- Alves dos Reis T, Zorzete P, Pozzi CR, da Silva VN, Ortega E, Correa B. Mycoflora and fumonisin contamination in Brazilian sorghum from sowing to harvest. *J Sci Food Agric*. 2010;90:1445–51.
- Arrus K, Blank G, Abramson D, Clear R, Holley RA. Aflatoxin production by *Aspergillus flavus* in Brazil nuts. *J Stored Prod Res*. 2005;41:513–27.
- Astoreca A, Vaamonde G, Dalcerro A, Marin S, Ramos AJ. Abiotic factors and their interactions influence on the co-production of aflatoxin B1 and cyclopiazonic acid by *Aspergillus flavus* isolated from corn. *Food Microbiol*. 2014;38:276–83.
- Asurmendi P, Barberis C, Pacual L, Dalcerro A, Barberis L. Influence of *Listeria monocytogenes* and environmental abiotic factors on growth parameters and aflatoxin B1 production by *Aspergillus flavus*. *J Stored Prod Res*. 2015;60:60–6.
- Bandopadhyay R, Butler DR, Chandrasekhar A, Reddy RK, Navi SS. Biology epidemiology and management of sorghum grain mold. In: Chandrasekhar A, Bandyopadhyay R, Hall AJ, editors. Technical and institutional options for sorghum grain mold management: proceedings of an international consultation. ICRISAT: Andhra Pradesh, India; 2000. p. 34–71.
- Baranyi J, Roberts TA. A dynamic approach to predicting bacterial growth in food. *Int J Food Microbiol*. 1994;23:277–94.
- Boutrif E, Jemmali M, Campbell AD, Pohland AE. Aflatoxin in Tunisian foods and foodstuffs. *Ann Nutr Aliment*. 1977;31:431–4.
- Dantigny P, Guilmar A, Bensoussan M. Basis of predictive mycology. *Int J Food Microbiol*. 2005;100:187–96.
- Da Silva JB, Dilkin P, Fonseca H, Correa B. Production of aflatoxins by *Aspergillus flavus* and of fumonisins by *Fusarium* species isolated from Brazilian sorghum. *Braz J Microbiol*. 2004;35:182–6.
- Diener UL, Davis ND. Aflatoxin production by isolates of *Aspergillus flavus*. *Phytopathology*. 1966;56:1390–3.
- Garcia D, Ramos AJ, Sanchis V, Marin S. Predicting mycotoxins in foods: a review. *Food Microbiol*. 2009;26:757–69.
- Garcia D, Ramos AJ, Sanchis V, Marin S. Intraspecific variability of growth and patulin production of *Penicillium*

- expansum* isolates at two temperatures. Int J Food Microbiol. 2011;151:195–200.
13. Garcia D, Ramos AJ, Sanchis V, Marin S. Modeling kinetics of aflatoxin production by *Aspergillus flavus* in maize-based medium and maize grain. Int J Food Microbiol. 2013;162:182–9.
  14. Ghali R, Hmaissia-khlifa K, Ghorbel H, Maaroufi K, Hedili A. Incidence of aflatoxins, ochratoxin A and zearalenone in Tunisian foods. Food Control. 2008;19:921–4.
  15. Ghali R, Hmaissia-khlifa K, Ghorbel H, Maaroufi K, Hedili A. Aflatoxin determination in commonly consumed foods in Tunisia. J Sci Food Agric. 2010;90:2347–51.
  16. Giorni P, Magan N, Pietri A, Battilani P. Growth and aflatoxin production of an Italian strain of *Aspergillus flavus*: influence of ecological factors and nutritional substrates. World Mycotoxin J. 2011;4:425–32.
  17. Godet M, Munaut F. Molecular strategy for identification in *Aspergillus* section *Flavi*. FEMS Microbiol Lett. 2010;304:157–68.
  18. Horn BW. Colonization of wounded peanut seeds by soil fungi: selectivity for species from *Aspergillus* section *Flavi*. Mycologia. 2005;97:202–17.
  19. Hunter JH [PhD Thesis] Growth and aflatoxin production in shelled corn by the *Aspergillus flavus* group as related to relative humidity and temperature. Purdue Univ D; 1969 [Abstr 30 1447 B].
  20. IARC. Monographs on the evaluation of the carcinogenic risk of chemicals to humans. Some naturally occurring substances. Food items and constituents, heterocyclic aromatic amines and mycotoxins, vol. 56. Lyon, France: IARC; 1993. p. 397–444.
  21. Isakeit T, Prom LK, Wheeler M, Puckhaber LS, Liu J. Mycotoxigenic potential of ten *Fusarium* species grown on sorghum and *in vitro*. Plant Pathol J. 2008;7:183–6.
  22. Marin S, Colom C, Sanchis V, Ramos AJ. Modelling of growth of aflatoxigenic *Aspergillus flavus* from red chilli powder as a function of water availability. Int J Food Microbiol. 2009;128:491–7.
  23. Marin S, Ramos AJ, Cano-Sancho G, Sanchis V. Mycotoxins occurrence, toxicology, and exposure assessment. Food Chem Toxicol. 2013;60:218–37.
  24. Marin S, Ramos AJ, Sanchis V. Modelling *Aspergillus flavus* growth and aflatoxin production in pistachio nuts. Food Microbiol. 2012;32:378–88.
  25. Marin S, Sanchis V, Saenz R, Ramos AJ, Vinas I, Magan N. Ecological determinants for germination and growth of some *Aspergillus* and *Penicillium* spp. from corn grain. J Appl Microbiol. 1998;84:25–36.
  26. Mohale S, Magan N, Medina A. Comparison on growth, nutritional utilization patterns, and niche overlap indices of toxigenic and atoxigenic *Aspergillus flavus* strains. Fungal Biol. 2013;117:650–9.
  27. Montani ML, Vaamonde G, Resnik SL, Buera P. Water activity influence on aflatoxin accumulation in corn. Int J Food Microbiol. 1988;6:349–53.
  28. Mousa W, Ghazali FM, Jinap S, Ghazali HM, Radu S. Modelling the effect of water activity and temperature on growth rate and aflatoxin production by two isolates of *Aspergillus flavus* on paddy. J Appl Microbiol. 2011;111:1262–74.
  29. Oueslati S, Blesa J, Molto JC, Ghorbel A, Mañes J. Presence of mycotoxin in sorghum and intake estimation in Tunisia. Food Addit Contam A. 2014;31:307–18.
  30. Oueslati S, Romero-Gonzalez R, Lasram S, Garrido-frenich A, Martinez-Vidal JL. Multi-mycotoxin determination in cereals and derived products marketed in Tunisia using ultra-high performance liquid chromatography coupled to triple quadrupole mass spectrometry. Food Chem Toxicol. 2012;50:2376–81.
  31. Pardo E, Marin S, Sanchis V, Ramos AJ. Prediction of fungal growth and ochratoxin A production by *Aspergillus ochraceus* on irradiated barley grain as influenced by temperature and water activity. Int J Food Microbiol. 2004;95:79–88.
  32. Pitt JI, Miscamble BF. Water relations of *Aspergillus flavus* and closely related species. J Food Prot. 1995;58:86–90.
  33. Proietti I, Frazzoli C, Mantovani A. Exploiting nutritional value of staple foods in the world's semi-arid areas. Risks, benefits, challenges and opportunities of sorghum. Healthcare. 2015;3:172–93.
  34. Rabie CJ, Smalley EB. Influence of temperature on the production of aflatoxin. In: Symposium on mycotoxins in foodstuffs. 1965. p. 18–29.
  35. Rodriguez-Herrera R, Waniska RD, Rooney WL. Antifungal proteins and grain mold resistance in sorghum with non-pigmented testa. J Agric Food Chem. 1999;47:4802–6.
  36. Samapundo S, Devlieghere F, Geeraerd AH, De Meulenaer B, Van Impe JF, Devere J. Modelling of the individual and combined effects of water activity and temperature on the radial growth of *Aspergillus flavus* and *A. parasiticus* on corn. Food Microbiol. 2007;24:517–29.
  37. Samson RA, Hoekstra ES, Frisvad JC, Filteborg O. Introduction to food-and airborne fungi. 6th ed. Utrecht: Centraal bureau voor schimmelcultures; 2000.
  38. Sautour M, Dantigny P, Divies C, Bensoussan M. A temperature-type model for describing the relationship between fungal growth and water activity. Int J Food Microbiol. 2001;67:63–9.
  39. Schindler AP, Palmer JG, Eisenberg WV. Aflatoxin production by *Aspergillus flavus* as related to various temperatures. Appl Microbiol. 1967;15:1006–9.
  40. Serrano AB, Font G, Ruiz MJ, Ferrer E. Co-occurrence and risk assessment of mycotoxin in food and diet from Mediterranean area. Food Chem. 2012;135:423–9.
  41. Waliyar F, Ravinder RC, Alur AS, Reddy BVS, Reddy AR, Gowda CLL. Management of grain mold and mycotoxins in sorghum. Patancheru, Andhra Pradesh, India: International Crops Research Institute for Semi-Arid Tropics; 2007, 32 pp.