



ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect.com)International Journal for Parasitology:
Parasites and Wildlifejournal homepage: www.elsevier.com/locate/ijppaw

Invited Review

A review of *Theileria* diagnostics and epidemiologyBen J. Mans^{a,b,c,*}, Ronel Pienaar^a, Abdalla A. Latif^{a,b}^a Parasites, Vectors and Vector-Borne Diseases, Agricultural Research Council-Onderstepoort Veterinary Institute, Onderstepoort 0110, South Africa^b The Department of Veterinary Tropical Diseases, University of Pretoria, Pretoria, South Africa^c Department of Life and Consumer Sciences, University of South Africa, South Africa

ARTICLE INFO

Article history:

Received 19 November 2014

Revised 18 December 2014

Accepted 19 December 2014

Keywords:

Diagnostics

Serology

Parasitaemia

PCR

Species

Theileria

ABSTRACT

An extensive range of serological and molecular diagnostic assays exist for most of the economically important *Theileria* species such as *T. annulata*, *T. equi*, *T. lestoquardi*, *T. parva*, *T. uilenbergi* and other more benign species. Diagnostics of *Theileria* is considered with regard to sensitivity and specificity of current molecular and serological assays and their use in epidemiology. In the case of serological assays, cross-reactivity of genetically closely related species reduces the use of the gold standard indirect fluorescent antibody test (IFAT). Development of antigen-specific assays does not necessarily address this problem, since closely related species will potentially have similar antigens. Even so, serological assays remain an important line of enquiry in epidemiological surveys. Molecular based assays have exploded in the last decade with significant improvements in sensitivity and specificity. In this review, the current interpretation of what constitute a species in *Theileria* and its impact on accurate molecular diagnostics is considered. Most molecular assays based on conventional or real-time PCR technology have proven to be on standard with regard to analytical sensitivity. However, consideration of the limits of detection in regard to total blood volume of an animal indicates that most assays may only detect >400,000 parasites/L blood. Even so, natural parasitaemia distribution in carrier-state animals seems to be above this limit of detection, suggesting that most molecular assays should be able to detect the majority of infected individuals under endemic conditions. The potential for false-negative results can, however, only be assessed within the biological context of the parasite within its vertebrate host, *i.e.* parasitaemia range in the carrier-state that will support infection of the vector and subsequent transmission.

© 2014 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	105
2. Diagnostics	106
2.1. Microscopy	106
2.2. Xenodiagnosis	106
2.3. Serological assays	106
2.4. Molecular assays	109
3. Specificity of molecular assays and the species concept	110
4. Sensitivity of molecular assays, carrier state and parasitaemia ranges	112
5. Epidemiology in the context of species	113
6. Conclusion	114
Acknowledgements	114
Conflict of interest	114
References	114

* Corresponding author. Parasites, Vectors and Vector-Borne Diseases, Agricultural Research Council-Onderstepoort Veterinary Institute, Onderstepoort 0110, South Africa. Tel.: +27 125299200; fax.: +27 12 838 2582.

E-mail address: mansb@arc.agric.za (B.J. Mans).

<http://dx.doi.org/10.1016/j.ijppaw.2014.12.006>

2213-2244/© 2014 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. A generalised lifecycle for the *Theileria* using *T. parva* as example.

1. Introduction

The phylum Apicomplexa comprises a large group of complex eukaryotic organisms known to be obligate parasites of vertebrates and invertebrates. These organisms share a common characteristic of having an apical complex which contains secretory organelles considered to be involved in invasion and/or establishment of the parasite in the mammalian or invertebrate host (Bishop et al., 2004). The phylum is divided into four principal groups; the Coccidia, Gregarinasina (gregarines), Haemospororida (haemosporidians) and the Piroplasmorida (piroplasmids) (Adl et al., 2012). The Piroplasmorida comprises two main genera (*Babesia* and *Theileria*) responsible for the economic important diseases of domestic and wild animals. New species of the piroplasmids are still being discovered and their full biology is not completely documented yet. Many of this order's parasites were formerly classified based on morphology, host cells in which schizogony occurs, the observation of piroplasms in the red blood cells associated with disease manifestation and host-vector specificity (Barnett, 1977; Uilenberg, 2006). The genus *Theileria* is distinguished by infection of leukocytes by sporozoites, maturation of schizonts into merozoites and subsequent infection of red blood cells to form piroplasms (Uilenberg, 2006).

A generalised lifecycle for the *Theileria* genus include secretion of infective sporozoites during tick feeding into the feeding site (Fig. 1). Sporozoites then infect leukocytes and multiply by merogony, after which merozoites are released, which invade red blood cells thereby establishing the piroplasm stage. During a next feeding cycle, larval or nymphal vector ticks ingest piroplasms and the released parasites undergo syngamy in the tick gut, forming a zygote, the only diploid stage. The zygote divides into motile kinetes that infect the tick gut epithelial cells and migrate to the haemolymph and subsequently infect the salivary glands. After moulting and commencement of feeding by the tick, sporogony results in the multiplication of sporozoites in the salivary gland acini before injection into the feeding site by nymphs or adult ticks (McKeever, 2009).

The *Theileria* species infect a wide range of both domestic and wild animals and are transmitted by ixodid ticks of the genera *Amblyomma*, *Haemaphysalis*, *Hyalomma* and *Rhipicephalus*. Most of these ticks are renowned for the large economic losses they cause to the agricultural industry due to disease outbreaks, mortalities, damage to hides and poor production in domestic animals (Bishop et al., 2004). The expansion of wildlife husbandry and conservation has also made *Theileria* of wildlife important subjects of study. The *Theileria* can be grouped into schizont “transforming” and

“non-transforming” species (Sivakumar et al., 2014). Transforming parasites all group in the *T. taurotragi* clade (Fig. 2) (Sivakumar et al., 2014), and uncontrolled proliferation of schizonts results in the pathologies associated with Corridor disease (*Theileria parva*), East Coast fever (*T. parva*), Tropical theileriosis (*T. annulata*) in cattle and malignant theileriosis (*T. lestoquardi*) in goats and sheep (Bishop et al., 2004; McKeever, 2009). However, *T. taurotragi*, *T. sp.* (buffalo) and *T. sp.* (bougasvlei) do not cause schizont associated pathology (Young et al., 1977; Bishop et al., 2004; Pienaar et al., 2014). *Theileria sp.* (sable), that group within the antelope *Theileria* (Fig. 2), causes lymphoid hyperplasia typically associated with the transforming *Theileria* (Nijhof et al., 2005). This and the ability to culture schizonts (Zweygarth et al., 2009a) indicate that not all transforming parasites share a monophyletic origin (Sivakumar et al., 2014), which suggests that transformation of schizonts may occur more widely than expected. The non-transforming *Theileria* are regarded as being benign but still able to cause disease as a result of anaemia induced by the piroplasm stage (Sivakumar et al., 2014). The last decade has seen an increase in the discovery of new *Theileria* species and genotypes and the realisation that even domestic animals may harbour an extensive array of mixed infections (Criado-Fornelio et al., 2004; Mans et al., 2011a; Sivakumar et al., 2014). This is compounded by mixed *Theileria* infections in both mammalian hosts and tick vectors where wildlife and livestock share the same habitat and common tick species (Lawrence et al., 1983; Kariuki et al., 2012). The means to discriminate benign and virulent forms as well as species responsible for disease outbreaks is becoming more important, both for diagnostic and epidemiological purposes.

2. Diagnostics

Diagnostic methods are used alone or in combination with other criteria to accurately diagnose disease. The use of assays as diagnostic tools needs to fulfil some pre-requisites albeit for human or veterinary use (Pfaffl, 2004; Bustin, 2010; Peeling et al., 2010). As such the basic performance characteristics of a test need to be determined against a reference standard. This would generate data on reproducibility, sensitivity and specificity of the particular assay, parasite prevalence, genetic variation and/or methodology used: type of antigen or antibody, automated or manual procedures, all contributing to the variables influencing any given assay's performance.

2.1. Microscopy

Historically most *Theileria* species were first described based on light microscopy examination after Koch's discovery of *T. parva* (Koch, 1898). It was usually performed on Giemsa stained blood smears, however, the limitation of light microscopy as a diagnostic tool was that the detection of carrier animals regularly went unnoticed and the discrimination between piroplasms of other *Theileria* species were difficult as they are morphologically very similar (Dschunkowsky and Luhs, 1904; Theiler, 1904; Lawrence, 1935, 1979; Uilenberg, 1981). Even so, discovery of novel species may still utilise morphology as part of their formal description (Nijhof et al., 2005; Clark and Spencer, 2007; Oosthuizen et al., 2009; Papparini et al., 2012), and remains a fast method to derive at an initial differential diagnosis of potential theileriosis in clinical cases (Nijhof et al., 2005; Izzo et al., 2010). In regard to routine diagnostics, microscopy is several orders of magnitude less sensitive than molecular methods (Criado-Fornelio, 2007), and generally unable to accurately detect all carrier-state animals (Lawrence, 1979; Zweygarth et al., 1997).

2.2. Xenodiagnosis

Xenodiagnosis (tick pickup and transmission linked with clinical disease) as demanded by Koch's postulate was used to confirm disease-causing species and tick vectors (Dschunkowsky and Luhs, 1904; Theiler, 1904; Young et al., 1977; Skilton et al., 2002). Direct injection of infected blood may also distinguish species amenable to proliferation in the piroplasm stage (Theiler, 1906). Xenodiagnosis remains a useful tool in basic parasite research and epidemiological investigations and may be the only means to confirm tick vector designation and clinical pathology of specific *Theileria* strains or as first step in parasite isolation (Ngumi et al., 1994; Steyl et al., 2012; Mbizeni et al., 2013). It is, however, not suitable for high-throughput or routine analysis and remains in the domain of specialised research groups.

2.3. Serological assays

Serological diagnosis by complement fixation (Lichtenheld, 1910) was followed by the more advanced indirect fluorescent antibody test (IFAT) (Schindler and Wokatsch, 1965; Burrige, 1971; Burrige and Kimber, 1972). Antigen may be prepared from schizont or piroplasm antigen, derived from infected animals or cell culture. Cell culture of *Theileria* parasites may be useful in diagnosis of carrier state animals (Zweygarth et al., 1997), to identify transforming parasites (Zweygarth et al., 2009a, 2009b), as means to generate IFAT antigen (Burrige and Kimber, 1972), and distinguish different species and investigate distinct parasite populations using monoclonal antibodies (Conrad et al., 1987; Bishop et al., 1994a; Alhassan et al., 2007a; Zweygarth et al., 2009b). The IFAT remains the gold standard assay recommended by the OIE for most economically important parasites (OIE, 2014). The IFAT has several drawbacks, namely, subjective operator-dependent interpretation of results, low throughput and difficulty in standardisation (Katende et al., 1998). However, the biggest problem with the IFAT is the significant cross-reactivity observed between closely related species. Cross-reactivity between *T. parva* and *T. taurotragi* antigen and anti-sera has been observed (de Vos and Roos, 1981; Jongejan et al., 1986). Similar observation has been made in regard to *T. parva* and *T. sp.* (buffalo) (Conrad et al., 1987; Pienaar et al., 2014). An IFAT assay for *T. lestoquardi* showed significant cross-reactivity with *T. annulata* and *T. parva* anti-sera and vice versa (Leemans et al., 1997). Detection by IFAT may therefore be a poor measure to assign parasites to species (Uilenberg, 1981; Uilenberg et al., 1985; Stewart et al., 1996), especially when closely related, such as the genotypes found in the *T. buffeli*, *T. mutans* and *T. velifera* clades (Fig. 2). Even so, IFAT may still be useful in epidemiological studies (Thompson et al., 2008; Mbizeni et al., 2013), or where certain species are absent in specific carrier hosts such as the case for *T. taurotragi* and African buffalo (*T. parva* carrier) (Mans et al., 2011a), or *T. annulata* in small ruminants (*T. lestoquardi* carrier) (Leemans et al., 1997).

The IFAT utilises whole-body antigen, but the realisation that a select number of antigens are responsible for the dominant immune response against most *Theileria* parasites has stimulated interest in enzyme-linked immunosorbent assays (ELISA) (Katende et al., 1998). These single molecule assays have the promise of more species specific antigens amenable to high throughput analysis. The polymorphic immunodominant molecule (PIM) of *Theileria parva* and the p32 antigen of *T. mutans* were used to develop ELISAs (Katende et al., 1990, 1998; Morzaria et al., 1999). Both were commercially marketed and shown to be more sensitive than the IFAT, but the tests were discontinued due to specificity issues (OIE, 2014). However, both are still being used in research (Swai et al., 2007; Bazarusanga et al., 2008; Kiara et al., 2014). The EMA-1 antigen for *T. equi* and a monoclonal antibody against this antigen were used to develop a competitive inhibition ELISA (Knowles et al., 1992) that is




Fig. 2. Phylogenetic analysis of *Theileria* genotypes. The 18S ribosomal RNA sequences were extracted from GenBank using text based queries and BLAST analysis. Final non-redundant dataset were obtained by phylogenetic analysis and manual curating. The dataset were trimmed to include the V4 hyper-variable region, aligned using MAFFT (auto, 200PAM/k = 2; [Kato and Standley, 2013](#)) and analysed using neighbour-joining (bootstrap = 1000, matrix = number of differences) with Mega5 ([Tamura et al., 2011](#)). Host and geographic data for genotypes are indicated and were extracted from GenBank files or literature in the case of reverse line blot analysis.




Fig. 2. (continued)

Table 1

A summary of diagnostic assays that exist for the *Theileria*. Indicated are *Theileria* species, tick vectors, hosts, diseases caused, serological and diagnostic assays developed.

<i>Theileria</i> spp.	Tick vector	Hosts	Disease	Serology	Molecular
<i>T. parva</i>	<i>Rhipicephalus appendiculatus</i> ; <i>R. zambeziensis</i>	Cattle, African buffalo	Fatal; East Coast fever, Corridor disease, Zimbabwean theileriosis (January disease)	1, 2, 3	4, 5, 6, 7, 8
<i>T. mutans</i>	<i>Amblyomma</i> sp.	African buffalo, cattle	Benign theileriosis	1, 9, 10	4, 8, 11
<i>T. taurotragi</i>	<i>Rhipicephalus</i> sp.	Cattle, eland (sheep)	Benign African theileriosis	9, 12	4, 8
<i>T. velifera</i>	<i>Amblyomma</i> sp.	African buffalo, cattle	Benign	None	4, 8
<i>T. buffeli</i>	<i>Haemaphysalis</i> sp.	African buffalo, cattle, Water buffalo	Oriental theileriosis	13, 14, 15, 16	4, 8, 17, 18, 19, 20
<i>T. sergenti</i>					
<i>T. orientalis</i>					
<i>T. sinesis</i>					
<i>T. lestoquardi</i>	<i>Hyalomma</i> sp.	Sheep, goats	Malignant sheep theileriosis	21, 22	23, 24, 25
<i>T. ovis</i>	<i>Hyalomma</i> sp. <i>Rhipicephalus</i> ?	Sheep, goats	Benign	None	23, 26, 27, 28, 29
<i>T. separata</i>	<i>Rhipicephalus</i> sp., <i>Hyalomma</i> sp.	Sheep, antelope	Benign/pathogenic	None	23, 24
<i>T. annulata</i>	<i>Hyalomma</i> sp.	Cattle, water/domestic buffalo	Fata; Tropical or Mediterranean theileriosis	30, 31, 32, 33	4, 17, 24, 26, 34, 35, 36, 37
<i>T. sp.</i> (buffalo); <i>T. sp.</i> (bougasvlei)	Not known	African buffalo (South Africa); African buffalo, cattle (East Africa)	Benign	38	8, 39
<i>T. sp.</i> (sable)	<i>R. evertsi</i> / <i>R. appendiculatus</i> ?	Sable	Malignant	None	40
<i>T. equi</i>	<i>Hyalomma</i> spp., <i>Rhipicephalus</i> spp.	Horses, donkeys, giraffes	Acute to chronic	41, 42, 43, 44	45, 46, 47, 48
<i>T. annae</i>	<i>Ixodes</i> sp., <i>Rhipicephalus</i> sp., <i>Dermacentor</i> sp.	Dogs, foxes	Severe, regenerative anaemia	None	49, 50
<i>T. uilenbergi</i>	<i>Haemaphysalis</i> sp.	Sheep, goats, sika, red deer	Cervine theileriosis	51	52, 53
<i>T. luwenshuni</i>	<i>Haemaphysalis</i> sp.	Sheep, goats, sika, red deer	Cervine theileriosis	51	26, 52, 54
<i>T. bicornis</i>	<i>R. evertsi evertsi</i>	Black, white and Indian rhinoceros	Benign	None	55
<i>T. capreoli</i>	<i>Ixodes ricinus</i>	Sika, red deer	Theileriosis?	None	55, 56
<i>T. cervi</i>	<i>Amblyomma americanum</i>	White-tailed deer, elk	Benign	57	58
<i>T. sp.</i> OT 3	<i>Haemaphysalis</i> sp.?	?	Theileriosis?	None	26, 54, 59
<i>T. sp.</i> MK	Not known	Goats, sheep	Benign	None	60

1. Complement-fixation assay (Schindler and Mehlitz, 1969). 2. Piroplasm and schizont IFAT and ELISA (Burrige, 1971; Burrige and Kimber, 1972; Gray et al., 1980). 3. PIM indirect ELISA (Katende et al., 1998). 4. RLB (Gubbels et al., 1999). 5. Conventional and nested PCR for p67, p104, Tpr (Bishop et al., 1992; Skilton et al., 2002; Odongo et al., 2010; Pienaar et al., 2011a). 6. Real-time PCR targeting 18S (Sibeko et al., 2008; Papli et al., 2011; Pienaar et al., 2013). 7. LAMP targeting PIM and p150 (Thekiso et al., 2010). 8. Pan-FRET real-time PCR for Cox III (Chaisi et al., 2013a). 9. Piroplasm IFAT (de Vos and Roos, 1981). 10. Indirect ELISA for p32 (Katende et al., 1990). 11. Conventional PCR based on non-coding region (Bishop et al., 1994b). 12. Piroplasm IFAT (Jongejan et al., 1986). 13. Piroplasm IFAT (Uilenberg et al., 1985; Papadopoulos et al., 1996). 14. Piroplasm extract ELISA (Shimizu et al., 1988). 15. Latex agglutination test for major piroplasm surface protein p33 (Jeong et al., 2005). 16. Indirect ELISA for p33 (Wang et al., 2010b). 17. Conventional PCR for β -tubulin, MPSP, p23, p32, p33, p34 (Tanaka et al., 1993; Kawazu et al., 1995; Kubota et al., 1996; Govaerts et al., 1998; Sarathaphan et al., 1999; Cacciò et al., 2000; Liu et al., 2010b, 2011; Ota et al., 2010). 18. Semi-nested PCR for 18S rRNA (Ghaemi et al., 2012). 19. Multiplex-tandem real-time PCR for the major piroplasm surface protein, 23-kDa piroplasm membrane protein and ITS1 (Perera et al., 2014). 20. LAMP for p33 and ITS (Wang et al., 2010a; Liu et al., 2013). 21. Schizont IFAT (Leemans et al., 1997). 22. Clone 5 ELISA (Bakheit et al., 2006). 23. RLB (Schnittger et al., 2004). 24. PCR-RFLP of 18S rRNA (Heidarpour Bami et al., 2009). 25. LAMP of clone 5 (Salih et al., 2012). 26. Suspension arrays (Ros-García et al., 2012a). 27. Nested PCR of 18S rRNA (Altay et al., 2005). 28. PCR-RFLP of 18S rRNA (Heidarpour Bami et al., 2009). 29. 5.8S RNA PCR (Zhang et al., 2014). 30. Piroplasm and schizont IFAT and ELISA (Gray et al., 1980; Kachani et al., 1992; Darghouth et al., 1996). 31. TaSP, Tams1 indirect ELISAs (Bakheit et al., 2004; Salih et al., 2005; Rajendran and Ray, 2014). 32. Competitive ELISA for TaSP (Renneker et al., 2008). 33. TaSP lateral flow device (Abdo et al., 2010). 34. Cytochrome b RLB (Bilgic et al., 2010). 35. Conventional PCR for cytochrome b, HSP70 and *Tams1* (d'Oliveira et al., 1995; Shayan et al., 1998; Kirvar et al., 2000; Criado et al., 2006). 36. Conventional and real-time PCR for 18S rRNA gene (Ilhan et al., 1998; Ros-García et al., 2012b). 37. LAMP for the 18S rRNA, ITS and TA04795 (Salih et al., 2008; Liu et al., 2012). 38. Schizont IFAT (Conrad et al., 1987). 39. Real-time LNA probe assay (Pienaar et al., 2014). 40. Real-time PCR (Pienaar, Personal communication). 41. Piroplasm IFAT and complement fixation test (Ogunremi et al., 2007). 42. Competitive ELISA of EMA-1 (Knowles et al., 1992). 43. Indirect ELISA for EMA-2 (Huang et al., 2003). 44. Immuno-chromatographic test for EMA-2 (Huang et al., 2004). 45. RLB (Butler et al., 2008). 46. Conventional PCR of EMA-1, β -tubulin, 18S rRNA (Cacciò et al., 2000; Alhassan et al., 2005; Heim et al., 2007; Salim et al., 2008). 47. Real-time PCR and high resolution melt analysis of 18S rRNA (Kim et al., 2008; Bhoora et al., 2010a; Salim et al., 2013). 48. LAMP of EMA-1 and 18S rRNA (Alhassan et al., 2007a, 2007b; Xie et al., 2013). 49. RLB (Yisaschar-Mekuzas et al., 2013). 50. Conventional PCR-RFLP of 18S rRNA (Jefferies et al., 2007). 51. TIHSP70, rTuIP ELISA (Miranda et al., 2006; Liu et al., 2014). 52. RLB (Niu et al., 2009). 53. Conventional PCR of 18S rRNA, RPS8 (Yin et al., 2008; Tian et al., 2013; Zhang et al., 2014). 54. RLB (Nagore et al., 2004). 55. RLB (Nijhof et al., 2003). 56. RLB (García-Sanmartín et al., 2007). 57. Conventional PCR of 18S rRNA (Li et al., 2014). 58. IFAT (Schaeffler, 1963). 59. PCR and sequencing of 18S rRNA (Chae et al., 1999a). 60. Conventional PCR of 18S rRNA (Tian et al., 2014). 61. RLB and conventional PCR of 18S rRNA (Altay et al., 2007, 2008).

available as a commercial kit and considered as replacement for the IFAT (OIE, 2014). The EMA-2 antigen for *T. equi* was also investigated in an indirect ELISA and as rapid immune-chromatographic test (Huang et al., 2003, 2004). Other ELISAs developed include the *T. annulata* merozoite surface 1 antigen (Tams1) and the *T. annulata* macroschizont stage protein (TaSP) (Gubbels et al., 2000a; Seitzer et al., 2007; Renneker et al., 2008), a heat-shock 70 antigen for *T. sp.* (China) (Miranda et al., 2006) and *T. uilenbergi* immunodominant protein (TuIP) (Liu et al., 2010a). TaSP was also used to develop a rapid lateral flow device assay for *T. annulata* that compares well with other serodiagnostic assays (Abdo et al., 2010). One of the biggest advantages of ELISA is in its use as a high throughput, cheap and fast method to screen and diagnose large numbers of samples. In this regard, serodiagnosis remains the workhorse for laboratories where molecular infrastructure does not exist.

2.4. Molecular assays

The molecular revolution saw an explosion of diagnostic assays (Criado-Fornelio, 2007), that is target specific genes and species (Table 1). This included conventional PCR followed by agarose gel electrophoretic analysis (Bishop et al., 1992; Pienaar et al., 2011a), PCR-RFLP methods (Bishop et al., 1992; Geysen et al., 1999; Heidarpour Bami et al., 2009; Zaemi et al., 2011), nested-PCR (Odongo et al., 2010), PCR followed by dot blotting, capillary blotting or slot-blotting and hybridisation using radio-isotope labelled probes (Bishop et al., 1992; Allsopp et al., 1993; Collins et al., 2002; Skilton et al., 2002). The latter was improved by the non-radioactive reverse line blot method that used chemiluminescence (Gubbels et al., 1999; Schnittger et al., 2004), with recent variation on this technique using a DNA bead-based suspension array

(Ros-García et al., 2012a, 2013). This was followed by probe based real-time PCR methods (Jeong et al., 2003; Kim et al., 2008; Sibeko et al., 2008; Bhoora et al., 2010a; Pienaar et al., 2011b, 2014; Ros-García et al., 2012b), SYBR green real-time PCR assays (Pienaar et al., 2013), loop-mediated isothermal amplification (LAMP) assays (Alhassan et al., 2007a, 2007b; Liu et al., 2008, 2012, 2013; Salihi et al., 2008, 2012; Thekisoe et al., 2010; Wang et al., 2010a; Xie et al., 2013), pan-FRET based assays (Chaisi et al., 2013a; Perera et al., 2014; Yang et al., 2014a) and high-resolution melt analysis (Salim et al., 2013). In all cases, detection by molecular methods allow for direct confirmation of the presence of parasite genomic material, with the inference that live parasites are present in the animal at the moment of sampling. Developments from conventional to nested to real-time PCR has allowed improvement in sensitivity, quantification and speed of detection, while methods such as reverse line blot, bead arrays, pan-FRET assays and high-resolution melt analysis hold the promise of detection of multiple species or genotypes at the same time. Real-time melting profile based assays also hold the advantage that variation in probe regions may be detected by differences in melting profiles that may be related to genotypic or species differences. Whereas many of these assays require specialised equipment, LAMP assays hold the advantage of functioning at isothermal conditions, with possible application under field conditions.

3. Specificity of molecular assays and the species concept

Most molecular assays depend on primers and/or probes that target small regions of genes with the implicit assumption that these regions are conserved through all members of a species or genus. In the case of protein genes, the degenerate nature of the genetic code makes the design of specific primers or probes quite difficult, as exemplified by their reduced sensitivity compared to more conserved ribosomal genes (Pienaar et al., 2011a, 2013). Protein genes on the other hand may provide much greater specificity due to more distant orthologous relationships (Odongo et al., 2010; Pienaar et al., 2011a). The challenge in the development of specific assays therefore lies in the identification of these unique regions within genes or a genome, working with often limited information regarding the diversity of a gene or the availability of total genome information. For example, the reverse line blot (RLB) was developed well before extensive genomic information was available for the *Theileria* genus or estimates regarding 18S rRNA diversity in domestic animals and wildlife (Gubbels et al., 1999). The RLB primers for *Theileria* and *Babesia* target the 18S rRNA regions that flank the V4 hyper-variable region and to date have been found to be conserved in all members of these genera (Gubbels et al., 1999). The probes used in the reverse line blot were assumed to be species-specific and to be able to detect all members of a species, based on the assumption that the 18S hyper-variable region is conserved within species. It is therefore also useful to detect new species in the case where only a *Theileria* catch-all probe is detected (Oosthuizen et al., 2009; Chaisi et al., 2013b, 2014). More recently, some of these assumptions have been challenged.

Diagnostic screening of buffalo and cattle samples using the hybridisation real-time PCR for *T. parva* (Sibeko et al., 2008) detected samples with aberrant melting curves that suggested variation in the probe region of this assay that occurs within the 18S rRNA V4 hyper-variable region (Mans et al., 2011a). Sequencing of these samples indicated single nucleotide polymorphisms, suggesting that variation in the V4 hyper-variable region may occur in individuals within the *T. parva* population (Mans et al., 2011a). In geographically isolated populations these polymorphisms may become fixed, leading to cases where species identity may be questioned. It was as such proposed that organisms with 1–2 single nucleotide polymorphisms within the V4 18S rRNA hyper-variable region be considered the same species (Mans et al., 2011a). Conversely, the

interesting phenomenon exists, that certain genotypes within the *Theileria* genus show extensive genotypic variation within the 18S rRNA hyper-variable region, but due to the fact that these groups within the same clade during phylogenetic analysis are considered to be members of the same species. As such, 12 main clades for *Theileria* can be distinguished (Fig. 2), comprising at least 91 unique genotypes which may or may not be unique species as will be discussed later. The main clades found include the *T. buffeli*, *T. equi*, *T. mutans*, *T. taurotragi*, *T. velifera* clades, three clades for antelope *Theileria*, a clade for marsupial *Theileria* and for *Theileria annae*.

The *T. taurotragi* clade comprises multiple well-recognised species, even though their inter-species genetic distances are very similar to other clades (Mans et al., 2011a). *Theileria taurotragi* has been found in antelope (bushbuck, eland) and cattle and occurs from East to Southern Africa (Martin and Brocklesby, 1960; Young et al., 1977; de Vos and Roos, 1981; Oura et al., 2011); *T. annulata* infects cattle and causes tropical theileriosis, while *T. lestoquardi* infects small ruminants such as sheep and goats and causes malignant theileriosis (Leemans et al., 1999); *T. parva* infects cattle and African buffalo (Norval et al., 1992); *T. sp.* (buffalo) infects African buffalo and has been detected in cattle (Allsopp et al., 1993; Zwegarth et al., 2009b; Mans et al., 2011a; Pienaar et al., 2011a; Githaka et al., 2014); *T. sp.* (bougasvlei) infects African buffalo (Zwegarth et al., 2009b; Mans et al., 2011a; Pienaar et al., 2011a, 2014). *Theileria annulata* and *T. lestoquardi* differ by 2 base pairs in the hyper-variable region, while differences in this region between *T. parva*, *T. sp.* (buffalo) and *T. sp.* (bougasvlei) range from 4 to 6 base pairs (Mans et al., 2011a). Even so, analyses using the nuclear S5 and mitochondrial COI genes distinguish these species quite well, while *T. sp.* (buffalo) and *T. sp.* (bougasvlei) also show disparate geographic distributions (Mans et al., 2011a; Pienaar et al., 2014). All members of this clade can transform lymphocytes (assumed for *T. sp.* (bougasvlei)), but this phenomenon is not associated with pathology for all members (Mans et al., 2011b; Sivakumar et al., 2014).

Theileria annae is a parasite of dogs and other canines and is related to *Babesia microti* (Zahler et al., 2000; Camacho et al., 2001; Najm et al., 2014). It has been assigned to the genus *Theileria* based on the contention that the small canine Babesias are more closely related to *Theileria* than *Babesia* (Zahler et al., 2000). However, molecular analyses of the 18S rRNA gene indicate a close relationship to the *Babesia* (Matjila et al., 2008; Sivakumar et al., 2014). Other *Theilerias* found incidentally in dogs include *T. annulata* and *T. equi* (Criado-Fornelio et al., 2003; Criado et al., 2006; Beck et al., 2009). However, a genotype identical to *T. sp.* (sable) has been found extensively in dogs surveyed in KwaZulu-Natal, South Africa (Nijhof et al., 2005; Matjila et al., 2008). *Theileria sp.* (sable) groups within the clade of Antelope *Theilerias* have been found in an extensive range of other antelope as well (Nijhof et al., 2005). Since most of these identifications were done by RLB analysis these hosts remain to be confirmed by sequencing. The antelope clade is extensive and comprised of various genotypes found in different antelope species. In many cases, multiple genotypes have been found in single antelope species and are not distinguished by their genotype designation, even though their phylogenetic position would suggest that these may be unique species. This includes genotypes for *T. sp.* OT3 (Nagore et al., 2004; Tian et al., 2014), *T. capreoli* (García-Sanmartín et al., 2007; Yang et al., 2014b), *T. cervi* (Chae et al., 1998, 1999a), giraffe *Theileria* (Oosthuizen et al., 2009; Githaka et al., 2013), waterbuck *Theileria* (Githaka et al., 2014), white-tailed deer *Theileria*, water deer *Theileria* (Han et al., 2009). Many of these genotypes have been identified in multiple antelope species, making designation according to host specificity difficult. One genotype from this clade has been identified in cheetah and shows a close relationship to a genotype found in giraffes (Githaka et al., 2012).

The *Theileria* found in marsupials all cluster together and comprise 7 unique genotypes. *Theileria fuliginosa* is found in the western

grey kangaroo (*Macropus fuliginosus*), *T. penicillata* in the woylie (*Bettongia penicillata ogilbyi*), *T. gilberti* in Gilbert's potoroo (*Potorous gilbertii*), *T. sp.* long-nosed potoroo in the long-nosed potoroo (*Potorous tridactylus*) and *T. sp.* K1 in the boodie (*Bettongia lesueur*) (Clark and Spencer, 2007; Lee et al., 2009; Papparini et al., 2012). Two distinct genotypes for *T. brachyuri* have been found in the quokka (*Setonix brachyurus*) (Clark and Spencer, 2007), suggesting that at least one is a different species. At least three other marsupial *Theileria* exist for which no molecular data are available; they include *T. tachyglossi* in the echidna, *T. peramelis* in the southern brown bandicoot (*Isodon obesulus*), and *T. ornithorhynchi* in the platypus (*Ornithorhynchus anatinus*) (Lee et al., 2009; Papparini et al., 2012). The monophyletic relationship within this clade would suggest a single introduction into Australia, with subsequent host adaptations and speciation.

The *T. buffeli* clade possesses the most extensive number of genotypes (13) and occurs in all the major continents of the world, where they infect cattle, African buffalo, water buffalo and Yak (Chaisi et al., 2013a; Sivakumar et al., 2014). A number of different species names have historically been assigned to members of this clade including *T. buffeli*, *T. orientalis* and *T. sergenti*, but have been proposed to represent a single species designated as *T. buffeli* (Stewart et al., 1996; Gubbels et al., 2000b, 2002). This builds on a similar proposal by Uilenberg (1981) and Uilenberg et al. (1985) that used morphological and serological cross-reactivity to assign all members to *T. orientalis*. Both *T. buffeli* and *T. orientalis* continue to be used in literature depending on the historical background of different scientific groups (Chaisi et al., 2014; Sivakumar et al., 2014). Another "species" that belongs to this group is *T. sinensis* (Bai et al., 2002). The genotypes from this clade were postulated to be in the process of speciation that has not been completed yet (Schnittger et al., 2003), while others considered members to be distinct species (Kawazu et al., 1992; Fujisaki et al., 1994). Most members of this group are considered to be benign, but a recent outbreak of a virulent genotype (Ikeda type) in Australia readdressed the possibility of unique species in this group (Izzo et al., 2010; Kamau et al., 2011). More recently, related genotypes (*T. buffeli* C, *T. sinensis*-like) were found in African buffalo but not in cattle from the same geographic regions in southern Africa (Mans et al., 2011a; Chaisi et al., 2014). All these genotypes are generally considered to be genotypic variants of *T. buffeli* (Chae et al., 1999b; Gubbels et al., 2000b, 2002; Chaisi et al., 2014), even though host specificity, clinical presentation and specific geographic distributions are different. The use of other genes such as the major piroplasm surface protein (MPSP) also generally groups different genotypes into separate clades that correspond with those obtained using the 18S gene (Kamau et al., 2011; Sivakumar et al., 2014). Serological and molecular assays that can distinguish different genotypes have been developed (Kawazu et al., 1992, 1995; Liu et al., 2010b, 2013; Perera et al., 2014). It thus seem surprising that this clade does not have well recognised species, which complicates development of specific diagnostic assays and possible interpretation of diagnostic results. The relative caution in molecular species designation within this clade that occurred more than 14 years ago (Chae et al., 1999b; Gubbels et al., 2000b) has also permeated to other *Theileria* clades such as *T. mutans*, *T. velifera*, *T. equi*, the *Theileria* antelope clade and members of the *T. taurotragii* clade (discussed in their respective sections), irrespective of the fact that no recombination experiments have yet been attempted between different genotypes, that may warrant the conclusion of a single *Theileria* species per clade.

The *T. mutans* clade is comprised of five recognised genotypes, namely *T. mutans* that infects cattle and buffalo, *T. mutans* MSD that infects cattle and buffalo, *T. mutans*-like 1, *T. mutans*-like 2 and *T. mutans*-like 3 that has thus far been found only in African buffalo (Chae et al., 1999b; Mans et al., 2011a; Chaisi et al., 2013b). Of interest is that the *T. mutans*-like 1–3 genotypes were not detected

by the *T. mutans* RLB probe, but is still considered to be genotypic variants of *T. mutans* (Chaisi et al., 2013b), even though they differ by 7–13 nucleotide as well as a gapped position (Mans et al., 2011a). However, to date *T. mutans*-like 1–3 have only been observed in African buffalo and occur at significantly higher prevalence and parasitaemia compared to *T. mutans* and *T. mutans* MSD, which occur extensively in cattle (B.J. Mans, personal observation). Host specificity would therefore suggest that at least two species exist in this clade.

The *T. velifera* clade is comprised of four genotypes, *T. velifera*, *T. velifera* A, *T. velifera* B and *T. sp.* RR-2012, an as yet undescribed genotype from Indian bovines. *T. velifera* and *T. velifera* A have been found in cattle and buffalo, while *T. velifera* B has only been found in buffalo (Mans et al., 2011a; Chaisi et al., 2013b). Significant similarity exists between the RLB probe for *T. sp.* (sable) and *T. velifera*, so that reports of the extensive occurrence of the former in cattle and African buffalo (Nijhof et al., 2005; Yusufmia et al., 2010) could be traced back to the presence of *T. velifera* (Mans et al., 2011a). This again underscores the pitfalls of the RLB approach, when used for screening of novel vertebrate species or animals from new geographic regions, for which the assay has not been validated.

The *T. equi* clade shows similar genetic diversity as the *T. buffeli* clade (twelve genotypes) and is also considered to be representative of a single species (Kim et al., 2008; Bhoora et al., 2009; Salim et al., 2010). This is supported by extensive cross-reactivity on IFAT (Bhoora et al., 2009). Some genotypes have thus far been confined to equids other than horses, such as donkeys and zebra, but have also been found in waterbuck (Bhoora et al., 2010b; Githaka et al., 2014). Whether these will infect horses remains to be determined, but the wide distribution observed for most genotypes suggests that none are host specific (Fig. 2). Multiple genotypes also generally occur in similar geographic localities which make the assessment of strain virulence difficult, with the general assumption that all strains present similar virulence and pathologies. A recently developed real-time PCR assay selected a region within the 18S ribosomal RNA gene, for which both primers and probes are conserved in the *Theileria* (Kim et al., 2008; Bhoora et al., 2010a). It will therefore detect all genotypes from *T. equi* when used to screen equids but will not be useful for screening other animals suspected to harbour *T. equi* such as waterbuck (Githaka et al., 2014). From an equid epidemiological control perspective this may be ideal if the clade is treated as a single species or virulent group. However, from a phylogeographic perspective where evolution of unique genotypes is considered, large-scale surveys that rely on diagnostic assays will be hampered.

Two genotypes that do not fit into any major clade include *T. bicornis* found in black, Indian and white rhinoceros (Nijhof et al., 2003; Otiende et al., 2014). The other was found in domestic cats in a Brazilian zoo and may be an incidental infection transmitted from another felid carrier (André et al., 2014).

The origin of the genotypic diversity observed should be considered. To date genomes for *T. annulata*, *T. equi*, *T. parva* and *T. orientalis* have been sequenced and all indicate the presence of two 18S ribosomal RNA genes (Gardner et al., 2005; Pain et al., 2005; Hayashida et al., 2012; Kappmeyer et al., 2012). In all cases these genes are highly conserved, most probably due to concerted evolution (Eickbush and Eickbush, 2007), with the implication that genotypic diversity observed within the 18S gene for the *Theileria* is not due to multiple gene copies within their genomes (Mans et al., 2011a). Genotypic diversity observed in the 18S genes will therefore be due to conventional divergent evolution, linked with mutation and insertion–deletion events.

The genotypic diversity observed within the *Theileria* has interesting implications from a diagnostic perspective. On the one hand, some species such as *T. annulata*, *T. lestoquardi*, *T. parva*, *T. sp.* (buffalo) and *T. sp.* (bougasvlei) seem to be highly conserved which allows

for the development of specific probe-based assays that may be used to detect all individuals within the species (Schnittger et al., 2003; Pienaar et al., 2011b, 2014; Ros-García et al., 2012b). Conversely, if it is assumed that some clades do represent single species with genotypic diverse members, they are so diverse that no single assay may be able to detect all genotypes and therefore all individuals within the species. This necessitates the development of multiple assays or probes for geographically diverse populations (Gubbels et al., 2000b; Chaisi et al., 2013a; Perera et al., 2014).

The same generalities are found for all clades in the literature, *i.e.* when different genotypes are observed from a single host species and genotypes group within a monophyletic clade, it is assumed that these are the same *Theileria* species. Conversely, finding a single genotype associated with a single host species, even if the sampling was small and the genotype has not previously been observed, will lead to the conclusion that this is a unique species. From an evolutionary perspective this situation is extremely confusing. It suggests recent speciation of genotypes such as *T. parva*, *T. sp.* (buffalo) and *T. sp.* (bougasvlei) that allows conservation of the 18S V4 hypervariable region (Mans et al., 2011a; Pienaar et al., 2014), even though evidence suggests a more ancient origin, since *T. parva* for example shows extensive genetic diversity in field populations (McKeever, 2009). It also implies that the 18S ribosomal genes in the *T. buffeli*, *T. equi* and *T. mutans* clades had undergone rapid evolution, contravening the ribosomal molecular clock (Lack et al., 2012), or that these clades have an ancient origin with diversification of lineages, while maintaining their species identity. The latter may occur via bottleneck events that result in clonal populations in which the 18S gene diverged (Tibayrenc and Ayala, 2002; Tibayrenc, 2006; McKeever, 2009). Why this would be more prone in certain clades is uncertain, while divergence of the whole genome may be expected. Designation of species then becomes arbitrary and retaining diverse genotypes in a single species may be a social and philosophical response to a complex problem (Tibayrenc, 2006). An alternative and more parsimonious scenario would posit dispersal and geographic isolation linked with speciation of vertebrate host species during the Middle–Late Miocene (~13–8 MYA) for the ancestral lineages to the Bubalina (African and Asiatic buffalo and cattle) (Hassanin et al., 2012) and the Pliocene (4–3 MYA) for the Bovina (cattle and bison) (Hassanin et al., 2012, 2013) and caballine and non-caballine equids (Vilstrup et al., 2013). This coincided with diversification of *Theileria* parasites within the different clades (Criado-Fornelio et al., 2003; Gou et al., 2013). Subsequent breakdown of geographic barriers could have resulted in co-infection of Bovini hosts by related genotypes unable to sexually recombine, but still able to infect similar vectors and hosts. In this case, the genotypic diversity observed within the respective clades should be interpreted as distinct lineages, *i.e.* species. With ~6000 named apicomplexan species, which comprise only 0.1% of the estimated number of apicomplexan species (1.2–10 million) (Morrison, 2009), it seems rather conservative that only 91 genotypes and ~40 species have been described thus far in the *Theileria*.

4. Sensitivity of molecular assays, carrier state and parasitaemia ranges

The theoretical limit of detection (LOD) of molecular PCR assays is three molecules per assay, which generally equates with a sensitivity cut-off of ~37 cycles (Bustin, 2010). In practice, however, this may range from 3 – > 100 molecules and a cut-off from 37 to 40 cycles (Burns and Valdivia, 2008; Bhoora et al., 2010a). This depends on a complex array of factors, including assay mixture, robustness of the polymerase enzyme, inhibitors in the sample, primer and probe design and thermodynamics, effective temperature cycling and detector sensitivity of PCR equipment. In the case of *Theileria*,

most assays detect genomic DNA and sensitivity will therefore also be influenced by DNA extraction efficiency (*i.e.* yield), gene copy number in the genome, number of parasites (genomic copies) per cell sampled and the parasitaemia level.

Sensitive quantitative real-time PCR assays have been developed for *T. parva* and *T. equi*. In all cases the analytical LOD ranges from 3 to 10 copies, with efficiencies that fall within the acceptable levels for a PCR assay (Kim et al., 2008; Sibeko et al., 2008; Bhoora et al., 2010a; Pienaar et al., 2011b). It is, however, of interest to put this into practical perspective with regard to the detection limit in carrier animals. For the *T. parva* test, the LOD for whole blood was ~10 parasites from the equivalent of 5 µL whole blood (Sibeko et al., 2008). This calculates back to ~2000 parasites/mL of blood and 19 million parasites in a 500 kg buffalo. Similarly, nested PCR based on the p104 gene, conventional PCR based on the p104 gene and reverse line blot for *T. parva* had LOD of 400–2000 parasites/mL of blood (Skilton et al., 2002; Oura et al., 2004; Odongo et al., 2010). For the *T. equi* assay, the LOD was 190 parasites from the equivalent of 5 µL whole blood (Bhoora et al., 2010a). This calculates to a detection limit of 376–1520 million parasites in a 500 kg horse if the average number of parasites varies from 1 to 4 per red blood cell. Similarly, tests for *T. annulata* (d'Oliveira et al., 1995; Kirvar et al., 2000; Ros-García et al., 2012b) and *T. buffeli* (Jeong et al., 2003) gave LOD of 23–54 and 108 million parasites, respectively, in cattle weighing 500 kg. A sensitive PCR-RLB assay for *T. annulata*, based on the cytochrome b gene, gave a LOD of 100,000 parasites per liter (Bilgic et al., 2010). A nested PCR for *T. ovis* gave a LOD of 1200 parasites/mL of blood (Altay et al., 2005), which calculates to a detection limit of 5.7 million parasites in an 80 kg sheep. It would thus seem that in general, the LOD in animals is >400,000 parasites/L blood, implying that assays for most *Theileria* parasites, even if sensitive from an analytical perspective, may fail to detect carrier animals from a diagnostic perspective. The potential for false-negative results can, however, only be assessed within the biological context of the parasite within its vertebrate host, *i.e.* parasitaemia ranges and the carrier-state that impacts on the ability to infect the tick vector and subsequent transmission to a new host.

Parasitaemia is generally expressed as percentage infected red blood cells. It would seem that all *Theileria* species have defined parasitaemia ranges (Young et al., 1986; Ueti et al., 2012). During clinical reactions, piroplasm parasitaemia may be as high as 1–45% depending on the species (Uilenberg and Schreuder, 1976; Young et al., 1978; de Waal, 1992; Latif et al., 2002; Schetters et al., 2010), but this generally drops well below 1% for the asymptomatic carrier state (Uilenberg and Schreuder, 1976; Young et al., 1986; Grootenhuys et al., 1987; Zaemi et al., 2011; Ueti et al., 2012). Quantitative real-time PCR has allowed estimation of parasitaemia ranges in hosts and the first interesting observation was that parasitaemia ranges approximate normal distributions (Pienaar et al., 2011a, 2014). These ranges will be determined by the specific biology of the parasite in the carrier state, *i.e.* ability to maintain and propagate in the vertebrate host (will be discussed later). It also implies that the specific parasitaemia range of a species may determine its practical LOD. As such, >95% of the *T. parva* population in African buffalo have parasitaemias >0.00001%, which is well above the determined detection limit (Pienaar et al., 2011a). The same holds for *T. sp.* (buffalo) and *T. sp.* (bougasvlei) (Pienaar et al., 2011a, 2014). It is difficult to make similar estimations for other *Theileria* species since the numbers of animals sampled do not allow construction of representative frequency distribution plots of published Ct values (Jeong et al., 2003; Kim et al., 2008; Bhoora et al., 2010a). However, from work performed in our own laboratories we may conclude that the parasitaemia range for *T. equi* is above the LOD in horses from endemic areas (personal observation). Similarly, most benign *Theileria* species generally have higher parasitaemia ranges in the carrier state and it may be concluded that real-time PCR assays for most

Theileria species should be able to detect the majority of infected animals in a herd.

Exceptions to this may occur where the sensitivity of the assay is affected by the presence of mixed infections. Sensitivity of the RLB is severely affected by the fact that universal primers are used and these are depleted by the predominant species present, thereby suppressing the signal from less abundant template (Pienaar et al., 2011a). The sensitivity of the hybridisation assay for *T. parva* (Sibeko et al., 2008) is similarly affected by the presence of *T. sp.* (buffalo)-like parasites, since the primers for this assay also amplify the latter parasite template (Pienaar et al., 2011a). Design of more specific primers and stringent conditions alleviated this suppression problem, at the cost of being slightly less sensitive in regard to analytical sensitivity, but with increased overall performance for field samples (Pienaar et al., 2011b). Other assays that may also be affected by sensitivity suppression include the cox III assay that uses universal primers and may be compounded by the implementation of a nested PCR strategy (Chaisi et al., 2013b). In regard to the latter, nested PCR may increase overall sensitivity (Schnittger et al., 2004; Odongo et al., 2010; Ueti et al., 2012), but may introduce contamination problems in the routine diagnostic laboratory, or if primers are not specific, still lead to PCR suppression by the most dominant genotypes. In addition, it is well accepted that parasitaemia can fluctuate and drop below detection limits, only to increase during relapse, the so-called sporadic carrier state (Norval et al., 1992). This generally occurs under conditions of stress, immune suppression or splenectomy (Hooshmand-Rad, 1976; Zobba et al., 2008). Fluctuation in parasitaemia from below the PCR LOD is, however, not extensively documented and may contribute little to herd surveillance, but may have a significant impact on disease epidemiology, precipitating in disease outbreaks. Multiple testing rounds are generally considered as solution for this eventuality (Ueti et al., 2012).

The carrier-state for piroplasmida may be defined as the long-term persistence of a parasite in its host, with the ability of transmission to other hosts via vector infection, to maintain a transmission cycle. Long-term persistence implies that the parasite can maintain itself and propagate in the vertebrate host while escaping the immune system. In most *Theileria*, it is accepted that the piroplasm stage is maintained in the host via asexual division and re-infection of red blood cells (Norval et al., 1992). In the case of *T. parva*, limited division of piroplasms has been observed (Conrad et al., 1986; Fawcett et al., 1987). However, it has been proposed that a reservoir of schizonts that maintains parasitaemia levels exists that escapes the immune system, especially since schizonts sampled 657 days after initial infection could be propagated in cell culture (Grootenhuys et al., 1987). Maintenance of parasitaemia levels is necessary, since red blood cells have a lifespan that range from 70 to 160 days in large mammals before being destroyed (Adili and Melizi, 2014). This would imply that parasites need to replicate and re-infect new red blood cells at least once in this period to maintain their parasitaemia levels. In this regard, remarkable stability in piroplasm parasitaemia levels has been observed in *T. parva* carrier buffalo kept under tick-free conditions where animals maintained parasitaemia levels for up to 20 years (Pienaar et al., 2014). It is therefore also accepted that once infected, animals may remain lifelong carriers of *Theileria* (de Waal, 1992; Norval et al., 1992). While this is generally true, fluctuations in parasitaemia and loss of carrier state has also been documented, especially in cases where true carrier-status cannot be established, such as the case for Corridor disease and East Coast Fever.

In the case of East Coast Fever, infection of animals using specific strains of Muguga resulted in fluctuating parasitaemia above and below the LOD of the assay, showing in principle that fluctuating parasitaemia was possible for *T. parva*. Animals infected with this strain will eventually clear it completely and will not be infective to ticks (Bishop et al., 1992; Skilton et al., 2002; Odongo et al.,

2010). Whether this is representative for field strains is not known, since this strain has been propagated under laboratory conditions for extensive time periods. In contrast, the Marikebuni strain maintained the carrier-state indefinitely well above the LOD (Bishop et al., 1992; Skilton et al., 2002). Prevalence of *T. parva* in cattle from East Coast fever endemic regions range between 37 and 42% using nested PCR (Odongo et al., 2010), compared to ~80% in buffalo (Pienaar et al., 2011a), suggesting that the former is not compatible with an endemic stable situation. In East Coast fever, the carrier-state may therefore be strain dependent and intermittent. For Corridor disease, cattle infected with *T. parva* that survive may maintain the infection for up to 3 months, but with decreasing parasitaemia until the infection is cleared, with no relapse observed yet (even in splenectomised cattle) and unable to infect ticks (Mbizeni et al., 2013). The latter case may represent the dynamics of a parasite not fully adapted to its host and where the carrier-state may only be intermittent. In this case, the seasonality of brown ear ticks in South Africa preclude the maintenance of a transmission cycle (Norval et al., 1991; Mbizeni et al., 2013), and buffalo-adapted *T. parva* may therefore not have true carrier status in cattle. The latter becomes an important consideration in epidemiological surveillance, since the presence of parasite DNA or positive serology may not be indicative of true host carrier status, but only exposure to parasites during tick transmission.

5. Epidemiology in the context of species

Theileria epidemiology considers parasite and vector distribution, mortality and morbidity of disease outbreaks, disease outbreak risk assessment and disease control measures, socio-economic factors, climate change, host resistance and susceptibility (Gachohi et al., 2012). While many of these factors may be addressed by clinical differential diagnosis, vector mapping, vaccination strategies and control policy, serological and molecular diagnosis remain central to confirmation of parasite identity, distribution, disease surveillance, assessment of vaccine and as tool for informative control policies (Bakheit et al., 2007). From a disease control standpoint, accurate diagnostics and communication is hampered by loose treatment of species concepts and lumping of genotypes into large groups collectively designated as single species. Examples of this are the emergence of different virulent “strains or genotypes” from “*T. buffeli*” that cause outbreaks in geographic diverse regions (Izzo et al., 2010; Kamau et al., 2011; Sivakumar et al., 2014), or the presence of *T. parva* in South Africa that causes buffalo-derived Corridor disease, but not cattle-derived East Coast fever (Mbizeni et al., 2013). In contrast to the *T. buffeli* clade where ample genetic evidence exists for diverse genotypes, the case for separating the *T. parva* sub-types based on genetic evidence is not clear and the trinomial system used to distinguish Corridor disease (*T. parva lawrencei*), East Coast fever (*T. parva parva*) and January disease (*T. parva bovis*) was abolished for more than 20 years ago (Norval et al., 1992; Perry and Young, 1993). The biological differences between these “strains” are, however, well documented (Lawrence, 1979; Perry and Young, 1993) and treatment as one species disregards the epidemiology of these parasites. The search for markers to distinguish between these strains has continued, but with little success (Sibeko et al., 2010, 2011). Again, the probability exists that these parasites are related via an evolutionary continuum, *i.e.* recent divergence that will eventually result in speciation due to host specificity.

The study of parasite epidemiology is further complicated by treatment of distinct genotypes as discrete units (species), since this may mask additive or competitive effects between genotypic populations. Treatment of *T. sp.* (buffalo) and *T. sp.* (bougasvlei) as one species (Chaisi et al., 2011; Mans et al., 2011b) could have missed their skewed geographic distribution with regard to each other (Pienaar et al., 2014). Similarly, genotypes from the *T. mutans*,

T. velifera and *T. buffeli* clades (Chaisi et al., 2013a, 2014) have skewed distributions in cattle and African buffalo in southern Africa based on screening data of ~1000 cattle and 1000 buffalo distributed across the region (B.J. Mans, personal observation). While these are anecdotal in regard to all *Theileria* genotypes, similar disparities in genotypic distributions are likely to be prevalent in different hosts and geographic regions. To ultimately understand the phylogeographic history of the *Theileria* genus, all genotypes need to be accounted for. In this regard, lumping genotypes into species does not unify, but rather obscures lineage specific life histories.

6. Conclusion

Extensive resources exist for diagnostics of *Theileria* parasites. However, many of the available assays are restricted by limitations in specificity and sensitivity, and a positive diagnosis may in some cases only be made by multiple assays or xenodiagnosis. Critical examination of the measurements of uncertainty inherent in all assays will allow future improvements in diagnostic capability and may also contribute towards a better understanding of parasite biology. This in turn should have a positive impact on disease epidemiology.

Acknowledgements

Funding for this review was provided by the Department of Agriculture, Forestry and Fisheries (DAFF) of South Africa (OV21/03/C148) and *Theileria* Diagnostics (15/08/1P01) of Parasites, Vectors and Vector-borne Diseases (PVVD), Onderstepoort Veterinary Institute, Agricultural Research Council, South Africa.

Conflict of interest

The authors declared that there is no conflict of interest.

References

- Abdo, J., Kristersson, T., Seitzer, U., Renneker, S., Merza, M., Ahmed, J., 2010. Development and laboratory evaluation of a lateral flow device (LFD) for the serodiagnosis of *Theileria annulata* infection. *Parasitol. Res.* 107, 1241–1248.
- Adili, N., Melizi, M., 2014. Preliminary study of the influence of red blood cells morphology on the species determination of domestic animals. *Vet. World* 7, 219–223.
- Adl, S.M., Simpson, A.G., Lane, C.E., Lukeš, J., Bass, D., Bowser, S.S., et al., 2012. The revised classification of eukaryotes. *J. Eukaryot. Microbiol.* 59, 429–493.
- Alhassan, A., Pumidonming, W., Okamura, M., Hirata, H., Battsetseg, B., Fujisaki, K., et al., 2005. Development of a single-round and multiplex PCR method for the simultaneous detection of *Babesia caballi* and *Babesia equi* in horse blood. *Vet. Parasitol.* 129, 43–49.
- Alhassan, A., Govind, Y., Tam, N.T., Thekisoe, O.M.M., Yokoyama, N., Inoue, N., et al., 2007a. Comparative evaluation of the sensitivity of LAMP, PCR and *in vitro* culture methods for the diagnosis of equine piroplasmiasis. *Parasitol. Res.* 100, 1165–1168.
- Alhassan, A., Thekisoe, O.M., Yokoyama, N., Inoue, N., Motloang, M.Y., Mbatia, P.A., et al., 2007b. Development of loop-mediated isothermal amplification (LAMP) method for diagnosis of equine piroplasmiasis. *Vet. Parasitol.* 143, 155–160.
- Allsopp, B.A., Baylis, H.A., Allsopp, M.T.P.E., Cavalier-Smith, T., Bishop, R.P., Carrington, D.M., et al., 1993. Discrimination between six species of *Theileria* using oligonucleotide probes which detect small subunit ribosomal RNA sequences. *Parasitology* 107, 157–165.
- Altay, K., Dumanli, N., Holman, P.J., Aktas, M., 2005. Detection of *Theileria ovis* in naturally infected sheep by nested PCR. *Vet. Parasitol.* 127, 99–104.
- Altay, K., Dumanli, N., Aktas, M., 2007. Molecular identification, genetic diversity and distribution of *Theileria* and *Babesia* species infecting small ruminants. *Vet. Parasitol.* 147, 161–165.
- Altay, K., Aktas, M., Dumanli, N., Aydin, M.F., 2008. Evaluation of a PCR and comparison with RLB for detection and differentiation of *Theileria* sp. MK and other *Theileria* and *Babesia* species of small ruminants. *Parasitol. Res.* 103, 319–323.
- André, M.R., Baccarim Denardi, N.C., Marques de Sousa, K.C., Gonçalves, L.R., Henrique, P.C., Grosse Rossi Ontivero, C.R., et al., 2014. Arthropod-borne pathogens circulating in free-roaming domestic cats in a zoo environment in Brazil. *Ticks Tick Borne Dis.* 5, 545–551.
- Bai, Q., Liu, G.Y., Yin, H., Zhao, Q.Z., Liu, D.K., Ren, J.X., et al., 2002. *Theileria sinensis* nov: study on molecular taxonomy. *Xu Mu Shou Yi Xue Bao* 33, 185–190.
- Bakheit, M.A., Schnitger, L., Salih, D.A., Boguslawski, K., Beyer, D., Fadl, M., et al., 2004. Application of the recombinant *Theileria annulata* surface protein in an indirect ELISA for the diagnosis of tropical theileriosis. *Parasitol. Res.* 92, 299–302.
- Bakheit, M.A., Seitzer, U., Ahmed, J.S., 2006. A new recombinant protein-based ELISA for the diagnosis of malignant theileriosis of sheep and goats. *Parasitol. Res.* 98, 145–149.
- Bakheit, M.A., Seitzer, U., Mbatia, P.A., Ahmed, J.S., 2007. Serological diagnostic tools for the major tick-borne protozoan diseases of livestock. *Parasitologia* 49, S53–S62.
- Barnett, S.F., 1977. *Theileria*. In: Kreier, J.P. (Ed.), *Parasitic Protozoa*, vol. IV. Academic Press, New York, USA, pp. 77–113.
- Bazarusanga, T., Geysen, D., Vercauteren, J., Marcotty, T., 2008. The sensitivity of PCR and serology in different *Theileria parva* epidemiological situations in Rwanda. *Vet. Parasitol.* 154, 21–31.
- Beck, R., Vojta, L., Mrljak, V., Marinculić, A., Beck, A., Zivicnjak, T., et al., 2009. Diversity of *Babesia* and *Theileria* species in symptomatic and asymptomatic dogs in Croatia. *Int. J. Parasitol.* 39, 843–848.
- Bhoora, R., Franssen, L., Oosthuizen, M.C., Guthrie, A.J., Zweygarth, E., Penzhorn, B.L., et al., 2009. Sequence heterogeneity in the 18S rRNA gene within *Theileria equi* and *Babesia caballi* from horses in South Africa. *Vet. Parasitol.* 159, 112–120.
- Bhoora, R., Quan, M., Franssen, L., Butler, C.M., van der Kolk, J.H., Guthrie, A.J., et al., 2010a. Development and evaluation of real-time PCR assays for the quantitative detection of *Babesia caballi* and *Theileria equi* infections in horses from South Africa. *Vet. Parasitol.* 168, 201–211.
- Bhoora, R., Buss, P., Guthrie, A.J., Penzhorn, B.L., Collins, N.E., 2010b. Genetic diversity of piroplasms in plains zebra (*Equus quagga burchellii*) and Cape mountain zebra (*Equus zebra zebra*) in South Africa. *Vet. Parasitol.* 174, 145–149.
- Bilgic, H.B., Karagenc, T., Shiels, B., Tait, A., Eren, H., Weir, W., 2010. Evaluation of cytochrome b as a sensitive target for PCR based detection of *T. annulata* carrier animals. *Vet. Parasitol.* 174, 341–347.
- Bishop, R., Sohanpal, B., Kariuki, D.P., Young, A.S., Nene, V., Baylis, H., et al., 1992. Detection of a carrier state in *Theileria parva*-infected cattle by the polymerase chain reaction. *Parasitology* 104, 215–232.
- Bishop, R., Musoke, A., Morzaria, S., Gardner, M., Nene, V., 2004. *Theileria*: intracellular protozoan parasites of wild and domestic ruminants transmitted by ixodid ticks. *Parasitology* 129, S271–S283.
- Bishop, R.P., Spooner, P.R., Kanhai, G.K., Kiarie, J., Latif, A.A., Hove, T., et al., 1994a. Molecular characterization of *Theileria* parasites: application to the epidemiology of theileriosis in Zimbabwe. *Parasitology* 109, 573–581.
- Bishop, R.P., Sohanpal, B.K., Morzaria, S.P., 1994b. Cloning and characterisation of a repetitive DNA sequence from *Theileria mutans*: application as a species-specific probe. *Parasitol. Res.* 80, 33–41.
- Burns, M., Valdivia, H., 2008. Modelling the limit of detection in real-time quantitative PCR. *Eur. Food Res. Technol.* 226, 1513–1524.
- Burridge, M.J., 1971. Application of the indirect fluorescent antibody test in experimental East Coast fever (*Theileria parva* infection in cattle). *Res. Vet. Sci.* 12, 338–341.
- Burridge, M.J., Kimber, C.D., 1972. The indirect fluorescent antibody test for experimental East Coast fever (*Theileria parva* infection of cattle). Evaluation of a cell culture schizont antigen. *Res. Vet. Sci.* 13, 451–455.
- Bustin, S.A., 2010. Why the need for qPCR publication guidelines? – the case for MIQE. *Methods* 50, 217–226.
- Butler, C.M., Nijhof, A.M., van der Kolk, J.H., de Haseth, O.B., Taoufik, A., Jongejan, F., et al., 2008. Repeated high dose imidocarb dipropionate treatment did not eliminate *Babesia caballi* from naturally infected horses as determined by PCR-reverse line blot hybridization. *Vet. Parasitol.* 151, 320–322.
- Cacciò, S., Cammà, C., Onuma, M., Severini, C., 2000. The beta-tubulin gene of *Babesia* and *Theileria* parasites is an informative marker for species discrimination. *Int. J. Parasitol.* 30, 1181–1185.
- Camacho, A.T., Pallas, E., Gestal, J.J., Guitián, F.J., Olmeda, A.S., Goethert, H.K., et al., 2001. Infection of dogs in north-west Spain with a *Babesia microti*-like agent. *Vet. Rec.* 149, 552–555.
- Chae, J., Lee, J., Kwon, O., Holman, P.J., Waghele, S.D., Wagner, G.G., 1998. Nucleotide sequence heterogeneity in the small subunit ribosomal RNA gene variable (V4) region among and within geographic isolates of *Theileria* from cattle, elk and white-tailed deer. *Vet. Parasitol.* 75, 41–52.
- Chae, J.S., Waghele, S.D., Craig, T.M., Kocan, A.A., Wagner, G.G., Holman, P.J., 1999a. Two *Theileria cervi* SSU rRNA gene sequence types found in isolates from white-tailed deer and elk in North America. *J. Wildl. Dis.* 35, 458–465.
- Chae, J.S., Allsopp, B.A., Waghele, S.D., Park, J.H., Kakuda, T., Sugimoto, C., et al., 1999b. A study of the systematics of *Theileria* spp. based upon small-subunit ribosomal RNA gene sequences. *Parasitol. Res.* 85, 877–883.
- Chaisi, M.E., Sibeko, K.P., Collins, N.E., Potgieter, F.T., Oosthuizen, M.C., 2011. Identification of *Theileria parva* and *Theileria* sp. (buffalo) 18S rRNA gene sequence variants in the African Buffalo (*Syncerus caffer*) in southern Africa. *Vet. Parasitol.* 182, 150–162.
- Chaisi, M.E., Janssens, M.E., Vermeiren, L., Oosthuizen, M.C., Collins, N.E., Geysen, D., 2013a. Evaluation of a real-time PCR test for the detection and discrimination of *Theileria* species in the African buffalo (*Syncerus caffer*). *PLoS ONE* 8, e75827.
- Chaisi, M.E., Collins, N.E., Potgieter, F.T., Oosthuizen, M.C., 2013b. Sequence variation identified in the 18S rRNA gene of *Theileria mutans* and *Theileria velifera* from the African buffalo (*Syncerus caffer*). *Vet. Parasitol.* 191, 132–137.
- Chaisi, M.E., Collins, N.E., Oosthuizen, M.C., 2014. Phylogeny of *Theileria buffeli* genotypes identified in the South African buffalo (*Syncerus caffer*) population. *Vet. Parasitol.* 204, 87–95.

- Clark, P., Spencer, P.B.S., 2007. Description of three new species of *Theileria* Bettencourt, Franca Borges, 1907 from Macropodoidea in Western Australia. *Trans. R. Soc. S. Aust.* 131, 100–106.
- Collins, N.E., Allsopp, M.T., Allsopp, B.A., 2002. Molecular diagnosis of theileriosis and heartwater in bovines in Africa. *Trans. R. Soc. Trop. Med. Hyg.* 96, S217–S224.
- Conrad, P.A., Denham, D., Brown, C.G.D., 1986. Intraerythrocytic multiplication of *Theileria parva* in vitro: an ultrastructural study. *Int. J. Parasitol.* 16, 223–229.
- Conrad, P.A., Stagg, D.A., Grootenhuys, J.G., Irvin, A.D., Newson, J., Njamunggeh, R.E., et al., 1987. Isolation of *Theileria* parasites from African buffalo (*Syncerus caffer*) and characterization with anti-schizont monoclonal antibodies. *Parasitology* 94, 413–423.
- Criado, A., Martinez, J., Buling, A., Barba, J.C., Merino, S., Jefferies, R., et al., 2006. New data on epizootiology and genetics of piroplasmids based on sequences of small ribosomal subunit and cytochrome b genes. *Vet. Parasitol.* 142, 238–247.
- Criado-Fornelio, A., 2007. A review of nucleic acid-based diagnostic tests for *Babesia* and *Theileria*, with emphasis on bovine piroplasmids. *Parasitologia* 49, S39–S44.
- Criado-Fornelio, A., Martinez-Marcos, A., Buling-Saraña, A., Barba-Carretero, J.C., 2003. *Babesia*, *Theileria* and *Hepatozoon* in southern Europe: part II. Phylogenetic analysis and evolutionary history. *Vet. Parasitol.* 114, 173–194.
- Criado-Fornelio, A., González-del-Río, M.A., Buling-Saraña, A., Barba-Carretero, J.C., 2004. The “expanding universe” of piroplasmids. *Vet. Parasitol.* 119, 337–345.
- de Vos, A.J., Roos, J.A., 1981. The isolation of *Theileria taurotragi* in South Africa. *Onderstepoort J. Vet. Res.* 48, 149–153.
- de Waal, D.T., 1992. Equine piroplasmidosis: a review. *Br. Vet. J.* 148, 6–14.
- d’Oliveira, C., van der Weide, M., Habela, M.A., Jacquet, P., Jongejan, F., 1995. Detection of *Theileria annulata* in blood samples of carrier cattle by PCR. *J. Clin. Microbiol.* 33, 2665–2669.
- Darghouth, M.E., Bouattour, A., Ben Miled, L., Sassi, L., 1996. Diagnosis of *Theileria annulata* infection of cattle in Tunisia: comparison of serology and blood smears. *Vet. Res.* 27, 613–621.
- Dschunkowsky, E., Luhs, J., 1904. Die piroplasmosen der rinder. *Zentralbl. Bakteriol. Mikrobiol. Hyg.* 35, 486–493.
- Eickbush, T.H., Eickbush, D.G., 2007. Finely orchestrated movements: evolution of the ribosomal RNA genes. *Genetics* 175, 477–485.
- Fawcett, D.W., Conrad, P.A., Grootenhuys, J.G., Morzaria, S.P., 1987. Ultrastructure of the intra-erythrocytic stage of *Theileria* species from cattle and waterbuck. *Tissue Cell* 19, 643–655.
- Fujisaki, K., Kawazu, S., Kamio, T., 1994. The taxonomy of the bovine *Theileria* spp. *Parasitol. Today* 10, 31–33.
- Gachohi, J., Skilton, R., Hansen, F., Ngumi, P., Kitala, P., 2012. Epidemiology of East Coast fever (*Theileria parva* infection) in Kenya: past, present and the future. *Parasit. Vectors* 5, 194.
- García-Sanmartín, J., Aurensetxe, O., Barral, M., Marco, I., Lavin, S., García-Pérez, A.L., et al., 2007. Molecular detection and characterization of piroplasmids infecting cervids and chamois in Northern Spain. *Parasitology* 134, 391–398.
- Gardner, M.J., Bishop, R., Shah, T., de Villiers, E.P., Carlton, J.M., Hall, N., et al., 2005. Genome sequence of *Theileria parva*, a bovine pathogen that transforms lymphocytes. *Science* 309, 134–137.
- Geysen, D., Bishop, R., Skilton, R., Dolan, T.T., Morzaria, S., 1999. Molecular epidemiology of *Theileria parva* in the field. *Trop. Med. Int. Health* 4, A21–A27.
- Ghaemi, P., Hoghooghi-Rad, N., Shayan, P., Eckert, B., 2012. Detection of *Theileria orientalis* in Iran by semi-nested PCR. *Parasitol. Res.* 110, 527–531.
- Githaka, N., Konnai, S., Kariuki, E., Kanduma, E., Murata, S., Ohashi, K., 2012. Molecular detection and characterization of potentially new *Babesia* and *Theileria* species/variants in wild felids from Kenya. *Acta Trop.* 124, 71–78.
- Githaka, N., Konnai, S., Skilton, R., Kariuki, E., Kanduma, E., Murata, S., et al., 2013. Genotypic variations in field isolates of *Theileria* species infecting giraffes (*Giraffa camelopardalis tippelskirchi* and *Giraffa camelopardalis reticulata*) in Kenya. *Parasitol. Int.* 62, 448–453.
- Githaka, N., Konnai, S., Bishop, R., Odongo, D., Lekolool, I., Kariuki, E., et al., 2014. Identification and sequence characterization of novel *Theileria* genotypes from the waterbuck (*Kobus defassa*) in a *Theileria parva*-endemic area in Kenya. *Vet. Parasitol.* 202, 180–193.
- Gou, H., Guan, G., Liu, A., Ma, M., Chen, Z., Liu, Z., et al., 2013. Coevolutionary analyses of the relationships between piroplasmids and their hard tick hosts. *Ecol. Evol.* 3, 2985–2993.
- Govaerts, M.M., Voet, M., Volckaert, G., Goddeeris, B.M., 1998. PCR amplification and sequence of the p33 piroplasm surface antigen gene of a *Theileria* species isolated from cattle in west Java, Indonesia. *Ann. N. Y. Acad. Sci.* 849, 126–136.
- Gray, M.A., Luckins, A.G., Rae, P.F., Brown, C.G., 1980. Evaluation of an enzyme immunoassay for serodiagnosis of infections with *Theileria parva* and *T. annulata*. *Res. Vet. Sci.* 29, 360–366.
- Grootenhuys, J.G., Leitch, B.L., Stagg, D.A., Dolan, T.T., Young, A.S., 1987. Experimental induction of *Theileria parva lawrencei* carrier state in an African buffalo (*Syncerus caffer*). *Parasitology* 94, 425–431.
- Gubbels, J.M., de Vos, A.P., van der Weide, M., Viseras, J., Schouls, L.M., de Vries, E., et al., 1999. Simultaneous detection of bovine *Theileria* and *Babesia* species by reverse line blot hybridization. *J. Clin. Microbiol.* 37, 1782–1789.
- Gubbels, M.J., d’Oliveira, C., Jongejan, F., 2000a. Development of an indirect Tams1 enzyme-linked immunosorbent assay for diagnosis of *Theileria annulata* infection in cattle. *Clin. Diagn. Lab. Immunol.* 7, 404–411.
- Gubbels, M.J., Hong, Y., van der Weide, M., Qi, B., Nijman, I.J., Guanyuan, L., et al., 2000b. Molecular characterisation of the *Theileria buffeli/orientalis* group. *Int. J. Parasitol.* 30, 943–952.
- Gubbels, M.J., Yin, H., Bai, Q., Liu, G., Nijman, I.J., Jongejan, F., 2002. The phylogenetic position of the *Theileria buffeli* group in relation to other *Theileria* species. *Parasitol. Res.* 88, S28–S32.
- Han, J.J., Jang, H.J., Lee, S.J., Na, K.J., 2009. High prevalence of *Theileria* sp. in wild Chinese Water Deer (*Hydropotes inermis argyropus*) in South Korea. *Vet. Parasitol.* 164, 311–314.
- Hassanin, A., Delsuc, F., Ropiquet, A., Hammere, C., Jansen van Vuuren, B., Matthee, C., et al., 2012. Pattern and timing of diversification of Cetartiodactyla (Mammalia, Laurasiatheria), as revealed by a comprehensive analysis of mitochondrial genomes. *C. R. Biol.* 335, 32–50.
- Hassanin, A., An, J., Ropiquet, A., Nguyen, T.T., Couloux, A., 2013. Combining multiple autosomal introns for studying shallow phylogeny and taxonomy of Laurasiatherian mammals: application to the tribe Bovini (Cetartiodactyla, Bovidae). *Mol. Phylogenet. Evol.* 66, 766–775.
- Hayashida, K., Hara, Y., Abe, T., Yamasaki, C., Toyoda, A., Kosuge, T., et al., 2012. Comparative genome analysis of three eukaryotic parasites with differing abilities to transform leukocytes reveals key mediators of *Theileria*-induced leukocyte transformation. *mBio* 3, e204–e212.
- Heidarpoor Bami, M., Haddadzadeh, H.R., Kazemi, B., Khazraiania, P., Bandehpour, M., Aktas, M., 2009. Molecular identification of ovine *Theileria* species by a new PCR-RFLP method. *Vet. Parasitol.* 161, 171–177.
- Heim, A., Passos, L.M., Ribeiro, M.F., Costa-Júnior, L.M., Bastos, C.V., Cabral, D.D., et al., 2007. Detection and molecular characterization of *Babesia caballi* and *Theileria equi* isolates from endemic areas of Brazil. *Parasitol. Res.* 102, 63–68.
- Hooshmand-Rad, P., 1976. The pathogenesis of anaemia in *Theileria annulata* infection. *Res. Vet. Sci.* 20, 324–329.
- Huang, X., Xuan, X., Yokoyama, N., Xu, L., Suzuki, H., Sugimoto, C., et al., 2003. High-level expression and purification of a truncated merozoite antigen-2 of *Babesia equi* in *Escherichia coli* and its potential for immunodiagnosis. *J. Clin. Microbiol.* 41, 1147–1151.
- Huang, X., Xuan, X., Xu, L., Zhang, S., Yokoyama, N., Suzuki, N., et al., 2004. Development of an immunochromatographic test with recombinant EMA-2 for the rapid detection of antibodies against *Babesia equi* in horses. *J. Clin. Microbiol.* 42, 359–361.
- Ilhan, T., Williamson, S., Kirvar, E., Shiels, B., Brown, C.G., 1998. *Theileria annulata*: carrier state and immunity. *Ann. N. Y. Acad. Sci.* 849, 109–125.
- Izzo, M.M., Poe, I., Horadagoda, N., De Vos, A.J., House, J.K., 2010. Haemolytic anaemia in cattle in NSW associated with *Theileria* infections. *Aust. Vet. J.* 88, 45–51.
- Jefferies, R., Ryan, U.M., Irwin, P.J., 2007. PCR-RFLP for the detection and differentiation of the canine piroplasm species and its use with filter paper-based technologies. *Vet. Parasitol.* 144, 20–27.
- Jeong, W., Kweon, C.H., Kang, S.W., Paik, S.G., 2003. Diagnosis and quantification of *Theileria sergenti* using TaqMan PCR. *Vet. Parasitol.* 111, 287–295.
- Jeong, W., Kweon, C.H., Kim, J.M., Jang, H., Paik, S.G., 2005. Serological investigation of *Theileria sergenti* using latex agglutination test in South Korea. *J. Parasitol.* 91, 164–169.
- Jongejan, F., Musisi, F.L., Moorhouse, P.D.S., Snacken, M., Uilenberg, G., 1986. *Theileria taurotragi* in Zambia. *Vet. Q.* 8, 261–263.
- Kachani, M., Spooner, R.L., Rae, P., Bell-Sakyi, L., Brown, C.G., 1992. Stage-specific responses following infection with *Theileria annulata* as evaluated using ELISA. *Parasitol. Res.* 78, 43–47.
- Kamau, J., de Vos, A.J., Playford, M., Salim, B., Kinyanjui, P., Sugimoto, C., 2011. Emergence of new types of *Theileria orientalis* in Australian cattle and possible cause of theileriosis outbreaks. *Parasit. Vectors* 4, 22.
- Kappmeyer, L.S., Thiagarajan, M., Herndon, D.R., Ramsay, J.D., Caler, E., Djikeng, A., et al., 2012. Comparative genomic analysis and phylogenetic position of *Theileria equi*. *BMC Genomics* 13, 603.
- Kariuki, E.K., Penzhorn, B.L., Horak, I.G., 2012. Ticks (Acari: Ixodidae) infesting cattle and African buffaloes in the Tsavo conservation area, Kenya. *Onderstepoort J. Vet. Res.* 79, 437.
- Katende, J., Morzaria, S., Toye, P., Skilton, R., Nene, V., Nkonge, C., et al., 1998. An enzyme-linked immunosorbent assay for detection of *Theileria parva* antibodies in cattle using a recombinant polymorphic immunodominant molecule. *Parasitol. Res.* 84, 408–416.
- Katende, J.M., Goddeeris, B.M., Morzaria, S.P., Nkonge, C.G., Musoke, A.J., 1990. Identification of a *Theileria mutans*-specific antigen for use in an antibody and antigen detection ELISA. *Parasite Immunol.* 12, 419–433.
- Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780.
- Kawazu, S., Sugimoto, C., Kamio, T., Fujisaki, K., 1992. Antigenic differences between Japanese *Theileria sergenti* and other benign *Theileria* species of cattle from Australia (*T. buffeli*) and Britain (*T. orientalis*). *Parasitol. Res.* 78, 130–135.
- Kawazu, S., Kamio, T., Sekizaki, T., Fujisaki, K., 1995. *Theileria sergenti* and *T. buffeli*: polymerase chain reaction-based marker system for differentiating the parasite species from infected cattle blood and infected tick salivary gland. *Exp. Parasitol.* 81, 430–435.
- Kiara, H., Jennings, A., Bronsvort, B.M., Handel, I.G., Mwangi, S.T., Mbole-Kariuki, M., et al., 2014. A longitudinal assessment of the serological response to *Theileria parva* and other tick-borne parasites from birth to one year in a cohort of indigenous calves in western Kenya. *Parasitology* 141, 1289–1298.
- Kim, C.M., Blanco, L.B., Alhassan, A., Iseki, H., Yokoyama, N., Xuan, X., et al., 2008. Diagnostic real-time PCR assay for the quantitative detection of *Theileria equi* from equine blood samples. *Vet. Parasitol.* 151, 158–163.
- Kirvar, E., Ilhan, T., Katzer, F., Hooshmand-Rad, P., Zweggarth, E., Gerstenberg, C., et al., 2000. Detection of *Theileria annulata* in cattle and vector ticks by PCR using the Tams1 gene sequences. *Parasitology* 120, 245–254.

- Knowles, D.P., Lowell, J.R., Kappmeyer, S., Stiller, D., Hennager, S.G., Perryman, L.E., 1992. Antibody to a recombinant merozoite protein epitope identifies horses infected with *Babesia equi*. *J. Clin. Microbiol.* 30, 3122–3126.
- Koch, R., 1898. Reiseberichte über Rinderpest, Bubonenpest in Indien und Africa, Tsetse und Surrakrankhet, Texasfieber, Tropische Malaria, Schwarzwasserfieber. Springer, Berlin.
- Kubota, S., Sugimoto, C., Kakuda, T., Onuma, M., 1996. Analysis of immunodominant piroplasm surface antigen alleles in mixed populations of *Theileria sergenti* and *T. buffeli*. *Int. J. Parasitol.* 26, 741–747.
- Lack, J.B., Reichard, M.V., Van Den Busche, R.A., 2012. Phylogeny and evolution of the Piroplasmida as inferred from 18S rRNA sequences. *Int. J. Parasitol.* 42, 353–363.
- Latif, A.A., Hove, T., Kanhai, G.K., Masaka, S., 2002. Buffalo-associated *Theileria parva*: the risk to cattle of buffalo translocation into the Highveld of Zimbabwe. *Ann. N. Y. Acad. Sci.* 969, 275–279.
- Lawrence, D.A., 1935. Report of the Director of Veterinary Research, Southern Rhodesia, for the Year 1934. Rhodesian Printing and Publishing Co, Salisbury, Rhodesia.
- Lawrence, J.A., 1979. The differential diagnosis of the bovine Theilerias of Southern Africa. *J. S. Afr. Vet. Assoc.* 50, 311–313.
- Lawrence, J.A., Norval, R.A.L., Uilenberg, G., 1983. *Rhipicephalus zambeziensis* as a vector of bovine Theileriae. *Trop. Anim. Health Prod.* 15, 39–42.
- Lee, J.Y., Ryan, U.M., Jefferies, R., McInnes, L.M., Forshaw, D., Friend, J.A., et al., 2009. *Theileria gilberti* n. sp. (Apicomplexa: Theileriidae) in the Gilbert's potoroo (*Potorus gilbertii*). *J. Eukaryot. Microbiol.* 56, 290–295.
- Leemans, I., Hooshmand-Rad, P., Uggla, A., 1997. The indirect fluorescent antibody test based on schizont antigen for study of the sheep parasite *Theileria lestoquardi*. *Vet. Parasitol.* 69, 9–18.
- Leemans, I., Brown, D., Hooshmand-Rad, P., Kirvar, E., Uggla, A., 1999. Infectivity and cross-immunity studies of *Theileria lestoquardi* and *Theileria annulata* in sheep and cattle: I. *In vivo* responses. *Vet. Parasitol.* 82, 179–192.
- Li, Y., Chen, Z., Liu, Z., Liu, J., Yang, J., Li, Q., et al., 2014. Molecular identification of *Theileria* parasites of northwestern Chinese Cervidae. *Parasit. Vectors* 7, 225.
- Lichtenheld, G., 1910. Preliminary communication on the fixing of complement in horse-sickness and East Coast fever. *Z. Immun. Forsch.* 8, 232–238.
- Liu, A., Guan, G., Liu, Z., Liu, J., Leblanc, N., Li, Y., et al., 2010b. Detecting and differentiating *Theileria sergenti* and *Theileria sinensis* in cattle and yaks by PCR based on major piroplasm surface protein (MPSP). *Exp. Parasitol.* 126, 476–481.
- Liu, A., Guan, G., Du, P., Liu, Z., Gou, H., Liu, J., et al., 2012. Loop-mediated isothermal amplification (LAMP) assays for the detection of *Theileria annulata* infection in China targeting the 18S rRNA and ITS sequences. *Exp. Parasitol.* 131, 125–129.
- Liu, A., Guan, G., Du, P., Gou, H., Zhang, J., Liu, Z., et al., 2013. Rapid identification and differentiation of *Theileria sergenti* and *Theileria sinensis* using a loop-mediated isothermal amplification (LAMP) assay. *Vet. Parasitol.* 191, 15–22.
- Liu, A.H., Guan, G.Q., Liu, J.L., Liu, Z.J., Leblanc, N., Li, Y.Q., et al., 2011. Polymorphism analysis of Chinese *Theileria sergenti* using allele-specific polymerase chain reaction of the major piroplasm surface protein gene. *J. Parasitol.* 97, 116–121.
- Liu, Z., Hou, J., Bakheit, M.A., Salih, D.A., Luo, J., Yin, H., et al., 2008. Development of loop-mediated isothermal amplification (LAMP) assay for rapid diagnosis of ovine theileriosis in China. *Parasitol. Res.* 103, 1407–1412.
- Liu, Z., Wang, Z., Yin, H., Luo, J., Zhang, B., Kullmann, B., et al., 2010a. Identification of *Theileria uilenbergi* immunodominant protein for development of an indirect ELISA for diagnosis of ovine theileriosis. *Int. J. Parasitol.* 40, 591–598.
- Liu, Z., Li, Y., Salih, D.E., Luo, J., Ahmed, J.S., Seitzer, U., et al., 2014. Validation of a recombinant protein indirect ELISA for the detection of specific antibodies against *Theileria uilenbergi* and *Theileria luwenshuni* in small ruminants. *Vet. Parasitol.* 204, 139–145.
- Mans, B.J., Pienaar, R., Latif, A.A., Potgieter, F.T., 2011a. Diversity in the 18S SSU rRNA V4 hyper-variable region of *Theileria* in bovines and African buffalo (*Syncerus caffer*) from southern Africa. *Parasitology* 138, 766–779.
- Mans, B.J., Pienaar, R., Potgieter, F.T., Latif, A.A., 2011b. *Theileria parva*, *T. sp.* (buffalo) and *T. sp.* (bougasvlei) 18S variants. *Vet. Parasitol.* 182, 382–383.
- Martin, H., Brocklesby, D.W., 1960. A new parasite of Eland. *Vet. Rec.* 72, 331–332.
- Matjila, P.T., Leisewitz, A.L., Oosthuizen, M.C., Jongejan, F., Penzhorn, B.L., 2008. Detection of a *Theileria* species in dogs in South Africa. *Vet. Parasitol.* 157, 34–40.
- Mbizeni, S., Potgieter, F.T., Troskie, C., Mans, B.J., Penzhorn, B.L., Latif, A.A., 2013. Field and laboratory studies on Corridor disease (*Theileria parva* infection) in cattle population at the livestock/game interface of uPhongolo-Mkuze area, South Africa. *Ticks Tick Borne Dis.* 4, 227–234.
- McKeever, D.J., 2009. Bovine immunity – a driver for diversity in *Theileria* parasites? *Trends Parasitol.* 25, 269–276.
- Miranda, J., Bakheit, M.A., Liu, Z., Yin, H., Mu, Y., Guo, S., et al., 2006. Development of a recombinant indirect ELISA for the diagnosis of *Theileria sp.* (China) infection in small ruminants. *Parasitol. Res.* 98, 561–567.
- Morrison, D.A., 2009. Evolution of the Apicomplexa: where are we now? *Trends Parasitol.* 25, 375–382.
- Morzaria, S.P., Katende, J., Musoke, A., Nene, V., Skilton, R., Bishop, R., 1999. Development of sero-diagnostic and molecular tools for the control of important tick-borne pathogens of cattle in Africa. *Parassitologia* 41, S73–S80.
- Nagore, D., García-Sanmartín, J., García-Pérez, A.L., Juste, R.A., Hurtado, A., 2004. Identification, genetic diversity and prevalence of *Theileria* and *Babesia* species in a sheep population from Northern Spain. *Int. J. Parasitol.* 34, 1059–1067.
- Najm, N.A., Meyer-Kayser, E., Hoffmann, L., Herb, I., Fensterer, V., Pfister, K., et al., 2014. A molecular survey of *Babesia* spp. and *Theileria* spp. in red foxes (*Vulpes vulpes*) and their ticks from Thuringia, Germany. *Ticks Tick Borne Dis.* 5, 386–391.
- Ngumi, P.N., Lesan, A.C., Williamson, S.M., Awich, J.R., 1994. Isolation and preliminary characterisation of a previously unidentified *Theileria* parasite of cattle in Kenya. *Res. Vet. Sci.* 57, 1–9.
- Nijhof, A.M., Penzhorn, B.L., Lynen, G., Molle, J.O., Morkel, P., Bekker, C.P., et al., 2003. *Babesia bicornis* sp. nov. and *Theileria bicornis* sp. nov.: tick-borne parasites associated with mortality in the black rhinoceros (*Diceros bicornis*). *J. Clin. Microbiol.* 41, 2249–2254.
- Nijhof, A.M., Pillay, V., Steyl, J., Prozesky, L., Stoltz, W.H., Lawrence, J.A., et al., 2005. Molecular characterization of *Theileria* species associated with mortality in four species of African antelopes. *J. Clin. Microbiol.* 43, 5907–5911.
- Niu, Q., Luo, J., Guan, G., Ma, M., Liu, Z., Liu, A., et al., 2009. Detection and differentiation of ovine *Theileria* and *Babesia* by reverse line blotting in China. *Parasitol. Res.* 104, 1417–1423.
- Norval, R.A., Lawrence, J.A., Young, A.S., Perry, B.D., Dolan, T.T., Scott, J., 1991. *Theileria parva*: influence of vector, parasite and host relationships on the epidemiology of theileriosis in southern Africa. *Parasitology* 102, 347–356.
- Norval, R.A., Perry, B.D., Young, A.S., 1992. The Epidemiology of Theileriosis in Africa. Academic Press Inc, London, UK, pp. 136–154.
- Odongo, D.O., Sunter, J.D., Kiara, H.K., Skilton, R.A., Bishop, R.P., 2010. A nested PCR assay exhibits enhanced sensitivity for detection of *Theileria parva* infections in bovine blood samples from carrier animals. *Parasitol. Res.* 106, 357–365.
- Ogunremi, O., Georgiadis, M.P., Halbert, G., Benjamin, J., Pfister, K., Lopez-Rebollar, L., 2007. Validation of the indirect fluorescent antibody and the complement fixation tests for the diagnosis of *Theileria equi*. *Vet. Parasitol.* 148, 102–108.
- OIE, 2014. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals, vol. 1–2, seventh ed. Office International Des Epizooties, Paris.
- Oosthuizen, M.C., Allsopp, B.A., Troskie, M., Collins, N.E., Penzhorn, B.L., 2009. Identification of novel *Babesia* and *Theileria* species in South African giraffe (*Giraffa camelopardalis*, Linnaeus, 1758) and roan antelope (*Hippotragus equinus*, Desmarest 1804). *Vet. Parasitol.* 163, 39–46.
- Ota, N., Mizuno, D., Kuboki, N., Igarashi, I., Nakamura, Y., Yamashina, H., et al., 2010. Epidemiological survey of *Theileria orientalis* infection in grazing cattle in the eastern part of Hokkaido, Japan. *J. Vet. Med. Sci.* 71, 937–944.
- Otiende, M.Y., Kivata, M.W., Jowers, M.J., Makumi, J.N., Runo, S., Obanda, V., et al., 2014. Three novel haplotypes of *Theileria bicornis* in black and white Rhinoceros in Kenya. *Transbound. Emerg. Dis. doi:10.1111/tbed.12242*.
- Oura, C.A., Bishop, R., Wampande, E.M., Lubega, G.W., Tait, A., 2004. The persistence of component *Theileria parva* stocks in cattle immunized with the 'Muguga cocktail' live vaccine against East Coast fever in Uganda. *Parasitology* 129, 27–42.
- Oura, C.A., Tait, A., Asiimwe, B., Lubega, G.W., Weir, W., 2011. *Theileria parva* genetic diversity and haemoparasite prevalence in cattle and wildlife in and around Lake Mburo National Park in Uganda. *Parasitol. Res.* 108, 1365–1374.
- Pain, A., Renauld, H., Berriman, M., Murphy, L., Yeats, C.A., Weir, W., et al., 2005. Genome of the host-cell transforming parasite *Theileria annulata* compared with *T. parva*. *Science* 309, 131–133.
- Papadopoulos, B., Perić, N.M., Uilenberg, G., 1996. Piroplasmids of domestic animals in the Macedonia region of Greece. 1. Serological cross-reactions. *Vet. Parasitol.* 63, 41–56.
- Paparin, A., Ryan, U.M., Warren, K., McInnes, L.M., de Tores, P., Irwin, P.J., 2012. Identification of novel *Babesia* and *Theileria* genotypes in the endangered marsupials, the woylie (*Bettongia penicillata ogilbyi*) and boodie (*Bettongia lesueur*). *Exp. Parasitol.* 131, 25–30.
- Papli, N.E., Landt, O., Fleischer, C., Koekemoer, J.O., Mans, B.J., Pienaar, R., et al., 2011. Evaluation of a TaqMan real-time PCR for the detection of *Theileria parva* in buffalo and cattle. *Vet. Parasitol.* 175, 356–359.
- Peeling, R.W., Smith, P.G., Bossuyt, P.M.M., 2010. A guide for diagnostic evaluations. *Nat. Rev. Microbiol.* 8, S2–S6.
- Perera, P.K., Gasser, R.B., Firestone, S.M., Smith, L., Roeber, F., Jabbar, A., 2014. Semi-quantitative multiplexed-tandem PCR for the detection and differentiation of four *Theileria orientalis* genotypes in cattle. *J. Clin. Microbiol. doi:10.1128/JCM.02536-14*.
- Perry, B.D., Young, A.S., 1993. The naming game: the changing fortunes of East Coast fever and *Theileria parva*. *Vet. Rec.* 133, 613–616.
- Pfaffl, M.W., 2004. Quantification strategies in real-time PCR. In: Bustin, S.A. (Ed.), A-Z of Quantitative PCR. International University Line (IUL), La Jolla, CA, USA, pp. 87–112.
- Pienaar, R., Potgieter, F.T., Latif, A.A., Thekisoe, O.M.M., Mans, B.J., 2011a. Mixed *Theileria* infections in free-ranging buffalo herds: implications for diagnosing *Theileria parva* infections in Cape buffalo (*Syncerus caffer*). *Parasitology* 138, 884–895.
- Pienaar, R., Potgieter, F.T., Latif, A.A., Thekisoe, O.M.M., Mans, B.J., 2011b. The HybridII assay: a sensitive and specific real-time hybridization assay for the diagnosis of *Theileria parva* infection in Cape buffalo (*Syncerus caffer*) and cattle. *Parasitology* 138, 1935–1944.
- Pienaar, R., Latif, A.A., Thekisoe, O.M.M., Mans, B.J., 2013. Protein gene candidates for the qualitative molecular detection of *Theileria parva* in African buffalo (*Syncerus caffer*) using the real-time SYBR green polymerase chain reaction. Proceedings of the 1st Annual Conference on Advances in Veterinary Science Research, Singapore.
- Pienaar, R., Potgieter, F.T., Latif, A.A., Thekisoe, O.M.M., Mans, B.J., 2014. Geographic distribution of *Theileria sp.* (buffalo) and *Theileria sp.* (bougasvlei) in Cape buffalo (*Syncerus caffer*) in southern Africa: implications for speciation. *Parasitology* 141, 411–424.
- Rajendran, C., Ray, D.D., 2014. Diagnosis of tropical bovine theileriosis by ELISA with recombinant merozoite surface protein of *Theileria annulata* (Tams1). *J. Parasit. Dis.* 38, 41–45.

- Renneker, S., Kullmann, B., Gerber, S., Dobschanski, J., Bakheit, M.A., Geysen, D., et al., 2008. Development of a competitive ELISA for detection of *Theileria annulata* infection. *Transbound. Emerg. Dis.* 55, 249–256.
- Ros-García, A., Juste, R.A., Hurtado, A., 2012a. A highly sensitive DNA bead-based suspension array for the detection and species identification of bovine piroplasms. *Int. J. Parasitol.* 42, 207–214.
- Ros-García, A., Nicolás, A., García-Pérez, A.L., Juste, R.A., Hurtado, A., 2012b. Development and evaluation of a real-time PCR assay for the quantitative detection of *Theileria annulata* in cattle. *Parasit. Vectors* 5, 171.
- Ros-García, A., Barandika, J.F., García-Pérez, A.L., Juste, R.A., Hurtado, A., 2013. Assessment of exposure to piroplasms in sheep grazing in communal mountain pastures by using a multiplex DNA bead-based suspension array. *Parasit. Vectors* 6, 277.
- Salih, D.A., Liu, Z., Bakheit, M.A., Ali, A.M., El Hussein, A.M., Unger, H., et al., 2008. Development and evaluation of a loop-mediated isothermal amplification method for diagnosis of tropical theileriosis. *Transbound. Emerg. Dis.* 55, 238–243.
- Salih, D.A., Ali, A.M., Liu, Z., Bakheit, M.A., Taha, K.M., El Imam, A.H., et al., 2012. Development of a loop-mediated isothermal amplification method for detection of *Theileria lestoquardi*. *Parasitol. Res.* 110, 533–538.
- Salih, D.E., Ahmed, J.S., Bakheit, M.A., Ali, E.B., El Hussein, A.M., Hassan, S.M., et al., 2005. Validation of the indirect TaSP enzyme-linked immunosorbent assay for diagnosis of *Theileria annulata* infection in cattle. *Parasitol. Res.* 97, 302–308.
- Salim, B., Bakheit, M., Kamau, J., Nakamura, I., Sugimoto, C., 2010. Nucleotide sequence heterogeneity in the small subunit ribosomal RNA gene within *Theileria equi* from horses in Sudan. *Parasitol. Res.* 106, 493–498.
- Salim, B., Bakheit, M.A., Sugimoto, C., 2013. Rapid detection and identification of *Theileria equi* and *Babesia caballi* by high-resolution melting (HRM) analysis. *Parasitol. Res.* 112, 3883–3886.
- Salim, B.O., Hassan, S.M., Bakheit, M.A., Alhassan, A., Igarashi, I., Karanis, P., et al., 2008. Diagnosis of *Babesia caballi* and *Theileria equi* infections in horses in Sudan using ELISA and PCR. *Parasitol. Res.* 103, 1145–1150.
- Sarataphan, N., Nilwarangkoon, S., Tananyutthawongse, C., Kakuda, T., Onuma, M., Chansiri, K., 1999. Genetic diversity of major piroplasm surface protein genes and their allelic variants of *Theileria* parasites in Thai cattle. *J. Vet. Med. Sci.* 61, 991–994.
- Schaeffler, W.F., 1963. Serologic tests for *Theileria cervi* in white-tailed deer and for other species of *Theileria* in cattle and sheep. *Am. J. Vet. Res.* 24, 784–791.
- Schettler, T.P., Arts, G., Niessen, R., Schaap, D., 2010. Development of a new score to estimate clinical East Coast Fever in experimentally infected cattle. *Vet. Parasitol.* 167, 255–259.
- Schindler, R., Mehlitz, D., 1969. Serologic studies on *Theileria mutans* infection in cattle. *Z. Tropenmed. Parasitol.* 20, 459–473.
- Schindler, R., Wokatsch, R., 1965. Versuche zur differenzierung der Theilerien speziez des rindes durch serologische untersuchungen. *Tropenmed. Parasitol.* 16, 85–87.
- Schnittger, L., Yin, H., Gubbels, M.J., Beyer, D., Niemann, S., Jongejan, F., et al., 2003. Phylogeny of sheep and goat *Theileria* and *Babesia* parasites. *Parasitol. Res.* 91, 398–406.
- Schnittger, L., Yin, H., Qi, B., Gubbels, M.J., Beyer, D., Niemann, S., et al., 2004. Simultaneous detection and differentiation of *Theileria* and *Babesia* parasites infecting small ruminants by reverse line blotting. *Parasitol. Res.* 92, 189–196.
- Seitzer, U., Bakheit, M.A., Salih, D.E., Ali, A., Haller, D., Yin, H., et al., 2007. From molecule to diagnostic tool: *Theileria annulata* surface protein TaSP. *Parasitol. Res.* 101, S217–S223.
- Shayan, P., Biermann, R., Schein, E., Gerdes, J., Ahmed, J.S., 1998. Detection and differentiation of *Theileria annulata* and *Theileria parva* using macroschizont-derived DNA probes. *Ann. N. Y. Acad. Sci.* 849, 88–95.
- Shimizu, S., Suzuki, K., Nakamura, K., Kadota, K., Fujisaki, K., Ito, S., et al., 1988. Isolation of *Theileria sergenti* piroplasms from infected erythrocytes and development of an enzyme-linked immunosorbent assay for serodiagnosis of *T. sergenti* infections. *Res. Vet. Sci.* 45, 206–212.
- Sibeko, K.P., Oosthuizen, M.C., Collins, N.E., Geysen, D., Rambritch, N.E., Latif, A.A., et al., 2008. Development and evaluation of a real-time polymerase chain reaction test for the detection of *Theileria parva* infections in Cape buffalo (*Syncerus caffer*) and cattle. *Vet. Parasitol.* 155, 37–48.
- Sibeko, K.P., Geysen, D., Oosthuizen, M.C., Matthee, C.A., Troskie, M., Potgieter, F.T., et al., 2010. Four p67 alleles identified in South African *Theileria parva* field samples. *Vet. Parasitol.* 167, 244–254.
- Sibeko, K.P., Collins, N.E., Oosthuizen, M.C., Troskie, M., Potgieter, F.T., Coetzer, J.A., et al., 2011. Analyses of genes encoding *Theileria parva* p104 and polymorphic immunodominant molecule (PIM) reveal evidence of the presence of cattle-type alleles in the South African *T. parva* population. *Vet. Parasitol.* 181, 120–130.
- Sivakumar, T., Hayashida, K., Sugimoto, C., Yokoyama, N., 2014. Evolution and genetic diversity of *Theileria*. *Infect. Genet. Evol.* 27, 250–263.
- Skilton, R.A., Bishop, R.P., Katende, J.M., Mwaura, S., Morzaria, S.P., 2002. The persistence of *Theileria parva* infection in cattle immunized using two stocks which differ in their ability to induce a carrier state: analysis using a novel blood spot PCR assay. *Parasitology* 124, 265–276.
- Stewart, N.P., Uilenberg, G., de Vos, A.J., 1996. Review of the Australian species of *Theileria*, with special reference to *Theileria buffeli* of cattle. *Trop. Anim. Health Prod.* 28, 81–90.
- Steyl, J.C.A., Prozesky, L., Stoltz, W.H., Lawrence, J.A., 2012. Theileriosis (Cytauxzoonosis) in Roan antelope (*Hippotragus equinus*): field exposure to infection and identification of potential vectors. *Onderstepoort J. Vet. Res.* 79, 367.
- Swai, E.S., Karimuribo, E.D., Kambarage, D.M., Moshy, W.E., Mbise, A.N., 2007. A comparison of seroprevalence and risk factors for *Theileria parva* and *T. mutans* in smallholder dairy cattle in the Tanga and Iringa regions of Tanzania. *Vet. J.* 174, 390–396.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Tanaka, M., Onoe, S., Matsuba, T., Katayama, S., Yamanaka, M., Yonemichi, H., et al., 1993. Detection of *Theileria sergenti* infection in cattle by polymerase chain reaction amplification of parasite-specific DNA. *J. Clin. Microbiol.* 31, 2565–2569.
- Theiler, A., 1904. Rhodesian tick fever. *Transvaal Agr. J.* 2, 421–438.
- Theiler, A., 1906. *Piroplasma mutans* (n. spec) of South African cattle. *J. Comp. Path. Ther.* 19, 292–300.
- Thekisoe, O.M., Rambritch, N.E., Nakao, R., Bazie, R.S., Mbati, P., Namangala, B., et al., 2010. Loop-mediated isothermal amplification (LAMP) assays for detection of *Theileria parva* infections targeting the PIM and p150 genes. *Int. J. Parasitol.* 40, 55–61.
- Thompson, B.E., Latif, A.A., Oosthuizen, M.C., Troskie, M., Penzhorn, B.L., 2008. Occurrence of *Theileria parva* infection in cattle on a farm in the Ladysmith district, KwaZulu-Natal, South Africa. *J. S. Afr. Vet. Assoc.* 79, 31–35.
- Tian, Z., Liu, G., Yin, H., Luo, J., Guan, G., Luo, J., et al., 2013. Discrimination between ovine *Babesia* and *Theileria* species in China based on the ribosomal protein S8 (RPS8) gene. *Vet. Parasitol.* 197, 354–359.
- Tian, Z., Liu, G., Yin, H., Xie, J., Wang, S., Yuan, X., et al., 2014. First report on the occurrence of *Theileria* sp. OT3 in China. *Parasitol. Int.* 63, 403–407.
- Tibayrenc, M., 2006. The species concept in parasites and other pathogens: a pragmatic approach? *Trends Parasitol.* 22, 66–70.
- Tibayrenc, M., Ayala, F.J., 2002. The clonal theory of parasitic protozoa: 12 years on. *Trends Parasitol.* 18, 405–410.
- Ueti, M.W., Mealey, R.H., Kappmeyer, L.S., White, S.N., Kumpula-McWhirter, N., Pelzel, A.M., et al., 2012. Re-emergence of the apicomplexan *Theileria equi* in the United States: elimination of persistent infection and transmission risk. *PLoS ONE* 7, e44713.
- Uilenberg, G., 1981. Theilerial species of domestic livestock. In: Irving, A.D., Cunningham, M.P., Young, A.S. (Eds.), *Advances in the Control of Theileriosis*. Martinus Nijhoff Publishers, The Hague, Boston, London, pp. 4–37.
- Uilenberg, G., 2006. *Babesia* – a historical perspective. *Vet. Parasitol.* 138, 3–10.
- Uilenberg, G., Schreuder, B.E.C., 1976. Studies on Theileriidae (Sporozoa) in Tanzania. I. Tick transmission of *Haematotaxenus veliferus*. *Tropenmed. Parasitol.* 27, 106–111.
- Uilenberg, G., Perié, N.M., Spanjer, A.A., Franssen, F.F., 1985. *Theileria orientalis*, a cosmopolitan blood parasite of cattle: demonstration of the schizont stage. *Res. Vet. Sci.* 38, 352–360.
- Vilstrup, J.T., Seguin-Orlando, A., Stiller, M., Ginolhac, A., Raghavan, M., Nielsen, S.C., et al., 2013. Mitochondrial phylogenomics of modern and ancient equids. *PLoS ONE* 8, e55950.
- Wang, L.X., He, L., Fang, R., Song, Q.Q., Tu, P., Jenkins, A., et al., 2010a. Loop-mediated isothermal amplification (LAMP) assay for detection of *Theileria sergenti* infection targeting the p33 gene. *Vet. Parasitol.* 171, 159–162.
- Wang, L.X., Zhao, J.H., He, L., Liu, Q., Zhou, D.N., Zhou, Y.Q., et al., 2010b. An indirect ELISA for detection of *Theileria sergenti* antibodies in water buffalo using a recombinant major piroplasm surface protein. *Vet. Parasitol.* 170, 323–326.
- Xie, J., Liu, G., Tian, Z., Luo, J., 2013. Development of loop-mediated isothermal amplification (LAMP) for detection of *Theileria equi*. *Acta Trop.* 127, 245–250.
- Yang, J.F., Li, Y.Q., Liu, Z.J., Liu, J.L., Guan, G.Q., Chen, Z., et al., 2014b. Molecular evidence for piroplasms in wild Reeves' muntjac (*Muntiacus reevesi*) in China. *Parasitol. Int.* 63, 713–716.
- Yang, Y., Mao, Y., Kelly, P., Yang, Z., Luan, L., Zhang, J., et al., 2014a. A pan-*Theileria* FRET-qPCR survey for *Theileria* spp. in ruminants from nine provinces of China. *Parasit. Vectors* 7, 413.
- Yin, H., Liu, Z., Guan, G., Liu, A., Ma, M., Ren, Q., et al., 2008. Detection and differentiation of *Theileria luwenshuni* and *T. uilenbergi* infection in small ruminants by PCR. *Transbound. Emerg. Dis.* 55, 233–237.
- Yisaschar-Mekuzas, Y., Jaffe, C.L., Pastor, J., Cardoso, L., Baneth, G., 2013. Identification of *Babesia* species infecting dogs using reverse line blot hybridization for six canine piroplasms, and evaluation of co-infection by other vector-borne pathogens. *Vet. Parasitol.* 191, 367–373.
- Young, A.S., Grootenhuys, J.G., Kimber, C.D., Kanhai, G.K., Stagg, D.A., 1977. Isolation of a *Theileria* species from Eland (*Taurotragus oryx*) infective for cattle. *Tropenmed. Parasitol.* 27, 185–194.
- Young, A.S., Purnell, R.E., Payne, R.C., Brown, C.G., Kanhai, G.K., 1978. Studies on the transmission and course of infection of a Kenyan strain of *Theileria mutans*. *Parasitology* 76, 99–115.
- Young, A.S., Leitch, B.L., Newson, R.M., Cunningham, M.P., 1986. Maintenance of *Theileria parva parva* infection in an endemic area of Kenya. *Parasitology* 93, 9–16.
- Yusufmia, S.B.A.S., Collins, N.E., Nkuna, R., Troskie, M., Van den Bossche, P., Penzhorn, B.L., 2010. Occurrence of *Theileria parva* and other haemoprotozoa in cattle at the edge of Hluhluwe-iMfolozi Park, KwaZulu-Natal, South Africa. *J. S. Afr. Vet. Assoc.* 81, 45–49.
- Zaemi, M., Haddadzadeh, H., Khazraiiina, P., Kazemi, B., Bandehpour, M., 2011. Identification of different *Theileria* species (*Theileria lestoquardi*, *Theileria ovis*, and *Theileria annulata*) in naturally infected sheep using nested PCR-RFLP. *Parasitol. Res.* 108, 837–843.
- Zahler, M., Rinder, H., Schein, E., Gothe, R., 2000. Detection of a new pathogenic *Babesia microti*-like species in dogs. *Vet. Parasitol.* 89, 241–248.

- Zhang, X., Liu, Z., Yang, J., Chen, Z., Guan, G., Ren, Q., et al., 2014. Multiplex PCR for diagnosis of *Theileria uilenbergi*, *Theileria luwenshuni*, and *Theileria ovis* in small ruminants. *Parasitol. Res.* 113, 527–531.
- Zobba, R., Ardu, M., Niccolini, S., Chessa, B., Manna, L., Cocco, R., et al., 2008. Clinical and laboratory findings in equine piroplasmiasis. *J. Equine Vet. Sci.* 28, 301–308.
- Zweygarth, E., Just, M.C., De Waal, D.T., 1997. In vitro cultivation of *Babesia equi*: detection of carrier animals and isolation of parasites. *Onderstepoort J. Vet. Res.* 64, 51–56.
- Zweygarth, E., Benade, J., Steyl, J., Prozesky, L., Koekemoer, O., Josemans, A.I., 2009a. In vitro cultivation of a *Theileria* species from a roan antelope (*Hippotragus equinus*). *Parasitol. Res.* 105, 1755–1757.
- Zweygarth, E., Koekemoer, O., Josemans, A.I., Rambritch, N., Pienaar, R., Putterill, J., et al., 2009b. *Theileria*-infected cell line from an African buffalo (*Syncerus caffer*). *Parasitol. Res.* 105, 579–581.