

Research Note

# Susceptibility of South African native conifers to the pitch canker pathogen, *Fusarium circinatum*

B. Porter, M.J. Wingfield, T.A. Coutinho \*

Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa

Received 19 August 2008; received in revised form 6 November 2008; accepted 11 November 2008

## Abstract

*Fusarium circinatum* is an economically important pathogen of *Pinus* species, which also occurs on Douglas fir (*Pseudotsuga menzeisii*). It causes pitch canker of mature trees and root and collar rot of seedlings and cuttings. In 2007, pitch canker was observed on *Pinus radiata* in the Western Cape and this was the first outbreak of this disease in South Africa. The Cape flora in this area includes native coniferous species of *Podocarpus* and *Widdringtonia*. In this study, seedlings of *P. latifolia*, *P. elongatus*, *P. henkelii*, *W. schwartzii*, *W. cederbergensis* and *W. nodiflora* were inoculated with a virulent isolate of *F. circinatum*, to assess their susceptibility. Six weeks after inoculation, there was no lesion development in any of the species inoculated. *Pinus patula* seedlings used as a positive control were severely affected. Results of this study show that species of *Podocarpus* and *Widdringtonia*, native to the Western Cape, are not threatened by the pitch canker fungus.

© 2008 SAAB. Published by Elsevier B.V. All rights reserved.

**Keywords:** Conifer; *Fusarium circinatum*; Pitch canker; *Podocarpus*; *Widdringtonia*

*Fusarium circinatum* (Nirenberg and O'Donnell) is a pathogen which has a serious economic impact on pine production in South Africa (Wingfield et al., 2002). The South African forestry industry contributes significantly to the economy through the production of solid timber and pulp products (Anonymous, 2005), and is based on exotic plantations of pines and eucalypts. The *Pinus* species most extensively planted are *Pinus patula* (Schl. et Cham.) and *P. elliottii* (Chapm.). *P. radiata* (D. Don.) is only grown in the Western Cape.

*Fusarium circinatum* was first reported from a pine production nursery in Mpumalanga (Viljoen and Wingfield, 1994) and is now endemic in nurseries throughout the country, where it causes root and collar rot of both seedlings and cuttings. In 2007, large, resinous cankers appeared on mature *P. radiata* trees in the Western Cape and this was the first report of pitch canker in South Africa (Coutinho et al., 2007). These plantations are situated in close proximity to the unique Cape flora, which includes native conifers such as *Podocarpus* and

*Widdringtonia*. Because *F. circinatum* is an exotic pathogen in South Africa, is known to infect numerous *Pinus* species in America and Europe (Storer et al., 1997; Hodge and Dvorak, 2000; Enebak and Stanosz, 2003), and can cause disease on a non-pine species, *Pseudotsuga menziesii* (Mirb.) (Douglas-fir) (Gordon et al., 2006), concern has been raised that it might threaten native conifers.

Inoculation trials with 20 seedlings each of *Podocarpus latifolia* [(Thunb.) R.Br.: Mirb.], *P. elongatus* [(Ait.) L'Herit.: Pers.], *P. henkelii* (Stapf. Ex Dallim. & Jacks.), *Widdringtonia schwartzii* (Endl.), *W. cederbergensis* (Marsh) and *W. nodiflora* [(L.) Powrie], were conducted using a known virulent strain of *F. circinatum* (FCC 3579). Inoculum was produced by culturing this isolate on half strength potato dextrose agar (20 g Potato dextrose agar powder [Merck, Biolab Diagnostics (Pty) Ltd.], 5 g Agarose bacteriological powder [Merck, Biolab Diagnostics (Pty) Ltd.], 1 l distilled water) for 15 days. Spores were washed from the cultures with 15% glycerol, excess mycelium was filtered through sterile cheese cloth, and the concentration adjusted to  $5 \times 10^4$  spores/ml using a haemocytometer. The seedlings were wounded by needle puncture of the main stem [Promex hypodermic syringe needles, 21 G, 0.80 × 40 mm], and

\* Corresponding author.

E-mail address: [teresa.coutinho@fabi.up.ac.za](mailto:teresa.coutinho@fabi.up.ac.za) (T.A. Coutinho).




Fig. 1. A. *Podocarpus henkelii* 12 weeks after inoculation, arrows denote inoculation points. Seedling '1' negative control, '2' and '3' inoculated with *Fusarium circinatum*. B. *Widdringtonia nodiflora* seedlings 12 weeks after inoculation, arrows denote inoculation points. Seedling '1' negative control, '2' inoculated with *Fusarium circinatum*. C. Local discoloration around inoculation point on *Podocarpus latifolia* seedling (arrow denotes inoculation point). D. Positive controls, *P. patula* seedlings 3 weeks after inoculation. Bark has been removed to expose lesion development. All *P. patula* seedlings dead at 12 weeks.

inoculated with a 10  $\mu$ l drop of spore suspension. This technique was similar to that used by Kuhlman (1987), McCain et al. (1987) and Blakeslee and Rockwood (1999). *Podocarpus* seedlings were purchased from indigenous tree nurseries and *Widdringtonia* seedlings were grown from seed purchased from the Kirstenbosch Botanical Gardens ([www.sanbi.org/frames/kirstfram.htm](http://www.sanbi.org/frames/kirstfram.htm)). Ten seedlings of each species were inoculated with sterile 15% glycerol as a negative control, and 20 *Pinus patula* seedlings were inoculated as a positive control. At the time of inoculation, all plants were 10 months old, planted in 2.5l black plastic bags containing composted bark medium, and maintained at a constant temperature of 25 °C in a greenhouse. Seedlings were evaluated for disease development at 3, 6 and 12 weeks after inoculation. The experiment was repeated once.

None of the conifers inoculated showed lesion development (Fig. 1-A, B), although some seedlings developed minor callusing and discoloration localized to the bark above the inoculation point (Fig. 1-C). All *P. patula* seedlings had distinct lesion development at 3 weeks after inoculation (Fig. 1-D), and by 12 weeks all had died. Thus far all plant species, other than *Pinus* spp. and *P. menziesii*, that have been tested for susceptibility to *F. circinatum* have been found to be highly resistant (McCain et al., 1987; Dick and Simpson, 2004). The resistance of *Podocarpus* and *Widdringtonia* spp. shown in this study is reassuring as *W. cederbergensis* is under threat from

over-exploitation and fire (Manders, 1986). *W. cederbergensis* exists as small scattered populations in rocky outcrops of the Cederberg, and in some localities is incapable of maintaining the wild population (Manders and Botha, 1989). An additional threat by an aggressive, exotic pathogen could be detrimental.

#### Acknowledgements

We thank the Forestry and Agricultural Biotechnology Institute (FABI), the Centre for Tree Health Biotechnology (CTHB), the Tree Protection Co-operative Program (TPCP), and the National Research Foundation (NRF), for funding.

#### References

- Anonymous, 2005. State of the worlds forests. Food and Agriculture Organization of the United Nations. FAO, Rome, pp. 134–142.
- Blakeslee, G.M., Rockwood, D.L., 1999. Variation in resistance to pitch canker in slash and loblolly pines. In: Devey, M.E., Matheson, A.C., Gordon, T.R. (Eds.), Current and potential impacts of pitch canker in radiata pine. Proceedings of the IMPACT Monterey workshop, California, USA, 30 November to 3 December 1998. (CSIRO, Forestry and forest Products No. 112: Canberra), pp. 35–39.
- Coutinho, T.A., Steenkamp, E.T., Mongwaketsi, K., Wilmot, M., Wingfield, M.J., 2007. First outbreak of pitch canker in a South African pine plantation. Australasian Plant Pathology 36, 256–261.
- Dick, M.A., Simpson, J.A., 2004. *Fusarium circinatum* as a cause of damping-off: Exploring the host range. Unpublished data.

- Enebak, S.A., Stanosz, G.R., 2003. Responses of conifer species of the Great Lakes region of North America to inoculation with the pitch canker pathogen *Fusarium circinatum*. *Forest Pathology* 33, 333–338.
- Gordon, T.R., Kirkpatrick, S.C., Aegerter, B.J., Wood, D.L., Storer, A.J., 2006. Susceptibility of Douglas fir (*Pseudotsuga menziesii*) to pitch canker caused by *Gibberella circinata* (anamorph=*Fusarium circinatum*). *Plant Pathology* 55, 231–237.
- Hodge, G.R., Dvorak, W.S., 2000. Differential responses of central American and Mexican pine species and *Pinus radiata* to infection by the pitch canker fungus. *New Forests* 19, 241–258.
- Kuhlman, E.G., 1987. Effects of inoculation treatment with *F. moniliforme* var *subglutinans* on dieback of slash and loblolly pine seedlings. *Plant Disease* 71, 161–162.
- Manders, P.T., 1986. An assessment of the current status of the Clanwilliam cedar (*Widdringtonia cedarbergensis*) and the reasons for its decline. *South African Forestry Journal* 139, 48–53.
- Manders, P.T., Botha, S.A., 1989. A note on establishment of *Widdringtonia cedarbergensis* (Clanwilliam cedar). *Journal of Applied Ecology* 26, 571–574.
- McCain, A.H., Koehler, C.S., Tjosvold, S.A., 1987. Pitch canker threatens California pines. *California Agriculture* November–December, pp. 22–23.
- Storer, A.J., Gordon, T.R., Wood, D.L., Bonello, P., 1997. Pitch canker disease of pines: current and future impacts. *Journal of Forestry* 95, 21–26.
- Viljoen, A., Wingfield, M.J., 1994. First report of *Fusarium subglutinans* f. sp. *pini* on pine seedlings in South Africa. *Plant Disease* 78, 309–312.
- Wingfield, M.J., Coutinho, T.A., Roux, J., Wingfield, B.D., 2002. The future of exotic plantation forestry in the tropics and southern Hemisphere: lessons from pitch canker. *Southern African Forestry Journal* 195, 79–82.