

- xeroderma pigmentosum skin *in vitro*: a model to study hypersensitivity to UV light. *Photochem Photobiol* 81:19–24
- Epp N, Fürstenberger G, Müller K *et al.* (2007) 12R-lipoxygenase deficiency disrupts epidermal barrier function. *J Cell Biol* 177:173–82
- Gache Y, Baldeschi C, Del RM *et al.* (2004) Construction of skin equivalents for gene therapy of recessive dystrophic epidermolysis bullosa. *Hum Gene Ther* 15:921–33
- García M, Larcher F, Hickerson RP *et al.* (2011) Development of skin-humanized mouse models of pachyonychia congenita. *J Invest Dermatol* 131:1053–60
- Green H (1978) Cyclic AMP in relation to proliferation of the epidermal cell: a new view. *Cell* 15:801–11
- Lampert IA (1985) Expression of class II MHC antigen on epithelia and autoimmunity. *Lancet* 2:1078
- Leigh I, Watt F (1994) *Keratinocyte Methods*. Cambridge: Cambridge University Press
- Mildner M, Ballaun C, Stichenwirth M *et al.* (2006) Gene silencing in a human organotypic skin model. *Biochem Biophys Res Commun* 348:76–82
- Mildner M, Jin J, Eckhart L *et al.* (2010) Knockdown of filaggrin impairs diffusion barrier function and increases UV sensitivity in a human skin model. *J Invest Dermatol* 130:2286–94
- O'Shaughnessy RF, Choudhary I, Harper JI (2010) Interleukin-1 alpha blockade prevents hyperkeratosis in an *in vitro* model of lamellar ichthyosis. *Hum Mol Genet* 19:2594–605
- Oji V, Eckl KM, Aufenvenne K *et al.* (2010) Loss of corneodesmosin leads to severe skin barrier defect, pruritus, and atopy: unraveling the peeling skin disease. *Am J Hum Genet* 87:274–81
- Rheinwald JG, Green H (1975) Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells. *Cell* 6:331–43
- Schäfer-Korting M, Bock U, Diembeck W *et al.* (2008) The use of reconstructed human epidermis for skin absorption testing: results of the validation study. *Altern Lab Anim* 36:161–87
- Thomas AC, Tattersall D, Norgett EE *et al.* (2009) Premature terminal differentiation and a reduction in specific proteases associated with loss of ABCA12 in Harlequin ichthyosis. *Am J Pathol* 174:970–8
- Tjabringa G, Bergers M, van Rens D *et al.* (2008) Development and validation of human psoriatic skin equivalents. *Am J Pathol* 173: 815–23
- Williams ML (1992) Ichthyosis: mechanisms of disease. *Pediatr Dermatol* 9:365–8

AKT Has an Anti-Apoptotic Role in ABCA12-Deficient Keratinocytes

Journal of Investigative Dermatology (2011) **131**, 1942–1945; doi:10.1038/jid.2011.132; published online 2 June 2011

TO THE EDITOR

Harlequin ichthyosis (HI) is a hereditary skin disorder characterized by severe hyperkeratosis and impaired skin barrier function (Moskowitz *et al.*, 2004; Akiyama *et al.*, 2005). We have identified the ATP-binding cassette transporter A12 (ABCA12) as the causative gene of HI and, furthermore, demonstrated that ABCA12 is essential for keratinocyte lipid transport (Akiyama *et al.*, 2005; Yanagi *et al.*, 2008). Loss of ABCA12 function causes lipid transport to be defective in keratinocytes of the upper spinous and granular layers, resulting in the deposition of numerous intracellular lipid droplets and malformation of intercellular lipid layers (Akiyama *et al.*, 2005; Yanagi *et al.*, 2010). Recently, we have shown that gangliosides accumulate in the differentiated keratinocytes of HI patients (Mitsutake *et al.*, 2010). On the basis of the evidence that lipid accumulation is involved in keratinocyte apoptosis (Wang *et al.*, 2001; Uchida *et al.*, 2010), we investigated apoptotic and anti-apoptotic parameters in skin samples from HI patients and *Abca12*^{-/-} HI model mice.

We studied the skin of two HI patients and that of *Abca12*^{-/-} mice. The ABCA12 mutations of the two HI patients have been previously reported: one patient has the homozygous splice acceptor site mutation c.3295-2A>G and the other has the homozygous nonsense mutation p.Arg434X (Akiyama *et al.*, 2005). The procedure for generating *Abca12*^{-/-} mice, the establishment of primary-cultured keratinocytes, immunofluorescence staining, immunoblotting, and real-time reverse transcriptase PCR analysis has been previously described (Yanagi *et al.*, 2008, 2010). First, we investigated the apoptosis of HI patient epidermis by hematoxylin–eosin stain and TUNEL assay (*In situ* Apoptosis Detection Kit, Takara Bio, Otsu, Japan). In the HI patients, the nuclei of the granular-layer keratinocytes were condensed (Figure 1b) and they show positive for TUNEL labeling (Figure 1d), although apoptotic nuclei are rare in the normal human epidermis (Figure 1a, c). The histopathological findings and results of TUNEL staining of the *Abca12*^{-/-} mice

were similar to those in the skin of the HI patients (Figure 1f and h). TUNEL staining in the epidermis of 18.5-day embryos indicated that the apoptosis of keratinocytes started during fetal skin development (Figure 1j).

We assessed the degree of AKT activation of *Abca12*^{-/-} skin and keratinocytes using anti-AKT antibody #4691 and anti-phosphorylated AKT (Ser473) #4060 antibody (Cell Signaling, Danvers, MA). By immunoblot analysis, differentiated primary-cultured keratinocytes and the epidermis of *Abca12*^{-/-} mice showed higher expression levels of Ser-473 phosphorylated AKT than those of the control wild-type mice (Figure 1o). Immunofluorescence staining detected phosphorylated AKT in the upper granular-layer keratinocytes of the *Abca12*^{-/-} mouse skin (Figure 1l), but not in the skin of control wild-type mouse (Figure 1k). Cell proliferation was assessed by Ki-67 immunofluorescence (Figure 1). Ki-67 stain was similar in the wild-type and the *Abca12*^{-/-} samples, indicating that the granular-layer keratinocytes of the *Abca12*^{-/-} neonatal mice showed no excessive cell proliferation. To clarify whether AKT activation has

Abbreviations: ABCA12, ATP-binding cassette transporter A12; HI, harlequin ichthyosis; PPAR, peroxisome proliferator-activated receptor; RXR, retinoid X receptor

anti-apoptotic effects on *Abca12*^{-/-} keratinocytes, we performed TUNEL staining of keratinocytes treated with AKT inhibitor, which blocks AKT phosphorylation (#124017; InSolution Akt Inhibitor VIII, Calbiochem, San Diego, CA). *Abca12*^{-/-} keratinocytes incubated with 10 μM #124017 AKT inhibitor showed a notably greater number of TUNEL-positive

cells than both wild-type keratinocytes with AKT inhibitor and *Abca12*^{-/-} keratinocytes without AKT inhibitor (Figure 2). These results suggest that AKT activation helps *Abca12*^{-/-} keratinocytes to avoid apoptosis. Furthermore, mRNA and protein levels of peroxisome proliferator-activated receptor (PPAR)-δ from *Abca12*^{-/-} epidermis were shown

to be significantly higher than those from wild-type epidermis (Taqman Gene Expression Assay, probe ID, Mm00803184_m1, Mm99999915_g1, Applied Biosystems, Carlsbad, CA; anti-PPAR-δ antibody H-74, Santa Cruz, Santa Cruz, CA; Supplementary Figure S1 online), which suggests upregulation of PPAR-δ as a candidate pathway for AKT activation.

Herein, we have suggested that apoptosis is involved in the pathomechanism of HI. Defective lipid transport due to loss of ABCA12 function leads to the accumulation of intracellular lipids, including glucosylceramides and gangliosides (Akiyama *et al.*, 2005; Mitsuake *et al.*, 2010). Studies by Wang *et al.* (2001) and Sun *et al.* (2002) showed that the elevation of ganglioside levels leads to keratinocyte apoptosis. Thus, we are able to speculate that the accumulation of gangliosides leads to the apoptosis of *Abca12*^{-/-} keratinocytes, although the exact mechanism of apoptosis in *Abca12*^{-/-} keratinocytes remains unclear.




Figure 1. ATP-binding cassette transporter A12-deficient keratinocytes show TUNEL-positive nuclei and AKT activation. (a-d) In the harlequin ichthyosis patients, the nuclei of the granular-layer keratinocytes are condensed (b, white arrows) and they show positive TUNEL labeling (d, white arrows), although apoptotic nuclei are rare in the normal human epidermis (a, c). Data shown are representative of those from the two harlequin ichthyosis patients. (e, f) Granular-layer keratinocytes of *Abca12*^{-/-} mice show more condensed nuclei (f, white arrows) than those of wild-type mice (e). (g-j) Granular-layer keratinocytes of *Abca12*^{-/-} mice, a neonate (h) and an 18.5-day embryo (j), show TUNEL-positive nuclei. No TUNEL-positive cells are seen in the epidermis of the control wild-type mice (g, i). Dotted lines indicate the basement membrane. Nonspecific staining is seen on the skin surface (white arrowheads). (k, l) By immunofluorescence staining, AKT activation (Ser-473 phosphorylated AKT; green) is observed in granular-layer keratinocytes of *Abca12*^{-/-} mice. (m, n) Immunofluorescence staining for the Ki-67-proliferation marker shows similar staining patterns of basal keratinocytes in wild-type (m) and *Abca12*^{-/-} (n) samples. (a, b, e, f; hematoxylin-eosin (H-E) stain. Bars of c, d, g, h, i, j, k, l, m, n = 20 μm. Bars of a, b, e, f = 5 μm.) (o) Immunoblot analysis shows that levels of serine-473-phosphorylated AKT (pAKT) in neonatal epidermis and differentiated keratinocytes of *Abca12*^{-/-} mice are higher than those of wild-type mice.




Figure 2. Inhibition of AKT activation leads to apoptosis of *Abca12*^{-/-} keratinocytes. (a) Immunoblot analysis indicates that the AKT inhibitor can inhibit AKT activation (phosphorylated AKT (pAKT) synthesis) in differentiated keratinocytes. (b–e) TUNEL staining of keratinocytes cultured under high Ca²⁺ condition treated with/without the AKT inhibitor. Neither wild-type cells (b) nor *Abca12*^{-/-} cells (c) are TUNEL positive. *Abca12*^{-/-} keratinocytes with the AKT inhibitor (#124017; 10 μM) show many TUNEL-positive nuclei (e), although only a small number of wild-type cells with the AKT inhibitor are TUNEL positive (d). (Bars = 20 μm.) (f) Percentage of TUNEL-positive keratinocytes. *Abca12*^{-/-} keratinocytes with AKT inhibitor shows a significantly greater number of TUNEL-positive nuclei than wild-type keratinocytes with/without the AKT inhibitor and *Abca12*^{-/-} keratinocytes without the AKT inhibitor. (n = 3, mean ± SD, *P < 0.05).

Although *Abca12*^{-/-} granular-layer keratinocytes show characteristics of apoptosis, including condensed nuclei and positive TUNEL labeling, they are able to form epidermal stratification. In several disorders involving keratinocyte apoptosis, e.g., toxic epidermal necrolysis, the apoptotic epidermal keratinocytes show not only TUNEL-positive nuclei but also defective epidermal stratification (Abe *et al.*, 2003). Thrash *et al.* (2006) reported that AKT1 activation is an essential signal for keratinocyte cell survival and stratification, by experiments with gene silencing and three-dimensional cell cultures. Thus, we hypothesized that the AKT pathway might work as a compensatory mechanism against apoptosis in *Abca12*^{-/-} keratinocytes. We have clearly shown that AKT activation occurs in *Abca12*^{-/-} granular-layer keratinocytes, which suggests that AKT activation serves to prevent the cell death of *Abca12*^{-/-} keratinocytes. By immunoblot analysis using anti-AKT1/2/3 antibodies (#2938/3063/3788, Cell Signaling), *Abca12*^{-/-} epidermis showed expression of AKT1 and AKT2, but not AKT3 (Supplementary Figure S2 online). Compared with wild-type epidermis, *Abca12*^{-/-} epidermis seemed to have more AKT1 than AKT2. From our data and the literature (Thrash *et al.*, 2006), we are able to speculate that AKT1 is the major isoform of phosphorylated AKT in *Abca12*^{-/-} epidermis.

We have shown that PPAR-δ is a candidate molecule in the upstream of the AKT activation pathway in *Abca12*^{-/-} keratinocytes. Di-Poi *et al.* (2002) reported that PPAR-δ has an anti-apoptotic role in keratinocytes via transcriptional control of the AKT1 signaling pathway. PPAR-δ also regulates the expression of ABCA12 (Jiang *et al.*, 2008). From these studies, we can speculate that upregulation of PPAR-δ is in response to apoptosis or decreased ABCA12 expression. To ascertain the function of PPAR-δ, we performed the experiments using a PPAR-δ-specific antagonist (GSK0660, Santa Cruz). Differentiated *Abca12*^{-/-} keratinocytes treated with 1 μM GSK0660 for 48 hours showed TUNEL-positive nuclei, from which we are able to speculate an anti-apoptotic role for

PPAR- δ in *Abca12*^{-/-} keratinocytes (Supplementary Figure S1 online). From our studies and the literature (Di-Poi et al., 2002), PPAR- δ has been shown to have at least an anti-apoptotic role in *Abca12*^{-/-} keratinocytes; however, it remains unclear whether the upregulation of PPAR- δ is in response to apoptosis or decreased ABCA12 expression.

Furthermore, we have measured the mRNA expression levels of other nuclear hormone receptors, including PPAR- α , PPAR- γ , retinoic acid receptor- α , liver X receptor- α , liver X receptor- β , RXR- α , and RXR- γ (Applied Biosystems). The mRNA level of RXR- α from *Abca12*^{-/-} epidermis was shown to be significantly higher than that from wild-type epidermis (Supplementary Figure S1 online). The interaction between the upregulation of RXR- α and AKT activation in keratinocytes has not been reported. However, Wang et al. (2011) reported that RXR- α ablation in the epidermis enhances UV-induced apoptosis, which suggests that RXR- α has an anti-apoptotic function in keratinocytes. Thus, upregulation of RXR- α may also have an anti-apoptotic function in *Abca12*^{-/-} keratinocytes.

In conclusion, the present data suggest that keratinocyte apoptosis is involved in the pathomechanisms of HI and that the AKT signaling pathway helps *Abca12*^{-/-} keratinocytes to survive during the keratinization process. In light of this, activation of the AKT signal pathway may be to our knowledge, previously unreported strategy for treating keratinization disorders, including ichthyosis.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

We thank Ms Aoyanagi for her technical assistance. This work was supported in part by a grant-in-aid from the Ministry of Education, Science, Sports and Culture of Japan (Kiban A 23249058: to MA), a grant from the Ministry of Health, Labor and Welfare of Japan (Health and Labor Sciences Research Grants; Research on Intractable Disease: H22-177: to MA), and a grant-in-aid from the Japan Society for the Promotion of Science Fellows (to TY).

Teruki Yanagi¹, Masashi Akiyama^{1,2}, Hiroshi Nishihara³, Yuki Miyamura¹, Kaori Sakai¹, Shinya Tanaka⁴ and Hiroshi Shimizu¹

¹Department of Dermatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan; ²Department of Dermatology, Nagoya University Graduate School of Medicine, Nagoya, Japan; ³Laboratory of Translational Pathology, Hokkaido University Graduate School of Medicine, Sapporo, Japan and ⁴Laboratory of Cancer Research, Department of Pathology, Hokkaido University Graduate School of Medicine, Sapporo, Japan
E-mail: makiyama@med.nagoya-u.ac.jp

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

REFERENCES

- Abe R, Shimizu T, Shibaki A et al. (2003) Toxic epidermal necrolysis and Stevens-Johnson syndrome are induced by soluble Fas ligand. *Am J Pathol* 162:1515-20
- Akiyama M, Sugiyama-Nakagiri Y, Sakai K et al. (2005) Mutations in lipid transporter ABCA12 in harlequin ichthyosis and functional recovery by corrective gene transfer. *J Clin Invest* 115:1777-84
- Di-Poi N, Tan NS, Michalik L et al. (2002) Antiapoptotic role of PPARbeta in keratino-

cytes via transcriptional control of the Akt1 signaling pathway. *Mol Cell* 10:721-33

- Jiang YJ, Lu B, Kim P et al. (2008) PPAR and LXR activators regulate ABCA12 expression in human keratinocytes. *J Invest Dermatol* 128:104-9
- Mitsutake S, Suzuki C, Akiyama M et al. (2010) ABCA12 dysfunction causes a disorder in glucosylceramide accumulation during keratinocyte differentiation. *J Dermatol Sci* 60:128-9
- Moskowitz DG, Fowler AJ, Heyman MB et al. (2004) Pathophysiologic basis for growth failure in children with ichthyosis: an evaluation of cutaneous ultrastructure, epidermal permeability barrier function, and energy expenditure. *J Pediatr* 145:82-92
- Sun P, Wang XQ, Lopatka K et al. (2002) Ganglioside loss promotes survival primarily by activating integrin-linked kinase/Akt without phosphoinositide 3-OH kinase signaling. *J Invest Dermatol* 119:107-17
- Thrash BR, Menges CW, Pierce RH et al. (2006) AKT1 provides an essential survival signal required for differentiation and stratification of primary human keratinocytes. *J Biol Chem* 281:12155-62
- Uchida Y, Houben E, Park K et al. (2010) Hydrolytic pathway protects against ceramide-induced apoptosis in keratinocytes exposed to UVB. *J Invest Dermatol* 130:2472-80
- Wang XQ, Sun P, Paller AS (2001) Inhibition of integrin-linked kinase/protein kinase B/Akt signaling: mechanism for ganglioside-induced apoptosis. *J Biol Chem* 276:44504-11
- Wang Z, Coleman DJ, Bajaj G et al. (2011) RXRalpha ablation in epidermal keratinocytes enhances UVR-induced DNA damage, apoptosis, and proliferation of keratinocytes and melanocytes. *J Invest Dermatol* 131:177-87
- Yanagi T, Akiyama M, Nishihara H et al. (2008) Harlequin ichthyosis model mouse reveals alveolar collapse and severe fetal skin barrier defects. *Hum Mol Genet* 17:3075-83
- Yanagi T, Akiyama M, Nishihara H et al. (2010) Self-improvement of keratinocyte differentiation defects during skin maturation in ABCA12-deficient harlequin ichthyosis model mice. *Am J Pathol* 177:106-18

See related commentary on pg 1790

Interpretation of Skindex-29 Scores: Cutoffs for Mild, Moderate, and Severe Impairment of Health-Related Quality of Life

Journal of Investigative Dermatology (2011) 131, 1945-1947; doi:10.1038/jid.2011.138; published online 19 May 2011

TO THE EDITOR

Health-related quality of life (HRQL) is commonly assessed by means of standar-

dized questionnaires and expressed in domain and overall HRQL scores. An important challenge is to interpret these

scores correctly. What does a given score really mean? Although there is no standard approach, several methods exist to facilitate the interpretation of HRQL scores.

In a recently published study (Prinsen et al., 2010), we identified

Abbreviation: HRQL, health-related quality of life