

CONCEPTUAL PAPERS

CP1

THE EVALUATION OF ECONOMIC METHODS TO ASSESS THE SOCIAL VALUE OF MEDICAL INTERVENTIONS FOR ULTRA-RARE DISORDERS (URDS)

Schländler M¹, Garattini S², Holm S³, Kolominsky-Rabas PL⁴, Nord E⁵, Persson U⁶, Postma MJ⁷, Richardson J⁸, Simoons S⁹, de Sola-Morales O¹⁰, Tolley K¹¹, Toumi M¹²
¹Institute for Innovation & Valuation in Health Care (InnoVal-HC), Wiesbaden, Germany, ²Mario Negri Institute for Pharmacological Research, Milano, Italy, ³University of Manchester, Manchester, UK, ⁴University of Erlangen, Erlangen, Germany, ⁵Norwegian Institute of Public Health, Oslo, Norway, ⁶The Swedish Institute for Health Economics (IHE), Lund, Sweden, ⁷University of Groningen, Groningen, The Netherlands, ⁸Monash University, Clayton, Victoria, Australia, ⁹KU Leuven, Leuven, Belgium, ¹⁰Sabirmedical, Barcelona, Spain, ¹¹Tolley Health Economics Ltd., Buxton, Derbyshire, UK, ¹²University Claude Bernard Lyon 1, Lyon, France

OBJECTIVES: To develop a set of criteria to critically appraise the strengths and weaknesses of health economic methods for the systematic valuation of interventions for ultra-rare disorders (URDs). **METHODS:** An international group of clinical and health economic experts met in conjunction with the Annual European ISPOR Congresses in Berlin/Germany and Dublin/Ireland, November 2012 and 2013, to deliberate and agree on a set of criteria to assess the potential of the various methods, which have been used or proposed to estimate the social value of medical interventions for URDs. **RESULTS:** The group identified a broad set of potential criteria, which may be grouped according to the following dimensions: theoretical foundations (normative premises, i.e., links to moral and economic theories, including - but not limited to - nonutilitarian consequentialist and deontological reasoning, definition and treatment of core concepts of economic thinking such as opportunity costs and efficiency), empirical underpinnings (social preferences related to attributes of the health condition or of the person afflicted with it), and pragmatic aspects (feasibility of implementation and potential for bias and misuse). For each of the dimensions, a set of criteria has been agreed upon, which in turn will need further scrutiny and justification. **CONCLUSIONS:** Previously, a need had been identified for modifications or alternatives to the conventional logic of cost effectiveness applying benchmarks for the maximum allowable cost per quality-adjusted life year (QALY). We propose a framework for the systematic assessment how well different evaluation approaches reflect prevalent social norms and value judgments. As a next step, the framework shall be applied on multi-criteria decision analysis methods and social cost value analysis, either using the person trade-off (PTO) or the relative social willingness-to-pay (RS-WTP) instrument.

CP2

VALUE IN THE MAKING: HARVESTING THE VALUE OF COMPLEX MEDICAL INNOVATIONS IN PRACTICE

Abrishami P¹, Boer A², Horstman K¹

¹Maastricht University, Maastricht, The Netherlands, ²National Health Care Institute, Diemen, The Netherlands

Rapid development of medical innovations in the face of rising health care costs have been calling for a more value-conscious adoption and diffusions of innovations. This conceptual paper departs from swift adoption of the da Vinci surgical robot in the Netherlands. It describes three challenges facing health care systems to evaluate promising, yet complex and often expensive medical innovations. Firstly, they are often adopted and diffused prior to their evidence-based superiority being proven. Secondly, formal evaluation frameworks are somehow detached from the dynamics of and incentives for adoption and diffusion of these innovations. Third, the real risks and benefits of these innovations are not easily amenable to an experimental inquiry. Unlike pharmaceuticals, whose impact is intrinsic to its biochemical components and thus can be subject to experiment, the value of complex surgical devices, imaging equipments, or targeted therapy interventions are inseparable from actual patterns of human practices and clinical pathways that utilize them. Multi-stakeholder (early) deliberation has often been proposed to better align value requirements with the adoption and diffusion processes. This article examines the importance of developing a shared value perspective on the implementation of complex innovations through early deliberation. Product developers, (potential) adopters (providers or patients), purchasers, and policy makers may engage in an upfront iterative deliberation on all the particularities and (pre)conditions that account for delivering value of a certain innovation during early adoption in a given care delivery setting. Such deliberation offers a cumulative learning as to how to reduce true-to-life uncertainties and risks 'along the way', thereby serving for value 'fulfillment' in practice. Implication of such situated deliberative platforms for technology (outcome) assessment and for the role of authorities is discussed. A concrete framework for multi-stakeholder deliberation applied to the case of the da Vinci surgical robot in the Netherlands is also proposed.

CP3

EVALUATING THE QUALITY OF EVIDENCE FROM A NETWORK META-ANALYSIS

Higgins JP¹, Del Giovane C², Chaimani A³, Caldwell DM⁴, Salanti G³

¹University of Bristol, Bristol, UK, ²University of Modena and Reggio Emilia, Modena, Italy, ³University of Ioannina School of Medicine, Ioannina, Greece

Systematic reviews that collate data about the relative effects of multiple interventions via network meta-analysis are highly informative for decision-making purposes. A network meta-analysis provides two types of findings for a specific outcome: the relative treatment effect for all pairwise comparisons, and a ranking of the treatments. It is important to consider the confidence with which these two types of results can enable clinicians, policy makers and patients to make informed decisions. We propose an approach to determining confidence in the output of a network meta-analysis, based on methodology developed by the Grading of Recommendations Assessment, Development and Evaluation (GRADE) Working Group for pairwise meta-analyses. The suggested framework for evaluating a network meta-analysis acknowledges (i) the key role of indirect comparisons (ii) the contributions of each piece of direct evidence to the network meta-analysis estimates of effect size; (iii) the importance of the transitivity assumption to the

validity of network meta-analysis; and (iv) the possibility of disagreement between direct evidence and indirect evidence. We illustrate the framework using a network meta-analysis of topical antibiotics without steroids for chronically discharging ears with underlying eardrum perforations.

CP4

AMENDING THE GUIDE TO METHODS OF TECHNOLOGY APPRAISAL AT NICE TO INCORPORATE TWO NEW VALUE ELEMENTS: BURDEN OF ILLNESS AND WIDER SOCIETAL IMPACT

Knight H¹, Boysen M¹, Stevens A², Longson C¹

¹National Institute for Health and Care Excellence, Manchester, UK, ²University of Birmingham, Birmingham, UK

BACKGROUND: In July 2013 the Department of Health referred terms of reference for value based assessment of health technologies to NICE. **OBJECTIVE:** We present the approach taken over the past 11 months to amending the Guide to Methods of Technology Appraisal to incorporate burden of illness and wider societal impact. **METHODS:** Given the time frame available, NICE built on prior work undertaken by the Department of Health (in the context of value based pricing) on the concepts of burden of illness and wider societal benefits by commissioning the NICE Decision Support Unit to review and critique this existing work. NICE reconvened the working party from the Guide to the Methods of Technology Appraisal review which took place in 2012, which included standing membership drawn from the stakeholder communities such as patient and professional organisations, academia and pharmaceutical industry. The working party considered the prior work undertaken by the Department of Health and the Decision Support Unit's critique for burden of illness and wider societal benefit, and provided advice on the incorporation of the 2 new value elements into NICE's current methods at 4 meetings. A consultation paper describing NICE's proposals and draft of the amended sections of the methods guide was published in March 2014, and consultation ran for 12 weeks. It is anticipated that the final amendment of the methods guide will be considered by the NICE Board in advance of the ISPOR conference. **RESULTS:** Key points drawn from the discussion at the working party and consultation responses regarding burden of illness and wider societal impact, will be discussed. **DISCUSSION:** Considering NICE's 'position' in the world of health technology assessment and appraisal, the conclusions from this latest amendment of the Guide to Methods of Technology Appraisal to incorporate value based assessment will be (highly) anticipated.

DIAGNOSTIC RESEARCH STUDIES

D11

COST-EFFECTIVENESS (CE) OF IMAGING-GUIDED STRATEGIES FOR THE DIAGNOSIS OF CORONARY ARTERY DISEASE (CAD): RESULTS FROM THE EVINCI STUDY

Lorenzoni V¹, Pierotti F¹, Bellelli S¹, Neglia D², Rovai D², Turchetti G¹

¹Scuola Superiore Sant'Anna, Pisa, Italy, ²National Research Council, Pisa, Italy

OBJECTIVES: To evaluate the cost-effectiveness (CE) of imaging-guided strategies for the diagnosis of significant coronary artery disease (CAD) in patients with intermediate pre-test likelihood. **METHODS:** Significant CAD was defined as invasive coronary angiography (ICA) as >50% stenosis in the left main or >70% stenosis in a major coronary vessel or 30-70% stenosis with fractional flow reserve ≤ 0.8 . Nine diagnostic strategies were compared using a CE analysis. Strategies included the use of one single or two combined non-invasive imaging tests (CTCA as first line test and then stress ECHO, CMR, PET or SPECT) followed by ICA in the case of positivity of the single test or both non-invasive examinations in the case of combinations. ICERs were obtained using per-patient data collected throughout the EVINCI multi-centre European study. Strategy costs were calculated using examination country-specific reimbursements, while effectiveness was defined as the percentage of correct diagnosis. All costs were converted to Euro 2012 and adjusted using PPP. A propensity-score adjustment was used in the analysis and 95%CI were obtained with non-parametric bootstrap. **RESULTS:** Among the strategies analysed only three resulted cost-effective for the diagnosis of significant CAD. These included stress ECHO and CTCA as single non-invasive test, CTCA first then ECHO, CTCA first and then stress PET, all followed by ICA when required. Stress ECHO approach was the least costly but also the least effective, while CTCA alone [ICER: 2345 (2287-2400)] or in combination with PET [ICER: 5227(5161-5296)] had increasingly higher effectiveness for a willingness to pay (WTP) exceeding 2,000 Euro and 5,000 Euro, respectively. **CONCLUSIONS:** Results from the health-economic analysis of the EVINCI study showed that stress ECHO guided diagnostic strategy could be cost-effective when the WTP is low. Strategies involving CTCA alone or as first line exam followed by stress PET could allow a more accurate diagnostic workflow for higher WTP.

D12

THE VALUE OF RISK-STRATIFIED INFORMATION IN THE NATIONAL LUNG CANCER SCREENING TRIAL

Soeteman DJ¹, Cohen JT¹, Neumann PJ², Wong JB¹, Kent DM¹

¹Tufts Medical Center/Tufts University School of Medicine, Boston, MA, USA, ²Tufts Medical Center, Boston, MA, USA

OBJECTIVES: Clinical guideline recommendations are generally informed by population-based evidence. However, interventions that are (cost-)effective on average may not be (cost-)effective for many (even for most) patients meeting trial inclusion criteria. This study aims to investigate the value of risk-stratified recommendations for lung cancer screening among current or former smokers between the ages of 55 and 74 years compared to a screen-all policy. **METHODS:** Using data from the National Lung Cancer Screening Trial (NLST), we calculated the costs and QALYs for low-dose computed tomography (CT) versus chest radiography (X-ray) from empirically observed health states and 6 years life expectancy. Based on Kovalchik's risk of lung cancer death prediction model, we stratified 53,454 NLST trial patients into quintiles. The expected value of individualized care (EVIC) was calculated to