

REVIEW ARTICLE

Perioperative cardiovascular evaluation: heads or tails?

GABRIEL A.L. CARMO¹, DANIELA CALDERARO², PAI C. YU², DANIELLE M. GUALANDRO², ANDRÉ C. MARQUES², CRISTINA S. BITTAR¹, ADRIANA F. PASTANA³, BRUNO CARAMELLI²

¹MD, Interdisciplinary Medicine in Cardiology Unit, Instituto do Coração (InCor), Hospital das Clínicas, Medical School, Universidade de São Paulo (USP), São Paulo, SP, Brazil

²PhD in Cardiology, Interdisciplinary Medicine in Cardiology Unit, InCor, Hospital das Clínicas, Medical School, USP, São Paulo, SP, Brazil

³BSBIO, Interdisciplinary Medicine in Cardiology Unit, InCor, Hospital das Clínicas, Medical School, USP, São Paulo, SP, Brazil

SUMMARY

When dealing with surgical patients, a perioperative evaluation is essential to anticipate complications and institute measures to reduce the risks. Several algorithms and exams have been used to identify postoperative cardiovascular events, which account for more than 50% of perioperative mortality. However, they are far from ideal. Some of these algorithms and exams were proposed before important advances in cardiology, at a time when pharmacological risk reduction strategies for surgical patients were not available. New biomarkers and exams, such as C-reactive protein, brain natriuretic peptide, and multislice computed tomography have been used in cardiology and have provided important prognostic information. The ankle-brachial index is another significant marker of atherosclerosis. However, specific information regarding the perioperative context of all these methods is still needed. The objective of this article is to evaluate cardiovascular risk prediction models after noncardiac surgery.

Keywords: Perioperative care; general surgery; brain natriuretic peptide; general anesthesia; cardiovascular risk; ankle-brachial index.

©2012 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

Study conducted at the Interdisciplinary Medicine in Cardiology Unit, Instituto do Coração (InCor), Hospital das Clínicas, Medical School, Universidade de São Paulo, São Paulo, SP, Brazil

Submitted on: 10/17/2011

Approved on: 04/09/2012

Correspondence to:
Interdisciplinary Medicine
in Cardiology Unit
Instituto do Coração (InCor),
Hospital das Clínicas
Universidade de São Paulo
Avenida Dr. Enéas Carvalho de
Aguiar, 44, Anexo II
Unidade de Medicina
Interdisciplinar em Cardiologia
Cerqueira César
São Paulo – SP, Brazil
CEP: 05403-000
Phone: +55 (11) 2661-5376
Fax: +55 (11) 2661-5535
gabriel_carmo@yahoo.com.br

Conflict of interest: None.

INTRODUCTION

Perioperative evaluation is an important step before referring a patient to surgery. The purpose is not just clearance for surgery, but to perform an overall evaluation of the clinical status, make recommendations, deal with cardiovascular risk factors or cardiac problems, and estimate the procedure's risk. Thereupon the medical team and patient can decide the best way to minimize complications, or even to postpone the operation¹. In addition to clinical experience and common sense, algorithms for perioperative evaluation have been proposed, tested, and validated to contribute to this decision. In general, sensitivity and specificity are quite similar among algorithms; however, they are unfortunately only somewhat better than a coin toss. In addition, most of the assessments have been based upon clinical characteristics of patients submitted to surgery over ten years prior. More recent perioperative evaluation guidelines include flowcharts suggesting tests or pharmacological interventions that may be a rather frustrating support for the algorithms, although much more acceptable to attending physicians.

On the other hand, population growth, developing countries' welfare, technological improvement, new surgical techniques allied to new materials and devices, and faster non-invasive strategies have led to an increase in the number of surgeries^{2,3}. Indeed, patients previously considered non-candidates for surgery may now have renewed hope. Consequently, a larger number of patients older than 75 years are being referred to surgery involving multiple cardiovascular risk factors bringing about increased postoperative mortality and duration of hospitalization⁴. Recent predictions estimate that, from 2010 to 2040, the aging population in the United States will increase the incidence of coronary heart disease by approximately 26%, as well as the costs related to care by 41%⁵. It is estimated that more than 40 million surgeries are performed annually in Europe⁶, and 240 million around the world⁷. In developing countries, the same scenario is observed. From 1997 to 2007, an increase of 20.42% was observed in the number of surgical procedures in Brazil². In the last few years, however, the same authors observed a 30% increase in the number of case-fatalities². Whether coincident or related events, both findings represent important challenges: how to anticipate and prevent the increase in perioperative complications in this apparently higher risk population without imposing a great economic burden?

CARDIOVASCULAR COMPLICATIONS AND PERIOPERATIVE ALGORITHMS

Cardiovascular complications are of special concern when dealing with surgical patients, since approximately 1% of them present with acute myocardial infarction (AMI) after the procedure. Generally it is associated with other

noncardiac events with an odds ratio > 6, significantly increasing duration of hospitalization⁸. Related cardiovascular mortality reaches 0.3% (1.2 million patients in Europe alone), accounting for more than 50% of postoperative deaths after vascular surgeries^{6,9}.

Perioperative cardiovascular risk can be estimated by assessing clinical status, functional capacity and intrinsic risk of the surgery³. However, sometimes it may be very difficult to estimate this risk in patients with subclinical presentation of diseases.

Methods and algorithms were developed some time ago, and are being used in clinical practice with a high frequency¹⁰⁻¹². However, some of these algorithms were proposed prior to the development of important advances in cardiology such as the use of aspirin and statins for coronary artery disease and acute coronary syndromes¹³⁻¹⁹. They were also developed before important pharmacological risk reduction strategies were available for surgical patients, such as beta-blockers and statins²⁰⁻²². Despite having been cautiously validated in the past, it remains to be investigated whether these algorithms are still applicable in different populations submitted to different treatments. Do they continue to merit confidence? Do they predict cardiovascular events or are they little better than as tossing a coin?

The American Society of Anesthesiologists (ASA) classification was first described in 1941²³ and revised in 1963 (Box 1). It was the first attempt to predict surgical complications and is still, by far, the most often used by anesthesiologists. No other preoperative index has achieved the same widespread use. It was first designed to estimate the physiological status with no need for clinical resources and, although it can predict postoperative complications, it has a limited capability to predict cardiovascular complications. Moreover, there is a major problem related to poor reproducibility even among anesthesiologists²⁴.

The Detsky Index¹¹ is a modified version of the original risk index¹⁰. In his study, Detsky showed that the accuracy of the original method proposed by Goldman et al. had dropped from 81% to 69%. After inclusion of angina pectoris severity, previous myocardial infarction, critical aortic stenosis, and alveolar pulmonary edema, the accuracy of the new index increased to 75%. Conversely, the authors observed many events in patients with low-risk scores (false negative) and a limited discrimination power in patients referred for vascular or other major surgery. Based upon the findings of Eagle et al.⁹ and Vanzetto et al.,²⁵ the American College of Physicians (ACP) published a guideline for assessment and management of perioperative risk, suggesting the Modified Cardiac Index for stratification of all patients prior to surgery (Box 1)^{26,27}. To minimize the limitations of this method, they recommended the use of a non-invasive cardiac ischemic test for individuals with a

Box 1 – Perioperative risk indexes

American Society of Anesthesiologists Score (ASA Grade) ²³	Revised Cardiac Risk Index (Lee et al., 1999) ¹²
I – Normal health patient II – Patient with mild systemic disease III – Patient with severe systemic disease IV – Patient with severe systemic disease that is a constant threat to life V – A moribund patient who is not expected to survive without the operation	1 – High risk type of surgery (intraabdominal, intrathoracic, or suprainguinal vascular procedures) 2 – Ischemic heart disease (includes any of the following: history of myocardial infarction, history of positive exercise test, current complaint of chest pain, i.e., considered to be secondary to myocardial ischemia, use of nitrate therapy, or electrocardiography with pathologic Q waves) 3 – Congestive heart failure 4 – History of cerebrovascular disease 5 – Preoperative treatment with insulin 6 – Preoperative serum creatinine > 2.0 mg/dL
Modified Cardiac Risk Index Index^{9,11}	
Prior myocardial infarction <ul style="list-style-type: none"> Last infarction within 6 months: 10 points Last infarction more than 6 months ago: 5 points Angina pectoris <ul style="list-style-type: none"> Canadian Angina Class 3: 10 points Canadian Angina Class 4: 20 points 	Arrhythmia <ul style="list-style-type: none"> Rhythm other than sinus or sinus with PACs: 5 points More than five premature ventricular beats: 5 points Poor general medical status: 5 points <ul style="list-style-type: none"> pO₂ < 60 mm Hg, pCO₂ > 50 mmHg, K < 3 mmol/L, blood urea nitrogen level > 50 mmol/L, creatinine 3mg/dL, bedridden
Alveolar pulmonary edema <ul style="list-style-type: none"> Pulmonary edema within one week: 10 points Pulmonary edema at any time: 5 points 	Age > 70 years: 5 points
Suspected critical aortic stenosis: 20 points	Emergency surgery: 10 points
Eagle and Vanzetto variables (low risk variables)^{b,9,25}	
Age > 70 years	History of myocardial infarction
History of angina	ST-segment ischemic abnormalities during resting electrocardiography
Diabetes mellitus	Hypertension with severe left ventricular hypertrophy
Q waves on electrocardiogram	History of congestive heart failure
History of ventricular ectopy	

^aClass I: 0 to 15 points (< 3% perioperative cardiac events); class II 20 to 30 points (3 to 15% perioperative cardiac events); class III: more than 30 points (> 15% perioperative cardiac events). ^bClass I patients still do not reliably identify low risk patients for perioperative cardiac events. For these patients, when two or more low risk variables are present, consider referring for noninvasive testing. PAC, premature atrial complex.

low-risk of cardiovascular complications who were to undergo vascular surgery and had more than two low-risk variables (Box 1). Although recommended by the ACP, the Detsky Index did not show any improvement in the prediction of cardiac risk when compared to other methods such as the Goldman, the ASA, or the New York Heart Association classification of angina²⁸. Thus, despite the increased number of variables collected (and more complicated to use), the ACP algorithm does not offer clear advantages over simpler methods for patient evaluation prior to noncardiac surgeries.

Comparing a new algorithm proposal with Detsky's and Goldman's, Lee et al. found even lower areas under the ROC curve for these two methods, and demonstrated a

good accuracy for the prediction of cardiovascular events with a different system (Table 1)¹². This approach is recommended by the new guidelines of the American College of Cardiology, American Heart Association, and European Society of Cardiology^{1,29}. Although very simple to use, this new index, called the Revised Cardiac Risk Index (RCRI – Box 1), did not perform well for patients undergoing abdominal aortic aneurism (AAA) repair. A later validation of the RCRI showed limited value for the prediction of cardiovascular mortality³⁰.

Indeed, Bertges et al. found that the RCRI did not perform well for vascular surgery patients, consistently underestimating the composite cardiac complications across all risk groups. Therefore, the authors suggested

Table 1 – Area under the ROC curve for the perioperative cardiac indices (Based on Lee et al., 1999)¹²

Index	Derivation Cohort (ROC area)	Validation Cohort (ROC area)
Original Cardiac Risk Index	0.606	0.701
Modified Cardiac Risk Index	0.545	0.582
Revised Cardiac Risk Index	0.759	0.806

a new method (Vascular Study Group of New England Cardiac Risk Index – VSG CRI) to predict cardiovascular complications. They added older age, chronic obstructive pulmonary disease, smoking, and long term β -blocker use as clinical variables³¹. This last method seems to reflect a higher risk population instead of merely reflecting the effect of the drug. With these modifications, the sensitivity, specificity, positive predictive value, and negative predictive value were 68%, 62%, 11%, and 97%, respectively. In another study, the RCRI was inferior to the new proposed score (Myocardial Infarction and Cardiac Arrest Score – MICA) across all studied patients, including AAA surgery³².

In a systematic review evaluating the ability to predict cardiac complications and mortality after major noncardiac surgery, the RCRI was only moderately good at discriminating between low-risk versus high-risk patients for cardiac events³³. The index was not adequate to predict cardiac events after vascular noncardiac surgery or to predict all-cause mortality³³.

Moreover, the RCRI does not take age into account as a risk factor, although the same group of researchers later showed that age > 70 confers a higher risk for perioperative cardiac and noncardiac complications⁴.

In summary, there are several algorithms for perioperative risk estimation, with a wide variation of accuracy for the prediction of cardiovascular complications. The most important are shown in Table 2.

IMAGE METHODS

Echocardiography is the cornerstone of complementary cardiological evaluation and the most prescribed preoperative noninvasive test³⁴. It is readily available, with no contraindications and does not require intravenous injection, radioactive isotopes, or exposure to radiation. It provides information about cardiac dimensions, valve areas and function, abnormal communications between the left and right side, and calculation of cardiac output and ejection fraction. Despite all these advantages, the request of routine echocardiogram does not seem to improve survival or detection of cardiac complications³⁵, and may increase length of hospitalization after major elective noncardiac surgery³⁶.

Coronary angiography (CA) is a well-established method for detecting coronary artery disease (CAD). It is invasive, but widely available, even in small hospitals. Although there are no data regarding its use in the

perioperative period, its indications, as well as revascularization, are similar to angiography indications in non-surgical settings^{1,29}.

Cardiac stress tests are unequivocally the most often performed exams in patients referred to surgeries for detection of CAD. In general, their use before elective major surgery is associated with improved survival in intermediate and high risk patients. This is probably due to the identification of those who could benefit from the introduction of perioperative drug therapy such as β -blockers³⁷. They have been used since the late 1980's after the study by Eagle et al. demonstrated the benefit of dipyridamole-thallium image scans for patients undergoing vascular operations⁹. Their ability to predict cardiovascular complications is similar to that of clinical variables alone. However, when used together, thallium redistribution can separate intermediate risk patients into low- and high- risk groups with postoperative cardiac ischemic events rates of 3.2% and 29.6%, respectively⁹. These findings were confirmed by another study, which also found that the larger the ischemic area, the higher the probability of major cardiovascular events²⁵. Dobutamine-atropine echocardiography is a safe exam and lower in cost than scintigraphy. A positive result greatly increases the likelihood ratio (LR) of a perioperative cardiac event and, above all, it effectively identifies a low-risk group with a very low incidence of complications³⁸. Meta-analysis of published data also showed a better diagnostic performance of stress echocardiography in comparison to myocardial perfusion scintigraphy, with a negative LR of 0.23 vs. 0.44 and a positive LR of 4.09 vs. 1.83 for detection of myocardial infarction or death, respectively^{39,40}.

Recently, multislice computed tomography (MSCT) was shown to be useful in the screening for CAD with a high sensitivity and negative predictive value⁴¹. There is only one study evaluating the role of MSCT in patients referred for surgery and, although it is a retrospective study with serious limitations, it increased the capability of detecting CAD when used together with scintigraphy. Furthermore, when compared to CA, MSCT may identify patients at high-risk of cardiovascular events with the same rate, however at a lower cost⁴².

Reactive hyperemia is a vasodilatation of arterial vessels that occurs after a period of tissue ischemia and depends on local production of vasodilators. It is related to traditional cardiovascular risk factors and markers of inflammation^{43,44}.

Table 2 – Algorithms for perioperative risk evaluation

Score	Year	Advantages	Disadvantages
ASA ²³	1963	Very simple to use	Poor reproducibility, not developed for cardiovascular complications and does not take into account age and complexity of the operation
Goldman ¹⁰	1977	Simple to use	Developed before important advances in cardiology care
Detsky ¹¹	1986	Simple to use and well validated in other studies	Poor discriminative power for patients classified as intermediate risk. Developed before important advances in medicine
Charlson ^{45,46}	1987	Takes into account several risk factors not considered in other scores such as cancer, AIDS and liver disease. Validated in several populations	Originally used to predict mortality in general population. Used a high risk population admitted in a hospital. It was not validated in a large general population referred for noncardiac surgery
RCRI ¹⁵	1999	Simple to use. Well validated in other populations	Poor performance for vascular surgeries (especially AAA repair) and for mortality prediction
Fleisher-Eagle ⁴⁷	2001	Simple to use. Similar to RCRI without history of cerebrovascular disease	The algorithm is a theoretical approach and was not validated in any population
EMAPO ⁴⁸	2007	Considers diseases not mentioned in previous guidelines and modern treatment options. Stratifies patients into five levels with a narrower interval of complications in each category	Complex execution due to the large number of items to consider. It was not validated in other studies and didn't show any improvement compared to Detsky's model
POSSUM ⁴⁹	1991	Allows comparative audit between different populations	Not validated specifically for risk stratification. Seems to over predict death by a factor greater than two ⁵⁰
VSG-CRI ³¹	2010	Developed specifically for patients undergoing vascular surgeries with a better performance than the RCRI	Not validated in other populations
MICA ³²	2011	Performed better than RCRI for myocardial infarction and cardiac arrest. Takes into account the type of surgery	Developed from a surgical database without active search for cardiac complications. Not validated in other populations

AAA, aortic aneurism repair; AIDS, acquired immune deficiency syndrome; ASA, American Society of Anesthesiologists; EMAPO, multicenter study of perioperative evaluation; MICA, myocardial infarction or cardiac arrest score; POSSUM, physiological and operative severity score for enumeration of mortality and morbidity; VSG-CRI, vascular study group of New England cardiac risk index; RCRI, Revised Cardiac Risk Index.

There are few studies on this technique for perioperative evaluation and results are controversial^{51,52}. More research is needed to define its role in this field.

BIOMARKERS

In perioperative evaluation, new advances have recently been made, especially regarding biomarkers. Brain natriuretic peptide (BNP) and N-terminal pro-BNP (NT-proBNP) are produced by myocytes in response to stress and are important prognostic indicators of heart failure. They can identify patients with high-risk of cardiovascular complications after noncardiac surgery despite the absence of

inducible ischemia and low left ventricular ejection fraction⁵³⁻⁵⁵, and they seem to be useful even in emergency noncardiac surgery⁵⁶. Cuthbertson et al. demonstrated that a BNP > 40 pg.mL⁻¹ has a sensitivity of 75% and a specificity of 70% for predicting perioperative death or myocardial injury, and also performed better than the RCRI⁵⁷. When meta-analyzed, elevated BNP or NT-proBNP were predictors of adverse cardiovascular outcomes at 30 and 90 days and at over six months⁵⁸⁻⁶⁰. However, they are most useful to identify event-free survival and may not accurately identify mortality with a high degree of certainty – high negative predictive value and low positive predictive value⁵⁹.

Inflammatory markers such as C-reactive protein (CRP), which is related to atherogenesis and atherosclerotic plaque instability, may identify patients with elevated coronary risk. However, their use in perioperative evaluation has not been completely defined. In 2010, Choi et al. demonstrated that higher levels of CRP are associated with an increased risk of AMI, pulmonary edema, and cardiovascular death after major noncardiac surgery⁶¹. Yet, when associated with NT-proBNP, it increased the relative risk by threefold compared to classical algorithms for clinical events⁶¹. High-sensitivity CRP and fibrinogen were also recently studied, and at elevated levels they seem to independently predict cerebral ischemic events after carotid endarterectomy in patients, whether symptomatic or not⁶².

ANKLE BRACHIAL INDEX

The ankle brachial index (ABI) is valuable for cardiovascular risk quantification and is perhaps the most promising source of information in the perioperative period, especially when compared to intima media thickness, coronary calcium score, and coronary and carotid atherosclerotic plaques^{63,64}. In comparison with these methods, it is less expensive, faster, and feasible in office care, with a good acceptance by patients and little intra- and inter-observer variability⁶⁵⁻⁶⁷. The procedure is simple and can be carried out by a trained nurse or other health care professional⁶⁴. It is easily done by measuring the systolic blood pressure (BP) with a portable Doppler ultrasound machine on each arm and on the dorsalis pedis and posterior tibial arteries of each ankle. The highest of the two arm pressures is selected, as is the highest of the two pressures of each ankle. The ABI is obtained by dividing the highest ankle BP in each leg by the highest arm pressure. The lowest value between the two indices (one for each leg) defines the risk of the patient. Normally, the BP is higher in the legs than in the arms^{68,69}. Thus, it is expected that a normal ABI would be > 0.9 and ≤ 1.3 . Values > 1.3 suggest a non-compressible calcified artery in the ankle, thus the ABI is not reliable in this situation, because of unreliable BP measurements^{68,69}. Values between 0.41 and 0.9 are associated with peripheral arterial disease (PAD) and severe PAD when ≤ 0.4 ⁶⁹. Asymptomatic patients with abnormal ABI and no prior cardiovascular disease, older than 70 years, or older than 50 years with an additional cardiac risk factor, should be treated as a patient with established PAD⁷⁰. Furthermore, an abnormal ABI and PAD are associated with atherosclerosis in coronary and cerebral arteries and a higher incidence of AMI, stroke, and cardiovascular and all-cause mortality^{63,70-72}.

Although there is a close relationship between abnormal ABI and atherosclerosis, including in coronary arteries, the association of ABI with other cardiovascular scores has been poorly studied; however, it appears to be useful. In a recent meta-analysis the ABI has proven to be an independent

cardiovascular risk factor, and when associated with the Framingham risk score (FRS), an abnormal result doubled all-cause mortality, cardiovascular mortality and major coronary events in all groups of the FRS⁶⁴. Further, after inclusion of the ABI, 20% of men and 33% of women had their Framingham risk category changed. In women, this effect was mainly a change from low-risk to a high-risk category. In another study, the addition of ABI to traditional cardiovascular risk factors such as diabetes, hypertension, smoking, and hypercholesterolemia improved the sensitivity, specificity, and predictive values for a future cardiovascular event⁷².

Flu et al., recently demonstrated that an asymptomatic low ABI (< 0.9) increases the risk of perioperative myocardial damage over twofold in patients undergoing AAA or carotid artery stenosis repair, even after adjusting for other cardiovascular risk factors⁷³. Unfortunately, there are no data about the use of ABI in other patients referred for non-vascular surgery, which comprises the majority of surgeries performed worldwide. Such use could add extensive information about this population, especially if associated with well-validated perioperative cardiovascular risk scores.

CONCLUSIONS

With such conflicting information regarding the perioperative evaluation and its algorithms and exams, a cautious and accurate clinical evaluation of the patient's cardiac risk is critical. The accuracy of methods available for predicting postoperative complications is better than random, but their performance is not ideal. Thus, measures and approaches that are widely available and easy to perform for the accurate prediction of postoperative cardiovascular events are urgently needed.

The ideal risk prediction model would be one that is simple, reproducible, accurate, objective, and available to all patients⁷⁴. Coronary calcium score, intima media thickness, pulse wave velocity, and ABI, which can predict a higher cardiovascular risk in the general population, appear to be promising for perioperative evaluation, adding important information to current guidelines and algorithms. The ABI, an easy to perform and widely available exam in office practice and points of care, appears to be the best approach and the most promising method, but evidence to support its use in the perioperative context, especially for patients undergoing non-vascular surgeries is still needed.

REFERENCES

1. Fleisher LA, Beckman JA, Brown KA, Calkins H, Chaikof E, Fleischmann KE, et al. ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (writing committee to revise the 2002 guidelines on perioperative cardiovascular evaluation for noncardiac surgery): developed in collaboration with the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, and Society for Vascular Surgery. *Circulation*. 2007;116:e418-99.

2. Yu PC, Calderaro D, Gualandro DM, Marques AC, Pastana AF, Prandini JC, et al. Non-cardiac surgery in developing countries: epidemiological aspects and economical opportunities - the case of Brazil. *Plos One*. 2010;5:e10607.
3. Castro I, Gualandro DM, Yu PC, Calderaro D, Marques AC, Pinho C, et al. II guidelines for perioperative evaluation of the Brazilian Society of Cardiology. *Arq Bras Cardiol*. 2011;96:1-68.
4. Polanczyk CA, Marcantonio E, Goldman L, Rohde LE, Orav J, Mangione CM, et al. Impact of age on perioperative complications and length of stay in patients undergoing noncardiac surgery. *Ann Intern Med*. 2001;134:637-43.
5. Odden M, Coxson P, Moran A, Lightwood J, Goldman L, Bibbins-Domingo K. The impact of the aging population on coronary heart disease in the United States. *Am J Med*. 2011;124:827-33.
6. Poldermans D, Hoeks SE, Feringa HH. Pre-operative risk assessment and risk reduction before surgery. *J Am Coll Cardiol*. 2008;51:1913-24.
7. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet*. 2008;372:139-44.
8. Fleischmann KE, Goldman L, Young B, Lee TH. Association between cardiac and noncardiac complications in patients undergoing noncardiac surgery: outcomes and effects on length of stay. *Am J Med*. 2003;115:515-20.
9. Eagle KA, Coley CM, Newell JB, Brewster DC, Darling RC, Strauss HW, et al. Combining clinical and thallium data optimizes preoperative assessment of cardiac risk before major vascular-surgery. *Ann Intern Med*. 1989;110:859-66.
10. Goldman L, Caldera DL, Nussbaum SR, Southwick FS, Krogstad D, Murray B, et al. Multifactorial index of cardiac risk in non-cardiac surgical procedures. *N Engl J Med*. 1977;297:845-50.
11. Detsky AS, Abrams HB, McLaughlin JR, Drucker DJ, Sasson Z, Johnston N, et al. Predicting cardiac complications in patients undergoing noncardiac surgery. *J Gen Intern Med*. 1986;1:211-9.
12. Lee TH, Marcantonio ER, Mangione CM, Thomas EJ, Polanczyk CA, Cook EF, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation*. 1999;100:1043-9.
13. ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17187 cases of suspected acute myocardial-infarction - ISIS-2. *Lancet*. 1988;2:349-60.
14. Thérroux P, Ouimet H, Mc Cans J, Latour JG, Joly P, Lévy G, et al. Aspirin, heparin, or both to treat acute unstable angina. *N Engl J Med*. 1988;319:1105-11.
15. Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet*. 1994;344:1383-9.
16. Shepherd J, Cobbe SM, Ford I, Isles CG, Lorimer AR, MacFarlane PW, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med*. 1995;333:1301-7.
17. Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med*. 1996;335:1001-9.
18. Downs JR, Clearfield M, Weis S, Whitney E, Shapiro DR, Beere PA, et al. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study. *JAMA*. 1998;279:1615-22.
19. The Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *N Engl J Med*. 1998;339:1349-57.
20. Durazzo AES, Machado FS, Ikeoka DT, De Bernoche C, Monachini MC, Puech-Leão P, et al. Reduction in cardiovascular events after vascular surgery with atorvastatin: a randomized trial. *J Vasc Surg*. 2004;39:967-75.
21. Mangano DT, Layug EL, Wallace A, Tateo I. Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. Multicenter Study of Perioperative Ischemia Research Group. *N Engl J Med*. 1996;335:1713-20.
22. Poldermans D, Boersma E, Bax JJ, Thomson IR, van de Ven LL, Blakensteijn JD, et al. The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography Study Group. *N Engl J Med*. 1999;341:1789-94.
23. Saklad M. Grading of patients for surgical procedures. *Anesthesiology*. 1941;1:281-4.
24. Wolters U, Wolf T, Stützer H, Schröder T. ASA classification and perioperative variables as predictors of postoperative outcome. *Br J Anaesth*. 1996;77:217-22.
25. Vanzetto G, Machecourt J, Blendea D, Fagret D, Borrel E, Magne JL, Gattaz F, et al. Additive value of thallium single-photon emission computed tomography myocardial imaging for prediction of perioperative events in clinically selected high cardiac risk patients having abdominal aortic surgery. *Am J Cardiol*. 1996;77:143-8.
26. Palda VA, Detsky AS, Thibault GE, Feussner JR, Audet AMJ, Freisinger GC Jr, et al. Clinical guideline .1. Guidelines for assessing and managing the perioperative risk from coronary artery disease associated with major noncardiac surgery. *Ann Intern Med*. 1997;127:309-12.
27. Palda VA, Detsky AS. Clinical guideline .2. Perioperative assessment and management of risk from coronary artery disease. *Ann Intern Med*. 1997;127:313-28.
28. Gilbert K, Larocque BJ, Patrick LT. Prospective evaluation of cardiac risk indices for patients undergoing noncardiac surgery. *Ann Intern Med*. 2000;133:356-9.
29. Poldermans D, Bax JJ, Boersma E, De Hert S, Eekhout E, Fowkes G, et al. Guidelines for pre-operative cardiac risk assessment and perioperative cardiac management in non-cardiac surgery. *Eur Heart J*. 2009;30:2769-812.
30. Boersma E, Kertai MD, Schouten O, Bax JJ, Noordzij P, Steyerberg EW, et al. Perioperative cardiovascular mortality in noncardiac surgery: validation of the Lee cardiac risk index. *Am J Med*. 2005;118:1134-41.
31. Bertges DJ, Goodney PP, Zhao Y, Schanzer A, Nolan BW, Likosky DS, et al. The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the revised cardiac risk index in vascular surgery patients. *J Vasc Surg*. 2010;52:674-83, 83.e1-83.e3.
32. Gupta PK, Gupta H, Sundaram A, Kaushik M, Fanf X, Miller WJ, et al. Development and validation of a risk calculator for prediction of cardiac risk after surgery. *Circulation*. 2011;124:381-7.
33. Ford MK, Beattie WS, Wijeyesundera DN. Systematic review: prediction of perioperative cardiac complications and mortality by the revised cardiac risk index. *Ann Intern Med*. 2010;152:26-35.
34. Wijeyesundera DN, Austin PC, Beattie WS, Hux JE, Laupacis A. Outcomes and processes of care related to preoperative medical consultation. *Arch Intern Med*. 2010;170:1365-74.
35. Halm EA, Browner WS, Tubau JF, Tateo IM, Mangano DT. Echocardiography for assessing cardiac risk in patients having noncardiac surgery. Study of Perioperative Ischemia Research Group. *Ann Intern Med*. 1996;125:433-41.
36. Wijeyesundera DN, Beattie WS, Karkouti K, Neuman MD, Austin PC, Laupacis A. Association of echocardiography before major elective non-cardiac surgery with postoperative survival and length of hospital stay: population based cohort study. *BMJ*. 2011;342:d3695.
37. Wijeyesundera DN, Beattie WS, Austin PC, Hux JE, Laupacis A. Non-invasive cardiac stress testing before elective major non-cardiac surgery: population based cohort study. *BMJ*. 2010;340:b5526.
38. Poldermans D, Arnesse M, Fioretti PM, Salustri A, Boersma E, Thomson IR, et al. Improved cardiac risk stratification in major vascular surgery with dobutamine-atropine stress echocardiography. *J Am Coll Cardiol*. 1995;26:648-53.
39. Kertai MD, Boersma E, Bax JJ, Heijnenbroek-Kal MH, Hunink MG, L'italien GJ, et al. A meta-analysis comparing the prognostic accuracy of six diagnostic tests for predicting perioperative cardiac risk in patients undergoing major vascular surgery. *Heart*. 2003;89:1327-34.
40. Beattie WS, Abdelnaem E, Wijeyesundera DN, Buckley DN. A meta-analytic comparison of preoperative stress echocardiography and nuclear scintigraphy imaging. *Anesth Analg*. 2006;102:8-16.
41. Mollet NR, Cademartiri F, van Mieghem CA, Runza G, McFadden EP, Baks T, et al. High-resolution spiral computed tomography coronary angiography in patients referred for diagnostic conventional coronary angiography. *Circulation*. 2005;112:2318-23.
42. Kaneko K, Ito M, Takahashi T, Hashizume E, Owashi K, Kaneko H, et al. Computed tomography and scintigraphy vs. cardiac catheterization for coronary disease screening prior to noncardiac surgery. *Intern Med*. 2010;49:1703-10.
43. Mitchell GF, Parise H, Vita JA, Larson MG, Warner E, Keane JF Jr, et al. Local shear stress and brachial artery flow-mediated dilation: the Framingham Heart Study. *Hypertension*. 2004;44:134-9.
44. Vita JA, Keane JF, Larson MG, Keyes MJ, Massaro JM, Lipinska I, et al. Brachial artery vasodilator function and systemic inflammation in the Framingham Offspring Study. *Circulation*. 2004;110:3604-9.
45. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40:373-83.
46. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol*. 1994;47:1245-51.
47. Fleisher LA, Eagle KA. Clinical practice. Lowering cardiac risk in noncardiac surgery. *N Engl J Med*. 2001;345:1677-82.
48. Pinho C, Grandini PC, Gualandro DM, Calderaro D, Monachini M, Caramelli B. Multicenter study of perioperative evaluation for noncardiac surgeries in Brazil (EMAPO). *Clinics*. 2007;62:17-22.
49. Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg*. 1991;78:355-60.
50. Prytherc DR, Whiteley MS, Higgins B, Weaver PC, Prout WG, Powell SJ. POSSUM and Portsmouth POSSUM for predicting mortality. Physiological and operative severity score for the enumeration of mortality and morbidity. *Br J Surg*. 1998;85:1217-20.
51. Huang AL, Silver AE, Shvenke E, Schopfer DW, Jahangir E, Titas MA, et al. Predictive value of reactive hyperemia for cardiovascular events in patients with peripheral arterial disease undergoing vascular surgery. *Arterioscler Thromb Vasc Biol*. 2007;27:2113-9.
52. Calderaro D, Monachini MC, Vieira CL, Yu PC, Gualandro DM, Marques AC, et al. Reactive hyperemia revisited. *Arterioscler Thromb Vasc Biol*. 2008;28:e23-4; author reply e5.

53. Feringa HH, Bax JJ, Elhendy A, Jonge R, Lindemans J, Schouten O, et al. Association of plasma N-terminal pro-B-type natriuretic peptide with postoperative cardiac events in patients undergoing surgery for abdominal aortic aneurysm or leg bypass. *Am J Cardiol.* 2006;98:111-5.
54. Dernellis J, Panaretou M. Assessment of cardiac risk before non-cardiac surgery: brain natriuretic peptide in 1590 patients. *Heart.* 2006;92:1645-50.
55. Leibowitz D, Planer D, Rott D, Elitzur Y, Chajek-Shaul T, Weiss AT. Brain natriuretic peptide levels predict perioperative events in cardiac patients undergoing noncardiac surgery: a prospective study. *Cardiology.* 2008;110:266-70.
56. Cuthbertson BH, Card G, Croal BL, McNeilly J, Hillis GS. The utility of B-type natriuretic peptide in predicting postoperative cardiac events and mortality in patients undergoing major emergency non-cardiac surgery. *Anaesthesia.* 2007;62:875-81.
57. Cuthbertson BH, Amiri AR, Croal BL, Rajagopalan S, Alosairi O, Brittenden J, et al. Utility of B-type natriuretic peptide in predicting perioperative cardiac events in patients undergoing major non-cardiac surgery. *Br J Anaesth.* 2007;99:170-6.
58. Karthikeyan G, Moncur RA, Levine O, Heels-Ansdell D, Chan MT, Alonso-Coelho P, et al. Is a pre-operative brain natriuretic peptide or N-terminal pro-B-type natriuretic peptide measurement an independent predictor of adverse cardiovascular outcomes within 30 days of noncardiac surgery? A systematic review and meta-analysis of observational studies. *J Am Coll Cardiol.* 2009;54:1599-606.
59. Lurati Buse GA, Koller MT, Burkhart C, Seeberger MD, Filipovic M. The predictive value of preoperative natriuretic peptide concentrations in adults undergoing surgery: a systematic review and meta-analysis. *Anesth Analg.* 2011;112:1019-33.
60. Ryding AD, Kumar S, Worthington AM, Burgess D. Prognostic value of brain natriuretic peptide in noncardiac surgery: a meta-analysis. *Anesthesiology.* 2009;111:311-9.
61. Choi JH, Cho DK, Song YB, Hahn JY, Choi S, Gwon HC, et al. Preoperative NT-proBNP and CRP predict perioperative major cardiovascular events in non-cardiac surgery. *Heart.* 2010;96:56-62.
62. Heider P, Poppert H, Wolf O, Liebig T, Pelisek J, Schuster T, et al. Fibrinogen and high-sensitive C-reactive protein as serologic predictors for perioperative cerebral microembolic lesions after carotid endarterectomy. *J Vasc Surg.* 2007;46:449-54.
63. Heald CL, Fowkes FG, Murray GD, Price JF, Collaboration ABI. Risk of mortality and cardiovascular disease associated with the ankle-brachial index: systematic review. *Atherosclerosis.* 2006;189:61-9.
64. Fowkes FG, Murray GD, Butcher I, Heald CL, Lee RJ, Chambless LE, et al. Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA.* 2008;300:197-208.
65. Leng GC, Fowkes FG, Lee AJ, Dunbar J, Housley E, Ruckley CV. Use of ankle brachial pressure index to predict cardiovascular events and death: a cohort study. *BMJ.* 1996;313:1440-4.
66. Aboyans V, Lacroix P, Lebourdon A, Preux PM, Ferrières J, Laskar M. The intra- and interobserver variability of ankle-arm blood pressure index according to its mode of calculation. *J Clin Epidemiol.* 2003;56:215-20.
67. Endres HG, Hucke C, Holland-Letz T, Trampisch HJ. A new efficient trial design for assessing reliability of ankle-brachial index measures by three different observer groups. *BMC Cardiovasc Disord.* 2006;6:33.
68. Hiatt WR. Medical treatment of peripheral arterial disease and claudication. *N Engl J Med.* 2001;344:1608-21.
69. Mohler ER. Peripheral arterial disease: identification and implications. *Arch Intern Med.* 2003;163:2306-14.
70. Doobay AV, Anand SS. Sensitivity and specificity of the ankle-brachial index to predict future cardiovascular outcomes: a systematic review. *Arterioscler Thromb Vasc Biol.* 2005;25:1463-9.
71. Newman AB, Shemanski L, Manolio TA, Cushman M, Mittelmark M, Polak JF, Powe NR, et al. Ankle-arm index as a predictor of cardiovascular disease and mortality in the Cardiovascular Health Study. The Cardiovascular Health Study Group. *Arterioscler Thromb Vasc Biol.* 1999;19:538-45.
72. Lee AJ, Price JF, Russell MJ, Smith FB, van Wijk MC, Fowkes FG. Improved prediction of fatal myocardial infarction using the ankle brachial index in addition to conventional risk factors: the Edinburgh Artery Study. *Circulation.* 2004;110:3075-80.
73. Flu WJ, van Kuijk JP, Voute MT, Kuiper R, Verhagen HJ, Bax JJ, Poldermans D. Asymptomatic low ankle-brachial index in vascular surgery patients: a predictor of perioperative myocardial damage. *Eur J Vasc Endovasc Surg.* 2010;39:62-9.
74. Barnett S, Moonesinghe SR. Clinical risk scores to guide perioperative management. *Postgrad Med J.* 2011;87:535-41.