

## QUERCETIN, AN ENERGY TRANSFER INHIBITOR IN PHOTOPHOSPHORYLATION

Yasuo MUKOHATA, Shoji NAKABAYASHI and Mitsuhiro HIGASHIDA

*Department of Biology, Faculty of Science, Osaka University, Toyonaka, 560 Japan*

Received 3 November 1977

### 1. Introduction

Quercetin, 3,3',4',5,7-pentahydroxyflavone, was found to bind to either the  $\alpha$ - or the  $\beta$ -subunits [1] of isolated chloroplast coupling factor 1 (CF<sub>1</sub>).

It was also shown to inhibit the ATPase of mitochondria [2] or chloroplasts [3], but not oxidative phosphorylation [2].

In this report, we show that quercetin inhibits photophosphorylation as an energy transfer inhibitor in competition with the substrate nucleotide. Two quercetin molecules cooperatively bind to sites on the  $\beta$ -subunits of CF<sub>1</sub> in situ and thus inhibit phosphorylation.

### 2. Materials and methods

Chloroplasts were isolated from market spinach leaves in a choline medium by the procedure described [4]. Chlorophyll was determined by the method [5]. Quercetin (Wako Pure Chemicals, Osaka) was purified by repeated precipitation from ethanol-water.

The electron transport activity from H<sub>2</sub>O to methylviologen (50  $\mu$ M) was measured as oxygen uptake with an oxygen electrode (Beckmann; Field Lab). Phosphorylation was assayed by the method in [6] using 20  $\mu$ M PMS. A typical reaction mixture contained: 0.1 M sucrose, 5 mM MgCl<sub>2</sub>, 10 mM Tricine-NaOH at pH 8.3 and chloroplasts containing 20  $\mu$ g chlorophyll/ml. ADP, ATP, GDP, IDP, 1, *N*<sup>6</sup>-ethenoAMP and P<sub>i</sub> (<sup>32</sup>P]P<sub>i</sub> for phosphorylation assay) were added as required, together with methyl-

**Abbreviations:** CF<sub>1</sub>, chloroplast coupling factor 1; PMS, phenazine methosulfate

viologen or PMS. The ethanol solutions of quercetin were added to the reaction mixture maintaining the ethanol concentration in the mixture constant and less than 5%. The reaction mixture was illuminated by white light ( $5 \times 10^4$  lux) for 12–60 s at 15°C.

### 3. Results and discussion

Figure 1 shows that quercetin inhibits electron transport under phosphorylation conditions (ADP + P<sub>i</sub>), but not in the presence of ATP and P<sub>i</sub> (i.e. 'basal electron transport' [7]). The electron transport activity in the absence of any adenylate is little affected by quercetin. Thus, quercetin can not take the place of nucleoside di- or tri-phosphate [7–11] as




Fig. 1. The effect of quercetin on the rate of electron transport from H<sub>2</sub>O to methylviologen. The activity was measured under phosphorylation conditions (200  $\mu$ M ADP + 1 mM P<sub>i</sub>) or basal electron transport conditions (200  $\mu$ M ATP + 1 mM P<sub>i</sub>) or without addition (none).




Fig.2. Inhibition of photophosphorylation by quercetin. The concentration of nucleotides was as indicated, with  $P_i$  and PMS at 2.5 mM and 20  $\mu$ M, respectively. (A) inhibition profiles. (B) Hill plots, where  $v_0$  is the rate of phosphorylation in the absence of quercetin, and the line is calculated for a slope of 2.

the effector which binds to the  $\alpha$ -subunits [12] of  $CF_1$  altering its conformation [13,14]. Electron transport in the presence of quercetin can be assayed from  $H_2O$  to methylviologen or from reduced dichlorophenolindophenol to methylviologen, but not with other electron acceptors such as dimethylquinone or ferricyanide because of the latter's reaction (oxidation) with quercetin.

Figure 2 shows that quercetin inhibits photophosphorylation of nucleoside diphosphate in a sigmoidal fashion with respect to inhibitor concentration, as expected from fig.1. This inhibition profile resembles that of (modified  $CF_1$ ) ATPase [3]. The inhibition was greater for GDP and IDP than ADP when these nucleotides were examined at the same concentration. However, when these nucleotides were used at a concentration of  $10 \times K_m$  (fig.2), the inhibition profiles became almost coincident with each other. The small difference in profiles may be due to inaccuracies in the  $K_m$  values used [11]. Figure 2B shows that the apparent Hill constant for the inhibition by quercetin is two for ADP and somewhat larger for GDP and IDP.

A slight enhancement of phosphorylation (and also coupled electron transport (fig.1)) was sometimes observed at low quercetin concentrations around 10  $\mu$ M. A small enhancement was also found when

dimethylsulfoxide [3] was used as the solvent of quercetin. At very low concentrations, the inhibitor may activate the reaction [15] by replacing the nucleotide. Also, the solvents themselves were found to give a small enhancement of phosphorylation.

Figure 3A shows that quercetin inhibition of ADP phosphorylation is competitive with ADP. Figure 3B shows that  $[ADP]/\text{rate of phosphorylation}$  (the slope of the lines in fig.3A) is linearly related to the square of quercetin concentration. This relationship obeys the equation [15]:

$$v/V_{\max} = [ADP]/(K_m + [ADP] + K_m [Q]^2/K_{i1} \cdot K_{i2})$$

or after rearrangement,

$$[ADP] (1/v - 1/V_{\max}) = K_m (1 + [Q]^2/K_{i1} \cdot K_{i2})/V_{\max}$$

where  $[ADP]$  and  $[Q]$  represent concentrations of ADP and quercetin, respectively, and  $K_{i1}$  and  $K_{i2}$  the (intrinsic) inhibition constants of the first and the second quercetin, respectively. Although  $K_{i1} \gg K_{i2}$  (fig.2 and 3), if we assume  $K_{i1} = K_{i2}$ , we can obtain the apparent inhibition constant,  $K_i = 33 \mu$ M, using  $K_m = 35 \mu$ M and  $V_{\max} = 333 \mu\text{mol/mg}\cdot\text{chl}\cdot\text{h}$  (fig.3A). This  $K_i$  value agrees with the dissociation constant of 34  $\mu$ M [1] which was determined physico-chemically




Fig.3. Competitive inhibition of phosphorylation by quercetin with respect to ADP. The concentration of quercetin was as indicated, those of  $P_i$  and PMS being 1 mM and 20  $\mu$ M, respectively. The apparent values of  $K_m$  for ADP and  $V_{\max}$  are 35  $\mu$ M and 333  $\mu\text{mol/mg}\cdot\text{chl}\cdot\text{h}$ , respectively. (A) Lineweaver-Burk plots. (B) Plots of the slopes in fig.3A versus the square of quercetin concentration.




Fig. 4. Non-competitive inhibition of phosphorylation by quercetin with respect to phosphate. The concentrations of GDP and quercetin were as indicated. PMS was used at 20  $\mu\text{M}$ . The apparent value of  $K_m$  for  $\text{P}_i$  is 330  $\mu\text{M}$ . (A) Inhibition profiles. (B) Lineweaver-Burk plots.

for quercetin binding to isolated  $\text{CF}_1$  in competition with 8-anilino-1-naphthalenesulfonate.

Figure 4 shows that quercetin is a non-competitive inhibitor of phosphorylation with respect to  $\text{P}_i$ . Therefore, contrary to phlorizin which was shown to compete with  $\text{P}_i$  [16], quercetin is an energy transfer inhibitor which competes with the substrate nucleoside diphosphate.

The binding sites of quercetin were assumed to be located on either the  $\alpha$ - or the  $\beta$ -subunits of  $\text{CF}_1$  [1]. 1,  $N^6$ -EthenoAMP is a competitive [10] of nucleotide(s) which bind to the so-called tight binding site(s) [8–11] on the  $\alpha$ -subunit(s) [12]. When 1,  $N^6$ -ethenoAMP was also present with quercetin, the sigmoidicity of the curves of fig. 2A was slightly decreased but the  $[\text{Q}]^2$  relationship still held. Furthermore, the negligible effect of quercetin on the electron transport activities either in the presence of ATP (effector for the  $\alpha$ -subunit(s)) or in the absence of nucleotide (fig. 1) implies that quercetin does not interact, at least functionally, with the effector binding site(s) on the  $\alpha$ -subunit(s). Therefore, the binding sites for quercetin are most likely located on the  $\beta$ -subunits. Two quercetin molecules may be considered to bind cooperatively to the sites on the  $\beta$ -subunits, block the binding of nucleoside diphosphate and so inhibit phosphorylation.

In the absence of quercetin, two nucleoside di-

phosphate molecules bind to these two sites [17] on the  $\beta$ -subunits, and the initial binding of these two nucleotides to vacant sites on the  $\beta$ -subunits would also be expected to be cooperative. Such cooperative binding of two nucleotides has indeed been shown with isolated  $\text{CF}_1$  [18]. A pair of nucleotide-binding sites also fits the mechanism [19] proposed for quasi-arsenylation [20] in which an ATP–AMP pair is considered to be functionally equivalent to an ADP–ADP pair. Preliminary experiments indicate that quercetin inhibits quasi-arsenylation in competition with AMP.

### Acknowledgement

We thank Dr M. Avron of Weizmann Institute, Israel, for his valuable suggestions during preparation of the manuscript.

### References

- [1] Cantley, L. C., jr and Hammes, G. G. (1976) *Biochemistry* 15, 1–8.
- [2] Lang, D. R. and Racker, E. (1974) *Biochim. Biophys. Acta* 333, 180–186.
- [3] Deters, D. W., Racker, E., Nelson, N. and Nelson, H. (1975) *J. Biol. Chem.* 250, 1041–1047.
- [4] Mukohata, Y., Yagi, T., Matsuno, A., Higashida, M. and Sugiyama, Y. (1974) *Plant Cell Physiol.* 15, 163–167.
- [5] Arnon, D. I. (1948) *Plant Physiol.* 24, 1–15.
- [6] Asada, K., Takahashi, M. and Urano, M. (1972) *Anal. Biochem.* 48, 311–315.
- [7] Mukohata, Y., Yagi, T., Sugiyama, Y., Matsuno, A. and Higashida, M. (1975) *Bioenergetics* 7, 91–102.
- [8] Avron, M., Krogmann, D. W. and Jagendorf, A. T. (1958) *Biochim. Biophys. Acta* 30, 144–153.
- [9] Vambutas, V. and Bertsch, W. (1975) *ibid.* 376, 169–179.
- [10] Mukohata, Y. and Sugiyama, Y. (1976) *Plant Cell Physiol.* 17, 733–742.
- [11] Yagi, T. and Mukohata, Y. (1977) *J. Bioenerg. Biomembr.* 9, 31–40.
- [12] Nelson, N., Deters, D. W., Nelson, H. and Racker, E. (1973) *J. Biol. Chem.* 248, 2049–2055.
- [13] McCarty, R. E., Fuhrman, J. S. and Tsuchiya, Y. (1971) *Proc. Natl. Acad. Sci. USA* 68, 2522–2526.
- [14] Telfer, A. and Evans, M. C. W. (1972) *Biochim. Biophys. Acta* 256, 625–637.
- [15] Segel, I. H. (1975) in: *Enzyme Kinetics*, p. 385, John Wiley, New York.

- [16] Izawa, S., Winget, G. and Good, N. E. (1966) *Biochem. Biophys. Res. Commun.* 22, 223-226.
- [17] Higashida, M. and Mukohata, Y. (1976) *J. Biochem. (Tokyo)* 80, 1177-1179.
- [18] Nelson, N. (1976) *Biochim. Biophys. Acta* 456, 314-338.
- [19] Mukohata, Y. (1976) *Proc. II<sup>nd</sup> Symp. Japan Bioenergetics Group*, pp. 91-93 (in Japanese; English version in preparation).
- [20] Mukohata, Y. and Yagi, T. (1975) *Bioenergetics* 7, 111-120.