

## 217

**Potassium canrenoate prevents aldosterone-induced hypoxia-reoxygenation injury in isolated human right atrial myocardium in vitro**

Damien Legallois (1), Sandrine Lemoine (2), Lan Zhu (2), Guillaume Coutance (1), Vladimir Saplacan (3), Laurent Coulbault (4), Stéphane Allouche (4), Fabio Cuttone (3), Massimo Massetti (3), Jean-Luc Hanouz (2), Alain Manrique (2), Paul Milliez (1)  
 (1) CHU de Caen, service de cardiologie, Caen, France – (2) CHU de Caen, EA 4650, Caen, France – (3) CHU de Caen, service de chirurgie cardiaque, Caen, France – (4) CHU de Caen, service de biochimie, Caen, France

**Introduction:** Atrial fibrillation (AF) is the most common complication after cardiac surgery and is responsible for significant morbidity and mortality. To assess whether preconditioning with renin-angiotensin-aldosterone system agents could limit myocardial tissue injury, we tested the force of contraction of human right atrial trabeculae during hypoxia-reoxygenation when exposed to aldosterone and/or potassium canrenoate.

**Materials and methods:** We studied the effect of aldosterone and/or mineralocorticoid receptor (MR) antagonist potassium canrenoate administration prior to hypoxia-reoxygenation and in normoxic condition on force of contraction in 51 human right atrial trabeculae obtained from patients scheduled for coronary artery bypass surgery or aortic valve replacement. We also tested if aldosterone exposure results in changes in levels of ERK1/2 phosphorylation.

**Results:** The force of contraction of trabeculae was significantly reduced by aldosterone administration under normoxic or hypoxia-reoxygenation conditions. The rapid onset of this effect – 25 minutes – suggests a nongenomic mechanism. When compared with controls, potassium canrenoate by itself was not able to induce cardioprotection. However, coadministration of aldosterone and potassium canrenoate preconditioned isolated human atrial myocardium exposed to hypoxia – reoxygenation. This effect could be MR-dependent or not because both aldosterone and potassium canrenoate have some nongenomic MR-independent actions. Preliminary results showed that aldosterone exposure increased ERK1/2 phosphorylation levels.

**Conclusion:** Potassium canrenoate-induced cardioprotection prevents the deleterious effect of aldosterone on human atrial myocardial tissue. Further research required to determine the mechanism involved and the potential beneficial in clinical practice in order to reduce the incidence of postoperative atrial fibrillation.




Figure – Experimental conditions

January 19<sup>th</sup>, Saturday 2013

## 218

**Can the “Bleeding Academic Research Consortium” (BARC) classification be applied to pulmonary embolism?**

Nicolas Meneveau (1), Youssef El Omri (1), Romain Chopard (2), Marie France Séronde (2), Sebastien Janin (2), Philoktimon Plataras (1), Yvette Bernard (2), Francois Schiele (2)

(1) CHU Jean Minjot, Besançon, cardiologie, Besançon, France – (2) CHU Besançon, cardiologie, Besançon, France

**Background:** Bleeding is associated with higher risk of adverse outcomes in pulmonary embolism (PE). The “Bleeding Academic Research Consortium” (BARC) developed a classification of events combining laboratory & clinical parameters, but based on acute coronary syndrome patients. We investigated whether this classification is applicable in the context of PE, where no standard bleeding definition exists.

**Methods:** Prospective, single-center registry of patients with confirmed PE. We excluded BARC types 1 or 4 bleeding, considered not to be applicable to the context of PE. BARC type 2 bleeds were defined as any overt bleeding requiring non-surgical or medical care, or leading to hospitalisation. Type 3 bleeds were defined as drop of >3 g/dL in haemoglobin, any transfusion, tamponade, intra-cranial hemorrhage, or bleeding requiring surgical intervention. Type 5 bleeds were defined as any fatal bleed.

**Results:** From 2007 to 2011, 666 patients with confirmed PE were included; average age 66±18 years; 52% women; 25% low-risk, 61% intermediate-risk and 14% high-risk PE. Treatment was: unfractionated heparin in 93 (14%), enoxaparin in 200 (30%), fondaparinux in 373 (56%). Thrombolysis was given in 167 (25%). Sixty patients (9%) experienced bleeding (n=13, 43, 4 for BARC types 2, 3, 5 respectively). Main in-hospital events are shown in table 1. By multivariate analysis, independent predictors of in-hospital death were: cardiogenic shock (OR 12.6 [4.8-20.8]); chronic obstructive pulmonary disease (OR 5.27 [2.25-8.43]); acute right ventricular dysfunction (OR 2.98 [1.25-6.96]) and any bleeding (BARC 2,3,5) (OR 3.15 [1.34-7.37]).

**Conclusion:** Our data suggest that the BARC classification can be applied to acute PE and that bleeding is associated with unfavourable in-hospital outcome. We suggest use of the BARC as the standard for classification of bleeding events in PE.

Table – Results

	BARC type 2 (A)	BARC type 3/5 (B)	No bleed (n=606) (C)	p (A) vs (C)	p (B) vs (C)
Death	2 (15.4%)	7 (15%)	19 (3.1%)	0.034	0.0014
Recurrent PE	0	4 (8.5%)	5 (0.9%)	0.67	<0.0001
Treatment escalation	1 (7.7%)	2 (4.2%)	4 (0.7%)	0.017	0.028

## 219

**Prevalence, awareness, treatment and control of hypertension in a self-selected sub-Saharan African urban population: A cross-sectional study**

Anastase Dzudie Tamdja (1), Hamadou Ba (2), Andre Pascal Kengne (3), Alain Menanga (4), Charles Kouam Kouam (5), Joseph Abah (6), Felicite Kamdem (1), Yves Monkam Mbouende (1), Henry Luma (1), Walinjom Muna (4), Samuel Kingue (4)

(1) Hôpital Général de Douala, médecine interne/cardiologie, Douala, Cameroun – (2) Hôpital Régional de Graoua, Garoua, Cameroun – (3) (4) Department of Medicine, University of Cape Town & Medical research Council, Cape Town, Afrique du Sud – (4) Faculté de médecine et des sciences biomédicales, Université de Yaoundé I, Yaoundé, Cameroun – (5) Hôpital Régional de Bafoussam, Bafoussam, Cameroun – (6) Hôpital Militaire de Bamenda, Bamenda, Cameroun

**Objectives:** Hypertension has been established as a major public health problem in Africa, but its specific contributions to disease burden are still incompletely understood. We quantified the burden and determinants of hypertension, detection, treatment and control rates among adults in major cities in Cameroon.

**Patients and methods:** This was a cross-sectional and community-based multicenter study in major cities in Cameroon. Participants were self-selected urban dwellers from the Center, Littoral, North-West and West Regions, who attended on May 17th 2011 a screening campaign advertised through mass media. Primary outcome measure was hypertension defined as systolic (and/or diastolic) blood pressure  $\geq 140$  (90) mmHg, or ongoing blood pressure (BP) lowering medications.

**Results:** In all, 2120 participants (1003 women) were included. Among them, 1007 (prevalence rate 47.5%) had hypertension, including 319 (awareness rate 31.7%) who were aware of their status. The prevalence of hypertension increased with age overall and by sex and region. Among aware hypertensive subjects, 191 (treatment rate 59.9%) were on regular BP lowering medication, and among those treated, 47 (controlled rate 24.6%) were at target BP levels (i.e. systolic (and diastolic) BP  $< 140$  (90) mmHg). In multivariable logistic regressions analysis, male gender, advanced age, parental history of hypertension, diabetes mellitus, elevated waist, and elevated body mass index (BMI) were the significant predictors of hypertension. Likewise, male gender, high BMI and physical inactivity were associated with poor control.

**Conclusions:** High prevalence of hypertension with low awareness, treatment and control were found in this urban population; these findings are significant and alarming with consideration to the various improvements in the access to healthcare and the continuing efforts to educate communities over the past decades.

## 220

### Control of hypertension in a large population of Blida, Algeria

Abdelghani Bachir Cherif, Abdelhalim Taleb, Zahia Dahdouh, Djamilia Riahi, Mohamed Halimouche, Mohamed Temmar, Mohamed Bouafia  
*CHU Franz Fanon, cardiologie, Blida, Algérie*

**Objective:** We are presenting the main results of the study regarding the hypertension control rate in a large sample of hypertensive patients at cardiology center of Blida. Algeria

**Methods:** In the cross-sectional questionnaire-based observational study we have included 1184 hypertensive patients (56.3% females, mean age  $54.0 \pm 11.1$  years) attending a routine visit in specialist care. To be included patients had to be  $\geq 20$  years old and had to be treated for at least 12 months with antihypertensive drugs. Blood pressure (BP) was measured twice in a seated position according to ESH/ESC guidelines and mean value was calculated. The controlled hypertension was defined as BP level below 140/90 mmHg. Selected demographical and clinical data were evaluated.

**Results:** Among studied patients hypertension control rate was 36.7%. There were no significant differences in control rates between primary care and specialist care patients and between men and women. Patients with obesity (BMI  $\geq 30$  kg/m<sup>2</sup>) or abdominal obesity (ESH/ESC 2007 criteria) had lower rates of controlled hypertension as compared with patients without obesity or abdominal obesity (47.3% vs 52.1%,  $p < 0.01$  et 46.2% vs 51.7%,  $p < 0.001$  respectively). Patients with diabetes as well as patients with coronary artery disease (CAD) or cerebrovascular disease (CVD) had lower hypertension control rates in comparison with patients without these diseases (32.6% vs 38.9%;  $p < 0.001$ , 25.3% vs 29.1%,  $p < 0.001$  and 23.8% vs 36.2%;  $p = 0.03$  for diabetes, CAD and CVD respectively).

**Conclusion:** Our results showed that 36.7% of hypertensive patients treated for at least 1 year achieved blood pressure control. Patients at higher cardiovascular risk (established cardiovascular disease, diabetes or obesity) had lower BP control rates.

## 221

### Peripheral artery disease and description of risk factors profile in ambulatory patients with metabolic syndrome

Cristina Florescu, Cristina Ramona Barsan, Catalina Mariana Cojocaru, Laura Popescu, Rodica Musetescu  
*Centre de cardiologie Craiova, Craiova, Roumanie*

**Purpose:** The metabolic syndrome, a cluster of cardiovascular risk factors associated with insulin resistance and obesity has been associated with an increase in incidence of coronary artery disease, stroke and cardiovascular mortality. Less data are available on the relationship between the metabolic syndrome and incident lower extremity peripheral artery disease. Our aim is to evaluate the prevalence of peripheral artery disease (PAD) in ambulatory patients aged 45-65 years with metabolic syndrome (MS).

**Methods:** We included in our study 400 ambulatory patients (200 men and 200 women) who fulfilled criteria of metabolic syndrome. The diagnosis of PAD was established if ankle-brachial index (ABI) was  $< 0.9$ . We analyzed risk factors profile in these patients: body mass index (BMI), physical inactivity, smoking, hypertension, dyslipidemia, abnormal glucose metabolism and subclinical inflammatory status.

**Results:** Smoking was found in 38% of males and 21% of females. Among these patients with MS, an HDL-Chol  $< 50$  mg/dl was the most prevalent MS trait: 87% in males and 83% in females. This trait was followed by elevated BMI in 80% of males and 84% of females. Also, elevated blood pressure was found in 79% of males and 75% of females. Elevated triglycerides were present in 80% of males and 74% of females. Dysglycemia was found in 29% of males and 24% of females.

Among patients with metabolic syndrome, the prevalence of PAD was of 12% in males and 6% in females, correlated with smoking (87.5% in males and 91.66% in females with PAD), physical inactivity (79.16% in males and 91.66% in females), dysglycemia (75% in males and 83.33% in females), age (66.66% in males aged  $> 55$  years and 83.33% in females aged  $> 55$  years, total cholesterol (79.16% in males and 83.33% in females) and high hs-CRP (70.83% in males and 75% in females).

**Conclusion:** MS is associated with a risk of PAD, an important role playing smoking, dysglycemia, dyslipidemia and subclinical inflammation.

## 222

### Relation between pulse pressure and vascular event among the elderly in Monastir: a population-based study

Sonia Hammami (1), Said Hajem (2), Nadia Koubaa (1), Amel Nkbi (1), Khaldoun Ben Hamda (3), Mohamed Hammami (1)  
*(1) Faculté de Médecine Monastir, nutrition humaine et désordres métaboliques, Monastir, Tunisie – (2) CHU F Bourguiba Monastir, Tunisie, Tunisie – (3) CHU F Bourguiba, Cardiologie, Monastir, Tunisie*

**Background:** The sixth Joint National Committee (JNC-VI) classification system of blood pressure emphasizes both systolic blood pressure (SBP) and diastolic blood pressure (DBP) for cardiovascular disease risk assessment. Pulse pressure (PP)=[SBP – DBP], may also be a valuable risk assessment tool.

**Objectives:** In this study, we examined the association of cardiovascular events with arterial pulse pressure in elderly as well as their correlations within diabetes.

**Patients and methods:** A Population based survey supported by WHO and FNUAP, was undertaken in 2008-2009 to investigate health behaviours and health status of elderly living in their home in Monastir. This study was carried out in a representative sample of elderly aged more than 65 years. Standardized techniques were used for blood pressure (BP). Hypertension was defined as BP more than 140/90 mmHg. Diabetes was defined as known diabetics. Regression analyses were used to examine the relationships among pulse pressure age, diabetes and cardiovascular events (myocardial infarction, arteritis and stroke) This project was approved by the research Ethics Committee, CHU F Bourguiba, university of Monastir

**Results:** The study included 598 participants (396 women and 202 men) who were aged  $\geq 65$  years.