



## Concerted action of activation-induced cytidine deaminase and uracil-DNA glycosylase reduces covalently closed circular DNA of duck hepatitis B virus

Sajeda Chowdhury, Kouichi Kitamura\*, Miyuki Simadu, Miki Koura, Masamichi Muramatsu

Department of Molecular Genetics, Kanazawa University, Graduate School of Medical Sciences, Kanazawa, Japan



### ARTICLE INFO

#### Article history:

Received 25 June 2013

Accepted 24 July 2013

Available online 15 August 2013

Edited by Hans-Dieter Klenk

#### Keywords:

AID  
APOBEC  
Base excision repair  
Hepatitis B virus  
cccDNA

### ABSTRACT

**Covalently closed circular DNA (cccDNA) forms a template for the replication of hepatitis B virus (HBV) and duck HBV (DHBV). Recent studies suggest that activation-induced cytidine deaminase (AID) functions in innate immunity, although its molecular mechanism of action remains unclear, particularly regarding HBV restriction. Here we demonstrated that overexpression of chicken AID caused hypermutation and reduction of DHBV cccDNA levels. Inhibition of uracil-DNA glycosylase (UNG) by UNG inhibitor protein (UGI) abolished AID-induced cccDNA reduction, suggesting that the AID/UNG pathway triggers the degradation of cccDNA via cytosine deamination and uracil excision.**

© 2013 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

### 1. Introduction

The covalently closed circular DNA (cccDNA) of the hepatitis B virus (HBV) plays an essential role in chronic hepatitis. cccDNA exists as an episome in the nucleus of infected hepatocytes and transcribes all viral RNAs, including pregenomic (pg) RNA. pgRNA is assembled into the nucleocapsid (NC) in the cytoplasm and is then converted into a relaxed circular DNA (rcDNA) by viral polymerase activity. Mature NCs are secreted after combining with envelope proteins or translocated to the nucleus. In the nucleus, the rcDNA is again converted into cccDNA for the next replication cycle. Only one or a few copies of cccDNA per infected cell are stably maintained by an unknown mechanism [1].

Despite the crucial role of cccDNA in viral persistence, little is known about the host factors that target this viral nuclear intermediate owing to the absence of an efficient experimental system that can produce HBV cccDNA at a level sufficient for analysis. In view of the limitation of HBV in vitro systems, the duck HBV (DHBV)

*Abbreviations:* AID, activation-induced cytidine deaminase; APOBEC, apolipoprotein B mRNA-editing enzyme catalytic subunit; UNG, uracil-DNA glycosylase; DHBV, duck hepatitis B virus; cccDNA, covalently closed circular DNA; 3D-PCR, differential denaturation DNA PCR

\* Corresponding author. Address: Department of Molecular Genetics, Kanazawa University, Graduate School of Medical Sciences, 13-1 Takara-machi, Kanazawa 920-8640, Japan. Tel.: +81 76 265 2176; fax: +81 76 234 4225.

E-mail address: [kkita@med.kanazawa-u.ac.jp](mailto:kkita@med.kanazawa-u.ac.jp) (K. Kitamura).

model has been commonly used to study HBV replication [2]. Recently, we reported that an anti-viral factor APOBEC3G (A3G) induces hypermutation on DHBV cccDNA [3]. A3G belongs to the APOBEC (apolipoprotein B mRNA-editing catalytic polypeptide) family of proteins. In humans, this family comprises at least 11 members including AID and APOBEC-1, -2, -3A, -3B, -3C, -3DE, -3F, -3G, -3H and -4. These AID/APOBEC proteins have enzymatic activity that can deaminate a cytosine base in DNA and/or RNA, thereby generating a uracil base. A3G is known to restrict the replication of retroviruses, including human immunodeficiency virus type 1 (HIV-1), HBV and retrotransposable elements [4–6]. AID was initially identified as an essential factor for class-switch recombination (CSR) and somatic hypermutation (SHM) of immunoglobulin genes in B cells [7,8]. It was also found that AID is expressed in response to cytokine stimulation in human hepatocytes [9] and HBV patients [10], induces HBV hypermutation in NC-DNA [10] and pgRNA [11] and restricts viral replication [12]. It remains to be established whether AID targets hepadnavirus cccDNA.

UNG is a host nuclear factor that removes uracil bases from uracilated DNA and initiates base excision repair (BER), resulting in DNA repair or cleavage [13]. The involvement of UNG activity in A3G-induced viral restriction has been vigorously studied in the HIV field, but remains a subject of debate [14–17]. We previously demonstrated that UNG repairs DHBV cccDNA hypermutation induced by A3G [3]. The involvement of UNG in the

anti-HBV activity of AID is of interest because AID is well known to initiate CSR and SHM by coupling with downstream UNG-directed BER pathways [13,18].

Here we investigated the effects of chicken AID on DHBV replication using an *in vitro* DHBV replication system with a chicken hepatoma cell line LMH, which we utilized to analyse cccDNA formation and function [2,19,20]. We found that the expression of chicken AID in LMH cells resulted in the reduction of DHBV NC-DNA and cccDNA. Furthermore, UNG inhibition abolished AID-induced reduction of cccDNA but not NC-DNA, suggesting that viral DNA degradation is induced by the AID/UNG pathway in the nucleus.

## 2. Materials and methods

### 2.1. Plasmids

The DHBV replicon plasmid pCSD3.5 $\Delta$ S [21] and UGI expression vector pIRES-UGI [3] were described previously. The pCSD3.5 $\Delta$ S plasmid produces a DHBV surface-deficient mutant that accumulates cccDNAs more efficiently than wild-type DHBV [20,21]. Other expression vectors are described in the [Supplementary methods](#).

### 2.2. Cell culture and transfection

LMH cells were maintained in Dulbecco's modified Eagle medium (DMEM; Sigma, St. Louis, MO, USA) containing 10% foetal bovine serum, 100 U/mL penicillin and 100  $\mu$ g/mL streptomycin. Plasmid transfection was performed using Fugene 6 (Roche, Basel, Switzerland), according to the manufacturer's instructions. The total DNA amount (6  $\mu$ g for a 60-mm dish) for each transfection was kept constant by adding mock vector as needed. The ER-fusion protein that was expressed as an inactive form was activated by tamoxifen (OHT, Wako Pure Chemical Industries, Osaka, Japan) treatment [22]. In the subcellular localization experiments of GFP fusion proteins, cells were treated with 10 ng/mL leptomycin B (LMB, Sigma) for 4 h.

### 2.3. Purification of viral DNAs

Cells were lysed in 0.5% NP40 lysis buffer, and nuclei were collected by centrifugation to separate the cytoplasmic and nuclear fractions. Cytoplasmic supernatants were treated with DNase I and RNase A and digested with proteinase K and sodium dodecyl sulphate (SDS) to extract viral NC-DNA. A modified Hirt extraction procedure was used to extract cccDNA from the nuclear fraction [3]. The nuclear pellet was lysed in 50 mM Tris-HCl (pH 7.5), 10 mM EDTA and 2% SDS. After a 20-min incubation at room temperature, 0.5 M KCl was added to the lysate and incubated at 4°C overnight. All purified DNA solutions were treated with DpnI restriction enzyme to digest any transfected plasmid DNA.

### 2.4. Mutation analysis

Hypermutation was detected using differential DNA denaturation PCR (3D-PCR) with DHBV cccDNA-selective primers. The specific amplification of cccDNA with cccDNA-selective primers was verified previously [3]. 3D-PCR was performed as follows: initial denaturation for 5 min at 94–83°C, followed by 35 cycles of 30 s at 94–83°C, 30 s at 55°C and 2 min at 72°C, with a final elongation for 7 min at 72°C. To determine the hypermutation frequency, PCR fragments were cloned into T vectors (Promega, Madison, WI, USA). Successfully recombined clones were randomly selected and sequenced using ABI PRISM 3130 (Applied Biosystems, Foster City, CA, USA).

### 2.5. Southern blotting

Purified DHBV DNAs were analysed by Southern blotting using double-stranded DHBV DNA probes spanning the entire viral genome. Probe labelling and signal development were performed using the AlkPhos direct labelling system (GE Healthcare, Piscataway, NJ, USA), and signals were detected using the LAS 1000 imager system (GE Healthcare).

### 2.6. Rolling circle amplification (RCA)

Specific amplification of DHBV cccDNA by RCA method was described previously [3,23]. The RCA concatemeric product was converted to the monomeric full-length DHBV genome by digestion with EcoRI, which cuts the DHBV cccDNA once. After electrophoresis, the DNA fragments were analysed by Southern blotting.

### 2.7. Western blotting

Transgene expression was analysed by Western blotting as described previously [3]. The following antibodies were used in this study: rabbit anti-GAPDH (G9545, Sigma), mouse anti-FLAG (M2, Sigma), rabbit anti-ER (HC-20, Santa Cruz Biotechnology, CA, USA) and anti-rabbit Igs-horseradish peroxidase (HRP) (AL13404, eBioscience, San Diego, CA, USA).

## 3. Results

### 3.1. Chicken AID induced cccDNA reduction via a UNG-dependent pathway

A DHBV replication system in a chicken hepatoma cell line LMH was utilized to analyse cccDNA. With this *in vitro* system, we could detect cccDNA by Southern blotting of nuclear Hirt extracts (Fig. 1A). The three DNA forms (re, dl and cccDNA) in the nuclear Hirt extract were all converted into dlDNA position as 3.0-kb bands after EcoRI digestion (Fig. 1A). To investigate the anti-DHBV activity of AID, a chicken AID expression vector and DHBV replicon plasmid pCSD3.5 were transfected into LMH cells. Concurrently, the effect of UNG inhibition was also tested by co-transfection of a UGI expression vector. UGI is an irreversible inhibitory protein that forms a stable complex with UNG. We previously demonstrated a 91% reduction of UNG activity under the conditions shown in Fig. 1B [3]. Six days after transfection, cytoplasmic nucleocapsid (NC)-DNAs and nuclear Hirt-extracted DNAs were purified. Southern blotting showed that AID overexpression reduced DHBV NC-DNA and cccDNA (Fig. 1B). UNG inhibition by UGI abolished AID-induced reduction of cccDNA but not NC-DNA. We confirmed that the protein level of chicken AID was not affected by UGI overexpression (Fig. 1C).

To determine cccDNA level at an earlier time point, we used RCA, a highly sensitive method widely used to measure the level of closed circular DNA of various viral species [24]. Three days after transfection, cccDNA from nuclear Hirt-extracted DNAs was amplified by RCA and digested with EcoRI to cleave the concatemer into monomers of full-length viral genomes (3.0 kb). Amplified cccDNA was detected by Southern blotting. Consistent with data obtained from day 6 (Fig. 1B), AID overexpression reproducibly suppressed both NC-DNA and cccDNA levels and UGI abolished the reduction of only cccDNA at day 3 (Fig. 2A and B).

### 3.2. Chicken AID induced cccDNA hypermutation

Next, we investigated the hypermutation activity of AID for DHBV cccDNA by 3D-PCR using cccDNA-specific primers




(Fig. 3A). 3D-PCR can detect hypermutated DNA by gradually decreasing the denaturation temperature [25]. Our results showed that only AID-UGI co-expression resulted in lower temperature amplification (83.9 and 83.0°C, indicated by triangles in Fig. 3B). Sequence analysis of DNA amplified at the lower denaturation temperature revealed G-to-A and C-to-T hypermutations (Fig. 3C and D and Fig. S1). Dinucleotide context analysis of the hypermutated cccDNA showed a preference for GpC and ApC dinucleotides as a mutation target (Fig. 3E), which is consistent with a previous report of human AID-induced HBV hypermutation [10].

### 3.3. Chicken AID is a protein that shuttles between the cytoplasm and nucleus


Chicken AID has a nuclear localization signal (NLS) in the N-terminus and a nuclear export signal (NES) in the C-terminus [26]. To determine the subcellular localization of chicken AID in chicken hepatocytes, an AID-GFP vector was transfected into LMH cells. Similar to human AID [27], chicken AID-GFP protein localized predominantly in the cytoplasm (Fig. 4A). Cells were treated with leptomycin B (LMB) to inhibit exportin 1-dependent nuclear export [27,28]. LMB treatment for 4 h changed the localization of AID from the cytoplasm to both cytoplasm and nuclear. Consistent with this observation, deletion of C-terminal residues containing NES (AID $\Delta$ NES in Fig. 4B, corresponding to JP8Bdel human mutant) resulted in the nuclear localization of AID with minor cytoplasmic retention (Fig. 4A). These observations indicate that chicken AID shuttles between the cytoplasm and nucleus in LMH cells and that the subcellular localization pattern of AID is conserved in mammals and chicken.

### 3.4. Nuclear retention of AID enhanced cccDNA-reduction activity

Because the majority of AID $\Delta$ NES-GFP protein was localized mainly to the nucleus, we examined the cccDNA-reduction activity of this mutant. Wild-type AID or AID $\Delta$ NES was co-transfected with a DHBV plasmid. Southern blotting following RCA showed that AID $\Delta$ NES had a stronger DHBV cccDNA-reduction activity than wild-type AID, whereas reduction of cytoplasmic NC-DNA levels by AID $\Delta$ NES was slightly weaker than that by wild-type AID



**Fig. 4.** Chicken AID is a protein that shuttles between the cytoplasm and nucleus. (A) Chicken AID-GFP or AID $\Delta$ NES-GFP was transfected into LMH cells. The nucleus was visualized by co-transfection of the DsRed-NLS vector. LMB was treated for 4 h to block nuclear export. LMB treatment caused accumulation of AID-GFP in the nucleus. (B) Domain structure of chicken AID. Numbers indicate amino acid residue positions. NLS, nuclear localization signal; NES, nuclear export signal.




**Fig. 5.** Deletion of NES domain enhanced the cccDNA-reduction activity of chicken AID. (A) Southern blotting for cytoplasmic NC-DNA and RCA-amplified cccDNA. Three days after transfection (OHT stimulation for 2 days), DHBV DNAs were analysed. (B) Exogenous protein expression of wild-type AID- and AID $\Delta$ NES-ER was detected by Western blotting with anti-ER antibody. Expression of GAPDH is also shown to demonstrate equivalent protein loading.

(Fig. 5A). Protein levels of AID $\Delta$ NES were equivalent to that of wild-type AID (Fig. 5B). These data suggest that nuclear-localized AID is responsible for cccDNA reduction.

## 4. Discussion

In this study, we overexpressed chicken AID and inhibited UNG activity in LMH cells to evaluate anti-viral roles of AID and UNG against DHBV. We found that AID reduced both DHBV NC-DNA and cccDNA levels, and UNG inhibition abolished the reduction of cccDNA but not NC-DNA (Figs. 1 and 2). The UNG inhibition also revealed the AID-induced G-to-A and C-to-T hypermutation in cccDNA (Fig. 3). Furthermore, a nuclear-localized AID mutant more strongly reduced cccDNA levels than wild-type AID (Fig. 5). These results suggest that AID introduces C-to-U conversion in both strands of cccDNA and that UNG-mediated BER may degrade uracilated cccDNA (Fig. 6).

This study suggests that the molecular mechanism of AID-induced NC-DNA reduction could be different from that for cccDNA because it was found to be UNG-independent. We previously showed a physical association between human AID and HBV core proteins [11], suggesting the encapsidation of AID proteins in NC as reported in the case of A3G. Unlike HIV, UNG incorporation was not detected in HBV NC [3], implying no or little contribution



**Fig. 6.** A proposed model for AID-induced DHBV cccDNA reduction. Chicken AID shuttles between the cytoplasm and nucleus. Cytoplasmic AID reduces NC-DNA levels, presumably by an UNG-independent mechanism. Nuclear AID deaminates cccDNA, and then uracilated cccDNA can be cleaved by the UNG-mediated BER pathway. Previous HBV studies demonstrated that human AID hypermutates both pgRNA and NC-DNA [10,11].



of BER-mediated DNA cleavage in NC-DNA restriction. Similar to A3G [29,30], AID may suppress NC-DNA formation through deaminase-independent activity such as inhibition of reverse transcription. We also obtained the result, suggesting that human P19-AID mutant carrying a mutation in the deaminase motif [31] has the ability to reduce HBV NC-DNA as efficiently as wild-type AID (unpublished). In contrast, when AID deaminates cccDNA in nucleus, UNG-mediated BER can efficiently excise uracils from cccDNA and cleave DNA, similar to the case of the immunoglobulin gene undergoing CSR [13,18]. This can be a reason why cccDNA reduction was abolished by UNG inhibition.

Our previous study revealed that A3G reduces both DHBV cccDNA and NC-DNA [3]. In contrast to AID, cccDNA reduction by A3G was not abolished when UNG was inhibited. It is likely that AID has a different activity from A3G to suppress cccDNA. cccDNA reduction by A3G may simply be reflected by the reduction of NC-DNA, whereas de novo cccDNA degradation could be triggered by nuclear-localized AID followed by UNG activity (Fig. 6). Enhancement of cccDNA-reduction activity in the AID $\Delta$ NES mutant (Fig. 5) supports the idea that AID is directly targeted to nuclear cccDNA. Furthermore, mutation analysis showed AID-induced hypermutation in both strands of cccDNA (Fig. 3), which is different from the A3G-induced hypermutation exclusively occurring in the minus strand DNA [3]. cccDNA reduction mediated by the AID/UNG pathway can be explained by double-strand breaks initiated by deamination of both strands. AID is proposed to recruit UNG to the immunoglobulin gene to cause CSR [32]. We speculate that such a property may also underlie the different outcomes in cccDNA reduction observed between AID and A3G.

Recently, Vartanian et al. clearly demonstrated that APOBEC3 proteins are major deaminases that cause HBV NC-DNA hypermutation in vivo using dinucleotide context analysis of hypermutated NC-DNA obtained from patients [10]. However, the contribution of AID to cccDNA reduction and deaminase-independent NC-DNA reduction was still unknown in vivo. Our data suggests cccDNA as a novel target of the anti-HBV activity of AID for the first time. Further studies are required to reveal the mechanism of AID/UNG-mediated HBV cccDNA degradation in vitro and in vivo.

## Acknowledgements

This work was supported by the Funding Program for Next Generation World-Leading Researchers (NEXT project), MEXT KAKENHI Grant Number 20012018 (to M.M.) and JSPS KAKENHI Grant Number 21790654, 23790780 (to K.K.) of Japan. We thank Dr. K. Kinoshita for helpful discussions.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.febslet.2013.07.055>.

## References

- [1] Zhang, Y.Y., Zhang, B.H., Theele, D., Litwin, S., Toll, E. and Summers, J. (2003) Single-cell analysis of covalently closed circular DNA copy numbers in a hepadnavirus-infected liver. *Proc. Natl. Acad. Sci. USA* 100, 12372–12377.
- [2] Funk, A., Mhamdi, M., Will, H. and Sirma, H. (2007) Avian hepatitis B viruses: molecular and cellular biology, phylogenesis, and host tropism. *World J. Gastroenterol.* 13, 91–103.
- [3] Kitamura, K., Wang, Z., Chowdhury, S., Simadu, M., Koura, M. and Muramatsu, M. (2013) Uracil DNA glycosylase counteracts APOBEC3G-induced hypermutation of hepatitis B viral genomes: excision repair of covalently closed circular DNA. *PLoS Pathog.* 9, e1003361.
- [4] Goila-Gaur, R. and Strebel, K. (2008) HIV-1 Vif, APOBEC, and intrinsic immunity. *Retrovirology* 5, 51.

- [5] Harris, R.S. and Liddament, M.T. (2004) Retroviral restriction by APOBEC proteins. *Nat. Rev. Immunol.* 4, 868–877.
- [6] Malim, M.H. (2009) APOBEC proteins and intrinsic resistance to HIV-1 infection. *Philos. Trans. R Soc. Lond. B Biol. Sci.* 364, 675–687.
- [7] Muramatsu, M., Nagaoka, H., Shinkura, R., Begum, N.A. and Honjo, T. (2007) Discovery of activation-induced cytidine deaminase, the engraver of antibody memory. *Adv. Immunol.* 94, 1–36.
- [8] Muramatsu, M., Kinoshita, K., Fagarasan, S., Yamada, S., Shinkai, Y. and Honjo, T. (2000) Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* 102, 553–563.
- [9] Endo, Y. et al. (2007) Expression of activation-induced cytidine deaminase in human hepatocytes via NF- $\kappa$ B signaling. *Oncogene* 26, 5587–5595.
- [10] Vartanian, J.P. et al. (2010) Massive APOBEC3 editing of hepatitis B viral DNA in cirrhosis. *PLoS Pathog.* 6, e1000928.
- [11] Liang, G. et al. (2013) RNA editing of hepatitis B virus transcripts by activation-induced cytidine deaminase. *Proc Natl Acad Sci U S A* 110, 2246–2251.
- [12] Jost, S., Turelli, P., Mangeat, B., Protzer, U. and Trono, D. (2007) Induction of antiviral cytidine deaminases does not explain the inhibition of hepatitis B virus replication by interferons. *J. Virol.* 81, 10588–10596.
- [13] Sousa, M.M., Krokan, H.E. and Slupphaug, G. (2007) DNA-uracil and human pathology. *Mol. Aspects Med.* 28, 276–306.
- [14] Kaiser, S.M. and Emerman, M. (2006) Uracil DNA glycosylase is dispensable for human immunodeficiency virus type 1 replication and does not contribute to the antiviral effects of the cytidine deaminase APOBEC3G. *J. Virol.* 80, 875–882.
- [15] Langlois, M.A. and Neuberger, M.S. (2008) Human APOBEC3G can restrict retroviral infection in avian cells and acts independently of both UNG and SMUG1. *J. Virol.* 82, 4660–4664.
- [16] Mbisa, J.L. et al. (2007) Human immunodeficiency virus type 1 cDNAs produced in the presence of APOBEC3G exhibit defects in plus-strand DNA transfer and integration. *J. Virol.* 81, 7099–7110.
- [17] Yang, B., Chen, K., Zhang, C., Huang, S. and Zhang, H. (2007) Virion-associated uracil DNA glycosylase-2 and apurinic/aprimidinic endonuclease are involved in the degradation of APOBEC3G-edited nascent HIV-1 DNA. *J. Biol. Chem.* 282, 11667–11675.
- [18] Di Noia, J.M. and Neuberger, M.S. (2007) Molecular mechanisms of antibody somatic hypermutation. *Annu. Rev. Biochem.* 76, 1–22.
- [19] Schultz, U., Grgacic, E. and Nassal, M. (2004) Duck hepatitis B virus: an invaluable model system for HBV infection. *Adv. Virus Res.* 63, 1–70.
- [20] Kock, J., Rosier, C., Zhang, J.J., Blum, H.E., Nassal, M. and Thoma, C. (2010) Generation of covalently closed circular DNA of hepatitis B viruses via intracellular recycling is regulated in a virus specific manner. *PLoS Pathog.* 6, e1001082.
- [21] Summers, J., Smith, P.M., Huang, M.J. and Yu, M.S. (1991) Morphogenetic and regulatory effects of mutations in the envelope proteins of an avian hepadnavirus. *J. Virol.* 65, 1310–1317.
- [22] Doi, T., Kinoshita, K., Ikegawa, M., Muramatsu, M. and Honjo, T. (2003) De novo protein synthesis is required for the activation-induced cytidine deaminase function in class-switch recombination. *Proc. Natl. Acad. Sci. USA* 100, 2634–2638.
- [23] Margeridon, S., Carrouee-Durantel, S., Chemin, I., Barraud, L., Zoulim, F., Trepo, C. and Kay, A. (2008) Rolling circle amplification, a powerful tool for genetic and functional studies of complete hepatitis B virus genomes from low-level infections and for directly probing covalently closed circular DNA. *Antimicrob. Agents Chemother.* 52, 3068–3073.
- [24] Johne, R., Muller, H., Rector, A., van Ranst, M. and Stevens, H. (2009) Rolling-circle amplification of viral DNA genomes using phi29 polymerase. *Trends Microbiol.* 17, 205–211.
- [25] Suspene, R., Henry, M., Guillot, S., Wain-Hobson, S. and Vartanian, J.P. (2005) Recovery of APOBEC3-edited human immunodeficiency virus G $\rightarrow$ A hypermutants by differential DNA denaturation PCR. *J. Gen. Virol.* 86, 125–129.
- [26] Wakae, K., Magor, B.G., Saunders, H., Nagaoka, H., Kawamura, A., Kinoshita, K., Honjo, T. and Muramatsu, M. (2006) Evolution of class switch recombination function in fish activation-induced cytidine deaminase, AID. *Int. Immunol.* 18, 41–47.
- [27] Ito, S., Nagaoka, H., Shinkura, R., Begum, N., Muramatsu, M., Nakata, M. and Honjo, T. (2004) Activation-induced cytidine deaminase shuttles between nucleus and cytoplasm like apolipoprotein B mRNA editing catalytic polypeptide 1. *Proc. Natl. Acad. Sci. USA* 101, 1975–1980.
- [28] Nishi, K., Yoshida, M., Fujiwara, D., Nishikawa, M., Horinouchi, S. and Beppu, T. (1994) Leptomycin B targets a regulatory cascade of crml, a fission yeast nuclear protein, involved in control of higher order chromosome structure and gene expression. *J. Biol. Chem.* 269, 6320–6324.
- [29] Nguyen, D.H., Gummuru, S. and Hu, J. (2007) Deamination-independent inhibition of hepatitis B virus reverse transcription by APOBEC3G. *J. Virol.* 81, 4465–4472.
- [30] Noguchi, C. et al. (2007) Dual effect of APOBEC3G on Hepatitis B vims. *J. Gen. Virol.* 88, 432–440.
- [31] Ta, V.T. et al. (2003) AID mutant analyses indicate requirement for class-switch-specific cofactors. *Nat. Immunol.* 4, 843–848.
- [32] Ranjit, S., Khair, L., Linehan, E.K., Ucher, A.J., Chakrabarti, M., Schrader, C.E. and Stavnez, J. (2011) AID binds cooperatively with UNG and Msh2–Msh6 to Ig switch regions dependent upon the AID C terminus. *J. Immunol.* 187, 2464–2475.