INTERNATIONAL JOURNAL OF MYCOBACTERIOLOGY 4 (2015) 98-99



Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO



Comparative evaluation of LAMP and Nested-PCR for the diagnosis of bovine paratuberculosis



Shahabeddin Safi ^a, Orkideh Heidarnejhad ^{b,*}, Nader Mosavari ^c, Mehdi Sakha ^d, Davoud Afshar ^e, Lale Moazami ^f, Mohse Meshkat ^f, Rohollah Keshavarz ^c, Mohammad Taheri ^c, Reza Aref Pajoohi ^c, Keyvan Tadayon ^c, Samerand Reshadi ^c, Shojaat Dashtipour ^c

- ^a Department of Pathobiology, Faculty of Specialized Veterinary Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran
- ^b Department of Veterinary, Institute of Applied Scientific Higher Education of Jihad-e Agriculture, Iran
- ^c Department of Tuberculosis, Razi Vaccine & Serum Research Institute, Tehran, Iran
- ^d Department of Clinical Sciences, Faculty of Specialized Veterinary Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran
- ^e Department of Pathobiology, School of Public Health, Tehran University of Medical Science, Tehran, Iran

ARTICLEINFO

Article history:
Received 6 September 2014
Received in revised form
11 September 2014
Accepted 13 September 2014
Available online 30 October 2014

Keywords:
Loop mediated isothermal
amplification (LAMP)
Mycobacterium avium subsp.
paratuberculosis
Nested-PCR
Paratuberculosis
Bovine
Johne's disease

ABSTRACT

Introduction: Mycobacterium avium subsp. paratuberculosis causes paratuberculosis (Johne's disease), a systemic infection and chronic inflammation of the intestine that affects many species, including bovine. Infection is widespread in livestock, and human populations are exposed. A possible association between MAP infection and Crohn's disease in humans has been also described. Effective control of paratuberculosis has hampered due to lake of rapid and accurate diagnostic test. Range of diagnostic tests is available, but all have inborn limitations. The present study was designed to develop a loop-mediated isothermal amplification (LAMP) assay for the rapid and simple detection of Mycobacterium avium subsp. paratuberculosis (MAP).

Materials and methods: Six primers were specially designed for recognizing eight distinct sequence of insertion sequence 900 (IS900). To determine the sensitivity of the LAMP assay, 10-fold serial dilutions were made from 431 ng/ μ l MAP stock solution and compared with Nested-PCR results obtained using similar templates at identical concentrations. Detection limit of the LAMP was defined as the last positive dilution and the reactions were performed four times to examine the reproducibility of the test. The specificity of the assays were evaluated by testing three Gram-positive bacteria including Mycobacterium bovis AN5, Mycobacterium tuberculosis DT and Mycobacterium avium avium.

Results: Sensitivity of this assay for detection of DNA of MAP was 4 fg/µl and the specificity was 100%. This assay successfully detected MAP not only in the bacterial cultures but also in clinical fecal samples and the specificity of both PCR was 100%. This LAMP method is performed under isothermal conditions and no special apparatus is needed. In addition, its reactivity is directly observed with the naked eye without electrophoresis either as

E-mail address: heidarnejhad@yahoo.com (O. Heidarnejhad). http://dx.doi.org/10.1016/j.ijmyco.2014.09.015

^f Iran Veterinary Organizations, Iran

^{*} Corresponding author.

turbidity or in the form of a color change when SYBR Green 1, a fluorescent dsDNA intercalating dye, is employed.

Conclusions: This assay is rapid which requires nearly 1 h for detection of MAP, low in cost and simple to perform, sensitive and practical tool for the detection of MAP and will be useful in facilitating the early diagnosis of paratuberculosis (Johne's disease) caused by the organism.

© 2014 Asian-African Society for Mycobacteriology. Published by Elsevier Ltd. All rights reserved.