

and a catalytically inactive MrpA–D sub-complex is formed in the absence of MrpE, F or G. Studies of *mrpE* point mutations further demonstrated the strong influence of small changes in MrpE on the antiporter properties of the whole complex although MrpA and D have been hypothesized to be the actual antiporter subunits.

doi:10.1016/j.bbabbio.2008.05.108

### S3.10 A role for uncoupling protein 1 in the formation of the mitochondrial permeability transition pore?

Paul G. Crichton, Nadeene Parker, Martin D. Brand

MRC, Dunn Human Nutrition Unit, Cambridge, UK

E-mail: [pgc@mrc-dunn.cam.ac.uk](mailto:pgc@mrc-dunn.cam.ac.uk)

The mitochondrial permeability transition pore (mPTP) is a non-specific channel that forms in the inner mitochondrial membrane in response to elevated levels of matrix calcium. The pore is generally believed to be comprised of the adenine nucleotide translocase (ANT), as well as several other mitochondrial proteins (e.g. VDAC and Cyclophilin D). Recent studies, however, indicate that the presence of ANT is not essential for permeability transition, which has led to the proposal that other members of the mitochondrial carrier protein family may be able to play a similar function to ANT in pore formation. To investigate this possibility we are studying the permeability transition properties of brown adipose tissue (BAT) mitochondria in which levels of the mitochondrial carrier protein, UCP1, can approach those of ANT. Using BAT mitochondria isolated from both wild-type and UCP1<sup>-/-</sup> mice we assess UCP1-specific, membrane potential-independent, influences on mPTP formation by studying their swelling properties under strictly de-energised conditions. UCP1-dependent contributions to mPTP formation will, therefore, allow us to determine if the mitochondrial carrier protein involvement in permeability transition is a more general property of this family of proteins or is more likely to be restricted to specific members such as the ANT.

doi:10.1016/j.bbabbio.2008.05.109

### S3.11 The mechanism of transport across the inner mitochondrial membrane: Clues from the PxW subfamily

Scott A. Lawrence, Richard G. Moran

Department of Pharmacology and Toxicology, Virginia Commonwealth University Richmond, Virginia, USA

E-mail: [lawrencesa@vcu.edu](mailto:lawrencesa@vcu.edu)

Mitochondrial carrier family members possess three conserved peptides that structural studies show to be linked by ionic bonds integral to opening the transport barrier into the mitochondrial matrix. These peptides contain the Px(D/E)xx(R/K) motif across family members, but a small subset of these carriers substitute a PxWxx(R/K) motif in the second peptide, and hence, possess new information about the mechanism of carrier opening. We have been applying site-directed mutagenesis to investigate the role of this PxWxx(R/K) motif in the mitochondrial folate transporter (MFT). We mutated MFT conserved sequence residue R249 and concluded that a  $\pi$ -cation interaction replaced one of the ionic bonds, which was consistent with previously proposed molecular modeling studies. Current results demonstrate that the MFT evolved a W (W142) substitution in place of the (D/E) in the second peptide

because an acidic residue in that position is incompatible with MFT function. Computational studies suggest that the aromatic phenyl ring of the folate substrate likely facilitates breakage of a  $\pi$ -cation interaction. A mutation predicted to form an additional  $\pi$ -cation interaction with W142 and stabilize the transport barrier, eliminated MFT function. Thus, the energetics of breakage of the transport barrier are delicate and tuned to the characteristics of the substrate.

doi:10.1016/j.bbabbio.2008.05.110

### S3.12 The ADP/ATP carrier of the apicomplexa *Cryptosporidium parvum*

John R. Mifsud, Edmund R.S. Kunji

Medical Research Council, Dunn Human Nutrition Unit, Cambridge CB2 0XY, UK

E-mail: [ek@mrc-dunn.cam.ac.uk](mailto:ek@mrc-dunn.cam.ac.uk)

The eukaryotic parasite *Cryptosporidium parvum* has mitochondria with reduced functions. Consistent with this notion, we have identified only eight putative transport proteins of the mitochondrial carrier family (MCF). Members of the MCF support mitochondrial metabolic energy generation, macromolecular synthesis, and amino-acid metabolism by linking biochemical pathways in the mitochondrial matrix with those in the cytosol. One of the *Cryptosporidium* proteins has been expressed functionally in the bacterium *Lactococcus lactis* and has been identified by whole cell and fused vesicle uptake studies as the ADP/ATP carrier. The transporter has a similar substrate specificity and inhibitor profile as the bovine and yeast ADP/ATP carrier. By comparative modelling, the internal cavity of the ADP/ATP carrier is more akin to the bovine ADP/ATP carrier than that of the ADP/ATP carrier of *Entamoeba histolytica*, which is the organism that causes amoebic dysentery.

doi:10.1016/j.bbabbio.2008.05.111

### S3.13 Determination of the dimensions and mass of membrane proteins in detergents by size exclusion chromatography

Marilyn Harding, Edmund R.S. Kunji

The Medical Research Council, Dunn Human Nutrition Unit, Hills Road, Cambridge, CB2 0XY, UK

E-mail: [ek@mrc-dunn.cam.ac.uk](mailto:ek@mrc-dunn.cam.ac.uk)

Determining the oligomeric state of a membrane protein is an important step in understanding its mechanism. Here, we apply a new approach based on size exclusion chromatography (SEC) of membrane proteins in the alkyl-maltoside and Cymal detergent series. The procedures will be illustrated by using the yeast mitochondrial ADP/ATP carrier, which was shown to be monomeric in detergent rather than dimeric. Several parameters of the detergent-solubilised protein were determined, such as its molecular mass, its Stokes' radius, its radius at the midpoint of the membrane, and its excluded volume. The procedures can be used to determine the mass and dimensions of other membrane proteins when the chromatographic behaviour in SEC is determined largely by the associated detergent micelle.

doi:10.1016/j.bbabbio.2008.05.112