



Redox characterization of usnic acid and its cytotoxic effect on human neuron-like cells (SH-SY5Y)

Thallita Kelly Rabelo^a, Fares Zeidán-Chuliá^b, Laura Milán Vasques^b, João Paulo Almeida dos Santos^b, Ricardo Fagundes da Rocha^b, Matheus Augusto de Bittencourt Pasquali^b, José Luiz Rybarczyk-Filho^b, Adriano Antunes Souza Araújo^a, José Claudio Fonseca Moreira^b, Daniel Pens Gelain^{b,*}

^aLaboratory of Pharmaceutical Assays and Toxicity, Federal University of Sergipe (LeFT/UFS), São Cristóvão, SE, Brazil

^bCenter of Oxidative Stress Research (CEEQ), Department of Biochemistry, Federal University of Rio Grande do Sul (UFRGS), Porto Alegre, RS, Brazil

ARTICLE INFO

Article history:

Received 20 July 2011

Accepted 2 December 2011

Available online 13 December 2011

Keywords:

Usnic acid

Free radicals

Cellular viability

Oxidative stress

SH-SY5Y cells

ABSTRACT

Usnic acid (UA) is the most common and abundant lichenic secondary metabolite with potential therapeutic application. Anti-inflammatory and antitumour properties have already been reported and UA-enriched extracts are widely used to treat several diseases in the folk medicine. First, we performed *in silico* evaluation of UA interactions with genes/proteins and important compounds for cellular redox balance and NO pathway. Then, we assessed UA redox properties against different reactive species (RS) generated *in vitro*, and evaluated its action on SH-SY5Y neuronal like cells upon hydrogen peroxide (H₂O₂), since no *in vitro* neurotoxicological data has been reported so far. Total reactive antioxidant potential index (TRAP) showed a significant antioxidant capacity of UA at the highest tested concentration; UA was also effective against hydroxyl radicals and reduced the formation of nitric oxide. *In vitro*, lipoperoxidation was enhanced by UA and changed the cellular viability at highest concentration of 20 µg/mL for 1 and 4 h, as well as 2 and 20 µg/mL for 24 h of treatment, according to MTT reduction assay. Moreover, UA did not display protective effects against H₂O₂-induced cell death in any case. Evaluation of intracellular RS production by the DCFH-based assay indicated that UA was able to induce changes in basal RS production at concentration of 20 µg/mL for 1 h and from 2 ng/mL to 20 µg/mL for 4 and 24 h. In conclusion, UA could display variable redox-active properties, according to different system conditions and/or cellular environment. Moreover, our results suggest that potential neurotoxicological effects of UA should be further studied by additional approaches; for instance, *in vivo* and clinical studies.

© 2011 Elsevier Ltd. Open access under the [Elsevier OA license](http://www.elsevier.com/locate/elsevier).

1. Introduction

Therapeutic properties displayed by some species of lichens are related to the production of secondary metabolites such as depsides, depsidones, dibenzofuranes, xanthenes, anthraquinones, aliphatic acids, atranorin and usnic acid (Ingólfssdóttir, 2002). Usnic acid (UA) is one of the most abundant secondary lichen metabolites and has been extensively studied. It has several biological activities, such as antibiotic (Cocchietto et al., 2002), antiviral (Campanella et al., 2002; Scirpa et al., 1999), analgesic and antipyretic (Okuyama et al., 1995), as well as anti-inflammatory properties (Vijayakumar et al., 2000). UA was reported to induce apoptosis of murine leukemia L1210 cells in a dose- and time-dependent manner (Bezivin et al., 2004), and exhibited anti-proliferative action against MCF7 breast cancer cells (Mayer

et al., 2005). These results highlighted UA as a potentially new precursor for novel chemotherapeutic agents.

Already, UA has shown a dualistic effect *in vivo* when compared to *in vitro* data. For instance, it exerts a significant gastroprotective effect in indomethacin-induced gastric ulcer rats, by reducing the formation of reactive species, and therefore, oxidative damage (Odabasoglu et al., 2006); but also it has shown to be a potent hepatotoxic agent that disrupts electron transport in mitochondria, inducing oxidative stress in cells (Han et al., 2004; Joseph et al., 2009). However, there exist many controversies about the toxic activities undertaken by this compound. All in all, better comprehension about UA-induced cellular mechanism would significantly contribute towards a safer therapeutic use.

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are involved in the pathogenesis of numerous diseases such as cancer, inflammatory diseases and neurodegenerative disorders (Seifried et al., 2007). Under normal physiological conditions, ROS/RNS participate as intracellular messengers and regulatory molecules. They are tightly regulated by balancing systems formed

* Corresponding author. Address: Rua Ramiro Barcelos, 2600-Anexo, CEP 90035-003, Porto Alegre, RS, Brazil. Tel.: +55 51 3308 5577; fax: +55 51 3308 5535.

E-mail address: dgelain@yahoo.com.br (D.P. Gelain).

by different antioxidants, antioxidant enzymes, and proteins (Kowaltowski et al., 2009). Non-enzymatic antioxidants coming from diet or other processes regulate oxidative and nitrosative reactions in the body, which prevents oxidative stress by removing both ROS and RNS (Maes et al., 2011).

Main actions of secondary metabolites in biological systems have also been linked to their redox properties. Our group has been studying the redox properties of atranorin, another lichenic secondary metabolite with potential therapeutic activity, similar to that exerted by UA. In that study, atranorin increased lipoperoxidation and displayed general antioxidant activity (Melo et al., 2011). The potential health-promoting effects of naturally occurring compounds are traditionally ascribed to a general antioxidant action (Aravindaram and Yang, 2010). In fact, the properties of UA are generally associated to its widespread antioxidant action, found in most phenolic compounds, since many therapeutic properties attributed to lichen extracts (and to UA itself) are intimately associated to oxidative stress; together with an unbalanced free radical production, mutagenicity, and inflammation (Halliwell and Gutteridge, 2007). However, only few works have studied the potential pro- or antioxidant properties of UA (Carlos et al., 2009; Jayaprakasha and Rao, 2000; Toledo Marante et al., 2003; Valencia-Islas et al., 2007). Interestingly, a number of phenolic compounds and other naturally derived substances, that were initially observed to act as antioxidants in mammalian cells, have later been described to unbalance the cellular redox state towards pro-oxidant states, depending on specific conditions (i.e., higher or lower drug concentrations) (Halliwell, 2008).

Since there are some studies showing that UA prevents oxidative stress by removing both ROS and RNS, we decide to deeper characterize *in silico* the general landscape of interactions of UA with genes/proteins, and compounds generally involved in both redox and NO pathways, by using system biology tools. Thereafter, we performed an *in vitro* characterization of the redox properties of UA against different reactive species, and evaluated its potential effects on cellular viability by using SH-SY5Y cells neuronal like cells since no *in vitro* neurotoxicological reports have been published yet.

2. Materials and methods

2.1. Chemicals

AAPH (2,2'-Azobis(2-methylpropionamide)dihydrochloride), Luminol (5-amino-2,3-dihydro-1,4-phthalazinedione), 2-deoxyribose, glycine, Griess reagent, SNP (sodium nitroprusside), TBA (2-thiobarbituric acid), (4,6-dihydroxypyrimidine-2-thiol), H₂O₂ (hydrogen peroxide), adrenaline, catalase, SOD (superoxide dismutase), MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide), DMSO (Dimethyl sulfoxide), DCFH-DA (2',7'-dichlorodihydrofluorescein diacetate) and (+)-Usnic acid were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Materials used in cell culture were acquired from Gibco®/Invitrogen (São Paulo, SP, Brazil) and from Rio de Janeiro Cell Bank (BCRJ, Rio de Janeiro, Brazil). The rest of reagents used in this study were of either analytical or HPLC grade. For each assay, UA (20 mg mL⁻¹) was dissolved in DMSO (100%) and serial dilutions were obtained from this stock solution. Therefore, at the highest concentration of UA in these assays (20 µg mL⁻¹), concentration of the vehicle DMSO would correspond to 0.1%.

2.2. Interaction networks of compounds and gene/proteins

In order to develop a model network for gene/protein and UA interaction, we first selected a number of gene/proteins involved

in redox and NO-related pathways and then, by using STITCH 2.0 (Kuhn et al., 2008, 2010) we screened the possible protein–protein and protein–compound interactions based on experimental knowledge and database (confident score = 0.4, medium). A list with gene symbols and Ensembl protein IDs is additionally provided (Supplementary Table 1).

The network connected 61 proteins and 5 compounds together, based on their possible interaction through “activation”, “catalysis”, “binding”, “inhibition”, and “reaction”; giving rise to a model for UA interactions through redox/NO pathways (MUA network).

2.3. Total reactive antioxidant potential (TRAP) and total antioxidant reactivity (TAR)

Total reactive antioxidant potential (TRAP) is utilized to estimate the non-enzymatic antioxidant capacity of samples *in vitro*. This method is based on the quenching of luminol-enhanced chemiluminescence (CL) derived from the thermolysis of AAPH as the free radical source (Lissi et al., 1992). Briefly, we prepared AAPH solution, added luminol (AAPH + luminol, radical generating system) and then, we waited for the system to stabilize for 2 h before the first reading. Different concentrations of UA were added and the luminescence produced by the free radical reaction was quantified in a liquid scintillator counter (Wallac 1409, Perkin-Elmer, Boston, MA, USA) for 60 min. The results were transformed in percentage and area under curve (AUC), and calculated by software (GraphPad software.® San Diego, CA; version 5.00) as previously described (Dresch et al., 2009).

Total antioxidant reactivity (TAR) was analyzed by using the same samples utilized for TRAP readings. TAR results were calculated as the ratio of light intensity in absence of samples (I_0)/light intensity right after UA addition. Although TAR and TRAP evaluations are obtained in the same experiment, they represent different observations, since the TAR is more related to the antioxidant quality (reactivity, the scavenging capacity in a short-term period) and TRAP is more related to the antioxidant amount and kinetic behavior (Lissi et al., 1995).

2.4. Hydroxyl radical-scavenging activity

The formation of ·OH (hydroxyl radical) from Fenton reaction was quantified by using the 2-deoxyribose oxidative degradation assay. The principle of the assay is the incubation of 2-deoxyribose together with a hydroxyl radical generation system, which produces malondialdehyde (MDA). This system is then incubated with 2-thiobarbituric acid (TBA), which reacts with MDA and forms a chromophore quantifiable by spectrophotometry (Lopes et al., 1999). Briefly, typical reactions were started by the addition of Fe²⁺ (FeSO₄ 6 µM final concentration) to solutions containing 5 mM 2-deoxyribose, 100 mM H₂O₂ and 20 mM phosphate buffer (pH 7.2).

To measure UA antioxidant activity against hydroxyl radicals, different concentrations of UA were added to the system before Fe²⁺ addition. Reactions were carried out for 15 min at room temperature and were stopped by the addition of 4% phosphoric acid (v/v) followed by 1% TBA (w/v, in 50 mM NaOH). Solutions were boiled for 15 min at 95 °C, and then cooled at room temperature. The absorbance was measured at 532 nm and results were expressed as percentage of TBARS formed.

2.5. Nitric oxide (NO·) scavenging activity

Nitric oxide was generated from spontaneous decomposition of sodium nitroprusside in 20 mM phosphate buffer (pH 7.4). Once it is generated, NO interacts with oxygen to produce nitrite ions, which were measured by the Griess reaction (Basu and Hazra,

2006). The reaction mixture (1 mL) containing both 10 mM sodium nitroprusside (SNP) in phosphate buffer (pH 7.4) and UA at different concentrations were incubated at 37 °C for 1 h. An aliquot of 0.5 mL was taken and homogenized with 0.5 mL Griess reagent. The absorbance of chromophore was measured at 540 nm. Results were expressed as percentage of nitrite formed by SNP alone.

2.6. Thiobarbituric acid reactive species (TBARS)

Thiobarbituric acid-reactive substances (TBARS) assay was employed to quantify lipid peroxidation (Draper and Hadley, 1990), and an adapted TBARS method was used to measure the antioxidant capacity of UA using egg yolk homogenate as lipid rich substrate (Melo et al., 2011). The principle of the method is based on spectrophotometric measurement of the color produced during the reaction of thiobarbituric acid (TBA) with lipoperoxidation products, such as malondialdehyde and 4-hydroxynonenal (Mansour and Mossa, 2009). Briefly, egg yolk was homogenized (1% w/v) in 20 mM phosphate buffer (pH 7.4), 1 mL of homogenate was sonicated and then homogenized with 0.1 mL of UA at different concentrations. Lipid peroxidation was induced by addition of 0.1 mL of AAPH solution (0.12 M). Control was just incubation medium without AAPH. Reactions were carried out for 30 min at 37 °C. Samples (0.5 mL) were centrifuged with 0.5 mL of trichloroacetic acid (15%) at 1200g for 10 min. An aliquot of 0.5 mL from supernatant was mixed with 0.5 mL TBA (0.67%) and heated at 95 °C for 30 min. After cooling, sample's absorbance was measured by using a spectrophotometer (UV-1800 Shimadzu) at 532 nm. The results were expressed as percentage of TBARS formed by AAPH alone (induced control).

2.7. Determination of superoxide dismutase-like activity (SOD)

The ability of UA to scavenge superoxide anion (“superoxide dismutase-like activity” or “SOD-like activity”) was measured as previously described (Misra and Fridovich, 1972). UA was mixed with 200 μ l glycine buffer (50 mM, pH 10.2) and 5 μ l of native catalase 100 U/mL. Superoxide generation was initiated by addition of adrenaline 2 mM, and adrenochrome formation was monitored at 480 nm for 5 min (32 °C). Superoxide production was determined by monitoring the reaction curves of samples, and measured as percentage of the rate of adrenaline auto-oxidation into adrenochrome (Bannister and Calabrese, 1987).

2.8. Determination of catalase-like activity (CAT)

The capacity of UA to degrade the hydrogen peroxide (H₂O₂) added in an incubation medium (“catalase-like activity” or “CAT-like activity”) was measured as previously described (Aebi, 1984). Briefly, H₂O₂ diluted in 0.02 M phosphate buffer (pH 7.0), to obtain a 5 mM final concentration, was added to microplate wells, in which different concentrations of UA were already placed. The plate was then scanned in a spectrophotometric plate reader (SpectraMax 190, Molecular Devices) at 240 nm every 15 s for 5 min at 37 °C. Catalase-like activity was monitored based on the rate decomposition of H₂O₂. Data were expressed as percentage of the rate decomposition of H₂O₂.

2.9. Cell culture

Exponential growing human neuroblastoma cell line SH-SY5Y, obtained from Rio de Janeiro Cell Bank (BCRJ, Rio de Janeiro, Brazil), were maintained in a mixture 1:1 of Ham's F12 and Dulbecco Modified Eagle Medium (DMEM) supplemented with 10% heat-inactivated FBS, 2 mM of glutamine, 0.28 μ g/ μ L of gentamicin and 250 μ g of amphotericin B, in a humidified atmosphere of 5%

of CO₂ in air at 37 °C. Cell medium was replaced each 2 days and cells were sub-cultured once they reached 90% confluence. All treatments were performed when cells were 70–90% confluence.

2.10. MTT assay

After treating with either different concentrations of UA alone or in the presence of H₂O₂ 1 mM for 1, 4, and 24 h SH-SY5Y cells viability was quantified by the MTT assay as previously described (Gelain and Moreira, 2008). This method is based on the ability of viable cells to reduce MTT (3-(4,5-dimethyl)-2,5-diphenyl tetrazolium bromide) and form a blue formazan product. MTT solution (sterile stock solution of 5 mg/mL) was added to the incubation medium in the wells at a final concentration of 0.2 mg/mL. The cells were left for 45 min at 37 °C in a humidified 5% CO₂ atmosphere.

Culture medium was then removed and plates were shaken with DMSO for 30 min. Optical density of each well was measured at 550 nm (test) and 690 nm. H₂O₂ 1 mM was used as positive control for the assay. Data were expressed as percentage of the formazan formation in untreated cells (control). Qualitative morphology of the cell cultures was also evaluated by phase-contrast light microscopy (Nikon Eclipse TE 300).

2.11. DCFH-DA assay

Intracellular reactive species production was determined by the DCFH-DA assay, as described by (Wang and Joseph, 1999). In the presence of reactive species (RS), DCFH is oxidized to highly fluorescent dichlorofluorescein (DCF) can be used as an index to quantify the overall ROS in cells. Briefly, 2×10^4 SH-SY5Y cells were seeded in 96-well plates and 100 μ M DCFH-DA dissolved in medium containing 1% FBS was added to each well and incubated for 2 h to allow cellular incorporation. After, the medium was discarded and cells were treated with different concentrations of UA alone or in the presence of H₂O₂ (1 mM) and DCF fluorescence was read at 37 °C during 1, 4 and 24 h in a fluorescence plate reader (Spectra Max M2, Molecular Devices, USA) with an emission wavelength set at 535 nm and an excitation wavelength set at 485 nm. The results were expressed as percentage DCF fluorescence. Hydrogen peroxide (1 mM) was used as positive control for intracellular reactive species production.

2.12. Statistical analysis

The *in vitro* procedures were carried out with $n = 3$ (i.e., 3 vials per group) while cell culture experiments were performed with $n = 6$ (i.e., 6 wells per group). Experiments were repeated four different times, and the results were expressed as mean \pm standard error of the mean (SEM) of four independent experiments. The differences among data were evaluated by one-way analysis of variance (ANOVA) followed by Tukey's *post hoc* test. In all cases differences were considered significant if $p < 0.05$. Data analyses were performed using the (GraphPad software,® San Diego, CA; version 5.00).

3. Results

In silico analysis gave rise to a network where exactly 61 proteins and 5 compounds where interconnected (“MUA network”) (Fig. 1A). Based on experimental data and database, UA directly interacts with hydroxyl radicals (Fig. 1F) and through them, with antioxidant enzymes (Fig. 1E), such as SOD1, SOD3, CAT, GPX1, and GPX2. UA interaction with hydroxyl radicals interconnects it with NO and proteins involved in NO biosynthesis (Fig. 1C); for in-




Fig. 1. Model for UA interactions through redox and NO pathways (A) *In silico* analysis based on experimental data and database gave rise to a network where exactly 61 proteins and 5 compounds were interconnected (“MUA” network). (B) Landscape of protein–protein and protein–compound interactions through activation processes. (C) Landscape of protein–protein and protein–compound interactions through catalysis processes. (D) Landscape of protein–protein and protein–compound interactions through binding processes. (E) Landscape of protein–protein and protein–compound interactions through inhibition processes. (F) Landscape of protein–protein and protein–compound interactions through diverse reactions.

stance, Nostrin, which is known to modulate nitric oxide release (Zimmermann et al., 2002). Landscape of protein–protein and protein–compound interactions through either activation or binding processes are additionally provided (Fig. 1B and D, respectively).

Redox properties of UA were first evaluated (TRAP and TAR assays) by using a method based on the quenching of luminol-enhanced chemiluminescence (CL) of AAPH. At 20 $\mu\text{g}/\text{mL}$ of UA, we observed an antioxidant capacity of UA by TRAP assay (Fig. 2A). Usnic acid at 20 $\mu\text{g}/\text{mL}$ also showed significant antioxidant capacity according to TAR measurements (Fig. 2B). In both methods, this concentration was able to efficiently reduce AAPH-induced CL, indicating peroxy/alkoxy non-enzymatic scavenging activity.

The 2-deoxyribose degradation assay was employed to investigate the ability of UA to scavenge *in vitro*-generated hydroxyl radicals by Fenton reaction. Fig. 3A shows that concentrations of UA from 2 $\mu\text{g}/\text{mL}$ to 20 $\mu\text{g}/\text{mL}$ inhibited hydroxyl radical-induced deoxyribose degradation, indicating an antioxidant effect against hydroxyl radicals. The capacity of UA to scavenge NO was measured by quantifying the production of nitrite derived from sodium nitroprusside (SNP) by the Griess reaction. Fig. 3B shows that all concentrations of UA tested (2 ng/mL to 20 $\mu\text{g}/\text{mL}$) significantly decreased SNP-derived nitrite formation.

The protective effect of UA against oxidative damage to lipids was measured by quantifying TBARS generated by AAPH in a




Fig. 2. TRAP and TAR measurements. (A) The total reactive antioxidant potential (TRAP) of UA at different concentrations. A free radical source (AAPH) generating system produces peroxy/alcoxy radicals at a constant rate, and the effect of different concentrations of UA on free radical-induced chemiluminescence is measured as area under a curve for 60 min. (B) Total antioxidant reactivity (TAR) values are calculated as the ratio of light intensity in absence of samples (I_0)/light intensity right after UA addition (I) and expressed as percent of inhibition. All groups denote samples in the presence of AAPH. Trolox (75 µg/ml) was used as standard antioxidant. The experiments were performed in triplicate, and bars represent mean \pm SEM of four different experiments. *** $p < 0.0001$ (1-way ANOVA followed by Tukey's *post hoc* test).

lipid-rich incubation medium. All the tested concentrations (2 ng/mL–20 µg/mL) of UA increased the AAPH-induced lipoperoxidation (Fig. 4), indicating that UA is either an enhancer of lipid peroxidation or somehow, it may facilitate chain reactions involved in the propagation of lipid peroxides.

The activity of UA against superoxide anions (SOD-like activity) was quantified by the inhibition of superoxide-dependent adrenaline auto-oxidation to adrenochrome. Moreover, we also assessed the ability of UA to decompose H_2O_2 *in vitro* (CAT-like activity). Respectively, Fig. 5A and B show that all tested concentrations of UA did not exert any significant variation in CAT and SOD-like activities.

In order to elucidate the potential effects that UA may exert in a cellular system (challenged by a pro-oxidant agent), we chose SH-SY5Y cells as *in vitro* neurotoxicological model. Phase contrast microscopy showed that cells treated with the highest concentration of UA (20 µg/mL) for 1 and 4 h, induced changes in their neuronal-like morphology (Fig. 6). At 1 h of treatment with 20 µg/mL of UA, neuritic processes seem to be stopped (Fig. 6a7) and cells looked more rounded. When treatment reached 4 h of incubation at the same concentration, cells already seem to lose their viability and start to detach from the culture plate (Fig. 6b7). In the pres-




Fig. 3. Hydroxyl and nitric oxide (NO) scavenging activities assays. (A) Hydroxyl radical-scavenging activity was quantified by using hydroxyl-mediated 2-deoxyribose oxidative degradation *in vitro*, which produces malondialdehyde (MDA) by condensation with 2-thiobarbituric acid (TBA). System is MDA production from 2-deoxyribose degradation with $FeSO_4$ and H_2O_2 alone (hydroxyl generating system). Other groups denote MDA production by $FeSO_4$ and H_2O_2 in the presence of different concentrations of UA. (B) NO scavenging assay. Nitric oxide was generated from spontaneous decomposition of sodium nitroprusside (SNP) in the presence of oxygen, producing nitrite ions, which were measured by the Griess reaction. Nitrite production by SNP alone was compared to nitrite production by SNP in the presence of different concentrations of UA. Trolox was used as standard antioxidant in both assays. The experiments were performed in triplicate, and bars represent mean \pm SEM of four different experiments. *** $p < 0.0001$. One-way ANOVA followed by Tukey's *post hoc* test was applied to all data.

ence of H_2O_2 (1 mM) together with UA (20 µg/mL), cells detached and seemed to lose their viability already at 1 h of treatment (Fig. 6a14). Interestingly, such effect looked stronger in comparison to cells that were treated only with 20 µg/mL of UA (Fig. 6a7). Moreover, even though clear morphological effects and potential loss of viability were only detectable at the highest concentration of UA (20 µg/mL) when treatment lasted 4 h (Fig. 6b7), 4 h of H_2O_2 and UA co-treatment showed an UA dose-dependent cellular detachment in comparison to those cells treated with 1 mM of H_2O_2 alone (positive control) (Fig. 6b8–b12). Strongest effect appeared at the highest concentration of UA (20 µg/mL) together with H_2O_2 (1 mM) (Fig. 6b12). For studying long-term UA effect on the cells, treatments were performed (in the presence of 1% of FBS) for 24 h. Results showed signs of cell detachment and loss of viability when cells were incubated with 2 µg/mL of UA (24 h) (Fig. 7f), and such effect was significantly stronger with 20 µg/mL of UA, with virtually 100% of the cells detached from the plate (Fig. 7g). 24 h co-treatment of UA together with 1 mM of H_2O_2 resulted in complete lack of cellular viability at any concentration




Fig. 4. Production of Thiobarbituric acid-reactive substances (TBARS) *in vitro*. A lipid-rich system was incubated with a free radical source (AAPH) and the effect of different concentrations of UA on the lipoperoxidation was measured by quantifying TBARS. Trolox (75 µg/ml) was used as standard antioxidant. The experiments were performed in triplicate, and bars represent mean \pm SEM of four different experiments. ** $p < 0.001$, *** $p < 0.0001$ (1-way ANOVA followed by Tukey's *post hoc* test).




Fig. 5. SOD-like and CAT-like activities. (A) Superoxide dismutase-like (SOD-like) activity was determined by following formation of adrenochrome in a SOD reaction buffer containing native catalase and adrenaline O₂ generator group. (B) Catalase-like (CAT-like) activity was measured in a catalase reaction buffer with H₂O₂. The experiments were performed in triplicate, and bars represent mean \pm SEM of four different experiments. One-way ANOVA followed by Tukey's *post hoc* test was applied to all data.

of UA (Fig. 7j–n). These results were confirmed when MMT assays showed that cells treated for 1 and 4 h with the highest concentra-

tion of UA (20 µg/mL) cellular viability was reduced in a 41.3% and 88.67% respectively in relation to control (100%) (Fig. 8A and B). During long-term treatments of 24 h, with concentrations of 2 µg/mL and 20 µg/mL, cellular viability was reduced in a 41.9% and 95.23% respectively, in relation to control (100%) (Fig. 8C).

Moreover, UA did not display protective effects in any case after 1, 4 or 24 h of treatment against 1 mM H₂O₂-induced cell death, which on the contrary, was able to significantly decrease cellular viability in a 88.4%, 92.06% and 95.96% respectively, in relation to control (100%) (Fig. 8A, B and C). This result suggests that UA is not able to prevent oxidative-mediated H₂O₂-induced cell death at any concentration.

Next, we evaluated if UA was able to induce intracellular ROS production in SH-SY5Y cells by the DCFH-DA assay. As shown in Fig. 9A, SH-SY5Y cells treated with UA (20 µg/mL) for 1 h increased in 76.73% when compared to control cells. After 4 and 24 h of treatment, levels of intracellular ROS were enhanced at any tested concentration of UA (Fig. 9B and C). Respectively: 2 ng/mL (44.99–62.71%), 20 ng/mL (60.63–78.20%), 200 ng/mL (79.56–90.32%), 2 µg/mL (53.69–64.60%) and 20 µg/mL (106.07–122.70%).

These results suggest that UA is able to trigger ROS production in SH-SY5Y cells and therefore, to induce an oxidative stress scenario. Besides, we also evaluated the effect on intracellular ROS production when cells were co-treated with UA together with H₂O₂ (1 mM) for 1, 4, and 24 h. At any concentration, UA appears to potentiate the effect of H₂O₂ when it comes to ROS production (Fig. 9A, B and C).

4. Discussion

Several studies have shown that redox activity associated with natural antioxidants is attributed to total content of phenolic compounds (Halliwell, 2008; Rice-Evans et al., 1995; Scalbert et al., 2005). Moreover, antioxidant and pro-oxidant biochemical agents have presented important role to design strategies for prevention and/or management of oxidative damage. Here, we aimed to characterize the redox properties of UA by using different approaches, in order to understand possible interactions of such compound with different types of reactive species.

The potential of UA scavenging peroxy radicals, by using TRAP/TAR assays, indicated a significant antioxidant capacity at the highest tested concentration. Moreover, UA was capable to quench hydroxyl radicals. Hydroxyl radical has a high oxidant power and it is probably the most reactive radical (Pastor et al., 2000). It is able to join DNA nucleotides and cause strand breakage, which contributes to carcinogenesis, mutagenesis, and cytotoxicity (Manian et al., 2008). The hydroxyl radical-scavenging capacity of any compound is directly related to its antioxidant activity (Babu et al., 2001). In addition, NO is an important mediator of acute and chronic inflammation. NO stimulates cyclooxygenase (COX) activity resulting in unbalanced production of pro-inflammatory prostaglandins (PG) (Salvemini et al., 1993). Other mechanism by which NO may modulate inflammatory processes is through its interaction with the Rel/nuclear transcription factor kappaB (NF-κB) family of transcription factors (Laroux et al., 2001). At high concentrations, NO may interact with superoxide radicals generating peroxynitrite (ONOO⁻), a potent oxidizing molecule capable of eliciting damage to proteins, lipids and DNA, which modulates COX (Zhang et al., 1994; Mollace et al., 2005; Halliwell and Gutteridge, 2007). Here, we observed that UA is able to reduce the production of nitrite, indicating a potential role as NO-scavenging agent and consequently, it may limit the action of these reactive species in biological systems.

ROS and RNS have also been related to act as pro-inflammatory signals *in vivo* by stimulating the activation of TNF-α, IL-1β and IL-




Fig. 6. Morphological effects of short and medium-term UA treatment on SH-SY5Y cells. Phase contrast microscopy of SH-SY5Y cells after 1 (A) and 4 h (B) treatment with different concentrations of UA, in the absence or presence of H₂O₂ (1 mM). Photomicrographies representative of three independent experiments.

6 genes through activation of the redox-sensitive transcription factor NF- κ B (Beauparlant and Hiscott, 1996). Thus, the anti-inflammatory and analgesic activities exerted by UA could also be a consequence of its NO-scavenging activity, which may prevent free radical-induced NF- κ B activation and consequent pro-inflammatory cytokine production; a cycle that would perpetuate inflammatory processes.

On the other hand, UA presented a pro-oxidant capacity in a lipid-rich system, enhancing TBARS formation induced by AAPH incubation. Lipid peroxidation has been shown to induce disturbance of membrane organization, functional loss as well as modification of proteins and DNA bases (Niki, 2009). Usnic acid is a lipophilic weak acid that can diffuse through mitochondrial membranes and cause proton leaking (uncoupling) (Joseph et al., 2009).




Fig. 7. Morphological effects of long-term UA treatment on SH-SY5Y cells. Phase contrast microscopy of SH-SY5Y cells after 24 h treatment with different concentrations of UA, in the absence or presence of H₂O₂ (1 mM). Photomicrographs representative of three independent experiments.

The uncoupling of mitochondrial oxidative phosphorylation by UA is similar to that observed by the classical uncoupler, 2,4-dinitrophenol (DNP). In mouse liver mitochondria, high doses of DNP resulted in increased lipid peroxidation due to higher oxidative stress (Futakawa et al., 2006). Odabasoglu et al. (2006) showed an antioxidant effect of UA when used against indomethacin-induced gastric ulcers in rats. They observed a significant inhibition of reactive species formation, a decrease in lipid peroxidation and increase of antioxidant enzymes such as glutathione peroxidase and superoxide dismutase.

Although, the ability of compounds to quench hydroxyl radicals is generally related to the prevention of lipid peroxidation propagation (Shukla et al., 2009), the capacity of free radical scavenging by antioxidants does not necessarily correlate with the capability to inhibit lipid peroxidation. It is well-described that lipid peroxidation may occur not only through free radical-mediated oxidations but also upon enzymatic oxidation as well as radical-independent non-enzymatic oxidations. Lipid peroxidation leads to cytotoxicity, excepting for those cases of sublethal concentrations where the result could be either diverse cellular adaptive responses or even up-regulation of antioxidant enzymes with increased tolerance to following oxidative stress. All this together, provides putative scenarios that support our results in SH-SY5Y cells (Niki et al., 2005; Niki, 2009). Besides, studies concerning the hepatotoxic effect of (+) UA in isolated rat hepatocytes and liver mitochondria using carbon tetrachloride (CCl₄), as the reference hepatotoxin, showed that UA seems to trigger the same cytotoxic mechanisms as CCl₄, increasing the production of MDA, a marker for lipid peroxidation (Pramyothin et al., 2004).

Then, we could speculate that pro-oxidant actions of UA observed in mitochondria and hepatocytes could be a result of the capacity that UA has to enhance free radical-induced lipoperoxidation; similarly to what we observed in our AAPH-generating lipoperoxidation system.

Usnic acid has several interesting biological properties. For instance, it is used in pharmaceutical preparations against infections,

bacterial eczema, mastitis, furunculosis and polydermy (Cocchietto et al., 2002). Thus, novel natural compounds isolated from lichens may represent a source of novel substances with selective biological action. The fact that UA is a non-genotoxic antineoplastic agent that works in a p53-independent manner (Yanmamoto et al., 1995) makes it a potential candidate for novel cancer therapies. However, to the best of our knowledge, nothing is known about UA effects in any *in vitro* neurotoxicological model. For this purpose, we decide to use SH-SY5Y cell line as well-characterized and accepted neuronal-like *in vitro* model because of their neuron-like properties, their capability to undergo neurite outgrowth as well as morphological changes, induced by oxidative stress (Cheung et al., 2009; Frota Junior et al., 2011; Navarra et al., 2010; Nicolini et al., 1998; Pahlman et al., 1984; Yu et al., 2011).

Overproduction of ROS can cause severe impairment of cellular functions. RS are involved in apoptotic mechanisms and may contribute to the apoptotic process found in several diseases (Jung et al., 2007). Our results have shown, that UA induced changes in their neuronal-like morphology, as well as, seems to have stopped the neuritic processes. The morphological changes, mainly in neurite outgrowth in SH-SY5Y, can be induced by increased intracellular ROS production (Jung et al., 2007). In addition, UA decreased cell viability with consequent cell death. This loss of cellular viability induced by UA can be due to increase the ROS.

Through DCFH-DA assay, our results showed that UA significantly increased intracellular ROS production when cells were treated for 1, 4 and 24 h, respectively. The advantage of this assay is that one could use intact living present a relevant redox-active action, acting as either pro-oxidant or antioxidant agent, depending on the specific radical that is being generated and the microenvironment provided by the system. As a matter of fact this dualistic effect is already known for some antioxidants that are able to auto-oxidize and therefore, to both generate reactive substances and act as pro-oxidants; strongly depending on the system composition (Moure et al., 2001). For instance, our laboratory has extensively characterized an apparently unexpected pro-oxidant effect of vita-




Fig. 8. MTT assay. Cultured SH-SY5Y cells were incubated with different concentrations of UA during 1 h (A), 4 h (B), and 24 h (C) at 37 °C in humidified 5% CO₂. MTT solution (sterile stock solution of 5 mg/mL) was added to the incubation medium in the wells at a final concentration of 0.2 mg/mL and incubated for 45 min. The medium was then replaced for DMSO and the optical density of each well was measured at 550 nm (test) and 690 nm (reference). H₂O₂ 1 mM was used as positive control for cell death. Data were expressed as percentage of the formazan formation in untreated cells (control). Bars represent mean ± SEM of data from four different experiments (six wells per group in each independent experiment). Data were analyzed by one-way ANOVA followed by Tukey's *post hoc* test. ****p* < 0.001, *****p* < 0.0001. # Different from control.

min A (and related carotenoids) both *in vitro* and *in vivo* at initially therapeutic conditions (Dal-Pizzol et al., 2000; Gelain et al., 2006, 2008; Klamt et al., 2003).

Our results suggest that UA displays variable redox-active properties, acting an antioxidant e pro-oxidant agent, according to different system conditions and/or cellular environment. These pro-oxidant properties in biological systems might be responsible by potential neurotoxicological effects of UA. Data




Fig. 9. Intracellular reactive species production (DCFH assay). Cultured SH-SY5Y cells were seeded in 96-well plates and 100 µM DCFH-DA dissolved in medium containing 1% FBS was added to each well and incubated for 2 h. Medium was then discarded and cells were treated with different concentrations of UA alone or in the presence of H₂O₂ (1 mM) and DCF fluorescence was read at 37 °C during 1 h (A), 4 h (B), and 24 h (C) (endpoint) in a fluorescence plate reader (Spectra Max M2, Molecular Devices, USA) with an emission wavelength set at 535 nm and an excitation wavelength set at 485 nm. Bars represent mean ± SEM of data from four different experiments (six wells per group in each independent experiment). ***p* < 0.001, *****p* < 0.0001 (1-way ANOVA followed by Tukey's *post hoc* test). # Different from control.

presented in this work, may contribute to elucidation of the bifunctional behaviour and neurotoxicological actions of UA in biological systems.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgements

The Brazilian agencies CNPq, FAPERGS (PqG #1008860), PRO-CAD/CAPES (#066/2007), FAPITEC and IBN-Net (#01.06.0842-00) supported this work.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tiv.2011.12.003.

References

- Aebi, H., 1984. Catalase in vitro. *Meth. Enzymol.* 105, 121–126.
- Aravindaram, K., Yang, N.S., 2010. Anti-inflammatory plant natural products for cancer therapy. *Planta Med.* 76, 1103–1117.
- Babu, B.H., Shylesh, B.S., Padikkala, J., 2001. Antioxidant and hepatoprotective effect of *Alanthus icicifolius*. *Fitoterapia* 72, 272–277.
- Bannister, J.V., Calabrese, L., 1987. Assays for superoxide dismutase. *Meth. Biochem. Anal.* 32, 279–312.
- Basu, S., Hazra, B., 2006. Evaluation of nitric oxide scavenging activity, in vitro and ex vivo, of selected medicinal plants traditionally used in inflammatory diseases. *Phytother. Res.* 20, 896–900.
- Beauparlant, P., Hiscott, J., 1996. Biological and biochemical inhibitors of the NF- κ B/Rel proteins and cytokine synthesis. *Cytokine Growth Factor Rev.* 7, 175–190.
- Bezivin, C., Tomasi, S., Rouaud, I., Delcros, J.G., Boustie, J., 2004. Cytotoxic activity of compounds from the lichen: *Cladonia convolute*. *Planta Med.* 70, 874–877.
- Campanella, L., Delfini, M., Ercole, P., Iacoangeli, A., Risuleo, G., 2002. Molecular characterization and action of usnic acid: a drug that inhibits proliferation of mouse polyomavirus in vitro and whose main target is RNA transcription. *Biochimie* 84, 329–334.
- Carlos, I.Z., Quilles, M.B., Carli, C.B., Maia, D.C., Benzatti, F.P., Lopes, T.I., et al., 2009. Lichen metabolites modulate hydrogen peroxide and nitric oxide in mouse macrophages. *Z. Naturforsch. [C]* 64, 664–672.
- Cheung, Y.T., Lau, W.K., Yu, M.S., Lai, C.S., Yeung, S.C., So, K.F., Chang, R.C., 2009. Effects of all trans-retinoic acid on human SH-SY5Y neuroblastoma as in vitro model in neurotoxicity research. *Neurotoxicology* 30, 127–135.
- Cocchietto, M., Skert, N., Nimis, P.L., Sava, G., 2002. A review on usnic acid, an interesting natural compound. *Naturwissenschaften* 89, 137–146.
- Dal-Pizzol, F., Klamt, F., Frota Jr., M.L., Moraes, L.F., Moreira, J.C., Benfato, M.S., 2000. Retinol supplementation induces DNA damage and modulates iron turnover in rat Sertoli cells. *Free Radic. Res.* 33, 677–687.
- Draper, H.H., Hadley, M., 1990. Malondialdehyde determination as index of lipid peroxidation. *Meth. Enzymol.* 186, 421–431.
- Dresch, M.T., Rossato, S.B., Kappel, V.D., Biegelmeyer, R., Hoff, M.L., Mayorga, P., Zuanazzi, J.A., Henriques, A.T., Moreira, J.C., 2009. Optimization and validation of an alternative method to evaluate total reactive antioxidant potential. *Anal. Biochem.* 385, 107–114.
- Frota Junior, M.L., Pires, A.S., Zeidán-Chuliá, F., Bristot, I.J., Lopes, F.M., de Bittencourt Pasquali, M.A., Zanotto-Filho, A., Behr, G.A., Klamt, F., Gelain, D.P., Moreira, J.C., 2011. In vitro optimization of retinoic acid-induced neurogenesis and TH endogenous expression in human SH-SY5Y neuroblastoma cells by the antioxidant Trolox. *Mol. Cell Biochem.* 358, 325–334.
- Futakawa, N., Kondoh, M., Ueda, S., Higashimoto, M., Takiguchi, M., Suzuki, S., Sato, M., 2006. Involvement of oxidative stress in the synthesis of metallothionein induced by mitochondrial inhibitors. *Biol. Pharm. Bull.* 29, 2016–2020.
- Gelain, D.P., Cammarota, M., Zanotto-Filho, A., de Oliveira, R.B., Dal-Pizzol, F., Izquierdo, I., et al., 2006. Retinol induces the ERK1/2-dependent phosphorylation of CREB through a pathway involving the generation of reactive oxygen species in cultured Sertoli cells. *Cell Signal.* 18, 1685–1694.
- Gelain, D.P., Moreira, J.C.F., 2008. Evidence of increased reactive species formation by retinol, but not retinoic acid, in PC12 cells. *Toxicol. in Vitro* 22, 553–558.
- Halliwell, B., Gutteridge, J.M.C., 2007. *Free Radicals in Biology and Medicine*, fourth ed. Oxford University Press, Oxford.
- Halliwell, B., 2008. Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and in vivo studies? *Arch. Biochem. Biophys.* 476, 107–112.
- Han, D., Matsumaru, K., Rettori, D., Kaplowitz, N., 2004. Usnic acid-induced necrosis of cultured mouse hepatocytes: inhibition of mitochondrial function and oxidative stress. *Biochem. Pharmacol.* 67, 439–451.
- Ingólfssdóttir, K., 2002. Usnic acid *Phytochemistry* 61, 729–736.
- Jayaparaksha, G.K., Rao, L.J., 2000. Phenolic constituents from the lichen *Parmotrema stippeum* (Nyl.) Hale and their antioxidant activity. *Z. Naturforsch. [C]* 55, 1018–1022.
- Joseph, A., Lee, T., Moland, C.L., Branham, W.S., Fuscoe, J.C., Leakey, J.E.A., Allaben, W.T., Lewis, S.M., Ali, A.A., Desai, V.G., 2009. Effect of (+)-usnic acid on mitochondrial functions as measured by mitochondria-specific oligonucleotide microarray in liver of B6C3F1 mice. *Mitochondrion* 9, 149–158.
- Jung, T.W., Lee, J.Y., Shim, W.S., Kang, E.S., Kim, S.K., Ahn, C.W., Lee, H.C., Cha, B.S., 2007. Rosiglitazone protects human neuroblastoma SH-SY5Y cells against MPP+ induced cytotoxicity via inhibition of mitochondrial dysfunction and ROS production. *J. Neurol. Sci.* 253, 53–60.
- Klamt, F., Dal-Pizzol, F., Bernard, E.A., Moreira, J.C., 2003. Enhanced UV-mediated free radical generation; DNA and mitochondrial damage caused by retinol supplementation. *Photochem. Photobiol. Sci.* 2, 856–860.
- Kowaltowski, A.J., de Souza-Pinto, N.C., Castilho, R.F., Vercesi, A.E., 2009. Mitochondria and reactive oxygen species. *Free Radic. Biol. Med.* 47, 333–343.
- Kuhn, M., von Mering, C., Campillos, M., Jensen, L.J., Bork, P., 2008. STITCH: interaction networks of chemicals and proteins. *Nucleic Acids Res.* 36, D684–D688.
- Kuhn, M., Szklarczyk, D., Franceschini, A., Campillos, M., von Mering, C., Jensen, L.J., Beyer, A., Bork, P., 2010. STITCH 2: an interaction network database for small molecules and proteins. *Nucleic Acids Res.* 38, D552–D556.
- Laroux, F.S., Pavlick, K.P., Hines, I.N., Kawachi, H., Harada, S., Bharwani, S., Hoffman, J.M., Grisham, M.B., 2001. Role of nitric oxide in inflammation. *Act. Physiol. Scand.* 173, 113–118.
- Lissi, E., Pascual, C., Del Castillo, M.D., 1992. Luminol luminescence induced by 2,2'-Azo-bis(2-amidinopropane) thermolysis. *Free Radic. Res. Commun.* 17, 299–311.
- Lissi, E., Salim-Hanna, M., Pascual, C., Del Castillo, M.D., 1995. Evaluation of total antioxidant potential (TRAP) and total antioxidant reactivity from luminal-enhanced chemiluminescence measurements. *Free Radic. Biol. Med.* 18, 153–158.
- Lopes, G.K., Schulman, H.M., Hermes-Lima, M., 1999. Polyphenol tannic acid inhibits hydroxyl radical formation from Fenton reaction by complexing ferrous ions. *Biochim. Biophys. Acta* 1472, 142–152.
- Maes, M., Galecki, P., Chang, Y.S., Berk, M., 2011. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the neurodegenerative processes in that illness. *Prog. NeuroPsychopharmacol. Biol. Psychiatry* 35, 676–692.
- Manian, R., Anusuya, N., Siddhuraju, P., Manian, S., 2008. The antioxidant activity and free radical scavenging potential of two different solvent extracts of *Camellia sinensis* (L.) O. Kuntz, *Ficus bengalensis* L. and *Ficus racemosa* L. *Food Chem* 107, 1000–1007.
- Mansour, S.A., Mossa, A.T.H., 2009. Lipid peroxidation and oxidative stress in rat erythrocytes induced by chlorpyrifos and the protective effect of zinc. *Pestic. Biochem. Physiol.* 93, 34–39.
- Mayer, M., O'Neill, M.A., Murray, K.E., Santos-Magalhães, N.S., Carneiro-Leão, A.M.A., Thompson, A.M., Appleyard, V.C.L., 2005. Usnic acid: a non-genotoxic compound with anti-cancer properties. *Anticancer Drugs* 16, 805–809.
- Melo, M.G.D., Santos, J.P.A., Serafini, M.R., Caregnato, F.F., Pasquali, M.A.B., Rabelo, T.K., Rocha, R.F., Júnior, L.Q., Araújo, A.A.S., Silva, A.S., Moreira, J.C.F., Gelain, D.P., 2011. Redox properties and cytoprotective actions of atranorin, a lichen secondary metabolite. *Toxicol. in Vitro* 25, 462–468.
- Misra, H.P., Fridovich, I., 1972. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.* 247, 3170–3175.
- Mollace, V., Muscoli, C., Masini, E., Cuzzocrea, S., Salvemini, D., 2005. Modulation of prostaglandin biosynthesis by nitric oxide and nitric oxide donors. *Pharmacol. Rev.* 57, 217–252.
- Moure, A., Cruz, J.M., Franco, D., Dominguez, J.M., Sineiro, J., Dominguez, H., Núñez, M.J., Parajó, J.C., 2001. Natural antioxidants from residual sources. *Food Chem.* 72, 145–171.
- Navarra, M., Celano, M., Maiuolo, J., Schenone, S., Botta, M., Angelucci, A., Bramanti, P., Russo, D., 2010. Antiproliferative and pro-apoptotic effects afforded by novel Src-kinase inhibitors in human neuroblastoma cells. *BMC Cancer* 10, 602–614.
- Nicolini, G., Miloso, M., Zoia, C., Di Silvestro, A., Cavaletti, G., Tredici, G., 1998. Retinoic acid differentiated SH-SY5Y human neuroblastoma cells: an in vitro model to assess drug neurotoxicity. *Anticancer Res.* 18, 2477–2481.
- Niki, E., Yoshida, Y., Saito, Y., Noguchi, N., 2005. Lipid peroxidation: mechanisms, inhibition, and biological effects. *Biochem. Biophys. Res. Commun.* 338, 668–676.
- Niki, E., 2009. Lipid peroxidation: physiological levels and dual biological effects. *Free Rad. Biol. Med.* 47, 469–484.
- Odobasoglu, F., Cakir, A., Suleyman, H., Aslan, A., Bayir, Y., Halici, M., Kazaz, C., 2006. Gastroprotective and antioxidant effects of usnic acid on indomethacin-induced gastric ulcer in rats. *J. Ethnopharmacol.* 103, 59–65.
- Okuyama, E., Umeyama, K., Yamazaki, M., Kinoshita, Y., Yamamoto, Y., 1995. Usnic acid and diffractaic acid as analgesic and anti-pyretic components of *Usnea diffracta*. *Planta Med.* 61, 113–115.
- Pahlman, S., Ruusala, A.I., Abrahamsson, L., Mattsson, M.E., Esscher, T., 1984. Retinoic acid induced differentiation of cultured human neuroblastoma cells: a comparison with phorbol ester-induced differentiation. *Cell Differ.* 14, 135–144.
- Pastor, N., Weinstein, H., Jamison, E., Brenowitz, M., 2000. A detailed interpretation of OH radical footprints in a TBP-DNA complex reveals the role of dynamics in the mechanism of sequence-specific binding. *J. Mol. Biol.* 304, 55–68.
- Pramyothin, P., Jantasoat, W., Pongnimitprasert, N., Phrukudom, S., Ruangrunsi, N., 2004. Hepatotoxic effect of (+)usnic acid from *Usnea siamensis* Wainio in rats, isolated rat hepatocytes and isolated rat liver mitochondria. *J. Ethnopharmacol.* 90, 381–387.
- Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Bramley, P.M., Pridham, J.B., 1995. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic. Res.* 22, 375–383.

- Salvemini, D., Misko, T.P., Seibert, K., Masferrer, J.L., Currie, M.G., Needleman, P., 1993. Nitric oxide activates cyclooxygenase enzymes. *Proc. Natl. Acad. Sci. USA* 90, 7240–7244.
- Seifried, H.E., Anderson, D.E., Fisher, E.J., Milner, J.A., 2007. A review of the interaction among dietary antioxidants and reactive oxygen species. *J. Nutr. Biochem.* 18, 567–579.
- Scalbert, A., Johnson, I.T., Saltmarsh, M., 2005. Polyphenols: antioxidants and beyond. *Am. J. Clin. Nutr.* 81, 215S–217S.
- Scirpa, P., Scambia, G., Masciullo, V., Battaglia, F., Foti, E., Lopez, R., Villa, P., Mancuso, Malecore M., 1999. A zinc sulphate and usnic acid preparation used as post-surgical adjuvant therapy in genital lesions by human Papillomavirus. *Minerva Ginecol.* 51, 255–260.
- Shukla, S., Mehta, A., Bajpai, V.K., Shukla, S., 2009. In vitro antioxidant activity and total phenolic content of ethanolic leaf extract of *Stevia rebaudiana* Bert. *Food Chem. Toxicol.* 47, 2338–2343.
- Toledo Marante, F.J., Garcia Castellano, A., Estevez Rosas, F., Quintana Aguiar, J., Bermejo Barrera, J., 2003. Identification and quantitation of allelochemicals from the lichen *Lethariella canariensis*: phytotoxicity and antioxidative activity. *J. Chem. Ecol.* 29, 2049–2071.
- Valencia-Islas, N., Zambrano, A., Rojas, J.L., 2007. Ozone reactivity and free radical scavenging behavior of phenolic secondary metabolites in lichens exposed to chronic oxidant air pollution from Mexico City. *J. Chem. Ecol.* 33, 1619–1634.
- Vijayakumar, C.S., Viswanathan, S., Reddy, M.K., Parvathavarthini, S., Kundu, A.B., Sukumar, E., 2000. Anti-inflammatory activity of (+)-usnic acid. *Fitoterapia* 71, 564–566.
- Wang, H., Joseph, J.A., 1999. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radic. Biol. Med.* 27, 612–616.
- Yanmamoto, Y., Miura, Y., Kinoshita, Y., Higuchi, M., Yamada, Y., Murakami, A., Ohigashi, H., Koshimuzi, K., 1995. Screening of tissue cultures and thalli and some of their active constituents for inhibition of tumor promoter-induced Epstein-Barr virus activation. *Chem. Pharm. Bull.* 43, 1388–1390.
- Yu, J., Ye, J., Liu, X., Han, Y., Wang, C., 2011. Protective effect of L-carnitine against H₂O₂-induced neurotoxicity in neuroblastoma (SH-SY5Y) cells. *Neurol Res.* 33, 708–716.
- Zhang, J., Dawson, V.L., Dawson, T.M., Snyder, S.H., 1994. NO activation of poly (ADP-ribose) synthetase in neurotoxicity. *Scienc. Wash. D.C.* 263, 687–689.
- Zimmermann, K., Opitz, N., Dedio, J., Renne, C., Muller-Esterl, W., Oess, S., 2002. NOSTRIN: a protein modulating nitric oxide release and subcellular distribution of endothelial nitric oxide synthase. *Proc. Natl. Acad. Sci. USA* 99, 17167–17172.