

Affine Shuffles, Shuffles with Cuts, the Whitehouse Module, and Patience Sorting

metadata, citation and similar papers at core.ac.uk

Department of Mathematics, Stanford University, Building 360, MC 2125, Stanford,
California 94305

E-mail: fulman@math.stanford.edu

Communicated by Michel Broué

Received October 15, 1999

Type A affine shuffles are compared with riffle shuffles followed by a cut. Although these probability measures on the symmetric group S_n are different, they both satisfy a convolution property. Strong evidence is given that when the underlying parameter q satisfies $\gcd(n, q - 1) = 1$, the induced measures on conjugacy classes of the symmetric group coincide. This gives rise to interesting combinatorics concerning the modular equidistribution by major index of permutations in a given conjugacy class and with a given number of cyclic descents. Using representation theoretic work on the Whitehouse module, a formula is obtained for the cycle structure of a riffle shuffle followed by a cut. It is proved that the use of cuts does not speed up the convergence rate of riffle shuffles to randomness. Generating functions for the first pile size in patience sorting from decks with repeated values are derived. This relates to random matrices. © 2000 Academic Press

Key Words: card shuffling; conjugacy class; sorting; random matrix; cycle structure.

1. INTRODUCTION

In an effort to study the way real people shuffle cards, Bayer and Diaconis [BaD] performed a definitive analysis of the Gilbert–Shannon–Reeds model of riffle shuffling. For an integer $k \geq 1$, a k -shuffle can be described as follows. Given a deck of n cards, one cuts it into k piles with probability of pile sizes j_1, \dots, j_k given by $\binom{n}{j_1, \dots, j_k} / k^n$. Then cards are dropped from the packets with probability proportional to the pile size at a given time (thus if the current pile sizes are A_1, \dots, A_k , the next card is dropped from pile i with probability $A_i / (A_1 + \dots + A_k)$). It



was proved in [BaD] that $\frac{3}{2} \log_2 n$ shuffles are necessary and suffice for a 2-shuffle to achieve randomness (the paper [A] established this result asymptotically in n). It was proved in [DMP] that if $k = q$ is a prime power, then the chance that a permutation distributed as a q -shuffle has n_i i -cycles is equal to the probability that a uniformly chosen monic degree n polynomial over the field F_q factors into n_i irreducible polynomials of degree i .

These results have recently been extended to other Coxeter groups and placed in a Lie theoretic setting. The paper [BeBe] defines hyperoctahedral shuffles using descent algebras and the paper [BiHR] relates Gilbert–Shannon–Reeds shuffles to hyperplane walks. The paper [F2] defines riffle shuffling for arbitrary real hyperplane arrangements, with convergence rates obtainable from the theory in [BiHR]. The results of [DMP] are given a Lie theoretic formulation and extension, at least for types A and B , in [F3] and [F4]. (Random polynomials are replaced by the semisimple orbits of the adjoint action of a finite group of Lie type on its Lie algebra, and even in type A restrictions on the characteristic are needed.) The paper [F1] considers the cycle structure of permutations after biased shuffles, and generalizations based on dynamical systems appear in [La1, La2].

It is worth commenting that the combinatorics of type A riffle shuffles is intimately connected to cyclic and Hochschild homology [Han, GerS], the Poincaré–Birkhoff–Witt theorem [BeW], free Lie algebras [Ga], and Hopf algebras (Section 3.8 of [SSt]). In recent work, Stanley [Sta] has related biased riffle shuffles to the representation theory of the symmetric group, thereby giving an elementary probabilistic interpretation of Schur functions and a different approach to some work in the random matrix community. He recasts and extends results of [BaD] and [F1] using quasisymmetric functions.

Using a construction of Cellini [Ce1, Ce2], the paper [F5] studies combinatorially much more intricate shuffles called affine shuffles (they are reviewed in Section 2). The conjectures of [F5] state in type A that the chance that a permutation distributed as an affine q shuffle has n_i i -cycles is equal to the probability that a uniformly chosen monic degree n polynomial *with constant term 1* over the field F_q factors into n_i irreducible polynomials of degree i . (The abstraction of these polynomials is semisimple conjugacy classes of finite groups of Lie type.) These conjectures are remarkable in the sense that (unlike the Lie algebra case [F3]) no restrictions on the characteristic are needed and there seems to be a reasonably natural way of associating to such a polynomial a permutation in the right conjugacy class, such that choosing the polynomial at random induces the affine shuffling measure. As emerges in [F5] (which gives an application to

dynamical systems and hints at number theoretic applications), this conjecture seems challenging.

The second type of shuffle to be studied in this paper is riffle shuffling followed by a cut at a uniformly chosen random position. Section 3 develops combinatorial preliminaries of shuffles followed by cuts. It is shown there that doing r “ k -shuffles followed by a cut” is the same as doing r k -shuffles and then a single cut (this is known for $k = 2$ from [Ce3]). It is proved that the total variation distance between a sequence of x riffle shuffles and y cuts (performed in any order) and the uniform distribution on S_n is at least the total variation distance between a sequence of x riffle shuffles on S_{n-1} and the uniform distribution on S_{n-1} . In this precise sense cuts do not help speed up riffle shuffles. This perhaps surprising fact can be contrasted with a result of Diaconis [D2], who used representation theory to show that although shuffling by doing random transpositions becomes random in $\frac{1}{2}n \log(n)$ steps, the use of cuts at each stage drops the convergence time to $\frac{3}{8}n \log(n)$ steps. It would be worthwhile and interesting to systematically study the effect of cuts on convergence rates of shuffling methods. Section 3 also shows that a riffle shuffle followed by a cut is at least as random (and sometimes more so) than a cut followed by a riffle shuffle.

Section 4 uses representation theoretic work on the Whitehouse module to obtain a formula for the cycle structure of a riffle shuffle followed by a cut. More precisely, it is shown that the chance that a permutation distributed as a q riffle shuffle followed by a cut has n_i i -cycles is equal to the probability that a uniformly chosen monic degree n polynomial with *non-zero constant term* over the field F_q factors into n_i irreducible polynomials of degree i . The connection with representation theory is illuminating; it follows for instance that the work of [DMP] on the cycle structure of riffle shuffles is equivalent to a representation theoretic result of Hanlon [Han]. The Whitehouse module is interesting in its own right and appears in many places in mathematics; the interested reader should see the transparencies for a talk on the Whitehouse module on Richard Stanley’s MIT website.

Section 5 gives strong evidence for the assertion that affine shuffles and shuffles followed by a cut, although they are different probability measures, coincide when lumped according to conjugacy classes provided that the prime power q satisfies $\gcd(n, q - 1) = 1$. This leads to fascinating combinatorics concerning the modular equidistribution by major index of permutations in a given conjugacy class and with a given number of cyclic descents.

Section 6 considers questions which turn out to be related to cycle structure in multiset permutations. To motivate things, we follow the recent preprint [AD] in its description of patience sorting (which can also

be viewed as a toy model for solitaire, a card game which unlike blackjack has been extremely difficult to analyze mathematically). The simplest case is that one starts with a deck of cards labelled $1, 2, \dots, n$ in random order. Cards are turned up one at a time and dealt into piles on the table according to the rule that a card is placed on the leftmost pile whose top card is of higher value. If no such pile exists, the card starts a new pile to the right. For example, the ordering

$$7513624$$

leads to the arrangement

$$\begin{array}{ccc} 1 & & \\ 5 & 2 & 4 \\ 7 & 3 & 6 \end{array}$$

The survey paper [AD] details connections of patience sorting with ideas ranging from random matrices and the Robinson–Schensted correspondence to interacting particle systems [AD2, BaiDeJ, Han, Rai]. For example, the number of piles in patience sorting is the length of the longest increasing subsequence of a random permutation. After an appropriate renormalization, this statistic has the same distribution as the largest eigenvalue of a random *GUE* matrix. The usefulness of [AD] is its insight that other functions of the random shape obtained through patience sorting have interesting structure. They give results for various pile sizes and suggest the search for analogous results for two variations of patience sorting, from decks with repeated values: “ties allowed” (i.e., a card may be placed on a card of the same value) and “ties forbidden.” Section 6 gives generating functions for the first pile size for these two variations and also for the case where each card is chosen at random from a finite alphabet. We remark that our approach also works for a game one could call *m*-Solitaire in which each card value may be placed on a given pile up to *m* times (ties allowed and ties forbidden corresponding to $m = \infty$ and $m = 1$, respectively). As these games are of less interest, we omit the details.

It may strike the reader that Section 6, although involving cards and cycle structure, is unrelated to card shuffling. However, there is a relation. To elaborate, suppose one picks a random word of length *n* from a totally ordered alphabet, where the probability of getting symbol *i* is x_i . The number of piles in patience sorting applied to the word is the length of the longest weakly increasing subsequence. Each word corresponds to a possible riffle shuffle and the longest increasing subsequence in the corresponding permutation is the longest weakly increasing subsequence in the word [Sta].

To close the introduction, we introduce some terminology that will be used throughout the paper. $C_r(m)$ will denote the Ramanujan sum

$\sum_l e^{2\pi i l m / r}$ where the sum is over all l less than and prime to r . An element w in the symmetric group S_n is said to have a descent at position i if $1 \leq i \leq n - 1$ and $w(i) > w(i + 1)$. The notation $d(w)$ will denote the number of descents of w . The major index of w , denote $\text{maj}(w)$, will be the sum of the positions i at which w has a descent. The permutation w is said to have a cyclic descent at n if $w(n) > w(1)$. Then $cd(w)$ is defined as $d(w)$ if w has no cyclic descent at n and as $d(w) + 1$ if w has a cyclic descent at n . As noted in [Ce1], the concepts of descents and cyclic descents have analogs for all Weyl groups. (The descents are the simple positive roots mapped to negative roots and w has a cyclic descent if it keeps the highest root positive.)

2. TYPE A AFFINE SHUFFLES

This section gives six definitions for what we call type A affine k -shuffles. The first two are due to Cellini [Ce1] and are the best in that they generalize to all Weyl groups. The next four are due to the author [F5] and are very useful for computational purposes. (The final definition is a “physical” description of affine type A 2-shuffles. A good problem is to extend the physical description to higher values of k .) In all of the definitions k is a positive integer. A seventh definition (which is conjectural but very conceptual and potentially valid for all Weyl groups) appears in the final section of [F5]. As it requires some effort to describe and will not be used here, we omit it.

DEFINITIONS OF AFFINE TYPE A k -SHUFFLES

1. Let W_k be the subgroup of the type A affine Weyl group generated by reflections in the n hyperplanes $\{x_1 = x_2, \dots, x_{n-1} = x_n, x_n - x_1 = k\}$. This subgroup has index k^{n-1} in the affine Weyl group and has k^{n-1} unique minimal length coset representatives, each of which can be written as the product of a permutation and a translation. Choose one of these k^{n-1} coset representatives uniformly at random and consider its permutation part. Define a type A affine k -shuffle to be the distribution on permutations thus obtained.

2. A type A affine k -shuffle assigns probability to $w^{-1} \in S_n$ equal to $1/k^{n-1}$ multiplied by the number of integer vectors (v_1, \dots, v_n) satisfying the conditions:

- (a) $v_1 + \dots + v_n = 0$
- (b) $v_1 \geq v_2 \geq \dots \geq v_n, v_1 - v_n \leq k$
- (c) $v_i > v_{i+1}$ if $w(i) > w(i + 1)$ (with $1 \leq i \leq n - 1$)
- (d) $v_1 < v_n + k$ if $w(n) > w(1)$.

3. A type A affine k -shuffle assigns probability to $w^{-1} \in S_n$ equal to $1/k^{n-1}$ multiplied by the number of partitions with $\leq n - 1$ parts of size at most $k - cd(w)$ such that the total number being partitioned has size congruent to $-\text{maj}(w) \pmod n$. Equivalently, it assigns probability equal to $1/k^{n-1}$ multiplied by the number of partitions with $\leq k - cd(w)$ parts of size at most $n - 1$ such that the total number being partitioned has size congruent to $-\text{maj}(w) \pmod n$.

4. A type A affine k -shuffle assigns probability to $w^{-1} \in S_n$ equal to

$$\frac{1}{nk^{n-1}} \sum_{r|n, k-cd(w)} \binom{n+k-cd(w)-r}{r, k-cd(w)} C_r(-\text{maj}(w))$$

if $k - cd(w) > 0$,

$$\frac{1}{k^{n-1}}$$

if $k - cd(w) = 0, \text{maj}(w) = 0 \pmod n$,

0 otherwise.

5. A type A affine k -shuffle assigns probability to $w^{-1} \in S_n$ equal to

$$\frac{1}{k^{n-1}} \sum_{r=0}^{\infty} \text{Coeff. of } q^{r \cdot n} \text{ in } \left(q^{\text{maj}(w)} \binom{k+n-cd(w)-1}{n-1} \right),$$

where $\binom{A}{B}$ denotes the q -binomial coefficient

$$\frac{(1-q) \cdots (1-q^A)}{(1-q) \cdots (1-q^B)(1-q) \cdots (1-q^{A-B})}.$$

6. A type A affine 2 shuffle has the following physical description for the symmetric group S_{2n} .

Step 1. Choose an even number between 1 and $2n$ with the probability of getting $2j$ equal to $\binom{2n}{2j}/2^{2n-1}$. From the stack of $2n$ cards, form a second pile of size $2j$ by removing the top j cards of the stack and then putting the bottom j cards of the first stack on top of them.

Step 2. Now one has a stack of size $2n - 2j$ and a stack of size $2j$. Drop cards repeatedly according to the rule that if stacks 1, 2 have sizes A, B at some time, then the next card comes from stack 1 with probability

$A/(A + B)$ and from stack 2 with probability $B/(A + B)$. (This is equivalent to choosing uniformly at random one of the $\binom{2n}{2j}$ interleavings preserving the relative orders of the cards in each stack).

The description of x_2 is the same for the symmetric group S_{2n+1} , except that at the beginning of Step 1 the chance of getting $2j$ is $\binom{2n+1}{2j}/2^{2n}$ and at the beginning of Step 2 one has a stack of size $2n + 1 - 2j$ and a stack of size $2j$.

An important property of these shuffles is the so-called ‘‘convolution property,’’ which says that a k_1 shuffle followed by a k_2 shuffle is equivalent to a $k_1 k_2$ shuffle. It is interesting that the type A riffle shuffles of [BaD] satisfy the same convolution property, as do some of the generalizations in [F2].

The conjecture of [F5] that relates the cycle structure of permutations distributed as affine q -shuffles to the factorization of polynomials with constant term 1 appears to be interesting. For instance, it is shown there that for the case of the identity conjugacy class of S_n , it amounts to the $m = 0$ case of following observation in ‘‘modular combinatorial reciprocity.’’ We recently found this observation in the paper [EJPa] which interprets it as a Hermite reciprocity statement in invariant theory.

COROLLARY. *For any positive integers x, y , the number of ways (disregarding order and allowing repetition) of writing m (mod y) as the sum of x integers of the set $0, 1, \dots, y - 1$ is equal to the number of ways (disregarding order and allowing repetition) of writing m (mod x) as the sum of y integers of the set $0, 1, \dots, x - 1$.*

Let $f_{n,k,d}$ be the coefficient of z^n in $((z^k - 1)/(z - 1))^d$ and let μ be the Moebius function. Let $n_i(w)$ be the number of i -cycles in a permutation w . Then ([F5]) the conjecture is equivalent to the truly bizarre assertion (which we intentionally do not simplify) that for all n, m, k ,

$$\begin{aligned} & \sum_{m=0 \bmod n} \text{Coeff. of } q^m u^n t^k \\ & \text{in } \sum_{n=0}^{\infty} \frac{u^n}{(1-tq) \cdots (1-tq^n)} \sum_{w \in S_n} t^{cd(w)} q^{\text{maj}(w)} \prod x_i^{n_i(w)} \\ & = \sum_{m=0 \bmod k-1} \text{Coeff. of } q^m u^n t^k \\ & \text{in } \sum_{k=0}^{\infty} t^k \prod_{i=1}^{\infty} \prod_{m=1}^{\infty} \left(\frac{1}{1 - q^m x_i u^i} \right)^{1/i \sum_{d|i} \mu(d) f_{m,k,i/d}}. \end{aligned}$$

3. SHUFFLES FOLLOWED BY A CUT

To begin we remark that although the paper [BaD] says it examined shuffles followed by a cut, it investigated a cut followed by a shuffle, which is different.

Let s be the element of the group algebra of S_n denoting a k -riffle shuffle. Let ζ be the cyclic permutation $(1 \cdots n)$ and let $c = \frac{1}{n} \sum_{i=0}^{n-1} \zeta^i$. Thus in this notation a shuffle followed by a cut is simply cs . The inverse of an element $\sum r_w w$ of the group algebra will be taken to mean $\sum r_w w^{-1}$.

It is useful to recall the following formula of Bayer and Diaconis.

THEOREM 1 [BaD]. *The coefficient of a permutation w in the element s is*

$$\frac{1}{k^n} \binom{n+k-d(w^{-1})-1}{n}$$

Theorem 2 derives an analogous formula for a shuffle followed by a cut.

THEOREM 2. *The coefficient of a permutation w in the element cs is*

$$\frac{1}{nk^{n-1}} \binom{n+k-cd(w^{-1})-1}{n-1}$$

Proof. Consider instead the coefficient of w in $s^{-1}c$. This coefficient is equal to

$$\frac{1}{n} \sum_{k=0}^{n-1} \text{Coeff. of } w\zeta^k \text{ in } s^{-1}.$$

The element $w\zeta^k$ maps i to $w(i+k \bmod n)$. Consequently, letting $cd(w)$ be the number of cyclic descents of w , there are $cd(w)$ values of k for which $w\zeta^k$ has $cd(w) - 1$ descents, and $n - cd(w)$ values of k for which $w\zeta^k$ has $cd(w)$ descents. Combining this with Theorem 1 shows that the coefficient of w in $s^{-1}c$ is

$$\frac{1}{nk^n} \left(cd(w) \binom{n+k-cd(w)}{n} + (n-cd(w)) \binom{n+k-cd(w)-1}{n} \right),$$

which simplifies to the formula in the statement of the theorem. ■

This yields the following combinatorial corollary.

COROLLARY 1. Let $B_{n,i}$ be the number of elements of S_n with i cyclic descents. Let $A_{n,i}$ be the number of elements of S_n with $i - 1$ descents. Then

1. $x^{n-1} = \sum_{1 \leq i \leq n-1} (B_{n,i}/n) \binom{n+x-i-1}{n-1}$.
2. If $n > 1$ then $B_{n,i} = nA_{n-1,i}$.

Proof. The first assertion is immediate from Theorem 2. The second assertion follows from the first together with the well-known facts that $A_{n,i} = A_{n,n+1-i}$ and that $A_{n,i}$ is the unique sequence satisfying Worpitzky's identity $x^n = \sum_{1 \leq i \leq n} A_{n,i} \binom{x+i-1}{n}$. ■

Theorem 3 appears in [Ce3] for the case $k = 2$. As noted there, it implies that $(cs)^h = cs^h$ for any natural number h . The proof given here is simpler.

THEOREM 3. $csc = cs$.

Proof. Taking inverses and using the fact that $c^{-1} = c$, it is enough to show that $cs^{-1}c = s^{-1}c$. The coefficient of w in $cs^{-1}c$ is

$$\frac{1}{n} \sum_{k=0}^{n-1} \text{Coeff. of } \zeta^k w \text{ in } s^{-1}c.$$

It is easy to see that $\text{cd}(\zeta^k w) = \text{cd}(w)$ for all k . The result now follows from Theorem 2. ■

Next recall the notion of total variation distance $\|P_1 - P_2\|$ between two probability distributions P_1 and P_2 on a finite set X . It is defined as

$$\frac{1}{2} \sum_{x \in X} |P_1(x) - P_2(x)|.$$

The book [D2] explains why this is a natural and useful notion of distance between probability distributions. $P_1 * P_2$ (the convolution) is defined by $P_1 * P_2(\pi) = \sum_{\tau \in S_n} P_1(\pi\tau^{-1})P_2(\tau)$, and $P_1 * \cdots * P_k$ is defined inductively. The following elementary lemma will be helpful.

LEMMA 1. Let P, Q , be any measures on a finite group G and let U be the uniform distribution. Then $\|P * Q - U\| \leq \|Q - U\|$.

Theorem 4 shows that cuts do not speed up the convergence rate of riffle shuffles.

THEOREM 4. (1) *Let $S^{(k)}, C, U$ denote the probability distribution corresponding to a k -riffle shuffle, a cut, and the uniform distribution, respectively. Then $\|C * S^{(k)} - U\| \leq \|S^{(k)} * C - U\|$ and the inequality can be strict. (In words, a shuffle followed by a cut is more random than a cut followed by a shuffle.)*

(2) *For $n > 1$, $\|C * S^{(k)} - U\|_{S_n} = \|S^{(k)} - U\|_{S_{n-1}}$.*

(3) *Let W be the convolution of any finite sequence of riffle shuffles and cuts. Let W' be the convolution of the same finite sequence, but with the cuts eliminated. (By abuse of notation, these can be viewed on any symmetric group.) Then*

$$\|W - U\|_{S_n} \geq \|W' - U\|_{S_{n-1}}.$$

Proof. For the first assertion, observe that Theorem 3 gives that $C * S^{(k)} = C * S^{(k)} * C$. Now use Lemma 1. Computations with the symmetric group S_4 show that the inequality can be strict.

For the second assertion, let $B(n, i)$ be the number of elements of S_n with i cyclic descents and let $A(n, i)$ be the number of elements of S_n with $i - 1$ descents. Observe that

$$\begin{aligned} \|C * S^{(k)} - U\| &= \frac{1}{2} \sum_{i=1}^{n-1} B(n, i) \left| \frac{\binom{k+n-i-1}{n-1}}{nk^{n-1}} - \frac{1}{n!} \right| \\ &= \frac{1}{2} \sum_{i=1}^{n-1} A(n-1, i) \left| \frac{\binom{k+n-i-1}{n-1}}{k^{n-1}} - \frac{1}{(n-1)!} \right| \\ &= \|S^{(k)} - U\|_{S_{n-1}}. \end{aligned}$$

The second equality is the second part of Corollary 1 and the final equality follows from Theorem 1.

For the third assertion, the inequality is clear if W has no cuts. Otherwise, combining the fact that $S^{(i)} * S^{(j)} = S^{(ij)}$ for any i, j with Theorem 3 shows that W is equivalent to a convolution of the form $S^{(k_1)} * C * S^{(k_2)}$ (with k_1 or k_2 possibly 0 and $S^{(0)}$ denoting the measure placing all mass on the identity). Now observe that

$$\begin{aligned} \|S^{(k_1)} * C * S^{(k_2)} - U\|_{S_n} &\geq \|C * S^{(k_1)} * C * S^{(k_2)} - U\|_{S_n} \\ &= \|C * S^{(k_1 k_2)} - U\|_{S_n} \\ &= \|S^{(k_1 k_2)} - U\|_{S_{n-1}}. \end{aligned}$$

The first equality is Lemma 1, the second equality comes from Theorem 3, and the third equality is the second part of this theorem. ■

A formula for a cut followed by a riffle shuffle appears in [BaD], although it is not evident how it could be used to prove Part 1 of Theorem 4.

As a final problem, we observe that the n -cycle $\zeta = (1 \cdots n)$ is a minimal length Coxeter element for type A . As there are analogs of shuffling for other finite Coxeter groups [BeBe, F2], it may be possible to extend the results of this paper to other Coxeter groups.

4. REPRESENTATION THEORY

This section uses representation theory to obtain a formula for the cycle structure of a riffle shuffle followed by a cut.

It is useful to recall the notion of a cycle index associated to a character of the symmetric group. Letting $n_i(w)$ be the number of i -cycles of a permutation w and N be a subgroup of S_n , one defines $Z_N(\chi)$ as

$$Z_N(\chi) = \frac{1}{|N|} \sum_{w \in N} \chi(w) \prod_i a_i^{n_i(w)}.$$

The cycle index stores complete information about the character χ . For a proof of the following attractive property of cycle indices, see [Fe].

LEMMA 2. *Let N be a subgroup of S_n and χ be a class function on N . Then*

$$Z_{S_n}(\text{Ind}_N^{S_n}(\chi)) = Z_N(\chi).$$

Next, recall that an idempotent e of the group algebra of a finite group G defines a character χ for the action of G on the left ideal KGe of the group algebra of G over a field K of characteristic zero. For a proof of Lemma 3, which will serve as a bridge between representation theory and computing measures over conjugacy classes, see [Han]. For its statement, let $e\langle w \rangle$ be the coefficient of w in the idempotent e .

LEMMA 3. *Let C be a conjugacy class of the finite group G , and let χ be the character associated to the idempotent e . Then*

$$\frac{1}{|G|} \sum_{w \in C} \chi(w) = \sum_{w \in C} e\langle w \rangle.$$

It is also convenient to define

$$Z_{S_n}(e) = \sum_{w \in S_n} e\langle w \rangle \prod_i a_i^{n_i(w)},$$

which makes sense for any element e of the group algebra. Note that one does not divide by the order of the group. When e is idempotent and χ is the associated character, Lemma 3 can be rephrased as

$$Z_{S_n}(\chi) = Z_{S_n}(e).$$

To proceed recall the Eulerian idempotents e_n^j , $j = 1, \dots, n$ in the group algebra QS_n of the symmetric group over the rationals. These can be defined [GerS] as follows: Let $s_{i, n-i} = \sum w$ where the sum is over all $\binom{n}{i}$ permutations w such that $w(1) < \dots < w(i)$, $w(i + 1) < \dots < w(n)$, and let $s_n = \sum_{i=1}^{n-1} s_{i, n-i}$. Letting $\mu_j = 2^j - 2$, the e_n^j are defined as

$$e_n^j = \prod_{i \neq j} \frac{s_n - \mu_i}{(\mu_j - \mu_i)}.$$

They are orthogonal idempotents which sum to the identity.

The following result, which we shall need, is due to Hanlon. The symbol μ denotes the Moebius function of elementary number theory.

THEOREM 5 [Han].

$$1 + \sum_{n=1}^{\infty} \sum_{i=1}^n k^i Z_{S_n}(e_n^i) = \prod_{i \geq 1} (1 - a_i)^{-1/i} \sum_{d|i} \mu(d) k^{i/d}.$$

THEOREM 6 [Ga].

$$\sum_{i=1}^n k^i e_n^i = \sum_{w \in S_n} \binom{n+k-d(w)-1}{n} w.$$

Remark. Combining Lemma 3 and Theorem 6, one sees that the formula for the cycle structure of a riffle shuffle [DMP] and Theorem 5 imply each other. It is interesting that both proofs used a bijection of Gessel and Reutenauer [GesR].

To continue, we let $\overline{e_n^j}$ denote the idempotent obtained by multiplying the coefficient of w in e_n^j by $\text{sgn}(w)$. Let λ_{n+1} be the $n + 1$ cycle $(1\ 2 \cdots n + 1)$ and $\Lambda_{n+1} = 1/(n + 1) \sum_{i=0}^n (\text{sgn } \lambda_{n+1}^i) \lambda_{n+1}^i$. Viewing $\overline{e_n^j}$ as in the group algebra of S_{n+1} , Whitehouse [Wh] proves that for $j = 1, \dots, n$

the element $\Lambda_{n+1} \overline{e_n^j}$ is an idempotent in the group algebra QS_{n+1} , which we denote by f_{n+1}^j . Whitehouse's main result is the following:

THEOREM 7 [Wh]. *Let $F_{n+1}^j, \overline{E_n^j}$ be the irreducible modules corresponding to the idempotents f_{n+1}^j and e_n^j . Then*

$$F_{n+1}^j \oplus \bigoplus_{i=1}^j \overline{E_{n+1}^i} = \bigoplus_{i=1}^j \text{Ind}_{S_n}^{S_{n+1}} \overline{E_n^i}.$$

As a final preparation for the main result of this section, we link the idempotent $\Lambda_{n+1} \overline{e_n^j}$ with riffle shuffles followed by a cut.

LEMMA 4. *The coefficient of w in $\sum_{j=1}^n k^j \Lambda_{n+1} \overline{e_n^j}$ is $\text{sgn}(w) \frac{1}{n+1} \binom{k+n}{n} - \text{cd}(w)$.*

Proof. Given Theorem 6, this is an elementary combinatorial verification. ■

Theorem 8 now derives the cycle structure of a permutation distributed as a shuffle followed by a cut. To simplify the generating functions, recall that $\sum_{d|i} \mu(d)$ vanishes unless $i = 1$.

THEOREM 8.

$$\begin{aligned} 1 + \sum_{n \geq 1} \sum_{w \in S_{n+1}} \frac{1}{(n+1)k^{n+1}} \binom{n+k-\text{cd}(w)}{n} \prod_i a_i^{n_i(w)} \\ = 1 - \frac{1}{k-1} - \frac{a_1}{k} + \frac{1}{k-1} \prod_{i \geq 1} \left(1 - \frac{a_i}{k^i}\right)^{-1/i \sum_{d|i} \mu(d) (k^{i/d} - 1)}. \end{aligned}$$

If $k = q$ is the size of a finite field, this says that the cycle type of a permutation distributed as a shuffle followed by the cut has the same law as the factorization type of a monic degree n polynomial over F_q with a non-vanishing constant term.

Proof. Replacing a_i by $a_i k^i (-1)^{i+1}$, it is enough to show that

$$\begin{aligned} 1 + \sum_{n \geq 1} \sum_{w \in S_{n+1}} \text{sgn}(w) \frac{1}{(n+1)} \binom{n+k-\text{cd}(w)}{n} \prod_i a_i^{n_i(w)} \\ = 1 - \frac{1}{k-1} - a_1 + \frac{1}{k-1} \prod_{i \geq 1} \left(1 - (-1)^{i+1} a_i\right)^{-1/i \sum_{d|i} \mu(d) (k^{i/d} - 1)}. \end{aligned}$$

Using Lemmas 2–4 and Theorem 7, one sees that

$$\begin{aligned}
 & 1 + \sum_{n \geq 1} \sum_{w \in S_{n+1}} \operatorname{sgn}(w) \frac{1}{(n+1)} \binom{n+k-\operatorname{cd}(w)}{n} \prod_i a_i^{n_i(w)} \\
 &= 1 + \sum_{n=1}^{\infty} \sum_{j=1}^n k^j Z_{S_{n+1}}(f_n^j) \\
 &= 1 + \sum_{n=1}^{\infty} \sum_{j=1}^n k^j \sum_{i=1}^j Z_{S_{n+1}}(\operatorname{Ind}_{S_n}^{S_{n+1}}(\overline{e_n^i})) - \sum_{n=1}^{\infty} \sum_{j=1}^n k^j \sum_{i=1}^j Z_{S_{n+1}}(\overline{e_{n+1}^i}) \\
 &= 1 + a_1 \sum_{n=1}^{\infty} \sum_{i=1}^n Z_{S_n}(\overline{e_n^i}) \left(\frac{k^{n+1} - k^i}{k-1} \right) \\
 &\quad - \sum_{n=1}^{\infty} \sum_{i=1}^n Z_{S_{n+1}}(\overline{e_{n+1}^i}) \left(\frac{k^{n+1} - k^i}{k-1} \right) \\
 &= 1 + a_1 k Z_{S_1}(\overline{e_1}) + \frac{a_1 k - 1}{k-1} \sum_{n=2}^{\infty} k^n \sum_{i=1}^n Z_{S_n}(\overline{e_n^i}) \\
 &\quad + \frac{1 - a_1}{k-1} \sum_{n=2}^{\infty} \sum_{i=1}^n k^i Z_{S_n}(\overline{e_n^i}).
 \end{aligned}$$

To simplify things further, recall that $\sum_{i=1}^n Z_{S_n}(\overline{e_n^i})$ is a_1^n since the $\overline{e_n^i}$'s sum to the identity. The above then becomes

$$1 - \frac{1}{k-1} - a_1 + \frac{1 - a_1}{k-1} \left(1 + \sum_{n=1}^{\infty} \sum_{i=1}^n k^i Z_{S_n}(\overline{e_n^i}) \right),$$

so the sought-after result follows from Theorem 5. ■

We observe that a combinatorial proof of Theorem 8 (which must exist) would give a new proof of Theorem 7 by reversing the steps. To close this section we remark that Theorem 8 gives analogs of all the results in Section 5 of [DMP] for k -riffle shuffles followed by a cut. For instance, the expected number of fixed points is 1 whereas for k -riffle shuffles it is $1 + 1/k + \dots + 1/k^{n-1}$. For fixed k as $n \rightarrow \infty$ the joint distribution of the number of i -cycles after a k -shuffle followed by a cut converges to independent negative binomials with parameters $1/i \sum_{d|i} \mu(d)(k^{i/d} - 1)$ and $1/k^i$. The large cycles have the same limit theory as in [DMP]. Proofs are at the author's web site.

5. CONJUGACY CLASSES

The aim of this section is to give evidence for the conjecture at the end of Section 2, in the case when $\gcd(q-1, n) = 1$. Note that under this assumption a uniformly chosen degree n polynomial with non-zero constant term and a uniformly chosen degree n polynomial with constant term 1 have the same chance of factoring into n_i i -cycles. Hence in this case the conjecture amounts to the assertion that affine shuffles and shuffles followed by a cut, although they are different probability measures, induce the same distribution on conjugacy classes. Before posing a problem which would explain why this should hold, some lemmas are needed.

LEMMA 5. *If $r > 1$, then $\sum_{j=0}^{r-1} C_r(j) = 0$.*

Proof. If l is relatively prime to r , then multiplication by l permutes the numbers $\{0, 1, \dots, r-1\} \bmod r$. Thus

$$\sum_{j=0}^{r-1} C_r(j) = \sum_{\substack{0 \leq l \leq r \\ \gcd(l, r) = 1}} \sum_{j=0}^{r-1} e^{2\pi i j l / r} = \phi(r) \sum_{j=0}^{r-1} e^{2\pi i j / r} = 0.$$

■

LEMMA 6. *For $n \geq 1$, let t be the largest divisor of n such that $\gcd(cd-1, t) = 1$. Suppose that $\gcd(n, q-1) = 1$ and that r divides n and $q - cd$. Then r divides t .*

Proof. Observe that $\gcd(r, cd-1) = 1$. Suppose there is some $a > 1$ dividing r and $cd-1$. Then a divides $q - cd$ and $cd-1$, hence $q-1$. Since a divides r and r divides n , it follows that a divides n . This contradicts the assumption that $\gcd(q-1, n) = 1$. ■

Next we pose the problem of determining whether or not the following statement holds.

Statement 1. For $n \geq 1$, let t be the largest divisor of n such that $\gcd(cd-1, t) = 1$. Then for every conjugacy class C of S_n , the set of permutations in C with cd cyclic descents has its major index equidistributed mod t .

Theorem 9 shows that if Statement 1 holds, then the conjecture about the cycle structure of permutations distributed as affine shuffles is correct. Some evidence in favor of Statement 1 is then given.

THEOREM 9. *Suppose that $\gcd(q-1, n) = 1$. If Statement 1 is correct, then affine shuffles and shuffles followed by a cut have the same distribution on conjugacy classes.*

Proof. Suppose that Statement 1 is correct and recall the third definition of affine q -shuffles in Section 2. If $q < \text{cd}(w)$ then both the affine q -shuffle and the q -riffle shuffle followed by a cut assign probability 0 to w . If $q = \text{cd}(w)$, then the affine q -shuffle assigns probability $1/q^{n-1}$ to w if $\text{maj}(w) = 0 \pmod n$, and 0 otherwise. If $q = \text{cd}(w)$, then the q -riffle shuffle followed by a cut associates probability $1/nq^{n-1}$ to w . Since $q = \text{cd}$, the t in Statement 1 is equal to n , which implies that for every conjugacy class C the set of permutations in C with cd cyclic descents has major index equidistributed mod n . Hence Statement 1 holds in this case.

The third and final case is that $q > \text{cd}(w)$. Suppose that $r > 1$ divides n and $q - \text{cd}$. Lemma 6 implies that r divides t . Hence, by Statement 1, for any conjugacy class C , the set of permutations with cd cyclic descents has its major index equidistributed mod r . Consequently (the second equality below holding by Lemma 5 and the equidistribution property mod r),

$$\begin{aligned} & \sum_{\text{cd}=1}^n \sum_{\substack{w \in C \\ \text{cd}(w) = \text{cd}}} \frac{1}{nq^{n-1}} \sum_{r|n, q-\text{cd}} \left(\frac{\binom{n+q-\text{cd}-r}{r}}{\binom{q-\text{cd}}{r}} \right) C_r(-\text{maj}(w)) \\ &= \sum_{\text{cd}=1}^n \frac{1}{nq^{n-1}} \sum_{r|n, q-\text{cd}} \left(\frac{\binom{n+q-\text{cd}-r}{r}}{\binom{q-\text{cd}}{r}} \right) \sum_{\substack{w \in C \\ \text{cd}(w) = \text{cd}}} C_r(-\text{maj}(w)) \\ &= \sum_{\text{cd}=1}^n \frac{1}{nq^{n-1}} \binom{n+q-\text{cd}-1}{n-1} \sum_{\substack{w \in C \\ \text{cd}(w) = \text{cd}}} 1 \\ &= \sum_{w \in C} \frac{1}{nq^{n-1}} \binom{n+q-\text{cd}(w)-1}{n-1}. \end{aligned}$$

From Theorem 2 (the formula for a q -riffle shuffle followed by a cut), Statement 1 follows. ■

Next we consider evidence in favor of the ideas of this section. Incidentally, given Section 4, Proposition 1 confirms conjecture 1 in [F5] in type A when n is prime and q is a power of n .

PROPOSITION 1. *Suppose that n is prime and that q is a power of n . Then type A affine q -shuffles are exactly the same as q -riffle shuffles followed by a cut.*

Proof. The probability that an affine q shuffle yields w is

$$\frac{1}{nq^{n-1}} \sum_{r|n, q-\text{cd}(w^{-1})} \left(\frac{\frac{n+q-\text{cd}(w^{-1})-r}{r}}{\frac{q-\text{cd}(w^{-1})}{r}} \right) C_r(-\text{maj}(w^{-1})).$$

Since $1 \leq \text{cd}(w) \leq n-1$ for any w in S_n , the assumptions on n and q imply that the only r dividing n and $q - \text{cd}(w^{-1})$ is $r = 1$. The result now follows from Theorem 2. ■

THEOREM 10. *Statement 1 holds for the identity conjugacy class and for the conjugacy class of simple transpositions.*

Proof. For the identity conjugacy class, use the third definition of affine q -shuffles in Section 2 together with the assumption that $\text{gcd}(n, q-1) = 1$.

Next consider the case of simple transpositions. Suppose that $n \geq 4$, the other cases being trivial. One checks that all simple transpositions (i, j) with $i < j$ have either 2 or 3 cyclic descents. The easy case is that of 2 cyclic descents. The possible values of (i, j) are then $(i, i+1)$ for $1 \leq i \leq n-2$, $(1, n)$, and $(n-1, n)$. The values of the major index thus obtained are $\{1, \dots, n\}$, and each value is hit once. Thus Statement 1 holds in this case.

The harder case is that of three cyclic descents. The relevant transpositions are (i, j) with $1 \leq i, j < n$ and $j \neq i+1$ (having major index $i+j-1$) and (i, n) with $2 \leq i < n-1$ (having major index $i+n-1$).

First suppose that n is odd. It suffices to prove that

$$\sum_{\substack{w=(i,j) \\ \text{cd}(w)=3}} x^{\text{maj}(w) \bmod n}$$

is a multiple of $(x^n - 1)/(x - 1)$. Calculating gives

$$\begin{aligned} & \sum_{\substack{w=(i,j) \\ \text{cd}(w)=3}} x^{\text{maj}(w) \bmod n} \\ &= \sum_{i=1}^{(n-3)/2} x^i \sum_{j=i+1}^{n-i-1} x^j + \frac{1}{x^n} \sum_{i=1}^{(n-3)/2} x^i \sum_{j=n-i}^{n-2} x^j \\ & \quad + \frac{1}{x^n} \sum_{i=(n-1)/2}^{n-3} x^i \sum_{j=i+1}^{n-2} x^j + \sum_{i=1}^{n-3} x^i \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{x-1} \left(\sum_{i=1}^{(n-3)/2} x^i(x^{n-i} - x^{i+1}) + \frac{1}{x^n} \sum_{i=1}^{(n-3)/2} x^i(x^{n-1} - x^{n-i}) \right) \\
 &\quad + \frac{1}{x-1} \left(\frac{1}{x^n} \sum_{i=(n-1)/2}^{n-3} x^i(x^{n-1} - x^{i+1}) + x^{n-2} - x \right) \\
 &= \frac{n-3}{2} \frac{x^n - 1}{x-1},
 \end{aligned}$$

as desired.

Next suppose that $n = 2^a$ with $a > 0$. It suffices to prove that

$$\sum_{\substack{w=(i,j) \\ cd(w)=3}} x^{\text{maj}(w) \bmod n}$$

is a polynomial multiple of $(x^{n/2^a} - 1)/(x - 1)$. Calculating as above (and omitting the steps analogous to the previous computation) gives that

$$\begin{aligned}
 &\sum_{\substack{w=(i,j) \\ cd(w)=3}} x^{\text{maj}(w) \bmod n} \\
 &= \frac{1}{x-1} (1 + x^2 + x^4 + \dots + x^{n-2} - x - x^3 - x^5 - \dots - x^{n-1}) \\
 &= -\frac{x^n - 1}{x^2 - 1} \\
 &= -\frac{x^{n/2^a} - 1}{x-1} \frac{x^{(2^a-1)n/2^a} + \dots + x^{n/2^a} + 1}{x+1}.
 \end{aligned}$$

Since $n/2^a$ is odd, it follows that $x^{(2^a-1)n/2^a} + \dots + x^{n/2^a} + 1$ is divisible by $x + 1$. ■

6. PATIENCE SORTING

Having described the motivation in the Introduction, we outline and then execute a strategy for obtaining generating function information for the first pile size in patience sorting from decks with repeated values. The first step is to apply ideas of Foata to obtain generating functions for multiset permutations by the number of cycles. The second step is to give a multiset records-to-cycles bijection (generalizing the one used in [AD]), which converts information about the distribution of cycles to information

about the distribution of records. The final step is to read information off of the generating function.

Some notation is needed. Let \mathbf{a} denote the vector (a_1, a_2, \dots) with $a_i \geq 0$ and $\sum a_i < \infty$. Let $\text{Mult}(\mathbf{a})$ denote the collection of all (a_1, a_2, \dots) words of length $\sum a_i$ formed from a_i i 's.

We recall Foata's theory of cycle structure for multisets [Fo], following Knuth's superb exposition [Kn]. Suppose that the elements of the multiset are linearly ordered. Then multiset permutations can be written in two-line notation,

$$\begin{pmatrix} a & a & a & b & b & c & d & d & d & d \\ c & a & b & d & d & a & b & d & a & d \end{pmatrix}.$$

Foata introduced an intercalation product $_T$ which multiplies two multiset permutations α and β by expressing α and β in two line notation, juxtaposing these two-line notations, and then sorting the columns in non-decreasing order of the top line. For example,

$$\begin{array}{ccccccccc} a & a & b & c & d & & a & b & d & d & d \\ c & a & d & a & b & & b & d & d & a & d \end{array} \quad {}_T \quad \begin{array}{ccccccccc} a & a & a & b & b & c & d & d & d & d & d \\ c & a & b & d & d & a & b & d & a & a & d \end{array}$$

Foata proved that if the elements of the multiset M are linearly ordered by the relation $<$, then the permutations π of M correspond exactly to the possible intercalations

$$\pi = (x_{11} \cdots x_{1n_1} y_1) {}_T (x_{21} \cdots x_{2n_2} y_2) {}_T \cdots {}_T (x_{t1} \cdots x_{tn_t} y_t),$$

with $y_1 \leq y_2 \leq \cdots \leq y_t$ and $y_i < x_{ij}$ for $1 \leq j \leq n_i$, $1 \leq i \leq t$. This defines a notion of cycle structure for multiset permutations by letting the cycles be the intercalation factors. Let $C_i(\pi)$ be the number of length i cycles of π and $C(\pi) = \sum C_i(\pi)$ be the total number of cycles. Let $C'_i(\pi)$ be the number of i -cycles of π , where cycles with the same minimum value y_j are counted at most once. Let $C'(\pi) = \sum C'_i(\pi)$. For example, the multiset permutation

$$(431) {}_T (231) {}_T (4)$$

satisfies $C_3(\pi) = 2$, $C'_3(\pi) = 1$, $C(\pi) = 3$, and $C'(\pi) = 2$.

Proposition 2 gives generating functions for multiset permutations.

PROPOSITION 2.

$$1 + \sum_{\mathbf{a}} \sum_{\pi \in \text{Mult}(\mathbf{a})} u^{C(\pi)} \prod_{i \geq 1} x_i^{a_i} = \prod_{k=1}^{\infty} \frac{1}{1 - x_k u / (1 - \sum_{j>k} x_j)}$$

$$1 + \sum_{\mathbf{a}} \sum_{\pi \in \text{Mult}(\mathbf{a})} u^{C'(\pi)} \prod_{i \geq 1} x_i^{a_i} = \prod_{k=1}^{\infty} \left(1 + u \frac{x_k / (1 - \sum_{j>k} x_j)}{1 - x_k / (1 - \sum_{j>k} x_j)} \right)$$

Proof. Both generating functions follow easily from Foata’s method of representing permutations by intercalations. The k ’s on the right-hand side index the letters of the alphabet. The point is that cycles are formed by fixing a smallest element k and specifying an ordered choice of elements larger than k ; permutations are ordered multisets of such cycles. ■

We remark that the generating functions of Proposition 2 are quasi-symmetric functions in the sense that for any $i_1 < \dots < i_n$ and $j_1 < \dots < j_n$ the coefficients of $x_{i_1}^{j_1} \dots x_{i_n}^{j_n}$ and $x_{j_1}^{i_1} \dots x_{j_n}^{i_n}$ are equal.

Theorem 11 converts information about the distribution of cycles to information about the distribution of records. Some further notation is needed for its statement. Let π^{rev} be the word obtained by reading from right to left the bottom line in the 2-line form of π . Recalling the definition of solitaire from the Introduction, let $P_i(\pi)$ and $P'_i(\pi)$ be the number of cards in pile i of solitaire with ties allowed and ties forbidden, respectively.

THEOREM 11. 1. For any given \mathbf{a} , there is a bijection $\Phi: \text{Mult}(\mathbf{a}) \mapsto \text{Mult}(\mathbf{a})$ such that if R_1, \dots, R_t are the positions of the left-to-right minima (ties are allowed) of π^{rev} , then $R_2 - R_1, R_3 - R_2, \dots, R_t - R_{t-1}, (\sum a_i) + 1 - R_t$ are the cycle lengths in Foata’s factorization of $\Phi(\pi)$.

2. $P_1(\pi) = C(\Phi(\pi))$ and $P'_1(\pi) = C'(\Phi(\pi))$.

Proof. For the first assertion, define $\Phi(\pi)$ as an intercalation of cycles formed by entries in the bottom line of π , with cycles (from left-to-right) having lengths $(\sum a_i) + 1 - R_t, R_t - R_{t-1}, \dots, R_2 - R_1$. The assertion is then evident, and the following example may help to untangle the notation. The multiset permutation $\pi = ddbcbdbbcabacdbd$ has $\pi^{\text{rev}} = dbdcabacbbdcbdd$ with $R_1(\pi^{\text{rev}}) = 1, R_2(\pi^{\text{rev}}) = 2, R_3(\pi^{\text{rev}}) = 5, R_4(\pi^{\text{rev}}) = 7$, and $\sum a_i + 1 = 16$. Thus forming from π cycles of lengths 9, 2, 3, 1 gives $\Phi(\pi)$ as the intercalation

$$(ddbcdbbca)_T (ba)_T (cdb)_T (d).$$

For the second assertion, we give the argument for the first equality, the argument for the second assertion being analogous. The point is that the

number of cards in pile 1 of solitaire with ties allowed applied to π is simply the number of left-to-right minima with ties allowed to π . The result now follows from the first assertion. ■

Proposition 3 shows that when one considers random words from a finite alphabet, there is a factorization for the full cycle structure vector, not only the number of cycles. Recent work of Tracy and Widom [TW] connects random words with random matrices chosen from the Laguerre ensemble. Let $\text{Word}_n(N)$ be the N^n words of length n from an alphabet on N letters (say $1, 2, \dots, N$). Each such word can be viewed as a multiset permutation.

PROPOSITION 3.

$$\begin{aligned}
 & 1 + \sum_n \frac{1}{N^n} \sum_{\pi \in \text{Word}_n(N)} \prod_{k=1}^N x_k^{a_k(\pi)} \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 &= \prod_{i \geq 1} \prod_{k=1}^N \left(1 / \left(1 - u_i \frac{x_k}{N} \left(\sum_{j=k+1}^N \frac{x_j}{N} \right)^{i-1} \right) \right). \\
 & (1-x) + \sum_n \frac{(1-x)x^n}{N^n} \sum_{\pi \in \text{Word}_n(N)} \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 &= \prod_{i \geq 1} \prod_{k=1}^N \frac{1 - \frac{x^i}{N} \left(\frac{N-k}{N} \right)^{i-1}}{1 - \frac{u_i x^i}{N} \left(\frac{N-k}{N} \right)^{i-1}}. \\
 & 1 + \sum_n \frac{1}{N^n} \sum_{\pi \in \text{Word}_n(N)} \prod_{k=1}^N x_k^{a_k(\pi)} \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 &= \prod_{i \geq 1} \prod_{k=1}^N \left(1 + u_i \frac{x_k}{N} \frac{1}{1 - \sum_{j=k+1}^N \frac{x_j}{N}} \right). \\
 & (1-x) + \sum_n \frac{(1-x)x^n}{N^n} \sum_{\pi \in \text{Word}_n(N)} \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 &= \prod_{i \geq 1} \prod_{k=1}^N \frac{1 + u_i \frac{x}{N} \frac{1}{1 - \sum_{j=k+1}^N \frac{x}{N}}}{1 + \frac{x}{N} \frac{1}{1 - \sum_{j=k+1}^N \frac{x}{N}}}.
 \end{aligned}$$

Proof. For the first assertion, note by Foata's representation of multi-set permutations as intercalations that each i -cycle is formed by fixing a

smallest element k and specifying an ordered choice of $i - 1$ elements larger than k to occupy the first $i - 1$ positions of the cycle. Since multiset permutations are ordered multisets of such cycles, one concludes that

$$\begin{aligned}
 1 + \sum_n \sum_{\pi \in \text{Word}_n(N)} \prod_{k=1}^N x_k^{a_k(\pi)} \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 = \prod_{i \geq 1} \prod_{k=1}^N \frac{1}{1 - u_i x_k \left(\sum_{j=k+1}^N x_j\right)^{i-1}}.
 \end{aligned}$$

Now replace each x_i by x_i/N .

To prove the second assertion, replacing each x_i by x in the first assertion yields the equation

$$\begin{aligned}
 1 + \sum_n \frac{1}{N^n} \sum_{\pi \in \text{Word}_n(N)} x^n \prod_{i \geq 1} u_i^{C_i(\pi)} \\
 = \prod_{i \geq 1} \prod_{k=1}^N \left(1 / \left(1 - \frac{u_i x^i}{N} \left(\frac{N-k}{N} \right)^{i-1} \right) \right).
 \end{aligned}$$

Setting all $u_i = 1$ and taking reciprocals shows that

$$1 - x = \prod_{i \geq 1} \prod_{k=1}^N \left(1 - \frac{x^i}{N} \left(\frac{N-k}{N} \right)^{i-1} \right).$$

The result follows by multiplying the previous two equations.

The arguments for the third and fourth assertions are analogous. ■

The second and fourth equations have probabilistic interpretations. For instance, in the second equation fix x such that $0 < x < 1$. The equation then says that if one picks n geometrically with probability $(1 - x)x^n$ and then picks $\pi \in \text{Word}_n(N)$ uniformly at random, the random variables $C_i(\pi)$ are sums of independent geometrics. In the fourth equation the C'_i become sums of independent binomials. Lemma 7 permits asymptotic statements in the $n \rightarrow \infty$ limit.

LEMMA 7. *If $f(1) < \infty$ and the Taylor series of f around 0 converges at $u = 1$, then*

$$\lim_{n \rightarrow \infty} [u^n] \frac{f(u)}{1 - u} = f(1).$$

Proof. Write the Taylor expansion $f(u) = \sum_{n=0}^{\infty} a_n u^n$. Then observe that $[u^n]f(u)/(1 - u) = \sum_{i=0}^n a_i$. ■

As a corollary, one sees for instance that as $n \rightarrow \infty$ the number of i -cycles of a random length- n word from the alphabet $\{1, \dots, N\}$ converges to a sum of independent geometrics with parameters $(1 - \frac{k}{N})^{i-1}$ as $k = 1, \dots, N$. For more on this type of factorization result and its applications, see [LIS] for the symmetric groups, [DS] for the compact classical groups, and [F6] for the finite classical groups.

Finally, we consider the application of Proposition 2 and Theorem 11 to patience sorting. As above, $P_1(\pi)$ and $P'_1(\pi)$ are the number of cards in pile 1 of patience sorting with ties allowed and ties forbidden respectively.

THEOREM 12. *Let π be chosen uniformly at random from the possible orderings of a deck of cards with a_i cards labelled i . Then $E(P_1) = \sum_k (a_k / (a_1 + \dots + a_{k-1} + 1))$ and $E(P'_1) = \sum_k (a_k / (a_1 + \dots + a_{k-1} + a_k))$.*

Proof. By Proposition 2 and Theorem 11,

$$1 + \sum_{\mathbf{a}} \sum_{\pi \in \text{Mult}(\mathbf{a})} u^{P_1(\pi)} \prod_{i \geq 1} x_i^{a_i} = \prod_{k=1}^{\infty} \left(1 / \left(1 - \frac{x_k u}{1 - \sum_{j>k} x_j} \right) \right).$$

Differentiating with respect to u and setting $u = 1$ implies that the sought expectation is

$$\begin{aligned} & \frac{1}{\binom{n}{a_1, a_2, \dots}} \text{Coeff. of } \prod x_i^{a_i} \text{ in } \sum_k \frac{x_k}{1 - \sum x_i} \frac{1}{1 - \sum_{j \geq k} x_j} \\ &= \frac{1}{\binom{n}{a_1, a_2, \dots}} \\ & \times \sum_{k: a_k > 0} \sum_{b_k, b_{k+1}, \dots \geq 0} \binom{a_1 + \dots + a_{k-1} + b_k + b_{k+1} + \dots}{a_1, \dots, a_{k-1}, b_k, b_{k+1}, \dots} \\ & \times \binom{a_k - 1 - b_k + a_{k+1} - b_{k+1} + \dots}{a_k - 1 - b_k, a_{k+1} - b_{k+1}, \dots} \\ &= \frac{1}{\binom{n}{a_1, a_2, \dots}} \sum_{k: a_k > 0} \frac{1}{a_1! \dots a_{k-1}! (a_k - 1)! a_{k+1}! \dots} \\ & \times \sum_{b_k, b_{k+1}, \dots \geq 0} \frac{\binom{a_k - 1}{b_k} \binom{a_{k+1}}{b_{k+1}} \binom{a_{k+2}}{b_{k+2}} \dots}{\binom{\sum a_i - 1}{a_1 + \dots + a_{k-1} + b_k + b_{k+1} + \dots}}. \end{aligned}$$

Letting $s = b_k + b_{k+1} + \dots$, this simplifies to

$$\begin{aligned} & \sum_{k : a_k > 0} \sum_{s=0}^{a_k + a_{k+1} + \dots - 1} \frac{1}{\sum a_i} \frac{\binom{a_k + a_{k+1} + \dots - 1}{s}}{\binom{\sum a_i - 1}{a_1 + \dots + a_{k-1} + s}} \\ &= \sum_{k : a_k > 0} \frac{1}{\sum a_i} \frac{\binom{a_k + a_{k+1} + \dots - 1}{a_1 + \dots + a_{k-1} + s}}{\binom{\sum a_i - 1}{a_1 + \dots + a_{k-1}}} \\ &= \sum_{k : a_k > 0} \frac{a_k}{a_1 + \dots + a_{k-1} + 1} \\ &= \sum_k \frac{a_k}{a_1 + \dots + a_{k-1} + 1}. \end{aligned}$$

The second calculation is similar. ■

As a final result, we study patience sorting applied to I_{2n} , the fixed point free involutions in the symmetric group S_{2n} . By [Rai], the number of piles in such a game relates to the eigenvalues of random symplectic and orthogonal matrices. Consequently, this restricted version of patience sorting merits further study. Proposition 4 shows that the generating function for the first pile size factors.

PROPOSITION 4.

$$\sum_{\pi \in I_{2n}} x^{P_1(\pi)} = \prod_{i=1}^n (x^2 + 2(i-1)).$$

Proof. The proposition is proved by induction, the base case being trivial. Suppose that the proposition holds for $I_{2(n-1)}$. Given $\pi \in I_{2n}$, let j be the symbol with which $2n$ is switched. If $j \neq 1$, then $P_1(\pi)$ is the same as $P_1(\pi')$ where π' is obtained by crossing the symbols j and $2n$ out of π . If $j = 1$, then $P_1(\pi) = P_1(\pi') + 2$, where $P_1(\pi')$ is obtained by crossing the symbols $1, 2n$ out of π . Consequently,

$$\sum_{\pi \in I_{2n}} x^{P_1(\pi)} = (2n - 2) \sum_{\pi \in I_{2n-2}} x^{P_1(\pi)} + x^2 \sum_{\pi \in I_{2n-2}} x^{P_1(\pi)}$$

and the result follows by induction. ■

ACKNOWLEDGMENTS

This research was supported by an NSF postdoctoral fellowship. The author thanks Persi Diaconis for ongoing discussions.

REFERENCES

- [A] A. Aldous, “Random Walk on Finite Groups and Rapidly Mixing Markov Chains,” Springer Lecture Notes in Mathematics, Vol. 986, pp. 243–297, Springer-Verlag, New York/Berlin, 1983.
- [AD] D. J. Aldous and P. Diaconis, Longest increasing subsequences: from patience sorting to the Baik–Deift–Johansson theorem, *Bull. Amer. Math. Soc.* **36** (1999), 413–432.
- [AD2] D. J. Aldous and P. Diaconis, Hammersley’s interacting particle process and longest increasing subsequences, *Probab. Theory Relat. Fields* **103** (1995), 199–213.
- [BaiDeJ] J. Baik, P. A. Deift, and K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations, *J. Amer. Math. Soc.* **12** (1999), 1119–1178.
- [BaD] D. Bayer and P. Diaconis, Trailing the dovetail shuffle to its lair, *Ann. Appl. Probab.* **2** (1992), 294–313.
- [BeBe] F. Bergeron and N. Bergeron, Orthogonal idempotents in the descent algebra of B_n and applications, *J. Pure Appl. Algebra* **79** (1992), 109–129.
- [BeW] N. Bergeron and L. Wolfgang, The decomposition of Hochschild cohomology and the Gerstenhaber operations, *J. Pure Appl. Algebra* **104** (1995), 243–265.
- [BiHR] P. Bidigare, P. Hanlon, and D. Rockmore, A combinatorial description of the spectrum of the Tsetlin library and its generalization to hyperplane arrangements, *Duke Math J.* **99** (1999), 135–174.
- [Ce1] P. Cellini, A general commutative descent algebra, *J. Algebra* **175** (1995), 990–1014.
- [Ce2] P. Cellini, A general commutative descent algebra. II. The case C_n , *J. Algebra* **175** (1995), 1015–1026.
- [Ce3] P. Cellini, Cyclic Eulerian elements, *Europ. J. Combin.* **19** (1998), 545–552.
- [D1] P. Diaconis, “Group Representations in Probability and Statistics,” Institute of Mathematical Statistics Lecture Notes Monograph Series, Vol. 11, Institute of Mathematical Statistics, Hayward, CA, 1995.
- [D2] P. Diaconis, Finite Fourier methods: Access to tools, in “Probabilistic Combinatorics and Its Applications,” Proceedings Symposia Applied Mathematics, Vol. 44, pp. 171–194, Amer. Math. Soc., Providence, RI, 1991.
- [DMP] P. Diaconis, M. McGrath, and J. Pitman, Riffle shuffles, cycles, and descents, *Combinatorica* **15** (1995), 11–20.
- [DS] P. Diaconis and M. Shahshahani, On the eigenvalues of random matrices, *J. Appl. Probab.* **31** (1994), 49–61.
- [EJPa] A. Elashvili, M. Jibladze, and D. Pataria, Combinatorics of necklaces and Hermite reciprocity, *J. Algebraic Combin.* **10** (1999), 173–188.
- [Fe] W. Feit, “Characters of Finite Groups,” Yale Univ. Press, New Haven, CT, 1965.
- [Fo] D. Foata, Etude algebrique de certains problemes d’analyse combinatoire et du calcul des probabilites, *Publ. Inst. Statist. Univ. Paris* **14** (1965), 81–241.

- [F1] J. Fulman, The combinatorics of biased riffle shuffles, *Combinatorica* **18** (1998), 173–184.
- [F2] J. Fulman, Descent algebras, hyperplane arrangements, and shuffling cards, *Proc. Amer. Math. Soc.*, to appear; also available at <http://math.stanford.edu/~fulman>.
- [F3] J. Fulman, Semisimple orbits of Lie algebras and card shuffling measures on Coxeter groups, *J. Algebra* **224** (2000), 151–165.
- [F4] J. Fulman, Counting semisimple orbits of finite Lie algebras by genus, *J. Algebra* **217** (1999), 170–179.
- [F5] J. Fulman, Cellini’s descent algebra, dynamical systems, and semisimple conjugacy classes of finite groups of Lie type, preprint available at <http://math.stanford.edu/~fulman>.
- [F6] J. Fulman, Cycle indices for the finite classical groups, *J. Group Theory* **2** (1999), 251–289.
- [Ga] A. Garsia, Combinatorics of the free Lie algebra and the symmetric group, in “Analysis, et cetera . . .” pp. 309–382, Academic Press, New York, 1990.
- [GerS] M. Gerstenhaber and S. D. Schack, A Hodge-type decomposition for commutative algebra cohomology, *J. Pure. Appl. Algebra* **38** (1987), 229–247.
- [GesR] I. Gessel and C. Reutenauer, Counting permutations with given cycle structure and descent set, *J. Combin. Theory Ser. A* **64** (1993), 189–215.
- [Ham] J. M. Hammersley, A few seedling of research, in “Proceedings, Sixth Berkeley Symposium on Mathematical Statistics and Probability,” Vol. 1, pp. 345–394, University of California Press, Berkeley, 1972.
- [Han] P. Hanlon, The action of S_n on the components of the Hodge decomposition of Hochschild homology, *Michigan Math. J.* **37** (1990), 105–124.
- [Kn] D. Knuth, “The Art of Computer Programming,” Vol. 3, Addison–Wesley, Reading, MA, 1973.
- [La1] S. Lalley, Cycle structure of riffle shuffles, *Ann. Probab.* **24** (1996), 49–73.
- [La2] S. Lalley, Riffle shuffles and their associated dynamical systems, preprint.
- [LIS] S. P. Lloyd and L. A. Shepp, Ordered cycle lengths in a random permutation, *Trans. Amer. Math. Soc.* **121** (1966), 340–357.
- [M] C. L. Mallows, Patience sorting, *Bull. Inst. Math. Appl.* **9** (1973), 216–224.
- [Rai] E. M. Rains, Increasing subsequences and the classical groups, *Electron. J. Combin.* **5** (1998), R12.
- [SSt] S. Shnider and S. Sternberg, “Quantum Groups,” Graduate Texts in Mathematical Physics, Vol. 2, International Press, Cambridge, MA, 1993.
- [Sta] R. Stanley, Generalized riffle shuffles and quasisymmetric functions, preprint, math.CO/9912025 at xxx.lanl.gov.
- [TW] C. A. Tracy and H. Widom, On the distributions of the lengths of the longest monotone subsequences in random words, preprint, <http://xxx.lanl.gov/math.CO/9904042>.
- [Wh] S. Whitehouse, The Eulerian representations of S_n as restrictions of representations of S_{n+1} , *J. Pure Appl. Algebra* **115** (1997), 309–320.